

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/135233>

Please be advised that this information was generated on 2018-06-25 and may be subject to change.

The strong nilpotency index of a matrix

Michiel de Bondt

Radboud University, Nijmegen, The Netherlands

Email: debondt@math.ru.nl

February 28, 2013

Abstract

It is known that strongly nilpotent matrices over a division ring are linearly triangularizable. We describe the structure of such matrices in terms of the strong nilpotency index. We apply our results on quasi-translation $x + H$ such that $\mathcal{J}H$ has strong nilpotency index two.

Keywords: strongly nilpotent matrices, linear triangularization, linear dependence, quasi-translation.

2010 Mathematics Subject Classification: 14R10; 14R15; 15A03; 15A04.

1 Introduction

The Jacobian conjecture asserts that a polynomial map F over a field of characteristic zero has a polynomial inverse G in case $\det \mathcal{J}F$ is a unit of the base field. In [BCW] and [Ya], it is shown that the Jacobian Conjecture holds if it holds for all polynomial maps $x + H$ of such that $\mathcal{J}H$ is nilpotent. This result is refined in [E, §6.2], where it is shown that for any polynomial map F of degree d in dimension n , the Jacobian conjecture holds if it holds for all polynomial maps $x + H$ of degree d in dimension $(d - 1)n$ such that $\mathcal{J}H$ is nilpotent, which subsequently holds if the Jacobian conjecture holds for all polynomial maps $x + H$ of degree d in dimension $(d - 1)n + 1$ such that H is homogeneous of degree d .

So it is natural to look at nilpotent Jacobian matrices. For quadratic maps, it is already known that the Jacobian conjecture holds, but since we have $(d - 1)n = n$ in that case, we see that reduction to the case that $F = x + H$ with $\mathcal{J}H$ nilpotent does not cost extra dimensions. But one may even assume that additionally, H is homogeneous. That is why more than twenty years ago, Gary Meisters and Czesław Olech looked at nilpotent Jacobian matrices of quadratic homogeneous H in low dimensions in [MO], and it appeared that in dimensions $n \leq 4$, the Jacobian matrix of H was so-called *strongly nilpotent*, but not necessarily in higher dimensions.

Somewhat later, but still more than fifteen years ago, polynomial maps $x+H$ over a field of characteristic zero such that $\mathcal{J}H$ is strongly nilpotent were classified as so-called *linearly triangularizable* polynomial maps, in both [EH] and [Y1]. Since linear triangularizable maps are invertible, the Jacobian conjecture holds for polynomial maps $x+H$ over a field of characteristic zero such that $\mathcal{J}H$ is strongly nilpotent. But there are no results about the index of strong nilpotency: just as with the nilpotency index for nilpotent matrices, one can define the strong nilpotency index for strongly nilpotent matrices as the minimum number of factors such that its product is zero.

Actually, there are two equivalent definitions for strongly nilpotent matrices over a field of characteristic zero, but both definitions no longer correspond when the base ring becomes arbitrary. Sometimes, the definitions only deal with matrices which are Jacobians, but such a restriction is unnecessary, see also equation (2.10) in the beginning of the proof of corollary 2.2 below. It is neither necessary to assume that the base ring is commutative or even a field. Although the definitions do not require that the base ring is unital, all results in this article are about rings R with unity.

The first of both definitions is given in [MO]. Write $f|_{x=g}$ for substituting x by g in f .

Definition 1.1 (Meisters and Olech in [MO]). Let $R[x]$ be the polynomial ring over a ring R (not necessarily commutative) in n indeterminates x_1, x_2, \dots, x_n . A square matrix M with entries in $R[x]$ is called *strongly nilpotent* if for some $r \in \mathbb{N}$, we have

$$M|_{x=v_1} \cdot M|_{x=v_2} \cdots \cdots M|_{x=v_r} = 0 \tag{1.1}$$

for all $v_i \in R^n$.

Actually, r is equal to the dimension of the matrix M in the original definition, but I consider that a consequence of restricting to fields of characteristic zero, or just to reduced commutative rings, see the later proposition 3.2.

Definition 1.2 (Van den Essen and Hubbers in [EH]). Let $R[x]$ be the polynomial ring over a ring R (not necessarily commutative) in n indeterminates x_1, x_2, \dots, x_n , which commute with each other and with elements of R . A square matrix M with entries in $R[x]$ is called *strongly nilpotent* if for some $r \in \mathbb{N}$, we have

$$M|_{x=y^{(1)}} \cdot M|_{x=y^{(2)}} \cdots \cdots M|_{x=y^{(r)}} = 0 \tag{1.2}$$

where the $y^{(i)}$ are other tuples of n indeterminates, which commute with each other and with elements of R .

Again, r is equal to the dimension of the matrix M in the original definition, which is given in [EH]. Over infinite integral domains, both definitions are the same. We call the minimum possible $r \in \mathbb{N}$ such that (1.1) or (1.2) respectively holds the *strong nilpotency index* of M .

In section 2, we consider definition 1.2 because it does not correspond to definition 1.1 when R is a finite field. This is because different polynomials

can be the same as a function, whence substituting all possible elements of R in the indeterminates will not distinguish both polynomials. For instance, the polynomials x_1 and x_1^q over \mathbb{F}_q are the same as functions, so they act equal on substituting an element of \mathbb{F}_q for x_1 . Over infinite integral domains, different polynomials are always different function as well. So definitions 1.2 and 1.1 are equivalent when R is an infinite integral domain.

In [Y1], definition 1.1 has been generalized to functions to R in general, not just multivariate polynomials, which we will discuss in section 3. Just like the main results in [EH] and [Y1] are similar, while the respective definitions 1.2 and the generalization of 1.1 are different, the refinements of these results by way of the strong nilpotency index, which are theorem 2.1 and theorem 3.1 respectively, are similar as well.

In the last section, we apply the results of section 2 on quasi-translations of strong nilpotency index two. For more information about quasi-translations, we refer to the last section and its references.

2 Strongly nilpotent matrices in the sense of [EH]

In [EH], the authors prove that strongly nilpotent Jacobians in the sense of 1.2 are linearly triangularizable, but there is no need to restrict to Jacobians. Instead, one can look at any matrix of polynomials over a division ring D . Notice that for finite fields, different polynomials might represent the same function. Therefore, the generalization to polynomial matrices is not a special case of the generalization of definition 1.1 in [Y1] that will follow in the next section.

Theorem 2.1. *Let $n \geq 0$ and $x = x_1, x_2, \dots, x_n$ be variables which commute with each other and with elements of a division ring D . Assume that $M \in \text{Mat}_m(D[x])$, where $m \geq 1$.*

Then M has strong nilpotency index r (in the sense of definition [EH]), if and only if there exists a $T \in \text{GL}_m(D)$ such that $T^{-1}MT$ is of the form

$$\begin{pmatrix} 0_{s_1} & & & & \emptyset \\ A_1 & 0_{s_2} & & & \\ & A_2 & \ddots & & \\ & & \ddots & 0_{s_{r-1}} & \\ * & & & A_{r-1} & 0_{s_r} \end{pmatrix} \quad (2.1)$$

where 0_{s_i} is the square zero matrix of size $s_i \geq 1$ and A_i has independent columns over D for each i . In particular, the strong nilpotency index of M does not exceed m in case M is strongly nilpotent.

Proof. Write $y^{(i)} = (y_1^{(i)}, y_2^{(i)}, \dots, y_n^{(i)})$ for each i .

⇒ Assume that M has nilpotency index r . Then

$$M|_{x=y^{(r)}} \cdot M|_{x=y^{(r-1)}} \cdots \cdots M|_{x=y^{(2)}} \cdot M|_{x=y^{(1)}} = 0 \quad (2.2)$$

but

$$M|_{x=y^{(r-1)}} \cdots \cdots M|_{x=y^{(2)}} \cdot M|_{x=y^{(1)}} \neq 0 \quad (2.3)$$

Thus for some term in $y^{(r-1)}, \dots, y^{(2)}, y^{(1)}$, the coefficient matrix $C \in \text{Mat}_n(D)$ on the left hand side of (2.3) does not vanish, and we obtain from (2.2) that $M|_{x=y^{(r)}} \cdot C = 0$. Hence also $M \cdot C = 0$ and the columns of M are dependent over D . Thus if we choose $T \in \text{GL}_n(D)$ and $s_r \in \mathbb{N}$ maximal, such that the last s_r columns of $M \cdot T$ are zero, then $s_r \geq 1$ and the first $m - s_r$ columns of $M \cdot T$ are independent over D .

Now replace M by $T^{-1}MT$. Then the first $m - s_r$ columns of M become independent over D and the last s_r columns of M become zero. Furthermore, by the maximality of s_r , we have that the first $m - s_r$ rows of the left hand side of (2.3) become zero, since otherwise we can find another linear dependence between the columns of both $M|_{x=y^{(r)}}$ and M , namely by taking for C the coefficient matrix of the left hand side of (2.3) with respect to a term in $y^{(r-1)}, y^{(r-2)}, \dots, y^{(2)}, y^{(1)}$, such that the first $m - s_r$ rows of C are not zero everywhere. Hence the left hand side of (2.3) is of the form

$$\begin{pmatrix} 0_{m-s_r} & \emptyset \\ \tilde{A} & 0_{s_r} \end{pmatrix} \quad (2.4)$$

where 0_{m-s_r} is the square zero matrix of size $m - s_r$ and $\tilde{A} \neq 0$ because of (2.3).

Let \tilde{M} be the leading principal minor matrix of size $m - s_r$ of M . Since the last s_r columns of M , and hence also of $M|_{x=y^{(i)}}$ for all $i \geq 2$, are zero, we have that the product

$$\tilde{M}|_{x=y^{(r-1)}} \cdot \tilde{M}|_{x=y^{(r-2)}} \cdots \cdots \tilde{M}|_{x=y^{(2)}} \cdot \tilde{M}|_{x=y^{(1)}} \quad (2.5)$$

is equal to the leading principal minor matrix of size $m - s_r$ of the left hand side of (2.3), which is the submatrix 0_{m-s_r} of (2.4). The product

$$\tilde{M}|_{x=y^{(r-2)}} \cdots \cdots \tilde{M}|_{x=y^{(2)}} \cdot \tilde{M}|_{x=y^{(1)}} \quad (2.6)$$

is nonzero, because the last s_r columns of $M|_{x=y^{(r-1)}}$ are zero and hence (2.6) is a factor of \tilde{A} .

So by induction on m , there exists a $\tilde{T} \in \text{GL}_{m-s_r}(D)$ such that $\tilde{T}^{-1}\tilde{M}\tilde{T}$ is of the form

$$\begin{pmatrix} 0_{s_1} & & & & \emptyset \\ A_1 & 0_{s_2} & & & \\ & A_2 & \ddots & & \\ & & \ddots & 0_{s_{r-2}} & \\ * & & & A_{r-2} & 0_{s_{r-1}} \end{pmatrix}$$

where A_i has independent columns over D for each i . Since the first $m - s_r$ columns of M are independent over D and the last s_r columns of M are zero, the desired result follows.

⇐ Assume that a $T \in \text{GL}_m(D)$ as in theorem 2.1 exists. Then one can prove by induction on j that for all $j \leq r$, the first $s_1 + s_2 + \dots + s_j$ rows of $(T^{-1}MT)|_{x=y^{(j)}} \cdot (T^{-1}MT)|_{x=y^{(j-1)}} \cdots (T^{-1}MT)|_{x=y^{(1)}}$ are zero. Hence the strong nilpotency index of M does not exceed r . Furthermore, the first $s_1 + s_2 + \dots + s_j$ rows of $(T^{-1}MT)|_{x=y^{(j-1)}} \cdots (T^{-1}MT)|_{x=y^{(1)}}$ are of the form

$$\begin{pmatrix} & & & \emptyset \\ A_{j-1}|_{x=y^{(j-1)}} & \cdots & A_1|_{x=y^{(1)}} & \end{pmatrix}$$

Since the columns of A_i are independent over D for each i , we obtain by looking at coefficient matrices in a similar manner as in \Rightarrow and by induction on j , that $A_{j-1}|_{x=y^{(j-1)}} \cdots A_1|_{x=y^{(1)}} \neq 0$ for all $j \leq r$. Consequently, r is the strong nilpotency index of M . \square

Application to Jacobian matrices

In addition to the result we get by taking for M a Jacobian matrix in theorem 2.1, we have the following.

Corollary 2.2. *Assume $H \in K[x]^n = K[x_1, x_2, \dots, x_n]^n$, where K is any field. Write $y^{(i)} = (y_1^{(i)}, y_2^{(i)}, \dots, y_n^{(i)})$ for all i .*

Then for each nonzero $r \in \mathbb{N}$, the following statements are equivalent.

- (1) $(\mathcal{J}H)|_{x=y^{(r)}} \cdot (\mathcal{J}H)|_{x=y^{(r-1)}} \cdots (\mathcal{J}H)|_{x=y^{(1)}} = 0$, i.e. the strong nilpotency index of $\mathcal{J}H$ (in the sense of definition 1.2) does not exceed r .
- (2) There exists a $T \in \text{GL}_n(K)$ such that the Jacobian of $T^{-1}H(Tx)$ is of the form

$$\begin{pmatrix} 0_{s_1} & & & \emptyset \\ & 0_{s_2} & & \\ & & \ddots & \\ * & & & 0_{s_{r'}} \end{pmatrix} \quad (2.7)$$

where 0_{s_i} is the square zero matrix of size $s_i \geq 1$ for each i and $r' = \min\{r, n\}$.

- (3) For all j with $1 \leq j \leq r$, we have that the Jacobian with respect to $y^{(1)}$ of

$$\begin{aligned} & (\mathcal{J}H)|_{x=y^{(r)}} \cdot (\mathcal{J}H)|_{x=y^{(r-1)}} \cdots (\mathcal{J}H)|_{x=y^{(j+1)}} \cdot \\ & t^{(j)} H \left(y^{(j)} + t^{(j-1)} H \left(\dots \left(y^{(2)} + t^{(1)} H(y^{(1)}) \right) \dots \right) \right) \end{aligned} \quad (2.8)$$

vanishes. (Thus (2.8) is constant with respect to $y^{(1)}$ when $\text{char } K = 0$.)

(4) There exists a j with $1 \leq j \leq r$, such that the Jacobian with respect to $y^{(1)}$ of

$$(\mathcal{J}H)|_{x=y^{(r)}} \cdot (\mathcal{J}H)|_{x=y^{(r-1)}} \cdots \cdots (\mathcal{J}H)|_{x=y^{(j+1)}} \cdot H\left(y^{(j)} + H\left(\cdots (y^{(2)} + H(y^{(1)})) \cdots\right)\right) \quad (2.9)$$

vanishes. (Thus (2.9) is constant with respect to $y^{(1)}$ when $\text{char } K = 0$.)

Proof. Since (3) \Rightarrow (4) is trivial, the following remains to be proved.

(1) \Rightarrow (2) Assume (1). From theorem 2.1, it follows that there exists a $T \in \text{GL}_n(K)$ such that $T^{-1} \cdot \mathcal{J}H \cdot T$ is of the form (2.7) for some $r' \leq r$ and hence also for $r' = \min\{r, n\}$. By the chain rule, we obtain

$$\mathcal{J}(T^{-1}H(Tx)) = T^{-1} \cdot (\mathcal{J}H)|_{x=Tx} \cdot T = (T^{-1} \cdot \mathcal{J}H \cdot T)|_{x=Tx} \quad (2.10)$$

which gives (2).

(2) \Rightarrow (1) Assume (2). From (2.10), it follows that $T^{-1} \cdot \mathcal{J}H \cdot T$ is of the form (2.7) as well. Subsequently, (1) can be proved in a similar manner as \Leftarrow of theorem 2.1.

(1) \Rightarrow (3) Assume (1). By taking the Jacobian with respect to $y^{(1)}$ of (2.8) and dividing by $t^{(j)}t^{(j-1)} \cdots t^{(1)}$, we obtain $(\mathcal{J}H)|_{x=y^{(r)}+\dots} \cdot (\mathcal{J}H)|_{x=y^{(r-1)}+\dots} \cdots \cdots (\mathcal{J}H)|_{x=y^{(1)}+\dots}$. This product is zero because it is a homomorphic image of the left hand side of the equality in (1), and (3) follows.

(4) \Rightarrow (1) Assume (4). By taking the Jacobian with respect to $y^{(1)}$ of (2.9), we obtain $(\mathcal{J}H)|_{x=y^{(r)}+\dots} \cdot (\mathcal{J}H)|_{x=y^{(r-1)}+\dots} \cdots \cdots (\mathcal{J}H)|_{x=y^{(1)}+\dots}$, which is zero on assumption. Furthermore, the left hand side of (1) is a homomorphic image of it, and (1) follows. \square

Since all differentiated factors are commuted to the right in the chain rule, this rule does not apply in a noncommutative context. Therefore, (3) and (4) of corollary 2.2 cannot be generalized to division rings. We will use corollary 2.2 in the last section.

Comparing regular and strong nilpotency index

Before constructing some maps H with strongly nilpotent Jacobians, such that the nilpotency index of the Jacobian is less than the strong nilpotency index, we formulate a proposition.

Proposition 2.3. *Let D be a division ring and suppose that $M \in \text{Mat}_m(D[x]) = \text{Mat}_m(D[x_1, x_2, \dots, x_n])$ is strongly nilpotent with index r (in the sense of definition 1.2), such that r exceeds the regular nilpotency index. Then $3 \leq r \leq m - 1$.*

Proof. If the regular nilpotency index equal one, then $M = 0$ and we have $r = 1$ as well, which contradicts the assumptions. So $r \geq 3$. Following the proof of \Leftarrow of theorem 2.1, we see that $M^{r-1} = 0$ gives $A_{r-1} \cdot A_{r-2} \cdots A_1 = 0$ for nonzero matrices A_i , which consist of only one entry when $r = m$. This is a contradiction, so $r \leq m - 1$. \square

Using (2) of corollary 2.2, one can construct a map H in dimension $n = 4$, such that $\mathcal{J}H$ has strong nilpotency index $r = 3$ (in the sense of definition 1.2) and regular nilpotency index two. Take for instance the (cubic) map

$$H = (0, x_1^2, x_1^3, 3x_2x_1^2 - 2x_3x_1)$$

in dimension four, or the (cubic) homogeneous map

$$H = (0, 0, x_1^3, x_1^2x_2, x_1x_2^2, x_3x_2^2 - 2x_4x_1x_2 + x_5x_1^2)$$

in dimension six. By proposition 2.3, we see that there is no other combination of strong nilpotency index r and matrix dimension m , such that $n \leq 4$ and r exceeds the regular nilpotency index.

The dimension $n = 6$ of the second homogeneous map H is minimal as well, at least if the base ring K is a field, but only for $r = 3$. The minimality of n for $r = 3$ can be seen as follows. If a homogeneous map H of degree d in dimension $n \leq 5$ with similar properties would exist, then by $\mathcal{J}H \cdot \mathcal{J}H = 0$, we have $\text{rk } \mathcal{J}H \leq \lfloor n/2 \rfloor \leq \lfloor 5/2 \rfloor = 2$. Hence $\text{rk } A_2 = \text{rk } A_1 = 1$. By homogeneity of H , the relations between the rows of A_1 decompose into linear relations, see e.g. the proof of [Ch, Lm. 3]. Consequently, the image of A_1 is determined by linear constraints and hence generated by vectors v over K .

Since $\text{rk } A_1 = 1$, the image of A_1 is generated by exactly one vector v over K . For this vector v , we have $C \cdot v = 0$ for all coefficient matrices C of A_2 in case $A_2 \cdot v = 0$. Following the proof of \Leftarrow of theorem 2.1, we see that $(\mathcal{J}H)^2 = 0$ gives $A_2 \cdot A_1 = 0$, so $A_2 \cdot v = 0$. Hence $C \cdot v = 0$ and $C \cdot A_1 = 0$ for all coefficient matrices C of A_2 . Following the proof of \Leftarrow of theorem 2.1 again, we get a contradiction with $r = 3$.

There is however a homogeneous map in dimension $n = 5$, from which the Jacobian M is strongly nilpotent with index $r = 4$ and regular nilpotency index three, namely

$$H = (0, x_1^d, x_1^{d-1}x_2, x_1^{d-2}x_2^2, 2x_1^{d-2}x_2x_3 - x_1^{d-1}x_4)$$

By proposition 2.3 and the study of the case $r = 2$ above, we see that there is no other combination of strong nilpotency index r and dimension n , such that M is a homogeneous Jacobian of a polynomial map over a field, $n \leq 5$ and r exceeds the regular nilpotency index.

Noncommutative polynomials

Notice that in theorem 2.1, one may also assume that the indeterminates of x are noncommutative, provided the indeterminates of $y^{(j)}$ are noncommutative for each j as well.

In [Y2], the author Jie-Tai Yu proves that a homogeneous nilpotent matrix M of polynomials over a field K is linearly triangularizable if there exist a (homogeneous) nilpotent matrix of noncommutative polynomials (free algebra) which corresponds to M . This result can be generalized as follows.

Theorem 2.4. *Let M be a matrix of size $m \times m$ of polynomials in noncommutative indeterminates $x = (x_i \mid i \in I)$ of (weighted) degree d over a division ring D .*

Then M has nilpotency index r , if and only if M has strong nilpotency index r (in the sense of definition 1.2), if and only if there exist a $T \in \text{GL}_m(D)$ such that $T^{-1}MT$ is of the form of (2.1) in theorem 2.1.

In the definition of strong nilpotency (in definition 1.2), for each i, j, k, l , it does not matter whether $y_i^{(j)}y_k^{(l)}$ and $y_k^{(l)}y_i^{(j)}$ are equalized (by commutativity assumptions) or not in case $j \neq l$. It is only necessary that $y_i^{(j)}y_k^{(j)}$ and $y_k^{(j)}y_i^{(j)}$ are not equalized (by commutativity assumptions). So $y_i^{(j)}y_k^{(j)}$ and $y_k^{(j)}y_i^{(j)}$ are equal, if and only if $i = k$.

Proof. For simplicity in writing, we assume that all entries of M are homogeneous of degree d instead of weighted homogeneous of degree d .

Suppose that $y^{(j)} = (y_i^{(j)} \mid i \in I)$ are partially noncommutative polynomial indeterminates satisfying the above noncommutativity properties. The noncommutative terms of degree rd in x correspond to (partially) noncommutative products of r terms of degree d in $y^{(r)}, y^{(r-1)}, \dots, y^{(1)}$, in that order, by

$$\begin{aligned} x_{i_1}x_{i_2} \cdots x_{i_{rd}} &\mapsto y_{i_1}^{(r)}y_{i_2}^{(r)} \cdots y_{i_d}^{(r)} \\ &\quad y_{i_{d+1}}^{(r-1)}y_{i_{d+2}}^{(r-1)} \cdots y_{i_{2d}}^{(r-1)} \\ &\quad \vdots \\ &\quad y_{i_{(r-1)d+1}}^{(1)}y_{i_{(r-1)d+2}}^{(1)} \cdots y_{i_{rd}}^{(1)} \end{aligned}$$

and each entry of $M^r = 0$ corresponds to the same entry of $M|_{x=y^{(r)}} \cdot M|_{x=y^{(r-1)}} \cdots M|_{x=y^{(1)}}$ in that manner, whence the latter matrix product vanishes as well as the former.

The rest of the proof is similar to that of theorem 2.1. \square

Taking the Jacobian of $H = (0, x_1x_2 - x_3, x_1^2x_2 - x_1x_3)$, we get

$$M = \begin{pmatrix} 0 & 0 & 0 \\ x_2 & x_1 & -1 \\ 2x_1x_2 & x_1^2 & -x_1 \end{pmatrix}$$

even if we see the components of H as noncommutative polynomials. Since the trace of $M \cdot M_{x=y}$ equals

$$0 + (x_1y_1 - y_2) - (x_2 + x_1y_1) = -y_2 - x_2 \neq 0$$

we see that M is not strongly nilpotent. But $M^3 = 0$ holds, regardless of whether we see the entries as commutative or noncommutative polynomials over \mathbb{C} . Hence the above theorem does not hold when M is not homogeneous.

3 Strongly nilpotent matrices in the sense of [MO]

In [Y1], the notion of strong nilpotency is generalized as follows. Instead of looking at Jacobians with polynomial entries, the author Jie-Tai Yu looks at arbitrary matrices of functions of a set S to a division ring D . Not all polynomials can be described as the functions they represent, e.g. taking the first variable over a finite field does not describe x_1 because the polynomial x_1^q corresponds to the identity as well over a finite field with q elements. But over an infinite field, different polynomials correspond to different functions.

For such matrices of functions, Yu proves that they are linearly triangularizable over D when M is so called *generalized strongly nilpotent*, which means that for some $r \in \mathbb{N}$,

$$M|_{v_1} \cdot M|_{v_2} \cdots M|_{v_r} = 0 \quad (3.1)$$

for all $v_i \in S$, where $f|_g$ means substituting g in the entries of f . Again, we call the minimum r such that (3.1) holds the *strong nilpotency index* of M . We can generalize Yu's result in a similar manner as in theorem 2.1.

Theorem 3.1. *Let $m \geq 1$ and $M \in \text{Mat}_m(D^S)$ be a matrix of functions from a set S to a division ring D .*

Then M has strong nilpotency index r (in the sense of definition 1.1, but with functions that do not need to be polynomials), if and only if there exists a $T \in \text{GL}_m(D)$ such that $T^{-1}MT$ is of the form

$$\begin{pmatrix} 0_{s_1} & & & & & \emptyset \\ A_1 & 0_{s_2} & & & & \\ & A_2 & \ddots & & & \\ & & \ddots & 0_{s_{r-1}} & & \\ * & & & A_{r-1} & 0_{s_r} & \end{pmatrix}$$

where 0_{s_i} is the square zero matrix of size $s_i \geq 1$ and A_i has independent columns over D for each i . In particular, the strong nilpotency index of M does not exceed m in case M is strongly nilpotent.

Proof. Since the proof is essentially similar to that of theorem 2.1, we only describe a part of the implication \Rightarrow .

Assume that M has strong nilpotency index r . Then

$$M|_{v_r} \cdot M|_{v_{r-1}} \cdots M|_{v_2} \cdot M|_{v_1} = 0 \quad (3.2)$$

for all $v_1, v_2, \dots, v_{r-1}, v_r \in S$, but

$$M|_{v_{r-1}} \cdots M|_{v_2} \cdot M|_{v_1} \neq 0 \quad (3.3)$$

for certain $v_1, v_2, \dots, v_{r-1} \in S$. Thus for these $v_1, v_2, \dots, v_{r-1} \in S$, the left hand side of (3.3) becomes a nonzero matrix $C \in \text{Mat}_m(D)$, and we obtain from (3.2) that $M|_{v_r} \cdot C = 0$ for all $v_r \in S$. \square

Notice that the strong nilpotency index does not exceed the dimension m of the matrix in both theorem 2.1 and theorem 3.1. We will prove below that this property still holds when the base division ring D is replaced by a reduced commutative ring R , see also [E, Exercise 7.4.2]. If R is a ring with unity which is not reduced, then there exists an $\epsilon \in R$ such that $\epsilon^2 = 0$, and for the matrix M defined by

$$M := \begin{pmatrix} \epsilon & 0 & \cdots & 0 & 0 \\ 1^2 & & & \emptyset & 0 \\ & 1^3 & & & \vdots \\ & & \ddots & & 0 \\ \emptyset & & & 1^m & 0 \end{pmatrix}$$

(where the powers of 1 are only taken to indicate that the size is m) we have that M^k is obtained from M by shifting all rows $k - 1$ places to below, where zero rows are shifted in above for each row that is shifted out below. Hence M^m is zero except for the lower left corner entry which equals ϵ , and the both the regular and the strong nilpotency index of M equal $m + 1$.

Proposition 3.2. *Let M be a square matrix of size m of functions from a set S to a reduced commutative ring R with unity. If there exists an $r \in \mathbb{N}$ such that $M|_{v_r} M|_{v_{r-1}} \cdots M|_{v_1} = 0$ for all $v_i \in S$, then $M|_{v_m} M|_{v_{m-1}} \cdots M|_{v_1} = 0$ for all $v_i \in S$.*

Proof. Suppose that there exists an $r \in \mathbb{N}$ such that $M|_{v_r} M|_{v_{r-1}} \cdots M|_{v_1} = 0$ for all $v_i \in S$. Since the intersection of all prime ideals of R is zero, it suffices to show that for every prime ideal \mathfrak{p} of R , all entries of $M|_{v_m} M|_{v_{m-1}} \cdots M|_{v_1}$ are contained in \mathfrak{p} for all $v_i \in S$. For that purpose, we look at residue classes of \mathfrak{p} and hence replace R by R/\mathfrak{p} , which is an integral domain. Let K be the field of fractions of R/\mathfrak{p} . By theorem 3.1, M is linearly triangularizable over K , which gives the desired result. \square

For reduced noncommutative rings, proposition 3.2 can also be reduced to the domain case, but domains can not always be embedded in division rings. Therefore, we cannot prove the following.

Conjecture 3.3. *Let M be a square matrix of size m of functions from a set S to a reduced noncommutative ring R . If there exists an $r \in \mathbb{N}$ such that $M|_{v_r} M|_{v_{r-1}} \cdots M|_{v_1} = 0$ for all $v_i \in S$, then $M|_{v_m} M|_{v_{m-1}} \cdots M|_{v_1} = 0$ for all $v_i \in S$.*

4 Quasi-translations with strong nilpotency index two

A quasi-translation is a polynomial map $x + H$ whose inverse is $x - H$. In 1876 in [GN], the authors proved that if $x + H$ is a quasi-translation such that H is homogeneous and $\text{rk } \mathcal{J}H \leq 2$, then $x + g^{-1}H$ is a quasi-translation such that

$\mathcal{J}(g^{-1}H)$ satisfies the second and hence any of the properties of corollary 2.2 with strong nilpotency index two, where $g = \gcd\{H_1, H_2, \dots, H_n\}$. Actually, the authors look at polynomial maps $x + H$ such that $\mathcal{J}H \cdot H = 0$, but that is the same as that $x + H$ is a quasi-translation on account of [Bo, Prop. 1.1]. In [W], the author proved a similar result as above for non-homogeneous quasi-translations in dimension three.

At the end of this section, we will show for two other types of quasi-translations $x + H$ that $\mathcal{J}H$ itself satisfies the properties of corollary 2.2 with strong nilpotency index two. Quasi-translations are important in the study of polynomials whose Hessian determinant vanishes, see [GN] or the beginning of section 2 of [BoE].

Theorem 4.1 below shows that theorem 2.1 and corollary 2.2 can be used to find equivalent properties for quasi-translations to have strong nilpotency index two. It was the study of such quasi-translations which caused the author to write this article.

Theorem 4.1. *Assume $x + H$ is a quasi-translation over a field K of characteristic zero. Then the following statements are all equivalent to $\mathcal{J}H \cdot \mathcal{J}H|_{x=y} = 0$.*

- (1) $\mathcal{J}H \cdot H(y) = 0$.
- (2) *There exists a $T \in \text{GL}_n(K)$ and an s with $0 \leq s < n$ such that for $\hat{H} := T^{-1}H(Tx)$, we have $\hat{H}_i = 0$ for all $i \leq s$ and $\hat{H}_i \in K[x_1, x_2, \dots, x_s]$ for all $i > s$.*
- (3) $H(x + tH(y)) = H$.
- (4) $H(x + H(y)) = H$.

Proof. By taking the Jacobian of (1) with respect to y , $\mathcal{J}H \cdot \mathcal{J}H|_{x=y} = 0$ follows. Conversely, if $\mathcal{J}H \cdot \mathcal{J}H|_{x=y} = 0$, then by integration with respect to y , we have $\mathcal{J}H \cdot H(y) = \mathcal{J}H \cdot H(0)$. By substituting y by x on both sides, we obtain $\mathcal{J}H \cdot H = \mathcal{J}H \cdot H(0)$, and both equalities combine to $\mathcal{J}H \cdot H(y) = \mathcal{J}H \cdot H$. Hence the equivalence of (1) and $\mathcal{J}H \cdot \mathcal{J}H|_{x=y} = 0$ follows from $\mathcal{J}H \cdot H = 0$, which is (2) in the proof of [Bo, Prop. 1.1]. By substituting $x = y^{(2)}$ and $y = y^{(1)}$, and vice versa, we see that $\mathcal{J}H \cdot \mathcal{J}H|_{x=y} = 0$ is equivalent to (1) of corollary 2.2 with $r = 2$. So (1) is equivalent to (1) of corollary 2.2 with $r = 2$.

We shall show that (2) in turn is equivalent to (2) of corollary 2.2 with $r = 2$. It is clear that (2) of corollary 2.2 with $r = 2$ follows from (2). Conversely, assume (2) of corollary 2.2 with $r = 2$. If $\mathcal{J}H = 0$, then we have (2) with $s = 0$, so assume $\mathcal{J}H \neq 0$. By (2) \Rightarrow (1) of corollary 2.2 and $\mathcal{J}H \neq 0$, we obtain that the strong nilpotency index of $\mathcal{J}H$ equals two. Hence we may assume that $\hat{H} := T^{-1}H(Tx)$ is of the form of (2.1) in theorem 2.1. By applying the fact that (2) of corollary 2.2 with $r = 2$ implies (1) (of this theorem), but on \hat{H} instead of H , we see that $\mathcal{J}\hat{H} \cdot \hat{H}(y) = 0$. Now the independence of the columns of A_1 subsequently gives $\hat{H}_1 = \hat{H}_2 = \dots = \hat{H}_{s_1} = 0$, so we have (2) with $s = s_1$ by definition of \hat{H} .

In order to show that (3) and (4) are equivalent to (1) and (2) as well, it suffices to show that (3) of corollary 2.2 with $r = 2$ implies (3), that (3) implies (4), and that (4) implies (4) of corollary 2.2 with $r = 2$. The latter follows by substituting $x = y^{(2)}$ and $y = y^{(1)}$, because the Jacobian matrix with respect to $y^{(1)}$ of $H(y^{(2)})$ is zero. That (3) implies (4) follows by substituting $t = 1$. So it remains to show that (3) of corollary 2.2 with $r = 2$ implies (3).

For that purpose, suppose we have (3) of corollary 2.2 with $r = 2$. By taking $j = 2$, we get $t^{(2)}H(y^{(2)} + t^{(1)}H(y^{(1)})) = H(y^{(2)} + t^{(1)}H(0))$, which gives $H(x + tH(y)) = H(x + tH(0))$ after suitable substitutions. By substituting y by x on both sides, we obtain $H(x + tH(x)) = H(x + tH(0))$, and both equalities combine to $H(x + tH(y)) = H(x + tH(x))$. Hence (3) follows from $H(x + tH) = H$, which is (1) in the proof of [Bo, Prop. 1.1]. \square

(2) \Rightarrow (3) of the following theorem was proved by E. Formanek in [F, Th. 4] for the case that H has no linear terms, but such a condition is unnecessary.

Theorem 4.2. *Assume $F = x + H$ is a Keller map in dimension n over a field K of characteristic zero, such that $\text{rk } \mathcal{J}H = 1$. Then F is invertible and for*

- (1) H has no linear terms,
- (2) $\det \mathcal{J}F = 1$,
- (3) $\mathcal{J}H \cdot \mathcal{J}H|_{x=y} = 0$,

we have (1) \Rightarrow (2) \Rightarrow (3).

Proof. By [F, Lm. 3], we have $H_i \in K[p]$ for some $p \in K[x]$, and by way of Gaussian elimination on the coefficients of the H_i with respect to p , we see that there exists a $T \in \text{GL}_n(K)$ and an $s \geq 0$ such that for $\hat{H} = T^{-1}H(Tx)$, we have $\hat{H}_i \in K$ for all $i \leq s$ and $0 < \deg \hat{H}_{s+1} < \deg \hat{H}_{s+2} < \dots < \deg \hat{H}_n$. Furthermore, $\det \mathcal{J}(x + \hat{H}) \in K^*$ and $\hat{H}_i \in K[\hat{p}]$ for all i , where $\hat{p} = p(Tx)$. We shall show below that $\mathcal{J}(x + \hat{H})$ is a lower triangular matrix (but not necessarily with ones on the diagonal). Consequently, $x + \hat{H}$ is a composition of elementary invertible polynomial maps and F is a tame invertible map, because $\det \mathcal{J}(x + \hat{H}) \in K^*$.

Notice that $\det \mathcal{J}(x + \hat{H}) - 1$ is the sum of the determinants of all principal minor matrices of $\mathcal{J}\hat{H}$. Since $\text{rk } \mathcal{J}\hat{H} = 1$, all minor determinants of size 2×2 or greater vanish. Hence $\text{tr } \mathcal{J}\hat{H} = \det \mathcal{J}(x + \hat{H}) - 1 \in K$. Take i maximal, such that $\frac{\partial}{\partial x_i} \hat{p} \neq 0$. If $i \leq s$, then we are in the situation of (2) of corollary 2.2 and we have (3) and hence also (2), because $\mathcal{J}H$ is nilpotent and $\det \mathcal{J}F - 1$ is the sum of the determinants of all principal minor matrices of $\mathcal{J}H$.

Thus assume that $i > s$. Then $\frac{\partial}{\partial x_j} \hat{H}_j = 0$ for all $j > i$, $\deg \frac{\partial}{\partial x_j} \hat{H}_j < \deg \hat{H}_j \leq \deg \hat{H}_i - \deg \hat{p}$ for all $j < i$, and $\deg \hat{H}_i - \deg \hat{p} \leq \deg \frac{\partial}{\partial x_i} \hat{H}_i$ because $\frac{\partial}{\partial x_i} \hat{H}_i$ is divisible by a polynomial in \hat{p} of sufficiently large degree. Hence $\deg \frac{\partial}{\partial x_j} \hat{H}_j < \deg \frac{\partial}{\partial x_i} \hat{H}_i$ for all $j \neq i$ and therefore $\deg \frac{\partial}{\partial x_i} \hat{H}_i = \deg \text{tr } \mathcal{J}\hat{H} \leq 0$. Consequently, $\frac{\partial}{\partial x_j} \hat{H}_j = 0$ for all $j \neq i$ and \hat{H}_i has degree one in \hat{p} , which in

turn has degree one in $x_i = x_{s+1}$ with leading coefficient in K as a polynomial over $K[x_1, x_2, \dots, x_s]$. This contradicts (1). By $\hat{H}_j \in K$ for all $j \leq s$ and $\hat{H}_j \in K[x_1, x_2, \dots, x_s, x_{s+1}]$ for all $j > s$, we get $\det \mathcal{J}(x + \hat{H}) = 1 + \frac{\partial}{\partial x_i} \hat{H}_i \neq 1$, which contradicts (2) by way of (2.10). \square

Corollary 4.3. *Assume $x + H$ is a quasi-translation over a field K of characteristic zero, such that $1 \in \{\deg H, \text{rk } \mathcal{J}H\}$. Then $\mathcal{J}H$ has strong nilpotency index two, i.e. $\mathcal{J}H \cdot \mathcal{J}H|_{x=y} = 0$.*

Proof. Suppose first that $\deg H = 1$. Since $\mathcal{J}H$ is a constant matrix, the strong nilpotency index and the nilpotency index correspond. Furthermore, the part of degree one of $\mathcal{J}H \cdot H$ equals $(\mathcal{J}H)^2 \cdot x$, and $(\mathcal{J}H)^2 = 0$ follows.

Suppose next that $\text{rk } \mathcal{J}H = 1$. By [Bo, Prop. 1.1], $\mathcal{J}H$ is nilpotent, so $\det \mathcal{J}H = 1$. Hence the desired result follows from (2) \Rightarrow (3) of theorem 4.2. \square

Acknowledgment. The author wishes to thank the referee for his careful reading and for several valuable comments.

This is a preprint of an article submitted for consideration in *Linear and Multilinear Algebra* © 2012 copyright Taylor & Francis.

References

- [BCW] Hyman Bass, Edwin H. Connell and David Wright, *The Jacobian conjecture: reduction of degree and formal expansion of the inverse*. Bull. Amer. Math. Soc. (N.S.) 7 (1982), no. 2, 287–330.
- [BoE] Michiel de Bondt and Arno van den Essen, *Singular Hessians*. J. Algebra 282 (2004), no. 1, 195–204.
- [Bo] Michiel de Bondt, *Quasi-translations and counterexamples to the homogeneous dependence problem*. Proc. Amer. Math. Soc. 134 (2006), no. 10, 2849–2856 (electronic).
- [Ch] Charles Ching-An Cheng, *Power linear Keller maps of rank two are linearly triangularizable*. J. Pure Appl. Algebra (2005), no. 2, 127–130.
- [E] Arno van den Essen, *Polynomial automorphisms and the Jacobian Conjecture*. Progress in Mathematics, Vol. 190, Birkhäuser, 2000.
- [EH] Arno van den Essen and Engelbert Hubbers, *Polynomial maps with strongly nilpotent Jacobian matrix and the Jacobian conjecture*. Linear Algebra Appl. 247 (1996), 121–132.
- [F] E. Formanek, *Two notes on the Jacobian conjecture*. Arch. Math. (Basel) 49 (1987), no. 4, 286–291.
- [GN] P. Gordan and M. Nöther, *Über die algebraischen Formen, deren Hesse'sche Determinante identisch verschwindet*. (German) Math. Ann. 10 (1876), no. 4, 547–568.

- [MO] Gary H. Meisters and Czesław Olech, *Strong nilpotence holds in dimensions up to five only*. Linear and Multilinear Algebra 30 (1991), no. 4, 231–255.
- [W] Zhiqing Wang, *Homogeneization of locally nilpotent derivations and an application to $k[X, Y, Z]$* . J. Pure Appl. Algebra, 196 (2005), no. 2-3, 323–337.
- [Ya] A.V. Yagzhev, *Keller’s problem*. Siberian Math. J. 21 (1980), no. 5, 747–754 (1981).
- [Y1] Jie-Tai Yu, *On generalized strongly nilpotent matrices*. Linear and Multilinear Algebra 41 (1996), no. 1, 19–22.
- [Y2] Jie-Tai Yu, *Noncommutative-nilpotent matrices and the Jacobian conjecture*. Linear and Multilinear Algebra 41 (1996), no. 2, 189–192.