

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/139565>

Please be advised that this information was generated on 2018-01-18 and may be subject to change.

# Measurement of the Forward-Backward Asymmetry in the Production of $B^\pm$ Mesons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

V.M. Abazov,<sup>31</sup> B. Abbott,<sup>67</sup> B.S. Acharya,<sup>25</sup> M. Adams,<sup>46</sup> T. Adams,<sup>44</sup> J.P. Agnew,<sup>41</sup> G.D. Alexeev,<sup>31</sup> G. Alkhazov,<sup>35</sup> A. Alton<sup>a</sup>,<sup>56</sup> A. Askew,<sup>44</sup> S. Atkins,<sup>54</sup> K. Augsten,<sup>7</sup> C. Avila,<sup>5</sup> F. Badaud,<sup>10</sup> L. Bagby,<sup>45</sup> B. Baldin,<sup>45</sup> D.V. Bandurin,<sup>73</sup> S. Banerjee,<sup>25</sup> E. Barberis,<sup>55</sup> P. Baringer,<sup>53</sup> J.F. Bartlett,<sup>45</sup> U. Bassler,<sup>15</sup> V. Bazterra,<sup>46</sup> A. Bean,<sup>53</sup> M. Begalli,<sup>2</sup> L. Bellantoni,<sup>45</sup> S.B. Beri,<sup>23</sup> G. Bernardi,<sup>14</sup> R. Bernhard,<sup>19</sup> I. Bertram,<sup>39</sup> M. Besançon,<sup>15</sup> R. Beuselinck,<sup>40</sup> P.C. Bhat,<sup>45</sup> S. Bhatia,<sup>58</sup> V. Bhatnagar,<sup>23</sup> G. Blazey,<sup>47</sup> S. Blessing,<sup>44</sup> K. Bloom,<sup>59</sup> A. Boehnlein,<sup>45</sup> D. Boline,<sup>64</sup> E.E. Boos,<sup>33</sup> G. Borissoy,<sup>39</sup> M. Borysova<sup>l</sup>,<sup>38</sup> A. Brandt,<sup>70</sup> O. Brandt,<sup>20</sup> R. Brock,<sup>57</sup> A. Bross,<sup>45</sup> D. Brown,<sup>14</sup> X.B. Bu,<sup>45</sup> M. Buehler,<sup>45</sup> V. Buescher,<sup>21</sup> V. Bunichev,<sup>33</sup> S. Burdin<sup>b</sup>,<sup>39</sup> C.P. Buszello,<sup>37</sup> E. Camacho-Pérez,<sup>28</sup> B.C.K. Casey,<sup>45</sup> H. Castilla-Valdez,<sup>28</sup> S. Caughron,<sup>57</sup> S. Chakrabarti,<sup>64</sup> K.M. Chan,<sup>51</sup> A. Chandra,<sup>72</sup> E. Chapon,<sup>15</sup> G. Chen,<sup>53</sup> S.W. Cho,<sup>27</sup> S. Choi,<sup>27</sup> B. Choudhary,<sup>24</sup> S. Cihangir,<sup>45</sup> D. Claes,<sup>59</sup> J. Clutter,<sup>53</sup> M. Cooke<sup>k</sup>,<sup>45</sup> W.E. Cooper,<sup>45</sup> M. Corcoran,<sup>72</sup> F. Couderc,<sup>15</sup> M.-C. Cousinou,<sup>12</sup> D. Cutts,<sup>69</sup> A. Das,<sup>42</sup> G. Davies,<sup>40</sup> S.J. de Jong,<sup>29,30</sup> E. De La Cruz-Burelo,<sup>28</sup> F. Déliot,<sup>15</sup> R. Demina,<sup>63</sup> D. Denisov,<sup>45</sup> S.P. Denisov,<sup>34</sup> S. Desai,<sup>45</sup> C. Deterre<sup>c</sup>,<sup>41</sup> K. DeVaughan,<sup>59</sup> H.T. Diehl,<sup>45</sup> M. Diesburg,<sup>45</sup> P.F. Ding,<sup>41</sup> A. Dominguez,<sup>59</sup> A. Dubey,<sup>24</sup> L.V. Dudko,<sup>33</sup> A. Duperrin,<sup>12</sup> S. Dutt,<sup>23</sup> M. Eads,<sup>47</sup> D. Edmunds,<sup>57</sup> J. Ellison,<sup>43</sup> V.D. Elvira,<sup>45</sup> Y. Enari,<sup>14</sup> H. Evans,<sup>49</sup> V.N. Evdokimov,<sup>34</sup> A. Fauré,<sup>15</sup> L. Feng,<sup>47</sup> T. Ferbel,<sup>63</sup> F. Fiedler,<sup>21</sup> F. Filthaut,<sup>29,30</sup> W. Fisher,<sup>57</sup> H.E. Fisk,<sup>45</sup> M. Fortner,<sup>47</sup> H. Fox,<sup>39</sup> S. Fuess,<sup>45</sup> P.H. Garbincius,<sup>45</sup> A. Garcia-Bellido,<sup>63</sup> J.A. García-González,<sup>28</sup> V. Gavrilov,<sup>32</sup> W. Geng,<sup>12,57</sup> C.E. Gerber,<sup>46</sup> Y. Gershtein,<sup>60</sup> G. Ginther,<sup>45,63</sup> O. Gogota,<sup>38</sup> G. Golovanov,<sup>31</sup> P.D. Grannis,<sup>64</sup> S. Greder,<sup>16</sup> H. Greenlee,<sup>45</sup> G. Grenier,<sup>17</sup> Ph. Gris,<sup>10</sup> J.-F. Grivaz,<sup>13</sup> A. Grohsjean<sup>c</sup>,<sup>15</sup> S. Grünendahl,<sup>45</sup> M.W. Grünewald,<sup>26</sup> T. Guillemain,<sup>13</sup> G. Gutierrez,<sup>45</sup> P. Gutierrez,<sup>67</sup> J. Haley,<sup>68</sup> L. Han,<sup>4</sup> K. Harder,<sup>41</sup> A. Harel,<sup>63</sup> J.M. Hauptman,<sup>52</sup> J. Hays,<sup>40</sup> T. Head,<sup>41</sup> T. Hebbeker,<sup>18</sup> D. Hedin,<sup>47</sup> H. Hegab,<sup>68</sup> A.P. Heinson,<sup>43</sup> U. Heintz,<sup>69</sup> C. Hensel,<sup>1</sup> I. Heredia-De La Cruz<sup>d</sup>,<sup>28</sup> K. Herner,<sup>45</sup> G. Hesketh<sup>f</sup>,<sup>41</sup> M.D. Hildreth,<sup>51</sup> R. Hirosky,<sup>73</sup> T. Hoang,<sup>44</sup> J.D. Hobbs,<sup>64</sup> B. Hoeneisen,<sup>9</sup> J. Hogan,<sup>72</sup> M. Hohlfeld,<sup>21</sup> J.L. Holzbauer,<sup>58</sup> I. Howley,<sup>70</sup> Z. Hubacek,<sup>7,15</sup> V. Hynek,<sup>7</sup> I. Iashvili,<sup>62</sup> Y. Ilchenko,<sup>71</sup> R. Illingworth,<sup>45</sup> A.S. Ito,<sup>45</sup> S. Jabeen<sup>m</sup>,<sup>45</sup> M. Jaffré,<sup>13</sup> A. Jayasinghe,<sup>67</sup> M.S. Jeong,<sup>27</sup> R. Jesik,<sup>40</sup> P. Jiang,<sup>4</sup> K. Johns,<sup>42</sup> E. Johnson,<sup>57</sup> M. Johnson,<sup>45</sup> A. Jonckheere,<sup>45</sup> P. Jonsson,<sup>40</sup> J. Joshi,<sup>43</sup> A.W. Jung,<sup>45</sup> A. Juste,<sup>36</sup> E. Kajfasz,<sup>12</sup> D. Karmanov,<sup>33</sup> I. Katsanos,<sup>59</sup> M. Kaur,<sup>23</sup> R. Kehoe,<sup>71</sup> S. Kermiche,<sup>12</sup> N. Khalatyan,<sup>45</sup> A. Khanov,<sup>68</sup> A. Kharchilava,<sup>62</sup> Y.N. Kharzheev,<sup>31</sup> I. Kiselevich,<sup>32</sup> J.M. Kohli,<sup>23</sup> A.V. Kozelov,<sup>34</sup> J. Kraus,<sup>58</sup> A. Kumar,<sup>62</sup> A. Kupco,<sup>8</sup> T. Kurča,<sup>17</sup> V.A. Kuzmin,<sup>33</sup> S. Lammers,<sup>49</sup> P. Lebrun,<sup>17</sup> H.S. Lee,<sup>27</sup> S.W. Lee,<sup>52</sup> W.M. Lee,<sup>45</sup> X. Lei,<sup>42</sup> J. Lellouch,<sup>14</sup> D. Li,<sup>14</sup> H. Li,<sup>73</sup> L. Li,<sup>43</sup> Q.Z. Li,<sup>45</sup> J.K. Lim,<sup>27</sup> D. Lincoln,<sup>45</sup> J. Linnemann,<sup>57</sup> V.V. Lipaev,<sup>34</sup> R. Lipton,<sup>45</sup> H. Liu,<sup>71</sup> Y. Liu,<sup>4</sup> A. Lobodenko,<sup>35</sup> M. Lokajicek,<sup>8</sup> R. Lopes de Sa,<sup>45</sup> R. Luna-Garcia<sup>g</sup>,<sup>28</sup> A.L. Lyon,<sup>45</sup> A.K.A. Maciel,<sup>1</sup> R. Madar,<sup>19</sup> R. Magaña-Villalba,<sup>28</sup> S. Malik,<sup>59</sup> V.L. Malyshev,<sup>31</sup> J. Mansour,<sup>20</sup> J. Martínez-Ortega,<sup>28</sup> R. McCarthy,<sup>64</sup> C.L. McGivern,<sup>41</sup> M.M. Meijer,<sup>29,30</sup> A. Melnitchouk,<sup>45</sup> D. Menezes,<sup>47</sup> P.G. Mercadante,<sup>3</sup> M. Merkin,<sup>33</sup> A. Meyer,<sup>18</sup> J. Meyer<sup>i</sup>,<sup>20</sup> F. Miconi,<sup>16</sup> N.K. Mondal,<sup>25</sup> M. Mulhearn,<sup>73</sup> E. Nagy,<sup>12</sup> M. Narain,<sup>69</sup> R. Nayyar,<sup>42</sup> H.A. Neal,<sup>56</sup> J.P. Negret,<sup>5</sup> P. Neustroev,<sup>35</sup> H.T. Nguyen,<sup>73</sup> T. Nunnemann,<sup>22</sup> J. Orduna,<sup>72</sup> N. Osman,<sup>12</sup> J. Osta,<sup>51</sup> A. Pal,<sup>70</sup> N. Parashar,<sup>50</sup> V. Parihar,<sup>69</sup> S.K. Park,<sup>27</sup> R. Partridge<sup>e</sup>,<sup>69</sup> N. Parua,<sup>49</sup> A. Patwa<sup>j</sup>,<sup>65</sup> B. Penning,<sup>45</sup> M. Perfilov,<sup>33</sup> Y. Peters,<sup>41</sup> K. Petridis,<sup>41</sup> G. Petrillo,<sup>63</sup> P. Pétroff,<sup>13</sup> M.-A. Pleier,<sup>65</sup> V.M. Podstavkov,<sup>45</sup> A.V. Popov,<sup>34</sup> M. Prewitt,<sup>72</sup> D. Price,<sup>41</sup> N. Prokopenko,<sup>34</sup> J. Qian,<sup>56</sup> A. Quadt,<sup>20</sup> B. Quinn,<sup>58</sup> P.N. Ratoff,<sup>39</sup> I. Razumov,<sup>34</sup> I. Ripp-Baudot,<sup>16</sup> F. Rizatdinova,<sup>68</sup> M. Rominsky,<sup>45</sup> A. Ross,<sup>39</sup> C. Royon,<sup>15</sup> P. Rubinov,<sup>45</sup> R. Ruchti,<sup>51</sup> G. Sajot,<sup>11</sup> A. Sánchez-Hernández,<sup>28</sup> M.P. Sanders,<sup>22</sup> A.S. Santos<sup>h</sup>,<sup>1</sup> G. Savage,<sup>45</sup> M. Savitskyi,<sup>38</sup> L. Sawyer,<sup>54</sup> T. Scanlon,<sup>40</sup> R.D. Schamberger,<sup>64</sup> Y. Scheglov,<sup>35</sup> H. Schellman,<sup>48</sup> C. Schwanenberger,<sup>41</sup> R. Schwienhorst,<sup>57</sup> J. Sekaric,<sup>53</sup> H. Severini,<sup>67</sup> E. Shabalina,<sup>20</sup> V. Shary,<sup>15</sup> S. Shaw,<sup>41</sup> A.A. Shchukin,<sup>34</sup> V. Simak,<sup>7</sup> P. Skubic,<sup>67</sup> P. Slattey,<sup>63</sup> D. Smirnov,<sup>51</sup> G.R. Snow,<sup>59</sup> J. Snow,<sup>66</sup> S. Snyder,<sup>65</sup> S. Söldner-Rembold,<sup>41</sup> L. Sonnenschein,<sup>18</sup> K. Soustruznik,<sup>6</sup> J. Stark,<sup>11</sup> D.A. Stoyanova,<sup>34</sup> M. Strauss,<sup>67</sup> L. Suter,<sup>41</sup> P. Svoisky,<sup>67</sup> M. Titov,<sup>15</sup> V.V. Tokmenin,<sup>31</sup> Y.-T. Tsai,<sup>63</sup> D. Tsybychev,<sup>64</sup> B. Tuchming,<sup>15</sup> C. Tully,<sup>61</sup> L. Uvarov,<sup>35</sup> S. Uvarov,<sup>35</sup> S. Uzunyan,<sup>47</sup> R. Van Kooten,<sup>49</sup> W.M. van Leeuwen,<sup>29</sup> N. Varelas,<sup>46</sup> E.W. Varnes,<sup>42</sup> I.A. Vasilyev,<sup>34</sup> A.Y. Verkheev,<sup>31</sup> L.S. Vertogradov,<sup>31</sup> M. Verzocchi,<sup>45</sup> M. Vesterinen,<sup>41</sup> D. Vilanova,<sup>15</sup> P. Vokac,<sup>7</sup> H.D. Wahl,<sup>44</sup> M.H.L.S. Wang,<sup>45</sup> J. Warchol,<sup>51</sup> G. Watts,<sup>74</sup> M. Wayne,<sup>51</sup> J. Weichert,<sup>21</sup> L. Welty-Rieger,<sup>48</sup>

M.R.J. Williams<sup>n,49</sup> G.W. Wilson,<sup>53</sup> M. Wobisch,<sup>54</sup> D.R. Wood,<sup>55</sup> T.R. Wyatt,<sup>41</sup> Y. Xie,<sup>45</sup> R. Yamada,<sup>45</sup> S. Yang,<sup>4</sup> T. Yasuda,<sup>45</sup> Y.A. Yatsunenkov,<sup>31</sup> W. Ye,<sup>64</sup> Z. Ye,<sup>45</sup> H. Yin,<sup>45</sup> K. Yip,<sup>65</sup> S.W. Youn,<sup>45</sup> J.M. Yu,<sup>56</sup> J. Zennamo,<sup>62</sup> T.G. Zhao,<sup>41</sup> B. Zhou,<sup>56</sup> J. Zhu,<sup>56</sup> M. Zielinski,<sup>63</sup> D. Zieminska,<sup>49</sup> and L. Zivkovic<sup>14</sup>

(The D0 Collaboration\*)

<sup>1</sup>LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

<sup>2</sup>Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>3</sup>Universidade Federal do ABC, Santo André, Brazil

<sup>4</sup>University of Science and Technology of China, Hefei, People's Republic of China

<sup>5</sup>Universidad de los Andes, Bogotá, Colombia

<sup>6</sup>Charles University, Faculty of Mathematics and Physics,  
Center for Particle Physics, Prague, Czech Republic

<sup>7</sup>Czech Technical University in Prague, Prague, Czech Republic

<sup>8</sup>Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

<sup>9</sup>Universidad San Francisco de Quito, Quito, Ecuador

<sup>10</sup>LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France

<sup>11</sup>LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3,  
Institut National Polytechnique de Grenoble, Grenoble, France

<sup>12</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

<sup>13</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

<sup>14</sup>LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France

<sup>15</sup>CEA, Irfu, SPP, Saclay, France

<sup>16</sup>IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France

<sup>17</sup>IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France

<sup>18</sup>III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany

<sup>19</sup>Physikalisches Institut, Universität Freiburg, Freiburg, Germany

<sup>20</sup>II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany

<sup>21</sup>Institut für Physik, Universität Mainz, Mainz, Germany

<sup>22</sup>Ludwig-Maximilians-Universität München, München, Germany

<sup>23</sup>Panjab University, Chandigarh, India

<sup>24</sup>Delhi University, Delhi, India

<sup>25</sup>Tata Institute of Fundamental Research, Mumbai, India

<sup>26</sup>University College Dublin, Dublin, Ireland

<sup>27</sup>Korea Detector Laboratory, Korea University, Seoul, Korea

<sup>28</sup>CINVESTAV, Mexico City, Mexico

<sup>29</sup>Nikhef, Science Park, Amsterdam, the Netherlands

<sup>30</sup>Radboud University Nijmegen, Nijmegen, the Netherlands

<sup>31</sup>Joint Institute for Nuclear Research, Dubna, Russia

<sup>32</sup>Institute for Theoretical and Experimental Physics, Moscow, Russia

<sup>33</sup>Moscow State University, Moscow, Russia

<sup>34</sup>Institute for High Energy Physics, Protvino, Russia

<sup>35</sup>Petersburg Nuclear Physics Institute, St. Petersburg, Russia

<sup>36</sup>Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Física d'Altes Energies (IFAE), Barcelona, Spain

<sup>37</sup>Uppsala University, Uppsala, Sweden

<sup>38</sup>Taras Shevchenko National University of Kyiv, Kiev, Ukraine

<sup>39</sup>Lancaster University, Lancaster LA1 4YB, United Kingdom

<sup>40</sup>Imperial College London, London SW7 2AZ, United Kingdom

<sup>41</sup>The University of Manchester, Manchester M13 9PL, United Kingdom

<sup>42</sup>University of Arizona, Tucson, Arizona 85721, USA

<sup>43</sup>University of California Riverside, Riverside, California 92521, USA

<sup>44</sup>Florida State University, Tallahassee, Florida 32306, USA

<sup>45</sup>Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

<sup>46</sup>University of Illinois at Chicago, Chicago, Illinois 60607, USA

<sup>47</sup>Northern Illinois University, DeKalb, Illinois 60115, USA

<sup>48</sup>Northwestern University, Evanston, Illinois 60208, USA

<sup>49</sup>Indiana University, Bloomington, Indiana 47405, USA

<sup>50</sup>Purdue University Calumet, Hammond, Indiana 46323, USA

<sup>51</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA

<sup>52</sup>Iowa State University, Ames, Iowa 50011, USA

<sup>53</sup>University of Kansas, Lawrence, Kansas 66045, USA

<sup>54</sup>Louisiana Tech University, Ruston, Louisiana 71272, USA

<sup>55</sup>Northeastern University, Boston, Massachusetts 02115, USA

<sup>56</sup>University of Michigan, Ann Arbor, Michigan 48109, USA

<sup>57</sup>Michigan State University, East Lansing, Michigan 48824, USA

<sup>58</sup>University of Mississippi, University, Mississippi 38677, USA

<sup>59</sup>University of Nebraska, Lincoln, Nebraska 68588, USA

<sup>60</sup>Rutgers University, Piscataway, New Jersey 08855, USA

<sup>61</sup>Princeton University, Princeton, New Jersey 08544, USA

<sup>62</sup>State University of New York, Buffalo, New York 14260, USA

<sup>63</sup>University of Rochester, Rochester, New York 14627, USA

<sup>64</sup>State University of New York, Stony Brook, New York 11794, USA

<sup>65</sup>Brookhaven National Laboratory, Upton, New York 11973, USA

<sup>66</sup>Langston University, Langston, Oklahoma 73050, USA

<sup>67</sup>University of Oklahoma, Norman, Oklahoma 73019, USA

<sup>68</sup>Oklahoma State University, Stillwater, Oklahoma 74078, USA

<sup>69</sup>Brown University, Providence, Rhode Island 02912, USA

<sup>70</sup>University of Texas, Arlington, Texas 76019, USA

<sup>71</sup>Southern Methodist University, Dallas, Texas 75275, USA

<sup>72</sup>Rice University, Houston, Texas 77005, USA

<sup>73</sup>University of Virginia, Charlottesville, Virginia 22904, USA

<sup>74</sup>University of Washington, Seattle, Washington 98195, USA

(Dated: December 22, 2014)

We present a measurement of the forward-backward asymmetry in the production of  $B^\pm$  mesons,  $A_{\text{FB}}(B^\pm)$ , using  $B^\pm \rightarrow J/\psi K^\pm$  decays in  $10.4 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  collected by the D0 experiment during Run II of the Tevatron collider. A nonzero asymmetry would indicate a preference for a particular flavor, i.e.,  $b$  quark or  $\bar{b}$  antiquark, to be produced in the direction of the proton beam. We extract  $A_{\text{FB}}(B^\pm)$  from a maximum likelihood fit to the difference between the numbers of forward- and backward-produced  $B^\pm$  mesons. We measure an asymmetry consistent with zero:  $A_{\text{FB}}(B^\pm) = [-0.24 \pm 0.41 (\text{stat}) \pm 0.19 (\text{syst})]\%$ .

PACS numbers: 13.25.Hw, 11.30.Er

Over the few past years there has been much interest in the forward-backward asymmetry in  $t\bar{t}$  production ( $A_{\text{FB}}^{t\bar{t}}$ ) [1], especially since initial experimental results were larger than standard model (SM) predictions [2, 3]. These observations prompted development of models beyond the SM that could explain the excess [4]. The corresponding asymmetry in  $b\bar{b}$  production,  $A_{\text{FB}}^{b\bar{b}}$ , has the same sources as  $A_{\text{FB}}^{t\bar{t}}$  but is expected to have a smaller magnitude in the SM, making it an important probe of these new physics models [5, 6].

The most recent D0 measurements of  $A_{\text{FB}}^{t\bar{t}}$  [7] agree with the SM [8]. A closely related quantity called the  $t\bar{t}$  charge asymmetry has been studied at the LHC [9, 10]. The LHCb collaboration has recently measured the charge asymmetry between  $b$  and  $\bar{b}$  jets in  $pp$  col-

lisions [11].

A forward-backward asymmetry in the production of heavy quark  $Q$  is primarily caused by interference between tree-level and loop diagrams for  $q\bar{q} \rightarrow Q\bar{Q}$  interactions, and also by interference between initial and final state gluon radiation [12]. We measure the forward-backward asymmetry using fully reconstructed  $B^\pm \rightarrow J/\psi(\rightarrow \mu^+\mu^-)K^\pm$  decays where the  $B^\pm$  directly identifies the quark flavor (i.e.,  $b$  or  $\bar{b}$ ). Compared to  $b$  jet reconstruction, this method has the advantages that the charge of the  $b$  quark is unambiguously determined, and there is no need to account for  $B^0/\bar{B}^0$  oscillations. The quantity  $A_{\text{FB}}(B^\pm)$  is sensitive to the same production asymmetries as  $A_{\text{FB}}^{b\bar{b}}$ . In  $p\bar{p}$  collisions, the forward category indicates a  $b$  ( $\bar{b}$ ) quark, or  $B^-$  ( $B^+$ ) meson, emitted with a longitudinal momentum component in the direction of the proton (antiproton) beam.

We reconstruct a  $B^\pm$  meson and categorize it as forward or backward with a variable  $q_{\text{FB}} = -q_B \text{sgn}(\eta_B)$ , where  $q_B$  is the  $B^\pm$  meson electric charge,  $\text{sgn}(x)$  is the sign function, and  $\eta_B$  is the  $B^\pm$  meson pseudorapidity [13]. The forward-backward asymmetry of the  $B^\pm$  mesons is

$$A_{\text{FB}}(B^\pm) = \frac{N(q_{\text{FB}} > 0) - N(q_{\text{FB}} < 0)}{N(q_{\text{FB}} > 0) + N(q_{\text{FB}} < 0)}. \quad (1)$$

Inclusive predictions of  $A_{\text{FB}}^{b\bar{b}}$  give positive asymmetries of  $\approx 0.5\%$  [5, 14], but the mass scales of the  $b\bar{b}$  pairs considered [ $M(b\bar{b}) > 35 \text{ GeV}$ , or  $p(b) > \approx 15 \text{ GeV}$ ] are more

---

\*with visitors from <sup>a</sup>Augustana College, Sioux Falls, SD, USA, <sup>b</sup>The University of Liverpool, Liverpool, UK, <sup>c</sup>DESY, Hamburg, Germany, <sup>d</sup>Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Mexico <sup>e</sup>SLAC, Menlo Park, CA, USA, <sup>f</sup>University College London, London, UK, <sup>g</sup>Centro de Investigacion en Computacion - IPN, Mexico City, Mexico, <sup>h</sup>Universidade Estadual Paulista, São Paulo, Brazil, <sup>i</sup>Karlsruher Institut für Technologie (KIT) - Steinbuch Centre for Computing (SCC), D-76128 Karlsruhe, Germany, <sup>j</sup>Office of Science, U.S. Department of Energy, Washington, D.C. 20585, USA, <sup>k</sup>American Association for the Advancement of Science, Washington, D.C. 20005, USA, <sup>l</sup>Kiev Institute for Nuclear Research, Kiev, Ukraine, <sup>m</sup>University of Maryland, College Park, Maryland 20742, USA and <sup>n</sup>European Organization for Nuclear Research (CERN), Geneva, Switzerland

relevant for a jet-based analysis. To make SM predictions tailored to our kinematics and selections, we produce next-to-leading-order Monte Carlo (MC) samples for QCD production of  $B^\pm$  in the process  $p\bar{p} \rightarrow b\bar{b}X$ . MC events are generated using MC@NLO [15] with parton distribution function (PDF) set CTEQ6M1 [16] and HERWIG [17] for parton showering and hadronization. Detector simulation is performed using GEANT3 [18].

The D0 experiment collected data at  $\sqrt{s} = 1.96$  TeV during Run II of the Fermilab Tevatron  $p\bar{p}$  collider, from 2002 until the Tevatron shutdown in 2011. The D0 detector is described in detail elsewhere [19]. For this analysis the most important detector elements are the central tracking and muon systems. The central tracking system consists of a silicon microstrip tracker and a central fiber tracker, both located within a 1.9 T superconducting solenoidal magnet, with designs optimized for tracking and vertex finding at pseudorapidities  $|\eta| < 3$  and  $|\eta| < 2.5$ , respectively. The muon system has a layer of tracking detectors and scintillation trigger counters outside a liquid argon sampling calorimeter and two similar layers outside a 1.8 T iron toroid [20], and covers the region  $|\eta_{\text{det}}| \approx 2$  where  $|\eta_{\text{det}}|$  is measured from the center of the detector. The solenoid and toroid magnet polarities were reversed approximately every two weeks giving nearly equal beam exposure to each polarity combination. The data used in this analysis were collected with a suite of single muon and dimuon triggers.

We select  $B^\pm \rightarrow J/\psi K^\pm$  candidates from the D0 Run II data set with an integrated luminosity of  $10.4 \text{ fb}^{-1}$ . Candidates are reconstructed by identifying a pair of oppositely charged muons (decay products of the  $J/\psi$  meson) produced along with a charged track (the  $K^\pm$  candidate) at a common vertex displaced from the  $p\bar{p}$  interaction vertex.

All tracks must lie within the pseudorapidity coverage of the muon and central tracking systems,  $|\eta| < 2.1$ . Selected muons have transverse momentum  $p_T > 1.5$  GeV, and  $K^\pm$  candidates have  $p_T > 0.7$  GeV. At least one muon must traverse both inner and outer layers of the muon detector. Both muons must match to tracks in the central tracking system. The  $J/\psi$  candidates with reconstructed invariant mass  $M(\mu^+\mu^-)$  between 2.7 and 3.45 GeV are accepted if their transverse decay length ( $L_{xy}$ ) uncertainty is less than 0.1 cm, where  $L_{xy}$  is the distance from the  $p\bar{p}$  vertex to a particle's decay vertex in the  $x$ - $y$  plane. The cosine of the pointing angle [21] must be greater than zero.

The combination of  $\mu^+$ ,  $\mu^-$ , and  $K^\pm$  tracks to form a  $B^\pm$  decay vertex must have  $\chi^2 < 16$  for 3 degrees of freedom, and the cosine of the  $B^\pm$  pointing angle must be above 0.8.  $B^\pm$  candidates are accepted if they are significantly displaced from the  $p\bar{p}$  vertex. Their transverse decay length significance (defined as  $L_{xy}$  divided by its uncertainty) must be greater than three. To calculate the  $B^\pm$  candidate mass we correct the muon momenta

by constraining  $M(\mu^+\mu^-)$  to the world average  $J/\psi$  meson mass [22]. The selected  $B^\pm$  mass range is 5.05 – 5.65 GeV.

Because definitions of forward and backward are tied directly to  $\text{sgn}(\eta_B)$ , the ambiguous region near  $\eta_B = 0$  is given special consideration. We compare  $\eta$  of the  $B^\pm$  mesons and their parent  $b$  quarks at the production and reconstruction levels in MC@NLO. Rejecting events with  $|\eta_B| < 0.1$  removes all  $B^\pm$  mesons reconstructed with incorrect  $q_{\text{FB}}$  without significantly affecting  $A_{\text{FB}}(B^\pm)$ . After the cut, more than 99.9% of  $B^\pm$  mesons give the same  $q_{\text{FB}}$  as the parent  $b$  quark, indicating minimal hadronization effects on  $A_{\text{FB}}(B^\pm)$ . The distribution of  $(\eta_b - \eta_B)$  has a rms width of 0.11.

Background rejection is improved using a boosted decision tree (BDT) [23] trained on a simulated MC signal sample and a background sample from data sidebands around the selected  $B^\pm$  mass range (4.0 – 5.05 and 5.65 – 7.0 GeV). Leading-order signal MC events are generated with PYTHIA [24] and processed through the same reconstruction code used for data. We weight MC events so that the  $p_T$  distributions of the muons match the distributions in data, which are affected by trigger inefficiencies. Additional weights are applied to match distributions of  $p_T(B^\pm)$ ,  $p_T(K^\pm)$ , and  $\chi^2$  of the  $B^\pm$  decay vertex fit to data distributions. Finally, we weight MC events so that the probability of reconstructing isolated muons or  $B^\pm$  candidates matches the probability measured in data. Isolated particles have no other tracks in a cone of size  $\Delta\mathcal{R} = 1$  around them, where  $\Delta\mathcal{R} = \sqrt{\Delta\phi^2 + \Delta\eta^2}$  is the angular separation between tracks. This weighting gives optimal agreement between data and simulation in all 40 BDT input variables, which include particle momenta, distances from the  $p\bar{p}$  vertex, decay lengths, pointing angles, isolation of the muons and  $B^\pm$  meson, and azimuthal angle separation for various particle pairs. A cut on the BDT discriminant is chosen to minimize the statistical uncertainty of  $A_{\text{FB}}(B^\pm)$ . After all cuts we find one  $B^\pm$  candidate in 98.5% of events, with the remainder having two or more candidates. All candidates are used independently in this analysis.

We extract  $A_{\text{FB}}(B^\pm)$  from a maximum likelihood fit incorporating a signal probability distribution and three background distributions (see below), which are functions of the reconstructed  $B^\pm$  mass  $m_{J/\psi K}$  and the kaon energy  $E_K$ . The signal distribution  $S(m_{J/\psi K}, E_K)$  is modeled by a double-Gaussian function with six parameters, where both Gaussians have the same mean but different widths. The widths have an exponential dependence on  $E_K$ . Signal parameters are allowed to differ for the  $\eta < -0.1$  and  $\eta > 0.1$  regions to account for slight differences in the magnetic field along the beam direction.

The background distribution  $P(m_{J/\psi K}, E_K)$  describes  $B^\pm \rightarrow J/\psi\pi^\pm$  events where the pion is assigned the kaon mass, creating an artificially high reconstructed  $B^\pm$  mass. Distribution  $P$  is a reflection of  $S$  with the mean



mass value shifted to account for the  $K/\pi$  mass difference and the widths scaled by a ratio of the mean mass values. Background distribution  $T(m_{J/\psi K})$  describes partially reconstructed decays of type  $B_x \rightarrow J/\psi h^\pm X$ , which have reconstructed mass lower than the  $B^\pm$  mass. Distribution  $T$  is empirically modeled using a threshold function with a floating inflection point and the slope fixed from MC simulation [25, 26]. Finally, the background distribution  $E(m_{J/\psi K}, E_K)$  describes combinatoric background and is modeled using an exponential function with three parameters, where the slope depends on  $E_K$ .

The unbinned fit minimizes LLH, the negative log of the likelihood function  $\mathcal{L}_n$  summed over  $N$  selected  $B^\pm$  candidates, each with weight  $w_n$  (defined below):

$$\text{LLH} = -2 \sum_{n=1}^N w_n \ln(\mathcal{L}_n). \quad (2)$$

Here  $\mathcal{L}_n$  is a function of the four probability density distributions, with each assigned sample fraction  $f_i$  and forward-backward asymmetry  $A_i$ . While systematic effects were studied, the  $A_i$  parameters were blinded by adding unknown random offsets. The likelihood  $\mathcal{L}_n$  has 26 parameters and is normalized to 1:

$$\begin{aligned} \mathcal{L}_n = & \alpha(E_K)[f_S(1 + q_{\text{FB}}A_S)S + f_P(1 + q_{\text{FB}}A_P)P \\ & + f_T(1 + q_{\text{FB}}A_T)T] + f_E(1 + q_{\text{FB}}A_E)E, \end{aligned} \quad (3)$$

where  $f_E = [1 - \alpha(E_K)(f_S + f_P + f_T)]$  and  $\alpha(E_K)$  uses three parameters to describe the dependence of the sample fractions on  $E_K$  [25].

Asymmetries in the detector material and  $J/\psi$  or  $K^\pm$  reconstruction between  $\eta < 0$  (the ‘‘north’’ side of the detector) and  $\eta > 0$  (the ‘‘south’’ side) can result in an apparent  $A_{\text{FB}}$ . A north-south asymmetry is defined as  $A_{\text{NS}} = (N_N - N_S)/(N_N + N_S)$ . Because  $B^+$  and  $B^-$  particles on the same side of the detector have opposite  $q_{\text{FB}}$ , corrections for north-south efficiency differences will generally cancel when determining  $A_{\text{FB}}(B^\pm)$ . We measure  $A_{\text{NS}}$  in data samples with no expected production asymmetries. Decays of  $\phi \rightarrow K^+K^-$  are used to measure  $A_{\text{NS}}(K^\pm)$ . Signal and background models are determined from MC simulation and a  $\chi^2$  minimization fit is performed simultaneously on north- and south-side data. We measure  $A_{\text{NS}}(K^\pm)$  in bins of leading kaon  $|\eta|$ ; there is no significant dependence on  $p_T$ . Integrated over all  $|\eta|$ ,  $A_{\text{NS}}(K^+) = (0.39 \pm 0.22)\%$  and  $A_{\text{NS}}(K^-) = (0.64 \pm 0.23)\%$ .

We measure  $A_{\text{NS}}(J/\psi)$  using prompt  $J/\psi \rightarrow \mu^+\mu^-$  decays.  $J/\psi$  mesons with significant  $L_{xy}$  are generally from  $B$  decays which could exhibit a north-south asymmetry due to  $A_{\text{FB}}(B^\pm)$ . To reduce the fraction of nonprompt  $J/\psi$  mesons to a negligible level we require the  $J/\psi$   $L_{xy}$  significance to be less than 1.5. Background events under the peak from 2.9 – 3.3 GeV are removed with a sideband subtraction, and  $A_{\text{NS}}(J/\psi)$  is calculated in bins of  $|\eta|$  and

$p_T$ . Integrated over all  $|\eta|$  and  $p_T$ ,  $A_{\text{NS}}(J/\psi) = (-0.41 \pm 0.04)\%$ .

Measured  $A_{\text{NS}}$  values are used to determine ‘‘efficiency weights’’  $w_{K^\pm}$  and  $w_{J/\psi}$  that equalize the relative reconstruction efficiencies on both sides of the detector. Applying these weights has a small effect on  $A_{\text{FB}}(B^\pm)$ : a shift of 0.06% from  $w_{K^\pm}$  and a shift of  $-0.01\%$  from  $w_{J/\psi}$ . Uncertainties on  $A_{\text{NS}}(J/\psi)$  and  $A_{\text{NS}}(K^\pm)$  contribute an uncertainty of 0.003% to  $A_{\text{FB}}(B^\pm)$ , determined using an ensemble test with 500 Gaussian variations of the  $A_{\text{NS}}$  values.

The total event weight is  $w_n = w_{\text{magnet}}w_{K^\pm}w_{J/\psi}$ , where  $w_{\text{magnet}}$  equalizes the number of events in eight settings of solenoid polarity, toroid polarity, and  $B^\pm$  charge. Equalizing the contribution from each magnet polarity combination removes tracking charge asymmetries to first order, since in one polarity a  $B^+$  is reconstructed with the same sign of curvature as a  $B^-$  in the opposite polarity. Also equalizing the number of  $B^+$  and  $B^-$  candidates eliminates the need to correct for different  $K^+$  and  $K^-$  interaction cross sections in the detector [27].

The weighted data sample contains 160360  $B^\pm$  candidates and the fit yields  $89328 \pm 349$   $B^\pm \rightarrow J/\psi K^\pm$  decays. Although the fit was unbinned, to visualize the data and fit quality, binned distributions of invariant mass  $M(J/\psi K)$  for the sum and the difference in the numbers of forward and backward  $B^\pm$  candidates with their projected fits are shown in Figs. 1 and 2. Over both mass distributions we obtain  $\chi^2/\text{ndf} = 249/214$ . We measure a signal asymmetry consistent with zero:  $A_{\text{FB}}(B^\pm) = [-0.24 \pm 0.41(\text{stat})]\%$ . The asymmetry is consistent over time and with  $B^+$  and  $B^-$  samples fitted separately. Asymmetries of the background distributions are also consistent with zero.

To determine systematic uncertainties on  $A_{\text{FB}}(B^\pm)$  a number of variations are made to the analysis. Data sample variations include training four alternative BDTs with different variables or input samples and using a range of BDT discriminant cuts. Fit variations include varying the  $B^\pm$  mass range, removing dependences on  $E_K$  from the distributions, allowing the slope of  $T(m_{J/\psi K})$  to float, and fixing the background asymmetry parameters to zero.

To estimate the systematic error from the reconstruction asymmetries we measure  $A_{\text{NS}}(J/\psi)$  and  $A_{\text{NS}}(K^\pm)$  using alternate data samples and calculations in different bins or with alternate fit parameters. Biases in the fitting procedure are explored with ensemble tests on randomized data, comparing input and fitted values of  $A_{\text{FB}}(B^\pm)$ . No bias is observed, and a systematic uncertainty is assigned based on the spread of results in the ensemble test. The total systematic uncertainty on the data measurement is 0.19%, as summarized in Table I.

To compare this measurement to the SM, the MC@NLO simulation is analyzed as described above, applying  $B^\pm \rightarrow J/\psi K^\pm$  selections and weights to correct for muon




FIG. 1: (color online) Invariant mass  $M(J/\psi K)$  of (forward + backward) events with fitted distributions. The lower pane shows the residuals.




FIG. 2: (color online) Invariant mass  $M(J/\psi K)$  of (forward - backward) events with fitted distributions which include the asymmetry parameters  $A_i$ .

trigger effects. Additionally, reconstructed muon and kaon tracks must match tracks from generated  $B^\pm \rightarrow J/\psi K^\pm$  decays. Since matching reconstructed and generated  $B^\pm$  mesons leaves no background events,  $A_{\text{FB}}^{\text{SM}}(B^\pm)$  is calculated directly according to Eq. (1).

The dominant systematic uncertainty on  $A_{\text{FB}}^{\text{SM}}(B^\pm)$  is due to renormalization and factorization energy scale choices. MC@NLO defines  $\mu_R$  and  $\mu_F$  for renormalization and factorization energy scales [15] as the square root of the average of  $m_T^2 = m^2 + p_T^2$  for the  $b$  and  $\bar{b}$  quarks [28], with  $b$  quark mass  $m$  set to 4.75 GeV. Since  $A_{\text{FB}}^{b\bar{b}}$  is zero at leading order, there is a large scale dependence in predictions at next-to-leading order [29]. Both scales are varied independently from  $\frac{1}{2}\mu_{R,F}$  to  $2\mu_{R,F}$  to estimate

TABLE I: Summary of uncertainties on  $A_{\text{FB}}(B^\pm)$  in data.

Source	Uncertainty
Statistical	0.41%
Alternative BDTs and cuts	0.17%
Fit variations	0.06%
Reconstruction asymmetries	0.05%
Fit bias	0.02%
Systematic uncertainty	0.19%
Total uncertainty	0.45%

an uncertainty due to uncalculated higher orders. Half the largest spread of variations gives a systematic uncertainty of 0.44%. The uncertainty on  $A_{\text{FB}}^{\text{SM}}(B^\pm)$  due to  $b$  quark fragmentation is estimated by weighting events so the distribution of  $p(B^\pm)_\parallel/p(b)$  matches a Bowler function [30] tuned to LEP data or SLD data, where  $p(B^\pm)_\parallel$  is the component of  $p(B^\pm)$  in the  $b$  quark direction. Half the largest spread of variations to  $A_{\text{FB}}^{\text{SM}}(B^\pm)$  is 0.25%. The negligible PDF uncertainty of 0.03% is calculated by varying the twenty CTEQ6M1 eigenvectors by their uncertainties and determining the standard deviation of the variations. We find  $A_{\text{FB}}^{\text{SM}}(B^\pm) = [2.31 \pm 0.34 (\text{stat}) \pm 0.51 (\text{syst})]\%$ . Combining all data and MC uncertainties in quadrature, the MC@NLO result differs from data by  $(2.55 \pm 0.76)\%$ , or 3.3 standard deviations.

Figure 3 shows measurements of  $A_{\text{FB}}(B^\pm)$  and  $A_{\text{FB}}^{\text{SM}}(B^\pm)$  versus transverse momentum and pseudorapidity. The fully reconstructed  $J/\psi K^\pm$  final state produces good kinematic agreement between reconstructed and generated  $B^\pm$  mesons, so corrections to recover the true  $B^\pm$  kinematics are unnecessary. The average  $p_T$  of the  $B^\pm$  mesons is 12.9 GeV. We find that  $A_{\text{FB}}(B^\pm)$  is systematically lower than  $A_{\text{FB}}^{\text{SM}}(B^\pm)$  for all pseudorapidities, and for  $p_T(B) = 9 - 30$  GeV. Considering the MC systematic uncertainties to be correlated (uncorrelated), Fig. 3 (a) has  $\chi^2 = 10.3$  (11.8) for three bins and Fig. 3 (b) has  $\chi^2 = 6.6$  (7.0) for seven bins.

In conclusion, we have measured the forward-backward asymmetry in the production of  $B^\pm$  mesons with  $B^\pm \rightarrow J/\psi K^\pm$  decays in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV. For  $B^\pm$  mesons with a mean  $p_T$  of 12.9 GeV, the result is  $A_{\text{FB}}(B^\pm) = [-0.24 \pm 0.41 (\text{stat}) \pm 0.19 (\text{syst})]\%$ , which is the first measurement of this quantity. The observed discrepancy of  $\approx 3$  standard deviations between our measurement and the MC@NLO estimate suggests that more rigorous determination of the standard model prediction is needed to interpret these results.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the Department of Energy and National Science Foundation (United States of America); Alternative Energies and Atomic Energy Commission and National Center for Scientific Research/National Institute of Nuclear and Particle Physics (France); Ministry of Education and Science of the Rus-




FIG. 3: (color online) Comparison of  $A_{FB}(B^\pm)$  and  $A_{FB}^{\text{SM}}(B^\pm)$  in bins of (a)  $|\eta_B|$  and (b)  $p_T(B)$ . Data points and MC bands include statistical uncertainties convoluted with systematic uncertainties.

sian Federation, National Research Center “Kurchatov Institute” of the Russian Federation, and Russian Foundation for Basic Research (Russia); National Council for the Development of Science and Technology and Carlos Chagas Filho Foundation for the Support of Research in the State of Rio de Janeiro (Brazil); Department of Atomic Energy and Department of Science and Technology (India); Administrative Department of Science, Technology and Innovation (Colombia); National Council of Science and Technology (Mexico); National Research Foundation of Korea (Korea); Foundation for Fundamental Research on Matter (The Netherlands); Science and Technology Facilities Council and The Royal Society (United Kingdom); Ministry of Education, Youth and Sports (Czech Republic); Bundesministerium für Bildung und Forschung (Federal Ministry of Education and Research) and Deutsche Forschungsgemeinschaft (German Research Foundation) (Germany); Science Foundation Ireland (Ireland); Swedish Research Council (Sweden); China Academy of Sciences and National Natural Science Foundation of China (China); and Ministry of Education and Science of Ukraine (Ukraine).

- [1] L.G. Almeida, G. Sterman, and W. Vogelsang, Phys. Rev. D **78**, 014008 (2008); J.H. Kühn and G. Rodrigo, J. High Energy Phys. 01 (2012) 063; J.A. Aguilar-Saavedra, D. Amidei, A. Juste, and M. Perez-Victoria, arXiv:1406.1798.
- [2] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **84**, 112005 (2011).
- [3] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **83**, 112003 (2011); Phys. Rev. D **87**, 092002 (2013); Phys. Rev. D **88**, 072003 (2013).
- [4] M.I. Gresham, I.-W. Kim, and K.M. Zurek, Phys. Rev. D **83**, 114027 (2011); J.A. Aguilar-Saavedra and M. Perez-Victoria, J. High Energy Phys. 09 (2011) 097; J.F. Kamenik, J. Shu, and J. Zupan, Eur. Phys. J. C **72**, 2102 (2012).
- [5] B. Grinstein and C.W. Murphy, Phys. Rev. Lett. **111**, 062003 (2013); Phys. Rev. Lett. **112**, 239901 (2014).
- [6] S. Ipek, Phys. Rev. D **87**, 116010 (2013).
- [7] V. M Abazov *et al.* (D0 Collaboration), Phys. Rev. D **87**, 011103(R) (2013); Phys. Rev. D **88**, 112002 (2013); Phys. Rev. D **90**, 072011 (2014).
- [8] W. Bernreuther and Z.-G. Si, Phys. Rev. D **86**, 034026 (2012); M. Czakon, P. Fiedler, and A. Mitov, arXiv:1411.3007 (2014).
- [9] S. Chatrchyan *et al.* (CMS Collaboration), J. High Energy Phys. 04 (2014) 191.
- [10] G. Aad *et al.* (ATLAS Collaboration), J. High Energy Phys. 02 (2014) 107.
- [11] R. Aaij *et al.* (LHCb Collaboration), Phys. Rev. Lett. **113**, 082003 (2014).
- [12] J.H. Kühn and G. Rodrigo, Phys. Rev. D **59**, 054017 (1999).
- [13] The D0 collaboration defines a coordinate system with the  $z$  axis along the proton beam direction, the  $x$  axis pointing away from the Tevatron center, and the  $y$  axis pointing upwards. Pseudorapidity is defined as  $\eta = -\ln[\tan(\theta/2)]$ , where  $\theta$  is the laboratory frame polar angle. Angle  $\phi$  is the azimuthal angle in the  $x$ - $y$  plane.
- [14] A.V. Manohar and M. Trott, Phys. Lett. B **711**, 313 (2012).
- [15] S. Frixione and B.R. Webber, J. High Energy Phys. 06 (2002) 029; S. Frixione, P. Nason, and B.R. Webber, J. High Energy Phys. 08 (2003) 007.
- [16] J. Pumplin, D.R. Stump, J. Huston, H.-L. Lai, P. Nadolsky, and W.-K. Tung, J. High Energy Phys. 07 (2002) 012; D. Stump, J. Huston, J. Pumplin, W.-K. Tung, H.-L. Lai, S. Kuhlmann, and J. Francis Owens, J. High Energy Phys. 10 (2003) 046.
- [17] G. Corcella, J. High Energy Phys. 01 (2001) 010.
- [18] R. Brun and F. Carminati, CERN Program Library W5013, 1993. We use GEANT version 3.15.
- [19] V.M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods A **565**, 463 (2006).
- [20] V.M. Abazov *et al.*, Nucl. Instrum. Methods A **552**, 372 (2005); V.M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods A **737**, 281 (2014).
- [21] The pointing angle is defined as the angle between a particle’s momentum vector and the vector from the  $p\bar{p}$  vertex to the particle’s decay vertex, with vectors defined in the  $x$ - $y$  plane.
- [22] K.A. Olive *et al.* (Particle Data Group), Chin. Phys. C



- 38**, 090001 (2014).
- [23] A. Hoecker *et al.*, Toolkit for multivariate Data Analysis, Proc. Sci. ACAT (2007) 040. We use version 4.1.0.
- [24] T. Sjöstrand, S. Mrenna, and P. Skands, J. High Energy Phys. 05 (2006) 026.
- [25] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **110**, 241801 (2013).
- [26] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **87**, 072006 (2013).
- [27] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **86**, 072009 (2012); Phys. Rev. D **82**, 032001 (2010).
- [28] S. Frixione (private communication).
- [29] J.M. Campbell and R.K. Ellis, FERMILAB-PUB-12-078-T (2012).
- [30] G. Corcella and V. Drollinger, Nucl. Phys. B **730**, 82 (2005).