THE STEREOCHEMISTRY OF THE FORMATION OF Δ^3 -1,3,4-THIADIAZOLINE-1-OXIDES
AND EPISULFOXIDES FROM SULFINES AND 2-DIAZOPROPANE¹

L. Thijs, A. Wagenaar², Miss E.M.M. van Rens and B. Zwanenburg³.

Department of Organic Chemistry, University of Nijmegen,

Toernooiveld, Nijmegen, The Netherlands.

(Received in UK 16 July 1973; accepted for publication 31 July 1973)

Recently it was shown^{4,5} that sulfines react readily with diazoalkanes to Δ^3 -1,3,4-thiadiazoline-1-oxides in a regiospecific cyclo-addition process. In one case⁵ an aliphatic sulfine gave with diazomethane an episulfoxide instead of a five-membered ring product. Although we were inclined to believe that the cyclization to thiadiazoline-oxides would be a stereospecific process, recent results with the 1,3-dipolar cyclo-addition reaction of sulfines with diphenylnitrilimine⁶ (a regiospecific, but non-stereospecific process) threw doubt on this anticipation. Therefore, the stereochemistry of the diazoalkane-sulfine cyclization reaction requires a closer examination.

On that account we studied the reaction of 2-diazopropane with the geometrical isomers of different types of sulfines. Treatment of these sulfines (see Table) with 2-diazopropane in ether or ether/dichloromethane at -20° - -30° resulted, after addition of pentane, in the crystallization of the desired 1:1 adducts in high yields. In all cases studied each of the geometrical isomers led to a single product which was distinctly different from that obtained from the other isomer (see Table). Particularly, the NMR spectra (CDCl_3) revealed that only one adduct was obtained from each of the isomeric sulfines. From the sulfines VI, VII and VIII only the *E*-isomer could be studied, since the *Z*-isomer was not accessible by oxidation of the corresponding dithioester. Each of these sulfines gave only one cyclo-adduct in good yield.

The data presented in the Table allow the conclusion that the spatial arrangement of the S=O group and the substituents R_1 and R_2 is retained in the product. Hence, the cyclo-addition is a stereospecific process and most likely the product formation takes place in a concerted manner.

The isomeric mesityl-phenylsulfonyl-sulfines XIa and XIb reacted smoothly with 2-diazopropane in benzene/ether (1:1) at -10° . However, to our surprise an episulfoxide was isolated in 72.5% yield, instead of a five-membered ring product. From either of these isomeric sulfone sulfines the same 1:1 mixture of diastereomeric episulfoxides (m.p. $85-87^\circ$) was obtained, thus, indicating a non-stereospecific process (see Scheme). The mixture could not be separated because the com-

TABLE

	R ₁	R ₂	m.p.*	%	δCH ₃	other NMR signals
Ia(E)	phenyl	<i>o</i> -tolyl	70 ⁰	83	1.27; 2.00; 2.41	6.58-7.74(m)
Ib(Z)	<i>o</i> -tolyl	phenyl	75 ⁰	88	1.47; 1.90; 2.19	6.99-7.72(m)
IIa(E)	phenyl	α-naphthyl	85 ⁰	87	1.60; 2.41	arom. H
IIb(Z)	α-naphthyl	phenyl	89 ⁰	91	1.63; 1.93	arom. H
IIIa(E)	<i>p</i> -tolyl	<i>p</i> -chlorophenyl	80 ⁰	88	1.05; 1.94; 2.29	6.85-7.68(m)
IIIb(Z)	<i>p</i> -chloro-phenyl	<i>p</i> -tolyl	76-77 ⁰	67	1.14; 2.02; 2.40	6.83-7.63(m)
IVa(E)	phenyl	chloro	72-80 ⁰	56	1.80; 1.93	7.52
IVb(Z)	chloro	phenyl	84 ⁰	81	1.19; 1.92	7.47
Va(E)	phenyl	phenylthio	65-67 ⁰	82	1.62; 1.80	6.93-7.62(m)
Vb(Z)	phenylthio	phenyl	75-77 ⁰	68	1.09; 1.89	6.85-7.67(m)
VIa(E)	anisyl	<i>p</i> -tolylthio	80 ⁰	83	1.54; 1.78; 2.30; 3.78	6.90+7.46(AB, J 9Hz) 7.02+7.28(AB, J 7Hz)
VIIa(E)	phenyl	phenylsulfonyl	97 ⁰	75	1.72; 2.03	7.20-7.67(m)
VIIIa(E)	anisyl	<i>p</i> -tolylsulfonyl	dec.	75	1.66; 2.00; 2.36; 3.78	6.83+7.44(AB, J 9Hz) 7.14+7.39(AB, J 9Hz)
IX	phenylthio	phenylthio	55 ⁰	58	1.37; 1.54 (-30 ⁰)	6.74-7.81(m)
X	chloro	chloro	70 ⁰	44	1.66; 1.85 (CCl ₄)	

* All compounds show vigorous decomposition during melting.

(Characteristic i.r. absorptions for these compounds were observed at 1060-1080 (ν_{S=O}) and 1560-1575 cm⁻¹ (ν_{N=N}))

pound could not withstand extensive chromatography.

The episulfoxide structure was assigned on the following grounds: a correct elemental analysis for C₁₉H₂₂O₃S₂, i.r. absorptions (in CS₂) at 1050 cm⁻¹ (ν_{S=O}), 1150 and 1325 cm⁻¹ (ν_{SO₂}) and signals in the NMR spectrum (CDCl₃) at δ 1.01, 1.40, 1.70 and 1.78 ppm for the methyl protons at C-2 (note the distinct different position of the methyl protons at C-2 in the thiadiazoline-oxide derived from VIIa), at δ 2.21 and 2.47 ppm for the methyls at C-2', at δ 2.16 and 2.34

ppm for those at C-4', at δ 6.62 and 6.96 ppm for the protons at C-3' and at δ 7.18-7.80 ppm for the phenyl protons. Furthermore, oxidation of the product with *m*-chloroperbenzoic acid in ether at 20° gave 1-mesityl-2-methyl-1-phenylsulfonyl-1-propene (m.p. 120-122°) in 46% yield (oxidation to episulfone with subsequent extrusion of SO₂).

SCHEME

Bonini and Maccagnani⁷ found that aromatic sulfines such as diphenylsulfine and thiofluorenone-S-oxide react with phenyldiazomethane to give a triaryl substituted episulfoxide as a mixture of diastereomers (Z/E ratio ranging from 1:4 to 2:3 for the different aryl substituents). Thus, again a non-stereospecific formation of the three-membered ring.

To explain this remarkable difference in stereochemistry in the formation of thiadiazoline-oxides and episulfoxides, we suggest that the episulfoxide does not come about *via* an initially formed thiadiazoline-oxide, but most likely *via* a two step process in which firstly a nucleophilic attack of the diazocarbon at the sulfine sulfur provides a zwitter ionic diazonium compound (see Scheme). Subsequently, an internal 1,3-displacement of nitrogen produces the episulfoxide. Inspection of molecular models clearly reveals that steric crowding prevents the formation of a five-membered ring adduct and favors the less congested three-membered ring.

The mechanism in the Scheme is supported by the fact that we never found any indication of an episulfoxide formation from the thiadiazoline-oxides. However, these five-membered ring adducts are thermally rather unstable. Usually a retro-cyclo-addition reaction to starting materials as observed for the adducts derived from Va, VIa and X takes place. In some cases a reverse retro-cyclo-addition reaction is observed as nicely exemplified by the adduct from IX. Warming this adduct in chloroform at 40° or at 25° in benzene/pentane, containing some silicagel, gave besides 60% of the sulfine IX a 30% yield of tetrakis(phenylthio)-ethene arising from bis(phenylthio)diazomethane *via* dimerization of bis(phenylthio)carbene⁸.

With other sulfines having a bulky substituent attached to the sulfine function a deviating reaction pattern was observed. *Z*-mesityl-phenylsulfine did not react at all with 2-diazopropane, whereas the *E*-isomer was isomerized quantitatively to the *Z*-form. Similarly, *E*-mesityl-phenylthio-sulfine isomerized to the *Z*-isomer, while the *Z*-form did not react. This isomerization can be rationalized by assuming the formation of a zwitter ionic intermediate (see Scheme) which then splits off 2-diazopropane to give the thermodynamically⁹ more stable sulfine isomer instead of forming the three-membered ring.

We conclude that the normal reaction of sulfines with diazoalkanes will be the concerted cyclo-addition to Δ^3 -1,3,4-thiadiazoline-1-oxides. Introduction of bulky substituents in either of the reactants will sterically hamper this cyclization to five-membered rings and give rise to alternative reaction routes of which the non-stereospecific formation of episulfoxides is the most interesting one.

References and notes

1. Part XXII in the series "Chemistry of Sulfines", part XXI, Tetrahedron Lett., submitted for publication.
2. Department of Organic Chemistry, University of Groningen, The Netherlands.
3. To whom correspondence should be addressed.
4. B.F. Bonini, G. Maccagnani, A. Wagenaar, L. Thijs and B. Zwanenburg, J.C.S. Perkin I, 1972, 2490.
5. B. Zwanenburg, A. Wagenaar, L. Thijs and J. Strating, J.C.S. Perkin I, 1973, 73.
6. Part XXI in this series, see ref. 1.
7. B.F. Bonini and G. Maccagnani, Tetrahedron Lett., accompanying paper.
8. U. Schöllkopf and E. Wiskott, Angew. Chem. 75, 725 (1963).
9. The *Z*-isomers were formed when the *E*-isomers were allowed to stand in the refrigerator for several months.