A New Hydridocarbonyl Complex of Osmium(II)

By F. G. MOERS

(Inorganic Chemistry Laboratory, University of Nijmegen, Nijmegen, The Netherlands)

Reprinted from

Chemical Communications 1971

The Chemical Society, Burlington House, London W1V 0BN

A New Hydridocarbonyl Complex of Osmium(II)

By F. G. MOERS

(Inorganic Chemistry Laboratory, University of Nijmegen, Nijmegen, The Netherlands)

Summary The reaction of K_2OsCl_6 with tricyclohexylphosphine in alcohols leads to the hydridocarbonyl complex $OsHCl(CO)(PCy_3)_2$.

THE hydridocarbonylphosphine complexes with the formula $MHX(CO)(PPh_3)_3$ where $M = Ru$ or Os and $X = Cl$ or Br may be obtained directly from the metal halides, triphenylphosphine, and alcohols.¹ Stephenson and Wilkinson² describe the preparation of the complex $RuCl_2(PPh_3)_3$ by refluxing a methanolic solution of PPh_3 and $RuCl_3$ under nitrogen.

I now report the isolation of the complex $OsHCl(CO)(PCy_3)_2$ (PCy_3 = tricyclohexylphosphine). On boiling a solution of PCy_3 and K_2OsCl_6 in 2-methoxyethanol in an atmosphere of nitrogen for 48 h, a red complex, $OsHCl(CO)(PCy_3)_2$, was obtained. The complex had satisfactory analytical data.

The complex decomposes at 210° and is slightly soluble in organic solvents. It is diamagnetic (measured between room temperature and -150°) indicating a low spin d^6 -configuration.

The i.r. spectrum (KBr) in the range 2100 – 1850 cm^{-1} shows two bands attributable to $\nu(CO)$ stretching modes at $1932s$ and $1887m\text{ cm}^{-1}$ and one band at 2012 cm^{-1} assigned to $\nu(M-H)$. On performing the reaction in deuteriated ethanol, the expected frequency shift for a covalent osmium-hydrogen band was found ($\nu(M-D)$ at $1508w\text{ cm}^{-1}$), the other two bands being unchanged. The split nature of the carbonyl bands, which is maintained in benzene solution, suggests that the complex is a mixture of geometrical isomers. So far, attempts to separate isomers have not succeeded. On shaking $OsHCl(CO)(PCy_3)_2$ in an excess of

pyridine a yellow complex, $OsHCl(CO)(PCy_3)_2Py$, was obtained. The pyridine complex shows two i.r. absorption bands (KBr) at $2119m$ and $1870s\text{ cm}^{-1}$ attributable to $\nu(CO)$ and one weak band at 2040 cm^{-1} assigned to $\nu(M-H)$.

In the mass spectrum there is a cluster, with the lowest fragment ion having m/e 808 which is in agreement with the formula $^{184}OsH^{36}Cl(CO)(PCy_3)_2$. The relative abundances of the peaks in this cluster agree with the calculated ratios. Fragment ions with higher masses than $OsHCl(CO)(PCy_3)_2$ could not be detected. Comparison of the mass spectra of $OsHCl(CO)(PCy_3)_2$ and its deuteriated derivative, prepared as described above, showed appreciable deuteration of PCy_3 indicated by the presence of peaks at m/e 280–290. As a result we found that the ratios of the cluster of peaks due to $OsDCl(CO)(PCy_3)_2$ differed from the calculated $OsDCl(CO)(PCy_3)_2$ ratios in which the PCy_3 is non-deuteriated. These data suggest an exchange reaction between the complex and deuterium during the preparation of the complex.

All attempts to prepare $OsHCl(CO)(PCy_3)_3$ by addition of an excess of PCy_3 were unsuccessful. The fact that this complex could not be isolated may be due to the steric hindrance of the cyclohexyl groups.³

The structure of this complex is probably related to that of the complexes $RuHX(PPh_3)_3$,⁴ the diamagnetic complex $RuX_2(SbPh_3)_3$,² and $RuCl_2(PPh_3)_3$. An X-ray study by La Placa and Ibers⁵ has shown a quasi-octahedral structure for $RuCl_2(PPh_3)_3$.

I thank Prof. Dr. Ir. J. J. Steggerda for discussions, Mr. J. P. Langhout for helpful assistance, and Dr. F. Gerhartl for recording the mass spectra.

(Received, November 18th, 1970; Com. 1994.)

¹ L. Vaska and J. W. Diluzio, *J. Amer. Chem. Soc.*, 1961, **83**, 1262; J. Chatt and B. L. Shaw, *Chem. and Ind.*, 1960, 931.

² T. A. Stephenson and G. Wilkinson, *J. Inorg. Nuclear Chem.*, 1966, **28**, 945.

³ T. S. Green and K. Saito, *Chem. Comm.*, 1969, 208; F. G. Moers and P. H. op het Veld, *J. Inorg. Nuclear Chem.*, 1970, **32**, 3225.

⁴ T. Ito, S. Kitazume, A. Yamamoto, and S. Ikeda, *J. Amer. Chem. Soc.*, 1970, **92**, 3011.

⁵ S. J. La Placa and I. A. Ibers, *Inorg. Chem.*, 1965, **4**, 778.