

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/102799>

Please be advised that this information was generated on 2019-02-19 and may be subject to change.



Alcohol and tobacco use and heart rate reactivity to a psychosocial stressor in an adolescent population

Brittany E. Evans^{a,b,*}, Kirstin Greaves-Lord^{a,1}, Anja S. Euser^{a,c,2}, Joke H.M. Tulen^d,
Ingmar H.A. Franken^{c,2}, Anja C. Huizink^{b,e,3}

^a Department of Child and Adolescent Psychiatry/Psychology, Erasmus University Medical Center/Sophia Children's Hospital, Rotterdam, The Netherlands

^b Department of Developmental Psychology, VU University, Amsterdam, The Netherlands

^c Department of Psychology, Erasmus University Rotterdam, Rotterdam, The Netherlands

^d Department of Psychiatry, Erasmus University Medical Center, Rotterdam, The Netherlands

^e Behavioral Science Institute, Radboud University, Nijmegen, The Netherlands

ARTICLE INFO

Article history:

Received 24 January 2012

Received in revised form 17 May 2012

Accepted 24 May 2012

Available online 21 June 2012

Keywords:

Adolescence
Alcohol use
Autonomic nervous system
Heart rate
Stress reactivity
Tobacco use

ABSTRACT

Background: Few studies have investigated physiological stress (re)activity in relation to substance use, especially in adolescents. Using substances is one way to stimulate physiological arousal, therefore inherent hypo-arousal may be associated with substance use in adolescents. The purpose of this study was to examine the relation of autonomic nervous system (ANS) activity with alcohol and tobacco use in adolescents.

Methods: ANS activity and perceived stress during a social stress procedure were examined in relation to substance use. 275 Dutch adolescents from a general population study provided complete data. Heart rate was recorded continuously during a pre-task rest period, a stressful task period and a post-task recovery period. Alcohol and tobacco use were self-reported.

Results: Adolescents who consumed a medium and high number of alcoholic drinks per week (more than two) exhibited lower heart rates during the entire stress procedure as compared to those who consumed a low number of alcoholic drinks. Adolescents who smoked every day portrayed blunted heart rate reactivity to stress as compared to adolescents who smoked less frequently or not at all. Perceived stress was not related to alcohol or tobacco use.

Conclusions: Overall lower heart rate in adolescents who drank more and blunted heart rate reactivity to stress in those who used tobacco every day may indicate inherent hypo-arousal of the ANS system in those vulnerable to use substances more often. These adolescents may actively seek out substances in order to achieve a more normalized state of arousal.

© 2012 Elsevier Ireland Ltd. Open access under the [Elsevier OA license](http://www.elsevier.com/locate/elsevier).

1. Introduction

Adolescence is a critical developmental period regarding exploration behavior toward substances, with adolescents showing increased experimentation with alcohol and tobacco (Hardin and Ernst, 2009). Adolescents have generally not used substances in large amounts or for long periods of time, which facilitates the discernment of vulnerability markers for substance use problems. As

it is less likely that the use of substances has caused lasting physical changes in this population, it is possible that differences found in those adolescents who are prone to use more alcohol and/or tobacco may be due to underlying, inherent factors.

Stress reactivity may be one potential vulnerability marker for the development of substance use disorders (SUDs). It has been related to SUDs in adults (for reviews see Goeders, 2003; Sinha, 2008). One view of this association describes the tendency of individuals to use substances in order to alleviate symptoms of stress, or hyper-arousal; self-medication hypothesis (Khantzian, 1985). A second hypothesis draws on the observation that individuals with high sensation seeking tendencies are more likely to engage in substance use (Creemers et al., 2009; Martin et al., 2002; Zuckerman and Kuhlman, 2000). These individuals may be inherently hypo-aroused and deliberately seek out substances in order to achieve a state of normalized arousal and thereby physiological comfort (Goeders, 2003; Majewska, 2002); stimulus-seeking

* Corresponding author at: Department of Developmental Psychology, VU University, Van der Boerhorststraat 1, Room 3B-65, 1081 BT Amsterdam. Tel.: +31 205988736.

E-mail address: b.evans@erasmusmc.nl (B.E. Evans).

¹ Wyttemaweg 8, Room KP-2861, 3015 CN Rotterdam, The Netherlands.

² Burgemeester Oudlaan 50, 3062 PA Rotterdam, The Netherlands.

³ Van der Boerhorststraat 1, 1081 BT Amsterdam, The Netherlands.

hypothesis (Huizink et al., 2006; Zuckerman and Neeb, 1979). This hypo-arousal may be associated with multifarious stimulus-seeking behaviors, and has frequently been linked to externalizing behaviors (Raine, 2002), of which substance use may be one (Liu et al., 2009).

Stress reactivity can be assessed by measuring the activation of the autonomic nervous system (ANS), which is responsible for the body's immediate response to stress and plays an important role in allostasis. In response to a stressor, the ANS prepares the body for action by increasing heart rate (HR), blood pressure and respiration. HR is a valid physiological index of stress (Porges, 1995) and in the field of SUD research, assessing HR reactivity to a psychosocial stressor is ideal due to its ecological validity.

In alcohol dependent patients, resting HR may be higher compared to social drinkers (Sinha et al., 2009) and controls (e.g., Ingjaldsson et al., 2003). In response to psychological stress, though, alcohol dependent patients could have lower HR reactivity (Panknin et al., 2002). Pertaining to tobacco use, the acute effect of smoking entails an increase in HR (Hasenfratz and Battig, 1992; James and Richardson, 1991; Pauli et al., 1993), though differences between habitual smokers and non-smokers in HR response to stress is unclear. Some studies reported no differences between smokers and non-smokers in resting HR (Kirschbaum et al., 1993; Perkins et al., 1992; Roy et al., 1994) or in response to psychosocial stress (Back et al., 2008; Childs and de Wit, 2009; Hughes and Higgins, 2010; Kirschbaum et al., 1993; Perkins et al., 1992; Tersman et al., 1991). Others reported increased resting HR in smokers (al'Absi et al., 2003; Phillips et al., 2009; Sheffield et al., 1997; Tsuda et al., 1996) and attenuated HR responses to psychological stressors in large community samples of men (Sheffield et al., 1997), in women (Girdler et al., 1997; Straneva et al., 2000) and with light as well as heavy smokers showing attenuated HR reactivity in comparison to non-smokers (Phillips et al., 2009; Roy et al., 1994).

As the above mentioned studies were performed in subjects who had already used substances heavily or were dependent on a substance, it is unclear whether the results point to underlying variation in the ANS, or whether substance use had affected this system directly. In order to minimize the possibility of heavy substance use dysregulating the ANS, it is important to perform studies in the general population, with subjects who have used substances relatively less. Studies that examined HR in response to stress in individuals with a family history (FH) of alcoholism, who had not yet developed problem drinking, led to differing results. One found that adults with a multigenerational FH of alcoholism, as compared to those with a unigenerational and negative FH, showed increased HR to unavoidable shock (Finn et al., 1992; Finn and Pihl, 1987). One study in boys found increased HR reactivity to a mental arithmetic task (Harden and Pihl, 1995). Contrastingly, another found blunted HR responses to psychosocial stress in adults with a FH of alcoholism as compared to those with a negative FH (Sorocco et al., 2006). Clearly, more research is needed to elucidate this relationship.

Furthermore, as almost all of the above-mentioned studies were performed in adults, little is known about the relation between HR and substance use in adolescents. Substance use can be viewed as a manifestation of externalizing problems (e.g., symptoms of oppositional defiant and conduct disorder; Krueger et al., 2002; Liu et al., 2009). The relation between externalizing problems and HR has been well established; low resting HR is the best-replicated correlate of antisocial behavior in children and adolescents, and attenuated HR in response to a stressor is also well-confirmed (Ortiz and Raine, 2004). Thus, literature on the relation between externalizing problems and HR may provide insight into the relation that could be found between substance use during adolescence and HR reactivity, i.e., low resting HR and attenuated HR response to stress. Of interest here is whether HR is related to externalizing




Fig. 1. Flow chart of available data. HR = heart rate.

problems in general, or whether it is related specifically to substance use.

The goal of this study was to examine the relation between adolescent alcohol and tobacco use and HR(re)activity during a psychosocial stressor. We expected to find that adolescents who drank more alcohol and adolescents who used more tobacco would portray low resting HR and an attenuated HR response to the stressor. By entering number of externalizing problems into the model, we aimed to examine whether any found relation is specific for alcohol and tobacco use. Physiological responses are generally postulated to reflect subjective, or perceived stress (PS), responses (Thayer, 1970), however, convincing experimental evidence of this is limited (Oldehinkel et al., 2011). Therefore, a second aim of this study was to examine whether HR and PS were related, and whether alcohol and tobacco use were related to PS. Based on findings in earlier studies in which HR did but PS responses did not vary by risk group (e.g., Fairchild et al., 2008; Finn and Pihl, 1987), we hypothesized that HR would be related to PS, but that PS would not be related to alcohol and tobacco use in adolescents.

2. Method

2.1. Participants

The current sample of 275 14–20-year-old ($M=17.22$; $SD=1.31$) adolescents is part of a larger sample that participated in the South Holland 2 study, a large Dutch general population study of youth aged 6–20 years. For this larger study, children and adolescents were randomly drawn from registers of 35 representative municipalities in the Dutch province of South Holland including both urban and rural areas. At the second assessment wave, 536 individuals were eligible for the present study on the adolescent group, being between the ages of 14 and 20. Of these, 330 individuals took part in a stress procedure between January 2006 and March 2009. Of this sub-sample, complete HR data was available for 288 individuals (13% missing data, which is not irregular for research using HR data, see e.g., Dietrich et al., 2007). This latter group did not differ from the 330 who participated in the stress procedure according to gender, SES or internalizing and externalizing symptoms, although participants with usable HR data were younger ($p<.01$). Complete HR as well as substance use data for the entire stress procedure was available for 275 adolescents. The latter group did not differ from the sample of 536 eligible individuals in terms of age, SES or internalizing symptoms, though female gender did significantly predict being included in the analysis ($p<.01$), and those included in the analysis reported fewer externalizing symptoms ($p<.05$). See Fig. 1 for a flow chart of available data.

2.2. Procedure

Stress procedure sessions began at approximately 12 pm or 3 pm and commenced with an explanation of the procedure by the experiment leader. After the completion of two questionnaires, the electrodes of the electrocardiogram were attached and participants were told to breathe normally and to relax. After a 10 min




Fig. 2. Stress procedure during which heart rate was measured continuously. PS = perceived stress; MAT = mental arithmetic task; PST = public speaking task; CT = computer task.

rest period, the psychosocial stress tasks began, entailing mental arithmetic, public speaking and computer mathematics tasks (see Dieleman et al., 2010 for full details on the procedure). The session ended with a 5 min recovery period and a relaxing nature documentary (25 min). Fig. 2 depicts the procedure schematically. Written informed consent was obtained from participating adolescents and their parents, and adolescents received a gift certificate. The study was approved by the Ethics Committee of the Erasmus University Medical Center.

2.3. Measures

2.3.1. Alcohol and tobacco use. Self-reported alcohol use consisted of a composite of questions pertaining to the number of days per week on which alcohol was usually drunk multiplied by the number of alcoholic drinks that was usually consumed per occasion. This led to a continuous variable denoting the average number of alcoholic drinks consumed per week. Subjects were divided into three groups according to this (based on third percentiles; Hillers and Massey, 1985; Murray et al., 2002) which led to the variable group of Number of Drinks per Week (gNDW). Those who drank two alcoholic drinks per week or less ($N=93$) were considered Low Quantity (per week) Drinkers; between three and six ($N=88$) Medium Quantity Drinkers; and seven or more ($N=69$) High Quantity Drinkers. Table 1 describes additional alcohol use history variables.

Frequency of Tobacco Use was based on one multiple choice self-report question (*Have you ever smoked cigarettes?*). Three groups were formed based on the answer categories (group of Frequency of Tobacco Use; gFTU): Non-smokers ($N=124$) had never smoked, Low Frequency Smokers ($N=107$) smoke once in a while, but not every day or have smoked 1 or 2 cigarettes ever and High Frequency Smokers ($N=38$) smoke every day. Participants who used to smoke, but have quit were excluded from the analysis ($N=6$).

2.3.2. Heart rate. Heart rate was measured using a three-lead electrocardiogram (ECG) and was monitored constantly throughout the entire stress procedure. The ECG was sampled at 512 Hz and stored on a flashcard by means of a portable digital recorder (Vitaport™ System; TEMEC Instruments B.V., Kerkrade, The Netherlands). After completion of the recording, all physiological data were imported and processed on a Personal Computer using a Vitascore™ software module (TEMEC Instruments BV, Kerkrade, The Netherlands). A customized software program calculated the interbeat intervals (IBI) of the ECG using R-top detection, resulting in IBI time series. This time series was inspected for detection and removal of artifacts. HR time series were calculated from these IBI time series and expressed in beats per

minute (bpm); the HR time series were subsequently averaged per period during the stress procedure. For purposes of the analyses, the stress procedure was consolidated into three periods: a pre-task rest period (Rest), the period during any of the three stress tasks that elicited the maximum HR response (Task), and a post-task recovery period (Recovery). As expected, the maximum HR response occurred for most participants during either the mental arithmetic task (33.8%) or the speech part of the public speaking task (49.8%).

2.3.3. Perceived stress. Self-reported perceived stress (Dieleman et al., 2010) was assessed after the rest period, each of the tasks and at the end of the procedure. Participants answered seven questions (e.g., 'Can you feel your heart beating?', 'Are you nervous?') using a visual thermometer ranging from 0 (not at all) to 8 (very much). The scores were summed to a total score of PS for each period/task, Task PS entailed the maximum PS score during any of the three stress tasks.

2.3.4. Potential covariates. In previous studies examining heart rate reactivity, age (Phillips et al., 2009), gender (Back et al., 2008), pubertal stage (Carroll et al., 2008), body mass index (BMI; Carroll et al., 2008), oral contraceptive (OC) use (Girdler et al., 1997), socioeconomic status (SES; Miller et al., 2009), internalizing and externalizing problems (Greaves-Lord et al., 2007; Ortiz and Raine, 2004), parental substance use (Finn et al., 1992) and time of test session (Sheffield et al., 1997) have been taken into account. We assessed pubertal stage using self-reported Tanner stages (Marshall and Tanner, 1970). SES was based on the higher occupational level of either parent (Statistics, 2010) and coded into low ($x=1$), average ($x=2$) and high ($x=3$) SES. Internalizing and externalizing problems were evaluated using the Youth Self-Report (YSR; Achenbach and Rescorla, 2001). Scores on subscales affective, anxiety and somatic disorders were summed, leading to number of internalizing problems. Number of externalizing problems was similarly achieved (using subscales attention deficit hyperactive, conduct and oppositional defiant disorders subscales). Parent substance use was self-reported usually by the mother and entails the average number of alcoholic drinks consumed per week. Age, gender (boy: $x=1$; girl: $x=2$) and OC use (no: $x=0$; yes: $x=1$) were assessed using a demographics self-report questionnaire. Height and weight were measured prior to the test session to calculate BMI. Time of test session was coded noon ($x=1$) or late afternoon ($x=2$).

2.4. Statistical analysis

First, a manipulation check was performed by way of repeated measures analysis of variance (RM-ANOVA) in the entire sample in order to confirm that the stressful

Table 1
History of alcohol use descriptive variables.

Frequency of alcohol use (times used)	N	Frequency of occurrence (%)				
		Never	1–5	5–10	10–20	>20
How often have you used alcohol						
In your lifetime?	268	8.2	10.8	9.3	10.8	60.8
In the past 12 months?	268	13.1	14.2	19.0	14.2	39.6
In the past 4 weeks?	268	23.1	43.7	21.6	6.7	4.9
How often have you been drunk or tipsy						
In your lifetime?	271	31.7	40.6	9.2	7.0	11.4
In the past 12 months?	267	39.7	42.7	7.5	4.1	6.0
In the past 4 weeks?	266	66.2	30.1	3.4	0.4	0.0
Begin alcohol use	N	Age (years)				
		Never	<9	10–12	13–15	>16
How old were you the first time you						
Drank at least one glass of alcohol?	234		1.3	18.4	69.7	10.7
Were drunk?	266	44.7	0.0	1.5	35.0	18.8

<9 = 9 years or younger; >16 = 16 years or older.

Table 2
Descriptive statistics for independent and dependent variables and covariates.

Variable	N	Mean (SD)	Range	Frequency (%)
Age	275	17.22 (1.31)	14.52–20.84	
Gender	275			
Boys				42.5
Girls				57.5
Body mass index	273	21.99 (3.30)	16.00–6.75	
Pubertal stage	254	4.31 (0.66)	2.50–.00	
Socioeconomic status	274			
Low				4.4
Average				50.0
High				45.6
Internalizing symptoms	243	0.93 (0.67)	0.00–3.72	
Externalizing symptoms	264	1.45 (0.70)	0.12–3.63	
Parental alcohol use	247	5.22 (6.71)	0.00–42.00	
Time of test session	274			
Noon				59.9
Late afternoon				40.1
Oral contraceptive use (girls only)	155			
Use				54.8
No use				45.2
gNDW	275			
Low Quantity Drinkers				37.5
Med. Quantity Drinkers				34.5
High Quantity Drinkers				28.0
gFTU	269			
Non-smokers				46.1
Low freq. smokers				39.8
High freq. smokers				14.1
Heart rate (bpm) during	275			
Rest		75.87 (9.99)	55.62–122.43	
Task		86.33 (12.34)	60.04–141.16	
Recovery		71.92 (9.06)	54.03–102.81	
Perceived stress during	275			
Rest		7.88 (5.95)	0–32	
Task		14.92 (8.93)	0–42	
Recovery		3.89 (4.98)	0–34	
Drinks per week	275			
Low Quantity Drinkers		5.36 (5.80)	0–39	
Med. Quantity Drinkers		0.68 (0.82)	0–2	
High Quantity Drinkers		4.56 (1.19)	3–6	
		12.59 (5.94)	7–39	
Number of cigarettes per day (only High Frequency Smokers)	38	10.97 (6.08)	2–30	

gNDW = group of Number of Drinks per Week; gFTU = group of Frequency of Tobacco Use; med. = medium; freq. = frequency; bpm = beat per minute.

tasks did induce an increase in HR and PS as compared to the Rest period. Age and gender were entered into all models as covariates. Prior to the main analysis additional covariates were examined and added to the main analysis if they correlated significantly with both independent and dependent variables. To investigate HR during the stress procedure, a $3 \times 3 \times 3$ RM-ANOVA was performed with period (Rest, Task, Recovery) as the within-subjects factor and gNDW (Low, Medium, High Quantity) and gFTU (Non-smokers, Low, High Frequency Smokers) as between-subjects factors. Interactions between period and the between-subjects variables as well as between period and the covariates were examined. Simple contrasts were performed in order to explore between-group differences. Univariate ANOVAs of the change score in HR between Rest and Task periods were performed when interaction effects were present. An identical analysis was performed with PS as the dependent variable measured across the three periods. In all analyses, Greenhouse–Geisser statistics are reported when necessary to correct departures from sphericity.

3. Results

3.1. Manipulation check

For the entire sample, the tasks produced physiological stress, as indicated by a significant within-groups effect of period

($F(1.32,357.37) = 589.66, p < .001$). Simple contrasts showed that average HR during the Task was significantly higher than during Rest ($F(1,271) = 412.99, p < .001$). PS also differed across the three periods ($F(1.51,414.42) = 403.01, p < .001$), with simple contrasts again showing PS to be higher during the Task as compared to Rest ($F(1,274) = 293.39, p < .001$).

3.2. Preliminary analyses

Descriptives of, and correlations between, dependent, independent and potential covariates are shown in Tables 2 and 3. Age and gender were controlled in all models. No other variables correlated with both dependent and independent variables, and therefore were not included as covariates in the models. PS and HR showed a small, significant positive correlation, specifically during Task ($R = .13, p < .05$) and Recovery ($R = .12, p < .05$), additionally PS Rest with HR Task ($R = .13, p < .05$) and HR Recovery ($R = .16, p < .01$). The HR response measure was not significantly correlated with the PS response measure.

Table 3
Correlations between independent and dependent variables and covariates.

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1. HR Rest	1																	
2. HR Task	.68***	1																
3. HR Rec	.92***	.76***	1															
4. gNDW	-.25***	-.23***	-.29***	1														
5. gFTU	-.08	-.21**	-.16*	.38***	1													
6. PS Rest	.10	.13*	.16**	-.04	-.08	1												
7. PS Task	.04	.13*	.09	-.02	-.10	.65***	1											
8. PS Rec	.07	.10	.12*	-.03	-.03	.68***	.49***	1										
9. Age	.07	.03	.04	.22***	.16*	-.10	-.11	-.00	1									
10. Gender	.06	.15*	.03	-.14*	.13*	-.13*	-.04	-.07	.04	1								
11. BMI	-.05	-.11	-.02	.05	.17**	-.05	-.06	.02	.07	.15*	1							
12. Puberty	.20**	.08	.15*	.07	.09	.02	-.02	.00	.32***	-.04	.22***	1						
13. SES	-.10	.07	-.09	-.03	-.06	.02	.01	-.02	-.04	.11	-.13*	-.05	1					
14. Intern	.09	.08	.13*	-.03	.12	-.01	.16*	.03	.03	.39***	.04	.03	-.08	1				
15. Extern	-.13*	-.14*	-.15*	.27***	.26***	-.02	.01	-.01	-.03	-.04	.01	.04	-.17**	.35***	1			
16. PAU	.05	.04	-.02	.12	.07	.07	.05	.03	.05	.02	-.08	-.03	.22***	-.01	-.08	1		
17. TTS	-.02	.06	.07	-.10	.01	-.12	-.08	-.04	-.03	.06	-.10	-.00	.05	.05	-.03	-.06	1	
18. OC use	.10	-.03	.10	.23**	.36***	-.06	-.06	-.15	.29***	Na	.01	.29***	-.07	.08	-.01	.03	.00	1

HR=heart rate; Rec=recovery; gNDW=group of Number of Drinks per Week; gFTU=group of Frequency of Tobacco Use; PS=perceived stress; BMI=body mass index; Puberty=pubertal stage; SES=socioeconomic status; Intern=internalizing; Extern=externalizing; PAU=parent alcohol use; TTS=time of test session; OC=oral contraceptive; Na=not applicable, as OC use was only present in girls.
***p<.001; **p<.01; *p<.05.

3.3. Alcohol and tobacco use: heart rate

A significant between-subjects effect was evident for gNDW ($F(2,244)=6.12, p<.01$). Simple contrasts showed that Low Quantity Drinkers had a significantly higher HR during the entire procedure compared to Medium Quantity (adjusted $p<.001$) and High Quantity Drinkers (adjusted $p<.01$). Medium Quantity Drinkers did not differ significantly from High Quantity Drinkers, as shown in Table 4. No interaction between period and gNDW was evident.

There was no main effect of period, nor a between-subjects effect for gFTU in the RM-ANOVA. However, an interaction effect of gFTU with period was evident ($F(2.62,319.57)=5.54, p<.01$). A three-way interaction effect between gNDW, gFTU and HR was not. To investigate the interaction between gFTU and period, a univariate analysis of change scores in HR from Rest to Task was performed and revealed a significant between-subjects effect ($F(2,258)=10.42, p<.001$), with simple contrasts showing that High

Frequency Smokers portrayed blunted HR reactivity to the tasks as compared to Low Frequency Smokers (adjusted $p<.001$) and Non-smokers (adjusted $p<.001$). Low Frequency Smokers did not differ significantly from Non-smokers. Results are depicted in Figs. 3 and 4 (gNDW) and (gFTU).

3.4. Alcohol and tobacco use: perceived stress

Predicting PS, a within-subjects effect of period over time was evident ($F(1.51,389.12)=7.66, p<.01$). No between-subjects effects of gNDW or gFTU or interactions effects were observed.

Table 4
Values for repeated measures ANOVA model.

Variable	F	p	η^2
Period	2.59	.10	.01
Between-subjects effects			
gNDW	6.12	<.01	.05
gFTU	0.61	.55	.01
Age	3.24	.07	.01
Gender	1.28	.26	.01
Interaction effects			
Period × gNDW	0.15	.96	.00
Period × gFTU	5.54	<.01	.04
Period × age	0.16	.75	.00
Period × gender	4.57	<.05	.02
Period × externalizing	0.11	.81	.00
Period × gNDW × gFTU	0.69	.64	.01

Variable	Contrast estimate	p
Contrast analysis gNDW		
Low vs. Med. Quantity Drinkers	6.08	<.001
Low vs. High Quantity Drinkers	5.84	<.01
Med. vs. High Quantity Drinkers	-.023	.89

All values are from the repeated measures ANOVA with number of externalizing symptoms included; N=256; gNDW=group of Number of Drinks per Week; gFTU=group of Frequency of Tobacco Use; Med.=medium.




Fig. 3. Heart rate during the stress procedure for each group of Low, Medium and High Quantity Drinkers, corrected for age, gender and externalizing problems. Group of Number of Drinks per Week, --- Low Quantity Drinkers, — Medium Quantity Drinkers, and - - High Quantity Drinkers. Note: Covariates appearing in the model are evaluated at the following values: age = 17.21, gender = 0.5, and number of externalizing problems = -.01.




Fig. 4. Heart rate during the stress procedure for each group of Non-smokers, Low and High Frequency Smokers, corrected for age, gender and externalizing problems. Group of Frequency of Tobacco use, --- Non-smokers, — Low Frequency Smokers, -- High Frequency smokers. Note: Covariates appearing in the model are evaluated at the following values: age = 17.21, gender = 0.59, and number of externalizing problems = -.01.

3.5. Covariates

In order to examine whether the found effects were specific to alcohol and tobacco use alone, number of externalizing problems was added to the model predicting HR. The interaction between number of externalizing problems and HR was not significant, and the results of the model did not change.

In the model predicting HR, there was no main effect of gender, though an interaction effect of period and gender was found ($F(1.31, 319.57) = 4.57, p < .05$). A univariate ANOVA analysis showed this interaction to be due to girls reacting more strongly to the stress procedure; the change in HR from Rest to Task was greater for girls than for boys ($F(1.273) = 4.06, p < .05$). No main or interaction effect of gender was significant in the model predicting PS. When both RM-ANOVAs were run again in female subjects only, controlling for OC use, there were no main or interaction effects of OC use.

4. Discussion

In a sample of 14–20-year old adolescents, we examined whether alcohol and tobacco use were related to heart rate (HR) during a psychosocial stress procedure. To our knowledge, this is the first study to examine autonomic nervous system (ANS) (re)activity in relation to substance use in adolescents from the general population. We found that those who drank a medium and high number of alcoholic drinks per week (more than 2) portrayed a lower HR during the entire stress procedure as compared to those who drank fewer alcoholic drinks per week. Also, those who used tobacco every day showed blunted HR reactivity to the stressful tasks as compared to those who smoked less frequently or not at all. Thus, two of our hypotheses were confirmed (i.e., adolescents who drank more alcohol portrayed lower resting HR, and those who used more tobacco portrayed blunted HR response to stress) and two were not (i.e., we expected to find that those who drank more alcohol would portray blunted HR response to stress, and those who used more tobacco would portray lower resting HR). Furthermore,

we examined whether HR responses were related to PS responses, and whether the groups of alcohol and tobacco users differed with respect to PS responses. As expected, HR and PS responses were positively and significantly correlated, and PS responses were not related to alcohol or tobacco use.

Previous research showed that drinking more during adolescence strongly predicts the prevalence of alcohol use disorders in adulthood (Bonomo et al., 2004). Therefore the adolescents in our sample who drank more may form a group of those at risk for later alcohol use problems. Though we were unable to examine causality in this study, these preliminary findings may provide support for the theory of an inherent hypo-arousal of the ANS in individuals more vulnerable to substance use problems. These individuals may deliberately seek out and use alcohol or tobacco in order to achieve a state of normalized arousal (Goeders, 2003; Majewska, 2002). Because this frequent use of substances as a way of seeking stimulation occurs at an early age, during adolescence, these individuals could be more vulnerable to SUDs later in life.

Interestingly, adolescents who drank medium and high quantities of alcohol per week showed a lower HR during the entire stress procedure; a between-subjects effect was evident in the analyses. However, no interaction effect was observed, as we had expected. All groups showed a relatively similar peak in HR in response to the tasks. Adolescents who drank more thus did not react physiologically differently to the stressful tasks; they portrayed a more general lowered HR. As this study was performed cross-sectionally, we are unable to differentiate whether this effect is due to underlying ANS variation, or whether use of alcohol has already affected the ANS in these adolescents. However, due to the young age of the adolescents in our sample, and that they have as yet used relatively little alcohol, we consider it unlikely that the observed differences of the ANS are due to the use of alcohol and thus are possibly due to an underlying difference in general ANS regulation. Our results suggest additionally that this difference may not lie in the immediate response to stress, rather in overall ANS activity across situations. The present results are supported by our previously reported findings, in the same sample, of lower hypothalamus-pituitary-adrenal (HPA) axis activity (indexed by salivary cortisol) during the Rest and Task period in adolescents who began drinking at an earlier age (Evans et al., 2012). HPA and ANS measures in this sample were related, as indicated by significant positive correlations between cortisol and HR during Rest ($R = .18, p < .01$) and Task ($R = .32, p < .001$), as well as the response measures ($R = .16, p < .05$), thus reinforcing our findings of low physiological arousal in those more prone to risky substance use. These findings are in line with earlier suggestions that physiological stress response dysregulation in adolescents may signal vulnerability to various kinds of psychopathology (Stroud et al., 2009).

This is the first study to examine the relation between alcohol use and HR in a general adolescent population, therefore, the results are preliminary and must be interpreted cautiously. Our finding that those who drank more portrayed a lower HR during the stress procedure is in line with one finding in adults with a FH of alcoholism (Sorocco et al., 2006), though in contrast to other similar studies which found increased HR in response to unavoidable shock (Finn et al., 1992; Finn and Pihl, 1987) and a mental arithmetic task (Harden and Pihl, 1995). Further research in this area is needed in order to clarify these contrasting findings.

We observed that PS was significantly and positively related to HR, which confirmed findings from a previous study in adolescents from the general population (Oldehinkel et al., 2011). We did not find a relation between PS and alcohol and tobacco use, corroborating earlier reports of no difference in PS between control subjects and those at risk for a SUD (Finn and Pihl, 1987) and those exhibiting more externalizing problems (Fairchild et al., 2008). This was in line with our expectations; physiological responses reflect

underlying, biological processes, and we would not necessarily expect similar relations to be found with the subjective experience of a stressor. Physiological and perceived stress are distinct constructs (Oldehinkel et al., 2011), which was substantiated in our finding of a significant and positive, but not strong, correlation between HR and PS.

Our observations indicate a relation between tobacco use and HR reactivity. Those who smoked every day showed a blunted HR response to the stressful tasks compared to those who smoked less frequently or not at all. This finding is in line with several findings on adult smokers (Girdler et al., 1997; Phillips et al., 2009; Roy et al., 1994; Sheffield et al., 1997; Straneva et al., 2000) though is in contrast to other studies (Back et al., 2008; Childs and de Wit, 2009; Hughes and Higgins, 2010; Kirschbaum et al., 1993; Perkins et al., 1992; Tersman et al., 1991). While two studies examining HR reactivity in low versus high frequency tobacco users found no difference between these groups (both portrayed attenuated responses), we found that adolescents who smoked less frequently did not differ significantly from those who had never smoked. It is possible that in adolescents, underlying variation of the ANS is only evident in those who use tobacco more frequently.

Attenuated HR during rest as well as in reaction to stress in those who exhibit more externalizing problems has been well established (Ortiz and Raine, 2004), as has the relation between externalizing problems and substance use (Krueger et al., 2002; Young et al., 2000; Zimmermann et al., 2007). Therefore, we examined whether hypo-arousal of the ANS was specific to substance use, as opposed to generally related to externalizing problems. When number of externalizing problems was controlled for in the model, the relations between HR and alcohol and tobacco use remained significant, thus providing evidence for a relation between HR hypo-arousal and substance use specifically.

The time at which the last cigarette was smoked prior to the stress procedure was not asked, thus it cannot be excluded that the observed blunted HR reactivity to stress was due to nicotine withdrawal; smokers may have been less able to concentrate and were therefore perhaps less able to engage in the stressful tasks (Phillips et al., 2009). However, in two studies, nicotine withdrawal did not influence the response to stress (al'Absi et al., 2003; Tsuda et al., 1996). Furthermore, smokers exhibited a blunted reaction to stress whether they did or did not wear a nicotine patch (Girdler et al., 1997). Moreover, it is unlikely that cigarettes that may have been smoked just prior to the session influenced HR during the stress procedure due to the fact that nicotine causes an increase in HR (Hasenfratz and Battig, 1992; James and Richardson, 1991; Pauli et al., 1993) while we observed no differences in pre-task HR in High Frequency Smokers as compared to Non- and Low Frequency Smokers.

With regard to this study, the following should be taken into account. As mentioned above, the nature of the study was cross-sectional. We are unable to exclude the possibility of alcohol and tobacco use affecting the ANS directly. While we believe it to be unlikely because they have used relatively little alcohol and tobacco, this remains to be elucidated. Secondly, possible effects of a third variable cannot be eliminated. It is viable that factors such as temperament or the experience of stressful events in the past influenced the relation between alcohol and tobacco use and physiological stress (re)activity in this study. Thirdly, we had no information regarding the time at which the last cigarette was smoked, and therefore cannot exclude possible direct effects of nicotine or nicotine withdrawal on HR. Fourthly, we based our alcohol use variable on third percentiles, as has been done in previous studies (Hillers and Massey, 1985; Murray et al., 2002), in order to obtain relatively equal groups. A more widely used variable, such as binge-drinking, would be interesting, but was not possible in a general population group of this size and of this age group in which

most adolescents have not yet begun using large amounts of alcohol. Future follow-up measurements of this group will allow a more comprehensive investigation of risky substance use in relation to physiological stress (re)activity.

In conclusion, lower HR during a psychosocial stress procedure was evident in those adolescents from the general population who drank more as compared to those who drank less. Also, adolescents who smoked every day showed a blunted HR response to the stressful tasks as compared to those who smoked less frequently or not at all. PS was significantly correlated with HR, and was not related to alcohol or tobacco use. Though data are preliminary and must be interpreted with caution, it is possible that adolescents prone to use more alcohol and tobacco may be characterized by hypo-arousal. Conceivably, these adolescents use substances in order to seek stimulation and achieve a more normalized state of arousal. As they use more alcohol and tobacco, they may be vulnerable to SUDs later in life. ANS activity during rest as well as in reaction to stress may thus be a potential vulnerability marker for SUDs in adolescents.

Role of funding source

This project is supported by ZonMW grant 3116.0002 and partly by ERAB grant 0609. Neither ZonMW nor ERAB had any role in the study design; in the collection, analysis and interpretation of data, in the writing of the report, or in the decision to submit the paper for publication.

Contributors

A.C. Huizink and I.H.A. Franken designed the study and wrote the protocol, and collected the funding for this study. A.C. Huizink supervised the whole project. B.E. Evans managed the literature searches and summaries of previous related work, undertook the statistical analysis, and wrote the first draft of the manuscript. K. Greaves-Lord and A.S. Euser contributed to the statistical analyses, the revision of the manuscript and the collection of data. All authors contributed to and have approved the final manuscript.

Conflict of interest

No conflict declared.

Acknowledgments

We are grateful to all participants of the South Holland 2 Study and their parents. We also thank Olga Husson for her assistance in data collection.

References

- Achenbach, T., Rescorla, L.A., 2001. Manual for the ASEBA School-Age Forms & Profiles. University of Vermont, Research Center for Children, Youth and Families, Burlington, VT.
- al'Absi, M., Wittmers, L.E., Erickson, J., Hatsukami, D., Crouse, B., 2003. Attenuated adrenocortical and blood pressure responses to psychological stress in ad libitum and abstinent smokers. *Pharmacol. Biochem. Behav.* 74, 401–410.
- Back, S.E., Waldrop, A.E., Saladin, M.E., Yeatts, S.D., Simpson, A., McRae, A.L., Upadhyaya, H.P., Sisson, R.C., Spratt, E.G., Allen, J., Kreek, M.J., Brady, K.T., 2008. Effects of gender and cigarette smoking on reactivity to psychological and pharmacological stress provocation. *Psychoneuroendocrinology* 33, 560–568.
- Bonomo, Y.A., Bowes, G., Coffey, C., Carlin, J.B., Patton, G.C., 2004. Teenage drinking and the onset of alcohol dependence: a cohort study over seven years. *Addiction* 99, 1520–1528.
- Carroll, D., Phillips, A.C., Der, G., 2008. Body mass index, abdominal adiposity, obesity, and cardiovascular reactions to psychological stress in a large community sample. *Psychosom. Med.* 70, 653–660.
- Childs, E., de Wit, H., 2009. Hormonal, cardiovascular, and subjective responses to acute stress in smokers. *Psychopharmacology* 203, 1–12.

- Creemers, H.E., Korhonen, T., Kaprio, J., Vollebergh, W.A.M., Ormel, J., Verhulst, F.C., Huizink, A.C., 2009. The role of temperament in the relationship between early onset of tobacco and cannabis use: the TRAILS study. *Drug Alcohol Depend.* 104, 113–118.
- Dieleman, G.C., van der Ende, J., Verhulst, F.C., Huizink, A.C., 2010. Perceived and physiological arousal during a stress task: can they differentiate between anxiety and depression? *Psychoneuroendocrinology* 35, 1223–1234.
- Dietrich, A., Riese, H., Sondejker, F.E.P.L., Greaves-Lord, K., van Roon, A.M., Ormel, J., Neeleman, J., Rosmalen, J.G.M., 2007. Externalizing and internalizing problems in relation to autonomic function: a population-based study in preadolescents. *J. Am. Acad. Child Adolesc. Psychiatry* 46, 378–386.
- Evans, B.E., Greaves-Lord, K., Euser, A.S., Franken, I.H.A., Huizink, A.C., 2012. The relation between hypothalamic–pituitary–adrenal (HPA) axis activity and age of onset of alcohol use. *Addiction* 107, 312–322.
- Fairchild, G., van Gozen, S.H.M., Stollery, S.J., Brown, J., Gardiner, J., Herbert, J., Goodyer, I.M., 2008. Cortisol diurnal rhythm and stress reactivity in male adolescents with early-onset or adolescence-onset conduct disorder. *Biol. Psychiatry* 64, 599–606.
- Finn, P.R., Earleywine, M., Pihl, R.O., 1992. Sensation seeking, stress reactivity, and alcohol dampening discriminate the density of a family history of alcoholism. *Alcohol. Clin. Exp. Res.* 16, 585–590.
- Finn, P.R., Pihl, R.O., 1987. Men at high-risk for alcoholism—the effect of alcohol on cardiovascular-response to unavoidable shock. *J. Abnorm. Psychol.* 96, 230–236.
- Girdler, S.S., Jamner, L.D., Jarvik, M., Soles, J.R., Shapiro, D., 1997. Smoking status and nicotine administration differentially modify hemodynamic stress reactivity in men and women. *Psychosom. Med.* 59, 294–306.
- Goeders, N.E., 2003. The impact of stress on addiction. *Eur. Neuropsychopharmacol.* 13, 435–441.
- Greaves-Lord, K., Ferdinand, R.F., Sondejker, F.E.P.L., Dietrich, A., Oldehinkel, A.J., Rosmalen, J.G.M., Ormel, J., Verhulst, F.C., 2007. Testing the tripartite model in young adolescents: is hyperarousal specific for anxiety and not depression? *J. Affect. Disord.* 102, 55–63.
- Harden, P.W., Pihl, R.O., 1995. Cognitive function, cardiovascular reactivity, and behavior in boys at high-risk for alcoholism. *J. Abnorm. Psychol.* 104, 94–103.
- Hardin, M.G., Ernst, M., 2009. Functional brain imaging of development-related risk and vulnerability for substance use in adolescents. *J. Addict. Med.* 3, 47–54.
- Hasenfrazt, M., Battig, K., 1992. Action profiles of smoking and caffeine—stroop effect, EEG, and peripheral physiology. *Pharmacol. Biochem. Behav.* 42, 155–161.
- Hillers, V.N., Massey, L.K., 1985. Interrelationships of moderate and high alcohol consumption with diet and health status. *Am. J. Clin. Nutr.* 41, 356–362.
- Hughes, B.M., Higgins, N.M., 2010. Habituation-sensitization of cardiovascular reactivity to repeated stress in smokers and non-smokers: an anthropometrically matched trial. *Int. J. Psychophysiol.* 76, 34–39.
- Huizink, A.C., Ferdinand, R.F., Ormel, J., Verhulst, F.C., 2006. Hypothalamic–pituitary–adrenal axis activity and early onset of cannabis use. *Addiction* 101, 1581–1588.
- Ingjaldsson, J.T., Laberg, J.C., Thayer, J.F., 2003. Reduced heart rate variability in chronic alcohol abuse: relationship with negative mood, chronic thought suppression, and compulsive drinking. *Biol. Psychiatry* 54, 1427–1436.
- James, J.E., Richardson, M., 1991. Pressor effects of caffeine and cigarette-smoking. *Br. J. Clin. Psychol.* 30, 276–278.
- Khantzian, E.J., 1985. The self-medication hypothesis of addictive disorders: focus on heroin and cocaine dependence. *Am. J. Psychiatry* 142, 1259–1264.
- Kirschbaum, C., Strasburger, C.J., Langkrär, J., 1993. Attenuated cortisol response to psychological stress but not to CRH or ergometry in young habitual smokers. *Pharmacol. Biochem. Behav.* 44, 527–531.
- Krueger, R.F., Hicks, B.M., Patrick, C.J., Carlson, S.R., Iacono, W.G., McGue, M., 2002. Etiologic connections among substance dependence, antisocial behavior, and personality: modeling the externalizing spectrum. *J. Abnorm. Psychol.* 111, 411–424.
- Liu, J.H., Raine, A., Wuerker, A., Venables, P.H., Mednick, S., 2009. The association of birth complications and externalizing behavior in early adolescents: direct and mediating effects. *J. Res. Adolesc.* 19, 93–111.
- Majewska, M.D., 2002. HPA axis and stimulant dependence: an enigmatic relationship. *Psychoneuroendocrinology* 27, 5–12.
- Marshall, W.A., Tanner, J.M., 1970. Variations in the pattern of pubertal changes in boys. *Arch. Dis. Child.* 45, 13–23.
- Martin, C.A., Kelly, T.H., Rayens, M.K., Brogli, B.R., Brenzel, A., Smith, W.J., Omar, H.A., 2002. Sensation seeking, puberty, and nicotine, alcohol, and marijuana use in adolescence. *J. Am. Acad. Child Adolesc. Psychiatry* 41, 1495–1502.
- Miller, G.E., Chen, E., Fok, A.K., Walker, H., Lim, A., Nicholls, E.F., Cole, S., Kobor, M.S., 2009. Low early-life social class leaves a biological residue manifested by decreased glucocorticoid and increased proinflammatory signaling. *Proc. Natl. Acad. Sci. U.S.A.* 106, 14716–14721.
- Murray, R.P., Connett, J.E., Tyas, S.L., Bond, R., Ekuma, O., Silversides, C.K., Barnes, G.E., 2002. Alcohol volume, drinking pattern, and cardiovascular disease morbidity and mortality: is there a U-shaped function? *Am. J. Epidemiol.* 155, 242–248.
- Oldehinkel, A.J., Ormel, J., Bosch, N.M., Bouma, E.M.C., Van Roon, A.M., Rosmalen, J.G.M., Riese, H., 2011. Stressed out? Associations between perceived and physiological stress responses in adolescents: the TRAILS study. *Psychophysiology* 48, 441–452.
- Ortiz, J., Raine, A., 2004. Heart rate level and antisocial behavior in children and adolescents: a meta-analysis. *J. Am. Acad. Child Adolesc. Psychiatry* 43, 154–162.
- Panknin, T.L., Dickensheets, S.L., Nixon, S.J., Lovallo, W.R., 2002. Attenuated heart rate responses to public speaking in individuals with alcohol dependence. *Alcohol. Clin. Exp. Res.* 26, 841–847.
- Pauli, P., Rau, H., Zhuang, P., Brody, S., Birbaumer, N., 1993. Effects of smoking on thermal pain threshold in deprived and minimally-deprived habitual smokers. *Psychopharmacology* 111, 472–476.
- Perkins, K.A., Grobe, J.E., Fonte, C., Breus, M., 1992. Paradoxical effects of smoking on subjective stress versus cardiovascular arousal in males and females. *Pharmacol. Biochem. Behav.* 42, 301–311.
- Phillips, A.C., Der, G., Hunt, K., Carroll, D., 2009. Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *Int. J. Psychophysiol.* 73, 273–278.
- Porges, S.W., 1995. Cardiac vagal tone: a physiological index of stress. *Neurosci. Biobehav. Rev.* 19, 225–233.
- Raine, A., 2002. Annotation: the role of prefrontal deficits, low autonomic arousal, and early health factors in the development of antisocial and aggressive behavior in children. *J. Child Psychol. Psychiatry* 43, 417–434.
- Roy, M.P., Steptoe, A., Kirschbaum, C., 1994. Association between smoking status and cardiovascular and cortisol stress responsivity in healthy young men. *Int. J. Behav. Med.* 1, 264–283.
- Sheffield, D., Smith, G.D., Carroll, D., Shipley, M.J., Marmot, M.G., 1997. The effects of recent food, alcohol, and tobacco intake and the temporal scheduling of testing on cardiovascular activity at rest and during psychological stress. *Psychophysiology* 34, 204–212.
- Sinha, R., 2008. Chronic Stress, Drug Use, and Vulnerability to Addiction. *Addiction Reviews* 2008. Blackwell Publishing, Oxford, pp. 105–130.
- Sinha, R., Fox, H.C., Hong, K.A., Bergquist, K., Bhagwagar, Z., Siedlarz, K.M., 2009. Enhanced negative emotion and alcohol craving, and altered physiological responses following stress and cue exposure in alcohol dependent individuals. *Neuropsychopharmacology* 34, 1198–1208.
- Sorocco, K.H., Lovallo, W.R., Vincent, A.S., Collins, F.L., 2006. Blunted hypothalamic–pituitary–adrenocortical axis responsivity to stress in persons with a family history of alcoholism. *Int. J. Psychophysiol.* 59, 210–217.
- Statistics, C.B.O., 2010. Standard Classification of Professions. Den Haag/Heerlen: Central Bureau of Statistics.
- Straneva, P., Hinderliter, A., Wells, E., Lenahan, H., Girdler, S., 2000. Smoking, oral contraceptives, and cardiovascular reactivity to stress. *Obstet. Gynecol.* 95, 78–83.
- Stroud, L.R., Foster, E., Papandonatos, G.D., Handwerker, K., Granger, D.A., Kivlighan, K.T., Niaura, R., 2009. Stress response and the adolescent transition: performance versus peer rejection stressors. *Dev. Psychopathol.* 21, 47–68.
- Tersman, Z., Collins, A., Eneroth, P., 1991. Cardiovascular-responses to psychological and physiological stressors during the menstrual-cycle. *Psychosom. Med.* 53, 185–197.
- Thayer, R.E., 1970. Activation states as assessed by verbal report and four psychophysiological variables. *Psychophysiology* 7, 86–94.
- Tsuda, A., Steptoe, A., West, R., Fieldman, G., Kirschbaum, C., 1996. Cigarette smoking and psychophysiological stress responsiveness: effects of recent smoking and temporary abstinence. *Psychopharmacology* 126, 226–233.
- Young, S.E., Stallings, M.C., Corley, R.P., Krauter, K.S., Hewitt, J.K., 2000. Genetic and environmental influences on behavioral disinhibition. *Am. J. Med. Genet.* 96, 684–695.
- Zimmermann, U.S., Blomeyer, D., Laucht, M., Mann, K.F., 2007. How gene-stress-behavior interactions can promote adolescent alcohol use: the roles of predrinking allostatic load and childhood behavior disorders. *Pharmacol. Biochem. Behav.* 86, 246–262.
- Zuckerman, M., Kuhlman, D.M., 2000. Personality and risk-taking: common biosocial factors. *J. Pers.* 68, 999–1029.
- Zuckerman, M., Neeb, M., 1979. Sensation seeking and psychopathology. *Psychiatry Res.* 1, 255–264.