

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/107990>

Please be advised that this information was generated on 2019-02-23 and may be subject to change.

# Genetic Overlap between Apparently Sporadic Motor Neuron Diseases

Marka van Blitterswijk<sup>1\*</sup>, Lotte Vlam<sup>1</sup>, Michael A. van Es<sup>1</sup>, W-Ludo van der Pol<sup>1</sup>, Eric A. M. Hennekam<sup>2</sup>, Dennis Dooijes<sup>2</sup>, Helenius J. Schelhaas<sup>3</sup>, Anneke J. van der Kooi<sup>4</sup>, Marianne de Visser<sup>4</sup>, Jan H. Veldink<sup>1,9</sup>, Leonard H. van den Berg<sup>1,9</sup>

**1** Department of Neurology, Rudolf Magnus Institute of Neuroscience, University Medical Center Utrecht, Utrecht, The Netherlands, **2** Department of Medical Genetics, University Medical Center Utrecht, Utrecht, The Netherlands, **3** Department of Neurology, Donders Institute for Brain, Cognition and Behavior, Radboud University Nijmegen Medical Center, Nijmegen, The Netherlands, **4** Department of Neurology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

## Abstract

Progressive muscular atrophy (PMA) and amyotrophic lateral sclerosis (ALS) are devastating motor neuron diseases (MNDs), which result in muscle weakness and/or spasticity. We compared mutation frequencies in genes known to be associated with MNDs between patients with apparently sporadic PMA and ALS. A total of 261 patients with adult-onset sporadic PMA, patients with sporadic ALS, and control subjects of Dutch descent were obtained at national referral centers for neuromuscular diseases in The Netherlands. Sanger sequencing was used to screen these subjects for mutations in the coding regions of superoxide dismutase-1 (*SOD1*), angiogenin (*ANG*), fused in sarcoma/translated in liposarcoma (*FUS/TLS*), TAR DNA-binding protein 43 (*TARDBP*), and multivesicular body protein 2B (*CHMP2B*). In our cohort of PMA patients we identified two *SOD1* mutations (p.D90A, p.I113T), one *ANG* mutation (p.K171), one *FUS/TLS* mutation (p.R521H), one *TARDBP* mutation (p.N352S), and one novel *CHMP2B* mutation (p.R69Q). The mutation frequency of these genes was similar in sporadic PMA (2.7%) and ALS (2.0%) patients, and therefore, our findings demonstrate a genetic overlap between apparently sporadic PMA and ALS.

**Citation:** van Blitterswijk M, Vlam L, van Es MA, van der Pol W-L, Hennekam EAM, et al. (2012) Genetic Overlap between Apparently Sporadic Motor Neuron Diseases. PLoS ONE 7(11): e48983. doi:10.1371/journal.pone.0048983

**Editor:** Udai Pandey, Louisiana State University Health Sciences Center, United States of America

**Received:** September 5, 2012; **Accepted:** October 3, 2012; **Published:** November 14, 2012

**Copyright:** © 2012 van Blitterswijk et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The research leading to these results has received funding from the European Community's Health Seventh Framework Program (FP7/2007–2013) (grant agreement no. 259867), the VSB fonds, The Brain Foundation of the Netherlands, Prinses Beatrix Fonds, Catharijne Stichting, H. Kersten and M. Kersten, J. R. van Dijk, and the Adessium Foundation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: L.H.vandenBerg@umcutrecht.nl

<sup>9</sup> These authors contributed equally to this work.

## Introduction

Motor neuron diseases (MNDs) are a heterogeneous group of disorders characterized by muscle weakness and/or spasticity due to degeneration of motor neurons. Progressive muscular atrophy (PMA) refers to a subgroup of the MND patients with rapidly or gradually developing muscle weakness. PMA accounts for 5–10% of adult-onset MNDs, and is caused by a progressive loss of lower motor neurons (LMNs) [1,2]. Differentiation of PMA from amyotrophic lateral sclerosis (ALS) is important, since the median survival of patients with PMA is significantly longer than that of patients with ALS [3].

The etiology of MNDs is complex. Most of the ALS cases, for instance, are sporadic in nature and thought to be caused by an interaction of genetic and environmental factors [4]. Currently, many genes appear to be involved in the pathogenesis of ALS, including chromosome 9 open reading frame 72 (*C9orf72*), superoxide dismutase-1 (*SOD1*), angiogenin (*ANG*), fused in sarcoma/translated in liposarcoma (*FUS/TLS*), TAR DNA-binding protein 43 (*TARDBP/TDP-43*), vesicle-associated membrane protein B (*VAPB*), optineurin (*OPTN*), valosin-containing protein (*VCP*), ubiquilin-2 (*UBQLN2*), sequestosome-1 (*SQSTM1*),

and profilin-1 (*PFN1*) [5,6,7,8,9,10,11]. We have recently shown that *C9orf72* repeat expansions can also be detected in apparently sporadic PMA, but at a lower frequency (1.6%) than in apparently sporadic ALS (6.1%) [12]. Moreover, we have demonstrated that mutations in four major MND-associated genes, *SOD1*, *ANG*, *FUS/TLS* and *TARDBP*, account for less than two percent of the sporadic ALS cases [13]. The combined mutation frequency of these four MND-associated genes is unknown for sporadic PMA patients. Mutations in charged multivesicular body protein 2B (*CHMP2B*) have, however, been reported in sporadic PMA patients [14,15].

The objective of this study is to determine the mutation frequency of major MND-associated genes in patients with apparently sporadic PMA. We compared their mutation frequencies to those in a large cohort of patients with apparently sporadic ALS, revealing a genetic overlap.

## Materials and Methods

### Patient Selection

We included 261 patients with apparently sporadic PMA and screened their DNA for mutations in *SOD1*, *ANG*, *FUS/TLS*,

*TARDBP*, and *CHMP2B*. PMA patients had already been screened for mutations in transient receptor potential vanilloid 4 (*TRPV4*) and repeat expansions in *C9orf72* [12,16]. Their diagnosis was based on LMN involvement on clinical and electrophysiological examination at time of referral. We excluded patients with a family history of PMA, a history of acute poliomyelitis, spinal radiculopathy, diabetic amyotrophy, thyrotoxicosis, or hyperparathyroidism, clinical signs of upper motor neuron (UMN) involvement, sensory signs on neurological examination, structural lesions on magnetic resonance imaging or computed tomography of head and spine, and motor conduction block on extensive standardized nerve conduction studies [2].

Cohorts of sporadic ALS patients had already been screened for mutations in *SOD1* (N = 451), *ANG* (N = 941), *FUS/TLS* (N = 1,192), *TARDBP* (N = 1,192), and *C9orf72* (N = 1,422) [12,13,17,18,19]. We screened 1,002 sporadic ALS patients for mutations in *CHMP2B*. ALS patients were recruited through the Dutch Prospective Population-based ALS registry; they were diagnosed according to the El Escorial Criteria at national referral centers for neuromuscular diseases (University Medical Center Utrecht, Academic Medical Center Amsterdam, or Radboud University Nijmegen Medical Center) [20,21].

Mutations in *SOD1* (N = 1,894), *ANG* (N = 1,582), *FUS/TLS* (N = 970), and *TARDBP* (N = 1,415) had previously been reported in Dutch control subjects [13,18,19]. We screened a total of 750 control subjects of Dutch descent for mutations in *CHMP2B*.

### Ethics Statement

All material was obtained with approval of the medical ethics committee for research in humans of the University Medical Center Utrecht, The Netherlands, and all participants gave written informed consent.

### Genetic Analysis

Coding regions of *SOD1* (NM\_000454.4), *ANG* (NM\_001145.4), *FUS/TLS* (NM\_004960.3, exon 5, 6, 14, 15), *TARDBP* (NM\_007375.3, exon 6), and *CHMP2B* (NM\_014043.3) were screened for mutations using touchdown PCR, as described previously [13,22]. Sanger sequencing and data analysis were performed with BigDye Terminator 3.1 sequencing kit (Applied Biosystems, Foster City, California), DNA Analyzer 3730XL (Applied Biosystems) and PolyPhred [23]. Each mutation was confirmed on genomic DNA and its impact on the structure, and function of the protein was predicted with PolyPhen-2 (PolyPhen-2 version 2.1.0; <http://genetics.bwh.harvard.edu/pph2/>) and PMut (<http://mmb.pcb.ub.es/PMut/PMut.jsp>).

### Genealogical Analysis

Lists of descendants were compiled for index patients. Based on these lists, civil records/registers, and church records of the Dutch population, pedigrees were generated (containing two parents, four grandparents, eight great-grandparents, etc.). This information was then used to determine whether index patients were related, and detailed family trees were constructed.

### Haplotype Analysis

Extended haplotype analysis, using six extragenic polymorphic markers flanking *TARDBP* (D1S1612, D1S503, D1S244 proximal of *TARDBP*, and D1S2667, D1S2740 and D1S1597 distal of *TARDBP*), was performed to construct a haplotype segregating with the identified p.N352S mutation in *TARDBP*. Validity of the constructed haplotype was determined by segregation analysis in families and patients whose DNA was available for testing.

### Statistical Analysis

A Fisher's exact test or Chi-square test was used to compare mutation frequencies, gender, site of onset, and current status (alive/deceased) between PMA and ALS patients; a Mann-Whitney test was used to compare age at onset and disease duration (GraphPad Prism version 5; <http://www.graphpad.com>). P-values below 0.05 were considered significant.

## Results

### Study Population

Baseline characteristics of the 261 sporadic PMA patients and 1,002 sporadic ALS patients are shown in Table 1. Patients with PMA were more likely to be male (72% versus 59%); furthermore, they lived longer (7.6 year versus 3.8 year), and had a lower age at onset (58.0 year versus 60.6 year) than patients with ALS.

### Mutation Frequencies

Table 2 summarizes the mutations found in patients and control subjects. In individual PMA patients we detected heterozygous mutations in *SOD1* (p.D90A [c.1078A>C], p.I113T [c.1147T>C]), *ANG* (p.K17I [c.122A>T]), *FUS/TLS* (p.R521H [c.1562G>A]) and *TARDBP* (p.N352S [c.1055A>G]), accounting for 2.3% of the patients. Previously, we showed that missense mutations in *SOD1*, *ANG*, *FUS/TLS* and *TARDBP* were present in 1.7% of the ALS patients, and 0.4% of the control subjects [13,17,18,19]. In our current study, we also identified four novel *CHMP2B* mutations, one of which was present in a PMA patient (p.R69Q [c.206G>A]), and three in ALS patients (p.R22Q [c.65G>A], p.N54T [c.161A>C], p.T83I [c.248C>T]). All four *CHMP2B* mutations are located in a domain that is important for the formation of multivesicular bodies (MVBs), involved in sorting of cargo proteins to intraluminal vesicles [24]. These mutations are located in well conserved areas (Figure S1) and predicted to be pathological (Table 2). They account for 0.38% of the sporadic PMA patients and 0.30% of the sporadic ALS patients. None of these *CHMP2B* mutations was present in our control subjects; however, in one control subject (0.13%) we did detect a mutation (p.S194L [c.581C>T]) that had previously been reported in a patient with frontotemporal dementia (FTD) [25].

### Clinical Characteristics

The average age at onset of PMA patients with missense mutations was 48 years, and five of them were male (71%). Although only one of these patients had died, their average disease duration already exceeded 114 months (range 37–316). These clinical characteristics of sporadic PMA patients with missense mutations were consistent with the characteristics of our entire PMA cohort. More detailed signs and symptoms are provided in Table 3.

### Genealogical- and Haplotype Analyses

Previously, we have shown that the p.N352S mutation in *TARDBP* is a founder mutation in the Dutch ALS population [13]. Hence, we performed a thorough genealogical analysis and demonstrated that our PMA patients with p.N352S mutations had common ancestors, dating back to the 17<sup>th</sup> century in the north of France (Figure S2). Haplotype analysis revealed that these patients also shared the haplotype that was reported in Dutch ALS patients [13].

**Table 1.** Baseline characteristics of study population.

Cohort	Number (N)	Male/female (N) (%)	Age at onset (y) (CI)	Alive/deceased (N) (%)	Duration (y) (CI)
PMA	261	187/74 (72/28)	58.0 (56.4–59.7)	137/116 (54/46)	7.6 (6.7–8.5)
ALS	1,002	593/409 (59/41)	60.6 (59.8–61.3)	135/854 (14/86)	3.8 (3.6–4.1)

Abbreviations: PMA = progressive muscular atrophy, ALS = amyotrophic lateral sclerosis, N = number, y = years, and CI = 95% confidence interval. Disease duration is defined as the interval between age at onset and age at death, or between age at onset and age last known to be alive. Patients with sporadic PMA are more likely to be male (p-value 0.001), to have a lower age at onset (p-value 0.010), to be alive (p-value <0.001), and to have a longer disease duration than patients with sporadic ALS (p-value <0.001).

doi:10.1371/journal.pone.0048983.t001

## Discussion

Patients with isolated LMN signs represent a subgroup of the patients with MND. To assess the mutation frequency of MND-associated genes in this subgroup, we compared 261 apparently sporadic PMA patients to apparently sporadic ALS patients. Our PMA patients were more likely to be male and lived significantly longer than ALS patients, as reported previously [3]. We detected two *SOD1* mutations (p.D90A, p.I113T), one *ANG* mutation (p.K17I), one *FUS/TLS* mutation (p.R521H), one *TARDBP*

mutation (p.N352S), and one novel *CHMP2B* mutation (p.R69Q) in individual PMA patients. For each of these genes we compared mutation frequencies between our PMA patients and ALS patients, and did not detect significant differences.

Clinical and pathological similarities between PMA and ALS have already been reported: more than twenty percent of the patients with isolated LMN signs will develop UMN signs within six years, especially in the first years after symptom onset [1,3,26]. Nonetheless, it can be difficult to diagnose these UMN signs due to LMN wasting and pathophysiological abnormalities caused by

**Table 2.** Missense mutations found in *SOD1*, *ANG*, *FUS/TLS*, *TARDBP*, and *CHMP2B*.

Gene	Variant	Exon	PMA	ALS	CON	Prediction PolyPhen-2	Prediction PMut
<i>SOD1</i>	p.D90A	4	1/261 <sup>a</sup>	1/451 [17]	3/1,894 [13]	Benign	Pathological
	p.I113T	4	1/261	0/451	0/1,894	Probably damaging	Pathological
	p.I99V	4	0/261	1/451	0/1,894	Benign	Neutral
	<b>Total (%)</b>		<b>2/261 (0.77)</b>	<b>2/451 (0.44)</b>	<b>3/1,894 (0.16)</b>		
<i>ANG</i>	p.G(-10)D	2	0/261 <sup>a</sup>	1/941 [18]	0/1,582 [18]	N/A	N/A
	p.K17I	2	1/261	3/941	2/1,582	Benign	Pathological
	p.T80S	2	0/261	1/941	0/1,582	Possibly damaging	Neutral
	p.F100I	2	0/261	1/941	0/1,582	Probably damaging	Neutral
	<b>Total (%)</b>		<b>1/261 (0.38)</b>	<b>6/941 (0.64)</b>	<b>2/1,582 (0.13)</b>		
<i>FUS/TLS</i>	p.S115N	5	0/261 <sup>a</sup>	1/1,192 [13]	0/970 [19]	Unknown	Neutral
	p.Q210H	6	0/261	0/1,192	1/970	Unknown	Neutral
	p.R487C	14	0/261	1/1,192	0/970	Probably damaging	Pathological
	p.R495X	14	0/261	1/1,192	0/970	N/A	N/A
	p.R521H	15	1/261	0/1,192	0/970	Probably damaging	Pathological
	<b>Total (%)</b>		<b>1/261 (0.38)</b>	<b>3/1,192 (0.17)</b>	<b>1/970 (0.10)</b>		
<i>TARDBP</i>	p.N352S	6	2/261 <sup>a</sup>	3/1,192 [13]	0/1,415 [13]	Benign	Pathological
	p.I383V	6	0/261	1/1,192	0/1,415	Benign	Neutral
	<b>Total (%)</b>		<b>2/261 (0.77)</b>	<b>4/1,192 (0.34)</b>	<b>0/1,415 (0.00)</b>		
<i>CHMP2B</i>	p.R22Q	2	0/261 <sup>a</sup>	1/1,002 <sup>a</sup>	0/750 <sup>a</sup>	Possibly damaging	Pathological
	p.N54T	3	0/261	1/1,002	0/750	Probably damaging	Neutral
	p.R69Q	3	1/261	0/1,002	0/750	Probably damaging	Pathological
	p.T83I	3	0/261	1/1,002	0/750	Probably damaging	Pathological
	p.S194L	6	0/261	0/1,002	1/750	Benign	Neutral
	<b>Total (%)</b>		<b>1/261 (0.38)</b>	<b>3/1,002 (0.30)</b>	<b>1/750 (0.13)</b>		
<b>Total (%)</b>		<b>7 (2.7)</b>	<b>18 (2.0)</b>	<b>7 (0.5)</b>			

Abbreviations: CON = control subjects, and N/A = not applicable. Mutations in *SOD1*, *ANG*, *FUS/TLS*, *TARDBP*, and *CHMP2B* were present in 2.7% of the PMA patients, 2.0% of the ALS patients, and 0.5% of the control subjects. No PMA patients were detected with mutations in multiple MND-associated genes. A Fisher's exact test or Chi-square test was used to compare mutation frequencies between patients with PMA and ALS for each gene; no significant differences were detected (data not shown for simplicity).

<sup>a</sup>Cohort described in the present study.

doi:10.1371/journal.pone.0048983.t002

**Table 3.** Clinical characteristics of newly identified patients with missense mutations.

Group	Gene	Variant	Gender	LMN <sup>a</sup> signs	UMN <sup>a</sup> signs	Age at onset (y)	Site of onset	Duration (m)
PMA	<i>SOD1</i>	p.D90A	M	1	0	17	Cervical	316
		p.I113T	F	2	0	48	Lumbosacral	108
	<i>ANG</i>	p.K17I	M	1	0	66	Lumbosacral	52
	<i>FUS/TLS</i>	p.R521H	M	3	0	47	Cervical	68
	<i>TARDBP</i>	p.N352S	F	2	0	68	Cervical	37
		p.N352S	M	4	0	61	Lumbosacral	101 <sup>b</sup>
	<i>CHMP2B</i>	p.R69Q	M	1	0	26	Cervical	116
ALS	<i>CHMP2B</i>	p.R22Q	M	3	2	57	Cervical	68
		p.N54T	F	3	2	68	Bulbar	28 <sup>b</sup>
		p.T83I	M	2	1	71	Cervical	75

Abbreviations: M = male, F = female, LMN = lower motor neuron, UMN = upper motor neuron, and m = months. Clinical characteristics of ALS patients with *SOD1*, *ANG*, *FUS/TLS* and *TARDBP* mutations have been described elsewhere [13,17,18,19].

<sup>a</sup>Number of affected body regions at time of diagnosis (maximum four: bulbar, cervical, thoracic or lumbosacral).

<sup>b</sup>Deceased.

doi:10.1371/journal.pone.0048983.t003

damaged motor pathways, motor neurons and interneurons [27]. Pathological studies have also revealed ubiquitinated inclusions and involvement of the corticospinal tract in PMA patients, which are typical for ALS patients and emphasize similarities between these diseases [28,29].

Previously, mutations in *SOD1*, *ANG*, *FUS/TLS*, *TARDBP* and *CHMP2B* have been identified in patients with a range of clinical phenotypes, including combinations of FTD, Parkinson's disease, and ALS [14,18,22,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45].

*CHMP2B* mutations have also been described in sporadic PMA patients, while mutations in MND-associated genes have been detected in familial ALS patients with predominantly LMN signs [15,19,28,46,47,48,49,50]. In addition, we have recently shown that *C9orf72* repeat expansions are present in sporadic PMA patients, but at a lower frequency than in sporadic ALS patients (1.6% versus 6.1%) [12]; other studies have shown that *C9orf72* repeat expansions were present in approximately 7% of white sporadic ALS patients from the USA, Europe and Australia, and that bulbar onset ALS is frequently encountered in patients with these expansions [51,52,53,54]. Our current findings demonstrate that mutations in *SOD1*, *ANG*, *FUS/TLS*, *TARDBP* and *CHMP2B* are also associated with apparently sporadic PMA, thus expanding the wide range of clinical phenotypes reported to date. Furthermore, the comparable mutation frequencies between PMA and ALS patients show that, apart from clinical and pathological similarities, these diseases demonstrate a genetic overlap as well, suggesting that PMA is a subtype of ALS.

We detected a *SOD1* mutation (p.D90A) in a patient with sporadic PMA, a patient with classical sporadic ALS, and control subjects. This is the most common *SOD1* mutation, and causes both autosomal dominant and recessive ALS [55,56]. Although it behaves dominantly in many families, it is a polymorphism in the Swedish population, primarily causing ALS when in the homozygous state [57]. Another *SOD1* mutation (p.I113T) was also present in a PMA patient; it is known for its clinical heterogeneity, including asymptomatic subjects, patients with mild fasciculations, patients with typical ALS, and patients with ALS-FTD and chorea [41,58]. Both these *SOD1* mutations appear to result in ALS through aggregation of mutant *SOD1* protein [59].

In addition, we identified an *ANG* mutation (p.K17I) in one PMA patient, and in two out of 1,582 Dutch control subjects. The

p.K17I mutation has already been reported in ALS patients and in control subjects [18]. Despite its presence in control subjects, it does affect the neuroprotective-, angiogenic- and ribonucleolytic activity of *ANG* [50,60,61]. It seems likely that this mutation raises ALS susceptibility and/or acts as a genetic modifier, a hypothesis supported by recent reports of families that harbor a p.K17I mutation in combination with *TARDBP*- or *FUS/TLS* mutations, and a large international collaborative study, which demonstrates that *ANG* mutations confer a substantial risk for ALS [13,18,62].

In one PMA patient, we identified a *FUS/TLS* mutation (p.R521H); one of the most common *FUS/TLS* mutations with a disease duration of approximately four years [63,64]. In two other PMA patients we detected a *TARDBP* mutation (p.N352S), which has been described in German and Japanese ALS patients [65,66,67]. We have recently reported that p.N352S is a founder mutation in the Dutch ALS population [13]. In the present study, we revealed that our PMA patients had common ancestors and shared a haplotype also detected in Dutch ALS patients.

The four *CHMP2B* mutations we detected (p.R69Q, p.R22Q, p.N54T, p.T83I) are novel, absent in control subjects, located in well conserved areas, and predicted to be pathogenic. One of these was identified in a patient with sporadic PMA (0.38%), three in patients with sporadic ALS (0.30%). These mutation frequencies demonstrate that *CHMP2B* mutations are not specific for PMA, but are present in patients with PMA, FTD, ALS-FTD, and ALS. We also detected one previously reported mutation (p.S194L) in a control subject [25]. Since this variant is located within an area of low complexity and predicted to have neutral effects, it probably represents a rare benign polymorphism.

Recently, we have provided evidence for an oligogenic etiology of familial ALS [13]. We reported five families with mutations in multiple MND-associated genes: *ANG* mutations were detected in combination with *FUS/TLS* and *TARDBP* mutations, and *C9orf72* repeat expansions in combination with *TARDBP*, *SOD1*, and *FUS/TLS* mutations (p-value  $1.57 \times 10^{-7}$ ). In our present study, we did not identify mutations in multiple MND-associated genes in a single patient; however, the high phenotypic variability that is seen amongst patients with mutations in these genes (ranging from FTD to Parkinson's disease and MNDs) does suggest that other genetic and/or environmental factors influence disease characteristics of sporadic MNDs.

To summarize, we have detected comparable mutation frequencies in patients with apparently sporadic PMA and ALS, indicating a genetic overlap between these two diseases. Thus, our findings favor the hypothesis that PMA is a subtype of ALS and not a distinct entity, broadening the disease spectrum of ALS.

## Supporting Information

**Figure S1 CHMP2B mutations and conservation.** Conservation of amino-acid residues across species was generated using ClustalW2 online tool, <http://www.ebi.ac.uk/Tools/msa/clustalw2/>. (DOC)

## References

- Visser J, van den Berg-Vos RM, Franssen H, van den Berg LH, Wokke JH, et al. (2007) Disease course and prognostic factors of progressive muscular atrophy. *Arch Neurol* 64: 522–528.
- Van den Berg-Vos RM, Visser J, Kalmijn S, Fischer K, de Visser MI, et al. (2009) A long-term prospective study of the natural course of sporadic adult-onset lower motor neuron syndromes. *Arch Neurol* 66: 751–757.
- Kim WK, Liu X, Sandner J, Pasmantier M, Andrews J, et al. (2009) Study of 962 patients indicates progressive muscular atrophy is a form of ALS. *Neurology* 73: 1686–1692.
- Wingo TS, Cutler DJ, Yarab N, Kelly CM, Glass JD (2011) The heritability of amyotrophic lateral sclerosis in a clinically ascertained United States research registry. *PLoS One* 6: e27985.
- DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, et al. (2011) Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 72: 245–256.
- Renton AE, Majounie E, Waite A, Simon-Sanchez J, Rollinson S, et al. (2011) A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 72: 257–268.
- Andersen PM, Al-Chalabi A (2011) Clinical genetics of amyotrophic lateral sclerosis: what do we really know? *Nat Rev Neurol* 7: 603–615.
- Johnson JO, Mandrioli J, Benatar M, Abramzon Y, Van Deerlin VM, et al. (2010) Exome sequencing reveals VCP mutations as a cause of familial ALS. *Neuron* 68: 857–864.
- Deng HX, Chen W, Hong ST, Boycott KM, Gorrie GH, et al. (2011) Mutations in UBQLN2 cause dominant juvenile and adult-onset ALS and ALS/dementia. *Nature* 477: 211–215.
- Fecto F, Yan J, Vemula SP, Liu E, Yang Y, et al. (2011) SQSTM1 mutations in familial and sporadic amyotrophic lateral sclerosis. *Arch Neurol* 68: 1440–1446.
- Wu CH, Fallini C, Ticozzi N, Keagle PJ, Sapp PC, et al. (2012) Mutations in the profilin 1 gene cause familial amyotrophic lateral sclerosis. *Nature* 488: 499–503.
- van Rheenen W, van Blitterswijk M, Huisman MH, Vlam L, van Doormaal PT, et al. (2012) Hexanucleotide repeat expansions in C9ORF72 in the spectrum of motor neuron diseases. *Neurology* 79: 878–882.
- van Blitterswijk M, van Es MA, Hennekam EA, Dooijes D, van Rheenen W, et al. (2012) Evidence for an oligogenic basis of amyotrophic lateral sclerosis. *Hum Mol Genet* 21: 3776–3784.
- Parkinson N, Ince PG, Smith MO, Highley R, Skibinski G, et al. (2006) ALS phenotypes with mutations in CHMP2B (charged multivesicular body protein 2B). *Neurology* 67: 1074–1077.
- Cox LE, Ferraiuolo L, Goodall EF, Heath PR, Higginbottom A, et al. (2010) Mutations in CHMP2B in lower motor neuron predominant amyotrophic lateral sclerosis (ALS). *PLoS One* 5: e9872.
- Vlam L, Schelhaas HJ, van Blitterswijk M, van Vught PW, de Visser M, et al. (2012) Mutations in the TRPV4 Gene Are Not Associated With Sporadic Progressive Muscular Atrophy. *Arch Neurol* 69: 790–791.
- van Es MA, Dahlberg C, Birve A, Veldink JH, van den Berg LH, et al. (2010) Large-scale SOD1 mutation screening provides evidence for genetic heterogeneity in amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* 81: 562–566.
- van Es MA, Schelhaas HJ, van Vught PW, Ticozzi N, Andersen PM, et al. (2011) Angiogenin variants in Parkinson disease and amyotrophic lateral sclerosis. *Ann Neurol* 70: 964–973.
- Groen EJ, van Es MA, van Vught PW, Spliet WG, van Engelen-Lee J, et al. (2010) FUS mutations in familial amyotrophic lateral sclerosis in the Netherlands. *Arch Neurol* 67: 224–230.
- Huisman MH, de Jong SW, van Doormaal PT, Weinreich SS, Schelhaas HJ, et al. (2011) Population based epidemiology of amyotrophic lateral sclerosis using capture-recapture methodology. *J Neurol Neurosurg Psychiatry* 82: 1165–1170.
- Brooks BR, Miller RG, Swash M, Munsat TL (2000) El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Motor Neuron Disord* 1: 293–299.
- Skibinski G, Parkinson NJ, Brown JM, Chakrabarti L, Lloyd SL, et al. (2005) Mutations in the endosomal ESCRTIII-complex subunit CHMP2B in frontotemporal dementia. *Nat Genet* 37: 806–808.

**Figure S2 Pedigree of two PMA patients with p.N352S mutations in TARDBP.** (DOC)

## Author Contributions

Conceived and designed the experiments: MvB LV MAvE WLvdP JHV LHvdB. Performed the experiments: MvB LV EAMH DD HJS AJvdK MdV JHV LHvdB. Analyzed the data: MvB LV MAvE WLvdP EAMH DD JHV LHvdB. Contributed reagents/materials/analysis tools: MvB LV MAvE WLvdP EAMH DD HJS AJvdK MdV JHV LHvdB. Wrote the paper: MvB. Revising manuscript for content: MvB LV MAvE WLvdP EAMH DD HJS AJvdK MdV JHV LHvdB. Statistical analysis: MvB LV JHV. Study supervision or coordination: JHV LHvdB.

- Nickerson DA, Tobe VO, Taylor SL (1997) PolyPhred: automating the detection and genotyping of single nucleotide substitutions using fluorescence-based resequencing. *Nucleic Acids Res* 25: 2745–2751.
- Horii M, Shibata H, Kobayashi R, Katoh K, Yorikawa C, et al. (2006) CHMP7, a novel ESCRT-III-related protein, associates with CHMP4b and functions in the endosomal sorting pathway. *Biochem J* 400: 23–32.
- Ghanim M, Guillot-Noel L, Pasquier F, Jornea L, Deramecourt V, et al. (2010) CHMP2B mutations are rare in French families with frontotemporal lobar degeneration. *J Neurol* 257: 2032–2036.
- Traynor BJ, Codd MB, Corr B, Forde C, Frost E, et al. (2000) Clinical features of amyotrophic lateral sclerosis according to the El Escorial and Airlie House diagnostic criteria: A population-based study. *Arch Neurol* 57: 1171–1176.
- Swash M (2012) Why are upper motor neuron signs difficult to elicit in amyotrophic lateral sclerosis? *J Neurol Neurosurg Psychiatry* 83: 659–662.
- Cudkovic ME, McKenna-Yasek D, Chen C, Hedley-Whyte ET, Brown RH, Jr. (1998) Limited corticospinal tract involvement in amyotrophic lateral sclerosis subjects with the A4V mutation in the copper/zinc superoxide dismutase gene. *Ann Neurol* 43: 703–710.
- Ince PG, Evans J, Knopp M, Forster G, Hamdalla HH, et al. (2003) Corticospinal tract degeneration in the progressive muscular atrophy variant of ALS. *Neurology* 60: 1252–1258.
- Van Langenhove T, van der Zee J, Slegers K, Engelborghs S, Vandenberghe R, et al. (2010) Genetic contribution of FUS to frontotemporal lobar degeneration. *Neurology* 74: 366–371.
- Blair IP, Williams KL, Warraich ST, Durnall JC, Thoeng AD, et al. (2010) FUS mutations in amyotrophic lateral sclerosis: clinical, pathological, neurophysiological and genetic analysis. *J Neurol Neurosurg Psychiatry* 81: 639–645.
- Borroni B, Archetti S, Del Bo R, Papetti A, Buratti E, et al. (2010) TARDBP mutations in frontotemporal lobar degeneration: frequency, clinical features, and disease course. *Rejuvenation Res* 13: 509–517.
- Borroni B, Bonvicini C, Alberici A, Buratti E, Agosti C, et al. (2009) Mutation within TARDBP leads to frontotemporal dementia without motor neuron disease. *Hum Mutat* 30: E974–983.
- Ticozzi N, Silani V, LeClerc AL, Keagle P, Gellera C, et al. (2009) Analysis of FUS gene mutation in familial amyotrophic lateral sclerosis within an Italian cohort. *Neurology* 73: 1180–1185.
- Benajiba L, Le Ber I, Camuzat A, Lacoste M, Thomas-Anterior C, et al. (2009) TARDBP mutations in motoneuron disease with frontotemporal lobar degeneration. *Ann Neurol* 65: 470–473.
- Broustal O, Camuzat A, Guillot-Noel L, Guy N, Millecamps S, et al. (2010) FUS mutations in frontotemporal lobar degeneration with amyotrophic lateral sclerosis. *J Alzheimers Dis* 22: 765–769.
- Mackenzie IR, Rademakers R, Neumann M (2010) TDP-43 and FUS in amyotrophic lateral sclerosis and frontotemporal dementia. *Lancet Neurol* 9: 995–1007.
- Yan J, Deng HX, Siddique N, Fecto F, Chen W, et al. (2010) Frameshift and novel mutations in FUS in familial amyotrophic lateral sclerosis and ALS/dementia. *Neurology* 75: 807–814.
- van Es MA, Diekstra FP, Veldink JH, Baas F, Bourque PR, et al. (2009) A case of ALS-FTD in a large FALS pedigree with a K17I ANG mutation. *Neurology* 72: 287–288.
- Quadri M, Cossu G, Saddi V, Simons EJ, Murgia D, et al. (2011) Broadening the phenotype of TARDBP mutations: the TARDBP Ala382Thr mutation and Parkinson's disease in Sardinia. *Neurogenetics* 12: 203–209.
- Lopate G, Baloh RH, Al-Lozi MT, Miller TM, Fernandes Filho JA, et al. (2010) Familial ALS with extreme phenotypic variability due to the I113T SOD1 mutation. *Amyotroph Lateral Scler* 11: 232–236.
- Borghero G, Floris G, Cannas A, Marrosu MG, Murru MR, et al. (2011) A patient carrying a homozygous p.A382T TARDBP missense mutation shows a syndrome including ALS, extrapyramidal symptoms, and FTD. *Neurobiol Aging* 32: 2327 e2321–2325.
- Kovacs GG, Murrell JR, Horvath S, Haraszti L, Majtenyi K, et al. (2009) TARDBP variation associated with frontotemporal dementia, supranuclear gaze palsy, and chorea. *Mov Disord* 24: 1843–1847.

44. Praline J, Vourc'h P, Guennoc AM, Veyrat-Durebex C, Corcia P (2012) Co-occurrence of progressive anarthria with an S393L TARDBP mutation and ALS within a family. *Amyotroph Lateral Scler* 13: 155–157.
45. Camdessanche JP, Belzil VV, Jousserand G, Rouleau GA, Creac'h C, et al. (2011) Sensory and motor neuropathy in a patient with the A382P TDP-43 mutation. *Orphanet J Rare Dis* 6: 4.
46. Suzuki M, Irie T, Watanabe T, Mikami H, Yamazaki T, et al. (2008) Familial amyotrophic lateral sclerosis with Gly93Ser mutation in Cu/Zn superoxide dismutase: a clinical and neuropathological study. *J Neurol Sci* 268: 140–144.
47. Cervenakova L, Protas II, Hirano A, Votjakov VI, Nedzved MK, et al. (2000) Progressive muscular atrophy variant of familial amyotrophic lateral sclerosis (PMA/ALS). *J Neurol Sci* 177: 124–130.
48. Restagno G, Lombardo F, Sbaiz L, Mari C, Gellera C, et al. (2008) The rare G93D mutation causes a slowly progressing lower motor neuron disease. *Amyotroph Lateral Scler* 9: 35–39.
49. Gellera C, Colombrita C, Ticozzi N, Castellotti B, Bragato C, et al. (2008) Identification of new ANG gene mutations in a large cohort of Italian patients with amyotrophic lateral sclerosis. *Neurogenetics* 9: 33–40.
50. Wu D, Yu W, Kishikawa H, Folkert RD, Iafate AJ, et al. (2007) Angiogenin loss-of-function mutations in amyotrophic lateral sclerosis. *Ann Neurol* 62: 609–617.
51. Millicamps S, Boillec S, Le Ber I, Seilhean D, Teyssou E, et al. (2012) Phenotype difference between ALS patients with expanded repeats in C9ORF72 and patients with mutations in other ALS-related genes. *J Med Genet* 49: 258–263.
52. Stewart H, Rutherford NJ, Briemberg H, Krieger C, Cashman N, et al. (2012) Clinical and pathological features of amyotrophic lateral sclerosis caused by mutation in the C9ORF72 gene on chromosome 9p. *Acta Neuropathol* 123: 409–417.
53. Bretschneider J, Van Deerlin VM, Robinson JL, Kwong L, Lee EB, et al. (2012) Pattern of ubiquilin pathology in ALS and FTL indicates presence of C9ORF72 hexanucleotide expansion. *Acta Neuropathol* 123: 825–839.
54. Majounie E, Renton AE, Mok K, Doppler EG, Waite A, et al. (2012) Frequency of the C9orf72 hexanucleotide repeat expansion in patients with amyotrophic lateral sclerosis and frontotemporal dementia: a cross-sectional study. *Lancet Neurol* 11: 323–330.
55. Robberecht W, Aguirre T, Van den Bosch L, Tilkin P, Cassiman JJ, et al. (1996) D90A heterozygosity in the SOD1 gene is associated with familial and apparently sporadic amyotrophic lateral sclerosis. *Neurology* 47: 1336–1339.
56. Andersen PM, Nilsson P, Ala-Hurula V, Keranen ML, Tarvainen I, et al. (1995) Amyotrophic lateral sclerosis associated with homozygosity for an Asp90Ala mutation in CuZn-superoxide dismutase. *Nat Genet* 10: 61–66.
57. Valdmans PN, Rouleau GA (2008) Genetics of familial amyotrophic lateral sclerosis. *Neurology* 70: 144–152.
58. Eisen A, Mezei MM, Stewart HG, Fabros M, Gibson G, et al. (2008) SOD1 gene mutations in ALS patients from British Columbia, Canada: clinical features, neurophysiology and ethical issues in management. *Amyotroph Lateral Scler* 9: 108–119.
59. Banci L, Bertini I, Boca M, Giroto S, Martinelli M, et al. (2008) SOD1 and amyotrophic lateral sclerosis: mutations and oligomerization. *PLoS One* 3: e1677.
60. Sebastia J, Kieran D, Breen B, King MA, Nettelband DF, et al. (2009) Angiogenin protects motoneurons against hypoxic injury. *Cell Death Differ* 16: 1238–1247.
61. Padhi AK, Kumar H, Vasaikar SV, Jayaram B, Gomes J (2012) Mechanisms of loss of functions of human angiogenin variants implicated in amyotrophic lateral sclerosis. *PLoS One* 7: e32479.
62. Millicamps S, Salachas F, Cazeneuve C, Gordon P, Bricka B, et al. (2010) SOD1, ANG, VAPB, TARDBP, and FUS mutations in familial amyotrophic lateral sclerosis: genotype-phenotype correlations. *J Med Genet* 47: 554–560.
63. Kwiatkowski TJ, Jr., Bosco DA, Leclerc AL, Tamrazian E, Vanderburg CR, et al. (2009) Mutations in the FUS/TLS gene on chromosome 16 cause familial amyotrophic lateral sclerosis. *Science* 323: 1205–1208.
64. Vance C, Rogelj B, Hortobagyi T, De Vos KJ, Nishimura AL, et al. (2009) Mutations in FUS, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6. *Science* 323: 1208–1211.
65. Kuhnlein P, Sperfeld AD, Vanmassenhove B, Van Deerlin V, Lee VM, et al. (2008) Two German kindreds with familial amyotrophic lateral sclerosis due to TARDBP mutations. *Arch Neurol* 65: 1185–1189.
66. Kamada M, Maruyama H, Tanaka E, Morino H, Wate R, et al. (2009) Screening for TARDBP mutations in Japanese familial amyotrophic lateral sclerosis. *J Neurol Sci* 284: 69–71.
67. Iida A, Kamei T, Sano M, Oshima S, Tokuda T, et al. (2012) Large-scale screening of TARDBP mutation in amyotrophic lateral sclerosis in Japanese. *Neurobiol Aging* 33: 786–790.