

Article 25fa pilot End User Agreement

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with explicit consent by the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available for no consideration following a reasonable period of time after the work was first published, provided that clear reference is made to the source of the first publication of the work.

This publication is distributed under The Association of Universities in the Netherlands (VSNU) 'Article 25fa implementation' pilot project. In this pilot research outputs of researchers employed by Dutch Universities that comply with the legal requirements of Article 25fa of the Dutch Copyright Act are distributed online and free of cost or other barriers in institutional repositories. Research outputs are distributed six months after their first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and/or copyrights owner(s) of this work. Any use of the publication other than authorised under this licence or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please contact the Library through email: copyright@ubn.ru.nl, or send a letter to:

University Library
Radboud University
Copyright Information Point
PO Box 9100
6500 HA Nijmegen

You will be contacted as soon as possible.

Normal Range of Emphysema and Air Trapping on CT in Young Men

Onno M. Mets¹
 Robert A. van Hulst^{2,3}
 Colin Jacobs^{4,5}
 Bram van Ginneken^{5,6}
 Pim A. de Jong¹

Keywords: air trapping, CT, healthy subjects, pulmonary emphysema, quantitative assessment

DOI:10.2214/AJR.11.7808

Received August 30, 2011; accepted after revision November 8, 2011.

¹Department of Radiology, University Medical Center Utrecht, Heidelberglaan 100, Postbus 85500, 3508GA Utrecht, The Netherlands. Address correspondence to O. M. Mets (o.m.mets@umcutrecht.nl).

²Diving Medical Center, Royal Netherlands Navy, Den Helder, The Netherlands.

³Department of Anaesthesiology, Laboratory of Experimental Intensive Care and Anaesthesiology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

⁴Fraunhofer MEVIS, Bremen, Germany.

⁵Diagnostic Image Analysis Group, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands.

⁶Image Sciences Institute, University Medical Center Utrecht, Utrecht, The Netherlands.

AJR 2012; 199:336–340

0361–803X/12/1992–336

© American Roentgen Ray Society

OBJECTIVE. The purpose of our study was to assess the normal range of CT measures of emphysema and air trapping in young men with normal lung function.

MATERIALS AND METHODS. A cohort of 70 young men with high-normal spirometry and body plethysmography underwent paired inspiratory and expiratory CT. Visual and quantitative scores of emphysema and air trapping were obtained. On CT, emphysema was defined as the 15th percentile of the attenuation curve (Perc₁₅), and as the percentage of inspiratory voxels below –950 (IN_{–950}) and below –960 (IN_{–960}) HU. On CT, air trapping was defined as the expiratory-to-inspiratory ratio of mean lung density (EI-ratio_{MLD}), and the percentage of voxels below –856 HU in expiration (EXP_{–856}). Means, medians, and upper limits of normal (ULN) are presented for the total population and for smokers and nonsmokers separately.

RESULTS. The mean age (± SD) of the subjects was 36.1 ± 9.3 years. Smoking history was limited (range, 0–11 pack-years). Spirometry was high normal, ranging from 113% to 160% of predicted for vital capacity (VC), and from 104% to 140% of predicted for forced expiratory volume in 1 second (FEV₁). The ULN was 2.73% for IN_{–950}, 0.87% for IN_{–960}, –936 HU for Perc₁₅, 89.0% for EI-ratio_{MLD}, and 17.2% for EXP_{–856}. Visual CT scores showed minimal emphysema in eight (11%), > 5 lobules of air trapping in five (7%), and segmental air trapping in three (4%) subjects. CT measures were similar for never- and ever-smokers.

CONCLUSION. We report the normal range of CT values for young male subjects with normal lung function, which is important to define pulmonary disease.

Visual and quantitative CT are increasingly used in research and clinical practice—including young individuals—and it is important to understand the range of normality to diagnose disease [1]. Quantitative CT has matured since it was introduced two decades ago [2], and it is commonly used in the assessment of emphysema and air trapping [3–5]. However, air trapping and emphysema can also be seen in nonsmokers and individuals without disease [6–9].

Furthermore, a limited bilateral subpleural reticular pattern, cystic airspaces, and increased bronchial wall thickness have been described as related to increasing age and independent of smoking history [10]. Parenchymal changes are also present in never-smoking men [11]. All these CT features may be part of the normal spectrum in smokers and lung aging. The normal spectrum of CT features in young men is not well known, and we had the opportunity to retrospective-

ly analyze paired inspiratory and expiratory volumetric CT scans obtained in a population of young male subjects with normal lung function. Therefore, the purpose of this study was to assess the range of visual and commonly used quantitative CT measurements of emphysema and air trapping in a population of young men.

Materials and Methods

This study retrospectively evaluated clinically obtained CT data and was approved by the local ethical review board of the University Medical Center Utrecht. Written informed consent was waived.

Study Population

Recently, the Diving Medical Center (DMC) of the Royal Netherlands Navy, Ministry of Defense in The Netherlands treated six pulmonary barotraumas in professional military divers and submarine personnel. These subjects shared high-normal spirometric and body plethysmographic values, together with abnormalities found on chest CT (i.e., air trapping,

CT of Emphysema and Air Trapping

blebs, and bullae). This initiated the clinical protocol as part of the annual diving examination, decided by the national military board, to evaluate all military divers and submariners with high-normal lung function values with inspiratory and expiratory chest CT to identify possible lung abnormalities and possibly prevent future pulmonary barotraumas.

For this descriptive study, we retrospectively collected the first 77 consecutive subjects who had been referred to our hospital by the DMC. Clinical examinations (i.e., CT and pulmonary function tests) were postponed when acute illness was present, and all investigations were performed in clinically stable subjects. The inclusion criteria for our analysis were an available volumetric inspiratory and expiratory chest CT, pulmonary function test results, demographic patient characteristics, and smoking history. Two subjects were excluded because of a failed expiratory CT. Another five subjects were excluded because of lung segmentation errors (described later). The final study population consisted of 70 subjects.

CT Protocol

Volumetric paired inspiratory and expiratory CT was performed in all subjects. CT examinations were performed using the same scanner (Brilliance 16P, Philips Healthcare) in a single center. Settings were 120 kVp at 130 mAs for inspiratory CT and 90 kVp at 20 mAs for expiratory CT in all subjects. Images were reconstructed with a slice thickness of 1 mm at 0.7-mm increments using a sharp reconstruction kernel (L- and E-filters, Philips Healthcare, for inspiratory and expiratory CT, respectively).

Quantitative CT Analysis

Quantitative measures were obtained after fully automatic segmentation of the lungs [12]. The segmentation process was visually checked in all subjects. As mentioned earlier, five subjects were excluded because of errors in the lung segmentation. Quantitative CT measures of pulmonary emphysema and air trapping were calculated in the segmented lung volumes. CT emphysema was defined as the percentage of voxels below -950 HU in inspiration (IN_{-950}) [13, 14], the percentage of voxels below -960 HU in inspiration (IN_{-960}) [15], and the 15th percentile of the attenuation curve in inspiration ($Perc_{15}$) [16, 17]. CT air trapping was defined as the expiratory-to-inspiratory ratio of mean lung density ($EI\text{-ratio}_{MLD}$) [18, 19] and the percentage of voxels below -856 HU in expiration (EXP_{-856}) [20]. $Perc_{15}$ is expressed in HU, all other quantitative measurements were expressed as a percentage. Calculations were performed after application of a noise reduction filter [21] to decrease the influence of image noise on the quantitative CT measurements.

Visual Assessment

All CT examinations were scored in consensus for the presence and extent of emphysema and air trapping by two observers with 10 and 2 years of experience in chest radiology. Emphysema was defined as focal regions of low attenuation [22], and scored as the percentage of lung parenchyma involved. Air trapping was defined as sharply demarcated areas of lung tissue with a lower than normal increase in attenuation and lack of volume reduction after expiration [22]. Air trapping extent was scored as the absolute number of secondary pulmonary lobules, segments, or lobes involved.

Pulmonary Function Testing

Pulmonary function was measured by qualified respiratory technicians using the Vmax Encore pulmonary function testing system (Cardinal Health), and performed according to the European Respiratory Society guidelines [23]. Calibration of the Vmax Legacy was performed according to the manufacturer's guidelines before every measurement. Reference values were calculated using validated prediction equations [24]. Vital capacity (VC), forced expiratory flow in 1 second (FEV_1), ratio of FEV_1 over forced vital capacity (FEV_1/FVC), forced expiratory flow at 50%

of expiration (FEF_{50}), total lung capacity (TLC), and ratio of residual volume over TLC (RV/TLC) were measured. VC, FEV_1 , FEF_{50} , and TLC were presented as percentage of predicted values and FEV_1/FVC and RV/TLC as percentages.

Data Analysis

The mean \pm SD, median (25th–75th percentile), and 95th percentile (i.e., upper limit of normal) of the quantitative CT measures were calculated for the total study population as well as for smokers and never-smokers separately. The difference in visual emphysema presence between smokers and never-smokers was tested with the Fisher exact test. The difference in visual air trapping presence between smokers and never-smokers was tested with the chi-square test. The difference in extent of air trapping according to smoking status was tested with the Fisher exact test. A Mann-Whitney U test was used to test for differences in age between subjects with and without CT abnormalities. The association between smoking status and quantitative CT measurements was tested using the Mann-Whitney U test. A p value < 0.05 was considered significant. All statistical analyses were performed using SPSS statistical software version 15.0. Data are presented as mean \pm SD unless indicated otherwise.

TABLE I: Characteristics of the 70 Male Divers and Submariners

Parameter	Value
Age (y)	36.1 \pm 9.3
Length (cm)	183 \pm 6.6
BMI (kg/m ²)	26.2 \pm 2.0
Smoking status	
Never-smokers	47 (67)
Smokers	23 (33)
Pack-years	
Never-smokers	
Smokers ^a	4.4 (interquartile range, 2.2–8.3)
Days between PFT and CT ^a	14 (interquartile range, 7–30)
Pulmonary function	
VC (% predicted)	129 \pm 9
FEV_1 (% predicted)	125 \pm 8
FEV_1/FVC %	76.6 \pm 5.6
FEF_{50} (% predicted)	105 \pm 20
TLC (% predicted)	118 \pm 10
RV/TLC (%)	21 \pm 4

Note—Data are mean \pm SD. Except where indicated otherwise, data in parentheses are percentages. BMI = body mass index, PFT = pulmonary function test, VC = vital capacity; FEV_1 = forced expiratory flow in 1 second, FEV_1/FVC = ratio of FEV_1 to forced vital capacity, FEF_{50} = forced expiratory flow at 50% of expiration, TLC = total lung capacity, RV/TLC ratio of residual volume to total lung capacity.

^aMedian (25th–75th percentile).




Fig. 1—Inspiratory CT scan of chest shows bilateral paraseptal emphysematous lesions in lung apices (*arrowheads*) seen as focal regions of low attenuation in 35-year-old man who is current smoker with 4.4 pack-years of smoking history.




Fig. 2—Expiratory CT image of chest shows lobular air trapping in right upper lobe (*arrow*) in 45-year-old man who is never-smoker.

Results

Study Population

Our study population contained 70 male subjects, with a mean (\pm SD) age of 36.1 ± 9.3 years (range, 19–53 years). Forty-seven (67%) were never-smokers, and 23 (33%) were current or ex-smokers, with a median of 4.4 pack-years (range, 0.2–11.0 pack-years). Study population characteristics and pulmonary function data are presented in Table 1.


Visual Assessment

Eight subjects (11%) had a few lobules with minimal centrilobular or paraseptal emphysema; less than 1% of total lung parenchyma was involved in all (Fig. 1). The prevalence of emphysema was similar between smokers and never-smokers ($p = 0.42$). Air trapping was detected in 56 subjects (80%), of whom 55 (98%) showed lobular air trapping (Fig. 2). Three subjects showed segmental air trapping (Fig. 3), which was additional to lobular air trapping in two. The median number of lobules involved was two, with 14 subjects showing more than three lobules and five subjects showing more than five lobules. The presence and amount of lobular air trapping were not different between smokers and nonsmokers ($p > 0.26$). Segmental air trapping was only found in smokers ($p = 0.03$). We did not find an association between age and CT abnormalities. The complete results of visual scoring are summarized in Table 2.

Quantitative Analysis

Results for the quantitative assessment of CT emphysema and CT air trapping are summarized in Table 3. The upper limits of normal for CT emphysema are 2.73% for IN_{-950} ,

Fig. 3—Expiratory CT image of chest shows segmental air trapping in right middle lobe (*arrowheads*) in 47-year-old man who is former smoker with 3.0 pack-years of smoking history.



0.87% for N_{-960} , and -936 HU for $Perc_{15}$. The upper limits of normal for CT air trapping are 89.0% for $EI\text{-ratio}_{MLD}$ or 17.2% for EXP_{-856} . Results for quantitative CT measurements were not significantly different according to smoking status. A possible pattern of higher lung density values in smokers was seen, although this did not reach statistical significance.

Discussion

In this study, we present the range of visual and quantitative CT emphysema and CT air trapping measurements in a population of young men. Because little is known regarding the normal range of imaging in such a population, reporting the range of CT features in these subjects is important. Knowledge of the results of visual and quantitative CT measures may be useful in establishing the normal spectrum and help in the detection of disease at an early stage.

Quantitative CT emphysema and air trapping are surrogate measures of disease. On the basis of the correlation between disease parameters and lung density, a continuous radiologic measure is calculated. Therefore, there is not a strict binomial distribution of presence and absence. Accordingly, we showed that in a healthy population a low percentage of lung voxels show attenuation below the CT emphysema threshold. The presence of disease can be regarded as a value beyond the upper limit of normal in the healthy population. However, it is important to note that densitometry values can be affected by biologic and technical factors—for example, interstitial lung disease, contrast medium injection, reconstruction algorithm, CT scanner, and radiation dose. Ideally, with current technology, quantitative reference values should be obtained for each CT protocol individually.

CT of Emphysema and Air Trapping

TABLE 2: Visual Scoring for Emphysema and Air Trapping in Young Male Subjects With Normal Spirometry

Parameter	All (n = 70)	Never-Smokers (n = 47)	Smokers (n = 23)	p
CT emphysema				
Extent < 1%	8 (11)	4 (9)	4 (17)	NS
CT air trapping				
Present	56 (80)	37 (79)	19 (83)	NS
Lobular air trapping	55	37	18	NS
Lobules involved ^a	2 (interquartile range, 1–4)	2 (interquartile range, 1–4)	1 (interquartile range, 1–3)	NS
Segmental air trapping	3	0	3	0.03

Note—Except where indicated otherwise, data in parentheses are percentages. NS = not significant.

^aMedian (25th–75th percentile).

Regarding quantitative CT measures, no studies were found that assessed the range of quantitative CT air trapping measures in healthy men, and few studies were found on the normal range of quantitative emphysema measures. Marsh et al. [25] evaluated 185 never-smoking subjects without respiratory disease (age 25–75 years) and calculated IN_{-950} in three slices in the upper part of the lungs. They found a median of 1.4% (interquartile range 0.6–2.9) for all three slices, which is somewhat higher than our 0.59% (interquartile range 0.28–1.30). Irion et al. [26] evaluated 30 young nonsmoking subjects (age 19–41 years) who were scanned to rule out lung disease and

calculated IN_{-950} . They found a mean and 95th percentile of 0.19% and 0.35%, respectively. These values are lower than our 0.97% and 2.73%, even though they performed low-dose scanning without noise filtering. Differences in study populations, slice thicknesses, and CT protocols may have contributed to the differences in results. Nevertheless, the observed differences are small, and all studies, including ours, indicate that a low percentage of CT emphysema is found in healthy subjects.

Regarding visual scores, several studies in healthy subjects have been reported; however, none of these studies obtained volumetric CT data. Nevertheless, our finding that lobular air

trapping is visually present to some extent in the majority of healthy subjects is in line with previous results [6–9], and we correspondingly did not find a difference in the presence of lobular air trapping between nonsmokers and smokers [6, 8]. In 250 volunteers, Mastora et al. [6] showed that segmental air trapping is more frequently seen in smokers, a finding that is supported by our data. One finding we could not replicate was a significant increase in air trapping extent with increasing smoking history [6, 7, 9]. This difference may be explained by the very low smoking history of our subjects, which is different from previous studies in which smoking history was substantial.

Smoking has to be taken into account when studying chest CT abnormalities. Apart from its association with air trapping and emphysema, smoking has other effects in the lung. It has been shown that areas of ground-glass attenuation and ill-defined micronodules are related to smoking and are common findings in smokers [11, 27, 28]. These inflammatory changes of the lung may raise the lung density and thereby mask pulmonary emphysema, a process that is supported by the fact that smoking cessation leads to a paradoxical decrease in lung density (increase of emphysema) [29]. In the current study, we found slightly higher lung density in smokers, although it did not reach statistical significance. This may well be due to the minimal smoking history and relatively small number of subjects in our study population.

Recently the effect of aging on the lung has received considerable attention, and aging has been shown to induce lung changes. Copley et al. [10] showed significant differences in CT features between younger and older individuals. Those authors showed that bronchial dimensions, air cysts, and limited subpleural reticular patterns were significantly more frequent in subjects more than 75 years old. Vilkren et al. [11] correspondingly showed that bronchial altera-

TABLE 3: Quantitative CT Measures for Emphysema and Air Trapping in Young Male Subjects With Normal Spirometry

Parameter	IN_{-950}	IN_{-960}	$Perc_{15}$	EXP_{-856}	$EI-Ratio_{MLD}$
All (n = 70)					
Mean ± SD	0.97 ± 1.10	0.35 ± 0.33	-919 ± 13	5.12 ± 7.98	76.2 ± 7.3
Median	0.59	0.25	-921	3.22	76.4
25th–75th Percentile	0.28–1.30	0.13–0.51	-913 to -929	0.69–6.23	70.1–81.3
95th Percentile	2.73	0.87	-936	17.2	89.0
Never-smokers (n = 47)					
Mean ± SD	1.11 ± 1.26	0.38 ± 0.36	-921 ± 12	5.86 ± 9.13	76.5 ± 7.4
Median	0.68	0.27	-921	3.57	76.4
25th–75th Percentile	0.28–1.98	0.14–0.57	-915 to -930	0.79–6.25	70.2–81.2
95th Percentile	3.21	1.13	-937	28.2	90.5
Smokers (n = 23)					
Mean ± SD	0.69 ± 0.62	0.27 ± 0.22	-917 ± 13.7	3.62 ± 4.68	75.6 ± 7.0
Median	0.52	0.19	-920	1.12	76.3
25th–75th percentile	0.15–1.13	0.08–0.35	-909 to -928	0.17–5.21	69.9–81.4
95th Percentile	2.31	0.80	-933	17.6	89.1
p^a	0.18	0.24	0.12	0.18	0.65

Note— IN_{-950} = emphysema score below -950 HU in inspiratory scan, IN_{-960} = emphysema score below -960 HU in inspiratory scan, $Perc_{15}$ = emphysema score as 15th percentile of attenuation distribution curve on inspiratory scan, EXP_{-856} = air trapping score below -856 HU in expiratory scan, $EI-Ratio_{MLD}$ = expiration-to-inspiration ratio of mean lung density,

^aComparison between smokers and never-smokers.

tions, septal lines, areas of ground-glass attenuation, and subpleural and parenchymal nodules are to some extent considered normal in men approximately 60 years old and that apart from emphysema none of the CT features was pathognomonic for smoking-induced disease [11].

That we did find emphysema in both smokers and nonsmokers, in contrast to previous reports, is likely due to our volumetric scanning technique. Lesions were often single and all very small. Thus, they can be easily missed when nonvolumetric scanning is applied. Lee et al. [7] showed that, in addition to parenchymal changes, the frequency and extent of air trapping also increase with age. We did not find a relationship between age and CT abnormalities, which is most likely due to the age-distribution of our study population, with more than 75% of the subjects under 45 years old.

The strength of our study is that it provides reference values for several quantitative CT measures in young men and we used a single CT scanner to perform volumetric scanning in both inspiration and expiration. A limitation of our study is that the study population consists of healthy military divers and submariners with high-normal lung function, representing the healthy worker employed by the military. The larger lung volumes may produce lower density values, and therefore our measurements might overestimate, but certainly do not underestimate, the normal limits. Furthermore, inherent to current CT technology, the quantitative measures can differ according to the noise-filtering, data reconstruction, and CT scanner used. Differences in scanning parameters should be carefully ascertained before generalizing quantitative results.

In conclusion, we have shown the normal spectrum for visual and quantitative CT measurements of emphysema and air trapping in a population of young male subjects with normal pulmonary function tests. These data are important to define disease in subjects with corresponding age and sex.

References

- Hansell DM. Thin-section CT of the lungs: the hinterland of normal. *Radiology* 2010; 256:695–711
- Müller NL, Staples CA, Miller RR, Abboud RT. "Density mask": an objective method to quantify emphysema using computed tomography. *Chest* 1988; 94:782–787
- Matsuoka S, Yamashiro T, Washko GR, Kurihara Y, Nakajima Y, Hatabu H. Quantitative CT assessment of chronic obstructive pulmonary disease. *RadioGraphics* 2010; 30:55–66
- Lynch DA, Newell JD. Quantitative imaging of COPD. *J Thorac Imaging* 2009; 24:189–194
- Coxson HO, Rogers RM. Quantitative computed tomography of chronic obstructive pulmonary disease. *Acad Radiol* 2005; 12:1457–1463
- Mastora I, Remy-Jardin M, Sobaszek A, Boulenguez C, Remy J, Edme JL. Thin-section CT finding in 250 volunteers: assessment of the relationship of CT findings with smoking history and pulmonary function test results. *Radiology* 2001; 218:695–702
- Lee KW, Chung SY, Yang I, Lee Y, Ko EY, Park MJ. Correlation of aging and smoking with air trapping at thin-section CT of the lung in asymptomatic subjects. *Radiology* 2000; 214:831–836
- Tanaka N, Matsumoto T, Miura G, et al. Air trapping at CT: high prevalence in asymptomatic subjects with normal pulmonary function. *Radiology* 2003; 227:776–785
- Verschakelen JA, Scheinbaum K, Bogaert J, Demedts M, Lacquet LL, Baert AL. Expiratory CT in cigarette smokers: correlation between areas of decreased lung attenuation, pulmonary function tests and smoking history. *Eur Radiol* 1998; 8:1391–1399
- Copley SJ, Wells AU, Hawtin KE, et al. Lung morphology in the elderly: comparative CT study of subjects over 75 years old versus those under 55 years old. *Radiology* 2009; 251:566–573
- Vikgren J, Boijens M, Andelid K, et al. High-resolution computed tomography in healthy smokers and never-smokers: a 6-year follow-up study of men born in 1933. *Acta Radiol* 2004; 45:44–52
- van Rikxoort EM, de Hoop B, Viergever MA, Prokop M, van Ginneken B. Automatic lung segmentation from thoracic computed tomography scans using a hybrid approach with error detection. *Med Phys* 2009; 36:2934–2947
- Gevenois PA, de Maertelaer V, De Vuyst P, Zanen J, Yernault JC. Comparison of computed density and macroscopic morphometry in pulmonary emphysema. *Am J Respir Crit Care Med* 1995; 152:653–657
- Gevenois PA, De Vuyst P, de Maertelaer V, et al. Comparison of computed density and microscopic morphometry in pulmonary emphysema. *Am J Respir Crit Care Med* 1996; 154:187–192
- Madani A, Zanen J, De Maertelaer V, Gevenois PA. Pulmonary emphysema: objective quantification at multi-detector row CT—comparison with macroscopic and microscopic morphometry. *Radiology* 2006; 238:1036–1043
- Parr DG, Stoel BC, Stolk J, Stockley RA. Validation of computed tomographic lung densitometry for monitoring emphysema in alpha-1-antitrypsin deficiency. *Thorax* 2006; 61:485–490
- Parr DG, Sevenoaks M, Deng C, Stoel BC, Stockley RA. Detection of emphysema progression in alpha-1-antitrypsin deficiency using CT densitometry: methodological advances. *Respir Res* 2008; 9:21
- O'Donnell RA, Peebles C, Ward JA, et al. Relationship between peripheral airway dysfunction, airway obstruction, and neutrophilic inflammation in COPD. *Thorax* 2004; 59:837–842
- Kubo K, Eda S, Yamamoto H, et al. Expiratory and inspiratory chest computed tomography and pulmonary function tests in cigarette smokers. *Eur Respir J* 1999; 13:252–256
- Regan EA, Hokanson JE, Murphy JR, et al. Genetic epidemiology of COPD (COPDGene) study design. *COPD* 2010; 7:32–43
- Schilham AM, van Ginneken B, Gietema H, Prokop M. Local noise weighted filtering for emphysema scoring of low-dose CT images. *IEEE Trans Med Imaging* 2006; 25:451–463
- Hansell DM, Bankier AA, MacMahon H, McLoud TC, Müller NL, Remy J. Fleischner Society: glossary of terms for thoracic imaging. *Radiology* 2008; 246:697–722
- Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J* 2005; 26:319–338
- Quanjer PH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R, Yernault JC. Lung volumes and forced ventilatory flows: Report Working Party Standardization of Lung Function Tests, European Community for Steel and Coal. Official Statement of the European Respiratory Society. *Eur Respir J Suppl* 1993; 16:5–40
- Marsh S, Aldington S, Williams MV, et al. Utility of lung density measurements in the diagnosis of emphysema. *Respir Med* 2007; 101:1512–1520
- Irion KL, Marchiori E, Hochegger B, et al. CT quantification of emphysema in young subjects with no recognizable chest disease. *AJR* 2009; 192:684; [web]W90–W96
- Remy-Jardin M, Remy J, Boulenguez C, Sobaszek A, Edme JL, Furon D. Morphologic effects of cigarette smoking on airways and pulmonary parenchyma in healthy adult volunteers: CT evaluation and correlation with pulmonary function tests. *Radiology* 1993; 186:107–115
- Remy-Jardin M, Edme JL, Boulenguez C, Remy J, Mastora I, Sobaszek A. Longitudinal follow-up study of smoker's lung with thin-section CT in correlation with pulmonary function tests. *Radiology* 2002; 222:261–270
- Shaker SB, Stavngaard T, Laursen LC, Stoel BC, Dirksen A. Rapid fall in lung density following smoking cessation in COPD. *COPD* 2011; 8:2–7