

Reviewing reasons for the decreased CSF Abeta₄₂ concentration in Alzheimer disease

Petra E. Spies¹, Marcel M. Verbeek^{2,3}, Thomas van Groen^{4,5}, Jurgen A.H.R. Claassen¹

¹Department of Geriatric Medicine, Radboud University Nijmegen Medical Center, Donders Institute for Brain, Cognition and Behaviour, and Alzheimer Center Nijmegen, Nijmegen, The Netherlands, ²Department of Neurology, Radboud University Nijmegen Medical Center, Donders Institute for Brain, Cognition and Behaviour, and Alzheimer Center Nijmegen, Nijmegen, The Netherlands, ³Department of Laboratory Medicine, Radboud University Nijmegen Medical Center, Donders Institute for Brain, Cognition and Behaviour, and Alzheimer Center Nijmegen, Nijmegen, The Netherlands, ⁴Department of Cell Biology, Center for Glial Biology, University of Alabama at Birmingham, Birmingham, USA, ⁵Department of Neurobiology, Center for Glial Biology, University of Alabama at Birmingham, Birmingham, USA

TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Abeta₄₂ metabolism under normal conditions
 - 3.1. Production of Abeta₄₂
 - 3.2. Degradation of Abeta₄₂
 - 3.2.1. Proteases
 - 3.2.2. Microglia
 - 3.3. Clearance of Abeta₄₂
 - 3.3.1. Active transport of Abeta from ISF to systemic circulation
 - 3.3.2. Drainage of Abeta with ISF to lymphatics
 - 3.3.3. Transport of Abeta from ISF to CSF
 - 3.3.4. Absorption of Abeta from CSF to systemic circulation
 - 3.3.5. Drainage of Abeta with CSF to lymphatics
4. Why is CSF Abeta₄₂ decreased in Alzheimer disease?
 - 4.1. Is Abeta₄₂ production reduced?
 - 4.2. Is Abeta₄₂ degradation increased?
 - 4.3. Is microglial uptake of Abeta₄₂ increased?
 - 4.4. Is Abeta₄₂ clearance affected?
 - 4.4.1. Is clearance of Abeta₄₂ from ISF increased?
 - 4.4.2. Is transport of Abeta₄₂ from ISF to CSF hampered?
 - 4.4.2.1. Hampered flow of ISF to CSF
 - 4.4.2.2. Perivascular deposition of Abeta
 - 4.4.2.3. Parenchymal aggregation of Abeta
 - 4.4.3. Is CSF Abeta₄₂ decreased because more Abeta₄₂ is cleared from CSF?
5. Perspective
6. Acknowledgements
7. References

1. ABSTRACT

Cerebrospinal fluid (CSF) amyloid beta₄₂ (Abeta₄₂) concentrations are decreased in patients with Alzheimer disease (AD). Consequently, low Abeta₄₂ is considered a positive biomarker for AD. Surprisingly, the mechanisms that underlie the decrease in CSF Abeta₄₂ remain speculative. Better understanding of this biomarker is an essential step to unravel AD pathophysiology and to develop and evaluate treatment. Therefore, we systematically examined the possible reasons for the decreased CSF Abeta₄₂ concentration in AD. Under normal conditions, Abeta₄₂ can be degraded by proteases, taken up by microglia, or cleared from the brain interstitial fluid across the blood brain barrier. Alternatively, it can be transported to the CSF and be cleared from there. Aggregation of Abeta₄₂ appears the most likely cause for the decreased CSF Abeta₄₂ concentration in AD: the aggregated state inhibits Abeta₄₂ from being transported from the ISF to the CSF. Evidence for other possibilities

such as a decreased production of Abeta₄₂, an increased proteolytic breakdown or microglial uptake of Abeta₄₂, or an increased clearance of Abeta₄₂ to the blood, is - at best - scarce or even absent.

2. INTRODUCTION

The characteristic plaques of Alzheimer disease (AD) were discovered by Alois Alzheimer in 1906, but it was not until the 1980s that it was demonstrated that these plaques consist of the amyloid beta protein (Abeta). (1-3) The realization that this protein was the same as the protein accumulating in brains of patients with Down syndrome led to the discovery of the amyloid precursor protein (APP), located on chromosome 21, as the precursor protein of Abeta. (4) Subsequently, pathogenic mutations in the APP gene were discovered in familial AD. (5-8) In the 1990s, Abeta could be measured in cerebrospinal fluid (CSF) for the first time. (9, 10) It was thereafter repeatedly shown that CSF Abeta₄₂ was decreased in AD. (11-14) Nowadays,




Figure 1. The non-amyloidogenic and amyloidogenic pathways of APP processing APP is cleaved by either alpha- or beta-secretase. Cleavage by alpha-secretase (the non-amyloidogenic pathway) generates sAPP-alpha and C83 (left). Cleavage by beta-secretase (the amyloidogenic pathway) generates sAPP-beta and C99 (right). C83 is cleaved by gamma-secretase, generating AICD and p3 (not shown). C99 is also cleaved by gamma-secretase, generating AICD and Abeta (right). (See text for abbreviations.) Republished with permission of the American Society for Clinical Investigation, from reference no. 22; permission conveyed through Copyright Clearance Center, Inc.

it is an accepted biomarker for AD (15, 16) that is frequently used in clinical practice. (17, 18) Surprisingly, the cause of its decrease has not yet been fully elucidated, although several explanations have been offered: changes in Abeta generation or degradation may affect the CSF concentration, or an alteration in the solubility of Abeta42 may diminish clearance of Abeta42 from the interstitial fluid (ISF) to CSF. (11-14) Determination of the CSF Abeta42 concentration in AD may be masked by its interaction with binding proteins, such as apolipoprotein J or E, or 11. E. Matsubara, B. Frangione and J. Ghiso, Characterization of apolipoprotein J-Alzheimer's Aβ interaction. *J. Biol. Chem.* 270 (1995), pp. 7563–7567. View Record in Scopus | Cited By in Scopus increased clearance of Abeta42 from CSF might explain the diminished levels of Abeta42 in the CSF of AD patients. (14)

In spite of all these suggestions, the actual explanation for the decreased CSF Abeta42 in AD remains largely unidentified. Here, we will review the evidence for each of the proposed explanations to create a better understanding of the underlying mechanisms that lead to its recognition as an important AD biomarker.

3. ABETA₄₂ METABOLISM

3.1. Production of Abeta₄₂

Abeta₄₂ is produced by sequential cleavage of the amyloid precursor protein (APP) (Figure 1). APP is a membrane-bound protein whose function remains unclear, although a role in cell adhesion, cell growth and synaptogenesis has been suggested. (19, 20) APP can be cleaved by either alpha- or beta-secretase, and subsequently by gamma-secretase. Whether Abeta is produced from APP depends on which of these enzymes first cleaves APP.

Cleavage of APP by alpha-secretase at the plasma membrane or in the trans-Golgi network generates N-terminal fragment soluble APP-alpha (sAPP-alpha) and C-terminal fragment C83. (21) sAPP-alpha is released into intracellular vesicles or extracellularly. C83 remains membrane-bound and is cleaved by gamma-secretase to produce p3 (Abeta17-40 and Abeta17-42/43) and APP intracellular domain (AICD) (CT57-59). (22-25) The latter is released into the cytoplasm. This pathway is called the non-amyloidogenic pathway since no Abeta40 or Abeta42 is produced.

Decreased CSF Abeta42 in Alzheimer disease

Cleavage of APP by beta-secretase, the amyloidogenic pathway, takes place in the trans-Golgi network and in endosomes (20, 26) and generates N-terminal fragment soluble APP-beta and C-terminal fragment C₉₉. sAPP-beta – just like sAPP-alpha – can be released into intracellular vesicles or extracellularly. C₉₉ remains membrane-bound and is subsequently cleaved by gamma-secretase to produce Abeta₄₂ or Abeta₄₀ and AICD. (22, 23) Cleavage by gamma-secretase takes place in the endoplasmic reticulum, trans-Golgi network, endosomes, and for a small part at the plasma membrane. (26) Other isoforms of Abeta have been identified as well, with heterogeneity either at the C- or N-terminus of the peptide, but for clarity this review will be limited to Abeta₄₂ and Abeta₄₀.

The amyloidogenic and non-amyloidogenic pathways are mutually exclusive. (22) APP molecules that are not cleaved at the cell membrane by alpha-secretase can be internalized by endocytosis and subsequently be cleaved by beta-secretase. (27) Redistribution of APP towards either the cell membrane or the cell interior can therefore influence the proteolytic pathway that APP undergoes. (22)

Abeta₄₂ and Abeta₄₀ are either retained in cells or secreted into the ISF that surrounds the brain cells. (23) The equilibrium between intra- and extracellular Abeta₄₂ and Abeta₄₀ is in part regulated by active transport via the receptor for advanced glycation end products (RAGE), (28) a multiligand receptor of the immunoglobulin family. (29)

3.2. Degradation of Abeta₄₂

3.2.1. Proteases

Several proteases can break down Abeta (both Abeta₄₂ and Abeta₄₀) *in vitro*; *in vivo* evidence stems mostly from animal studies. Insulin-degrading enzyme (IDE) and neprilysin have received most attention. IDE is a zinc-metalloproteinase present in the cytosol, in intracellular membranes and at the cell surface. (25, 30-32) It is secreted by neurons (32) and has also been found in CSF. (25) It can therefore degrade both intracellular and extracellular Abeta. IDE only degrades soluble, monomeric Abeta. (25, 30, 31) The cleavage products are not neurotoxic and are not prone to accumulate in plaques. (25) Neprilysin is an axonal and synaptic membrane-anchored protein with its catalytic site facing the cell exterior (31, 32) and is therefore primarily involved in the breakdown of extracellular Abeta₄₂. It can degrade monomeric as well as oligomeric Abeta. (32)

Other proteases that have been reported to degrade Abeta *in vitro* include matrix metalloproteinase-9, angiotensin converting enzyme, cathepsin D, plasmin and endothelin converting enzyme-1. (32, 33) Abeta aggregates can stimulate tissue-type and urokinase-type plasminogen activator to generate plasmin. Plasmin can degrade both monomeric Abeta and Abeta fibrils *in vitro*, although degradation of fibrils is far less efficient. Its role in degrading Abeta *in vivo* is unclear. (33)

3.2.2. Microglia

Microglia are the macrophages of the brain. They take up soluble Abeta by macropinocytosis and possibly by

binding of Abeta to the low-density lipoprotein receptor-related protein (LRP). (34) Fibrillar forms of Abeta interact with the microglia cell surface and bind to the CD36 receptor expressed by microglia. CD36, a major pattern recognition receptor, mediates the microglial response to Abeta, leading to an intracellular signaling cascade that stimulates phagocytosis. (35) Ablation of microglia in mice led to an increase in soluble Abeta₄₀ and Abeta₄₂, but had no influence on number and size of Abeta plaques. (36) This would support the idea that microglia are inefficient in degrading fibrillar Abeta. (34) However, others have found that upon activation, for example by cerebral ischemia or after vaccination with Abeta specific antibodies, microglia can degrade fibrillar Abeta. (37-39)

3.3. Clearance of Abeta₄₂

Abeta₄₂ and Abeta₄₀ can also be cleared from the extracellular compartment. It can be cleared from the ISF across the blood-brain barrier towards the circulation, or it can be transported from the ISF to the CSF, and be cleared from there. Alternatively, Abeta can be transported with the ISF or CSF to the peripheral lymphatic system (Figure 2).

3.3.1. Active transport of Abeta from ISF to systemic circulation

Both Abeta₄₂ and Abeta₄₀ can be removed from ISF to the blood by crossing the blood-brain barrier. The blood-brain barrier is primarily formed by endothelial cells in brain capillaries that are closely connected via tight junctions which exclude the transfer of many macromolecules from ISF into blood. (40) Astrocytes and pericytes may also play a role in the control of transport across the blood-brain barrier. Abeta₄₂ and Abeta₄₀ are primarily transported across the blood-brain barrier by LRP. (29) LRP is a multiligand lipoprotein receptor that mediates endocytosis of secreted proteins. It provides a rapid transport of brain-derived Abeta into the blood. (41) LRP-mediated transport is more efficient for Abeta₄₀ when compared to Abeta₄₂ (29, 31), probably related to Abeta₄₀'s lower propensity to aggregate. Abeta₄₂ and Abeta₄₀ that is produced in the periphery can enter the brain ISF from the systemic circulation by active transport across the blood-brain barrier via RAGE, which is expressed on the luminal surface of brain vessels. (29)

3.3.2. Drainage of Abeta with ISF to lymphatics

One of the routes that ISF can follow is drainage along axonal tracts and through perivascular spaces. ISF travels along these perivascular spaces towards the surface of the brain and exits at the base of the skull, where it drains to regional lymph nodes in the neck. (42) The idea that Abeta is transported together with ISF along this pathway is supported by the fact that Abeta has been found in the basement membranes of capillary and arterial walls, but was not detectable further downstream in the walls of the carotid arteries. (43)

3.3.3. Transport of Abeta from ISF to CSF

Another route for ISF is towards the CSF, where it constitutes 10-30% to the total CSF production. (44) ISF mixes with CSF at the ventricles and presumably also in the subarachnoid compartment. (40, 44) Whether Abeta




Figure 2. Schematic view of Abeta₄₂ and Abeta₄₀ clearance, Abeta can be Abeta₄₂ or Abeta₄₀. Solid arrows depict active transport of Abeta; dashed arrows depict passive transport of Abeta together with ISF or CSF. BBB = blood brain barrier, BCB = blood CSF barrier at choroid plexuses, AV = arachnoid villi

reaches the CSF via this route is uncertain. The glia limitans and the pia mater separate ISF from CSF in the subarachnoid space, however, the extent to which these layers block the passage of solutes and ISF into CSF is unknown. They may be fully permeable to fluid and small molecules, (29) but on the other hand, the presence of Abeta deposits in the glia limitans in AD suggests that transport of Abeta from ISF to CSF is limited in the subarachnoid compartment. (43)

The concentration of Abeta₄₀ in CSF is about 6-fold greater than the concentration of Abeta₄₂; (45-47) in AD, the decreased concentration of CSF Abeta₄₂ makes the CSF Abeta₄₀ concentration at least 10-fold that of CSF Abeta₄₂. (46-48)

3.3.4. Absorption of Abeta from CSF to systemic circulation

Labeled Abeta injected into CSF of rats was detectable in blood a few minutes after injection. (49) From cell culture research it was inferred that Abeta can be absorbed from CSF at the choroid plexuses. (50) The endothelium of the blood vessels within the choroid plexus is leaky, but the epithelium between blood vessels and ventricular CSF contains tight junctions and forms the blood-CSF barrier (which should not be confused with the blood-brain barrier). Small molecules can enter or leave the brain. (40) Uptake of Abeta peptides at the choroid plexus was rapid and occurred as a non-diffusional, yet unclear uptake process. Efflux of Abeta from CSF to blood was favored over influx from blood into CSF. (50) In addition to the blood-CSF barrier, CSF drains to blood in the subarachnoid compartment, where CSF drains into the

major venous sinuses via arachnoid granulations. Tight junctions in these granulations prevent passage of proteins, (51) and it is therefore unclear whether Abeta can be cleared from CSF to blood in this way.

3.3.5. Drainage of Abeta with CSF to lymphatics

The perineural sheath along the olfactory nerve represents another pathway of CSF drainage. The olfactory nerve fibers enter the nasal mucosa in the roof of the nasal cavity. At this point CSF drains from the perineural subarachnoid space into the extracellular matrix, where it is absorbed by blind-ended lymphatic capillaries, and drained to the regional lymph nodes that serve the nasopharynx. (44, 51) CSF also flows along the perineuronal sheaths of the cranial and spinal nerves to regional lymphatics. (40, 51, 52) It seems plausible that Abeta is cleared together with CSF via this pathway, but studies to confirm this are currently lacking.

4. WHY IS CSF ABETA₄₂ DECREASED IN ALZHEIMER DISEASE?

Several morphological changes that are related to the fate of the Abeta₄₂ peptide have been observed in AD. Upon post-mortem examination, Abeta plaques are found: diffuse plaques that contain only Abeta₄₂, and classic plaques that consist of both Abeta₄₂ and Abeta₄₀. (53, 54) In most AD cases, at least a mild degree of cerebral amyloid angiopathy is found as well, that consists of Abeta deposits in the walls of leptomeningeal and cortical arteries and less frequently in capillaries. (55) In contrast to the plaques, these vascular deposits contain mostly Abeta₄₀. (56) The concentration of both Abeta₄₂ and Abeta₄₀ in the AD brain

Decreased CSF Abeta42 in Alzheimer disease

Table 1. Overview of arguments supporting or contradicting the hypothetical mechanisms that could explain the decreased CSF Abeta₄₂ concentration in AD

Hypothetical mechanism	Supporting arguments	Contradicting arguments
Decreased production of Abeta ₄₂ (4.1.)	- Could be the result of a reduction in viable neurons due to neurodegeneration	- An increased brain Abeta ₄₂ load is found post-mortem - Abeta ₄₀ shares the same production pathway as Abeta ₄₂ but CSF Abeta ₄₀ is not decreased - Disorders with an increased production of Abeta have decreased concentrations of CSF Abeta ₄₂ - The production of Abeta ₄₂ in AD did not differ from controls in a labeling study
Decreased secretion of Abeta ₄₂ from neurons (4.1.)	- Intraneuronal Abeta aggregation has been described in Tg mice	- The typical Abeta plaques of AD are found extracellularly
Increased degradation of Abeta ₄₂ (4.2.)		- An increased brain Abeta ₄₂ load is found post-mortem - Increased degradation would probably affect Abeta ₄₀ as well, yet the CSF Abeta ₄₀ concentration is unchanged - Levels of neprilysin (one of the proteases involved in Abeta degradation) decrease with aging
Increased microglial uptake of Abeta ₄₂ (4.3.)	- Activated microglia are found in the vicinity of Abeta plaques	- An increased brain Abeta ₄₂ load is found post-mortem - Microglia become dysfunctional with aging
Increased clearance of Abeta ₄₂ from ISF to blood (4.4.1.)	- Expression of LRP (which promotes transport of Abeta from ISF to blood across the blood brain barrier) by perivascular cells increased in reaction to Abeta ₄₂	- Perivascular cells expressing LRP degenerated after uptake of Abeta - LRP expression is reduced in AD - No increase in plasma Abeta ₄₂ is found - An increased brain Abeta ₄₂ load is found post-mortem
Hampered transport of Abeta ₄₂ with ISF to CSF (4.4.2.1.)	- Dilated perivascular spaces may indicate obstruction of ISF flow	- Dilated perivascular spaces are often seen in the elderly and their clinical relevance is unclear - Other proteins (including Abeta ₄₀) still reach the CSF
Perivascular deposition of Abeta ₄₂ (4.4.2.2.)	- Abeta deposits are found perivascularly	- Perivascular Abeta deposits contain mostly Abeta ₄₀
Parenchymal aggregation and deposition of Abeta ₄₂ (4.4.2.3.)	- An increased brain Abeta ₄₂ load is found post-mortem - Abeta ₄₂ is more prone to aggregation than Abeta ₄₀	
Increased clearance of Abeta ₄₂ from CSF to the blood (4.4.3.)		- Uptake of Abeta from CSF decreased with aging - No increase in plasma Abeta ₄₂ is found

The numbers behind each mechanism refer to the corresponding section in the text.

is thus increased. (57, 58) However, the CSF concentration of Abeta₄₂ is decreased in AD patients, while the CSF Abeta₄₀ concentration is unaltered. (14, 48) We will focus on reasons why CSF Abeta₄₂ is decreased, while keeping in mind the other changes that occur in AD. Table 1 provides an overview of these reasons and the arguments supporting or contradicting them.

4.1. Is Abeta₄₂ production reduced?

Theoretically, a decreased concentration of Abeta₄₂ in CSF could simply be the result of a decreased production of Abeta₄₂, for example by a reduction in the number of viable neurons that produce Abeta. However, several arguments make this unlikely. First, a decreased production does not logically connect to an increased Abeta₄₂ load in the brain. Second, if the production of Abeta₄₂ were decreased, the production of Abeta₄₀ would probably be decreased as well, since Abeta₄₂ and Abeta₄₀ likely share the same production pathway. This should then lead to a decreased CSF Abeta₄₀ concentration, which, however, is not observed in AD. Third, a decreased concentration of CSF Abeta₄₂ is also found in familial AD and in Down syndrome, disorders in which overproduction of Abeta₄₂ has been clearly demonstrated. (59-62) Decreased Abeta₄₂ thus exists despite overproduction, which implies that factors other than production play a role in causing the decreased CSF concentration.

An alternative theory is that Abeta₄₂ secretion from the cell is decreased, leaving less Abeta₄₂ available in the ISF for transport to the CSF. Although intraneuronal Abeta aggregation has been described in triple transgenic mouse models, (63) Abeta predominantly accumulates extracellularly in human AD. This implies that the secretion of Abeta from the cell to the exterior is not diminished and that a decreased secretion of Abeta₄₂ by the cell is unlikely to be the cause of the decreased CSF Abeta₄₂ concentration.

Supporting these arguments against a decreased production or secretion of Abeta₄₂ is a study showing that the production rate of Abeta₄₂ and Abeta₄₀, as measured in CSF using labeled Abeta₄₂ and Abeta₄₀, did not differ between controls and AD patients. (64)

4.2. Is Abeta₄₂ degradation increased?

It can be hypothesized that an increase in proteolytic breakdown of Abeta₄₂ leads to a decreased CSF Abeta₄₂ concentration. However, an increase in proteolytic breakdown would clear the brain from not only Abeta₄₂, but also from Abeta₄₀ (65) and would preclude formation of plaques. Understandably, research has focused on finding defects in proteolytic degradation of Abeta rather than overactivity of these proteases to explain the elevated cerebral levels of Abeta. (32, 33) It appears that levels of

Decreased CSF Abeta₄₂ in Alzheimer disease

neprilysin decrease with aging (66) and that low levels of neprilysin make certain brain areas more vulnerable to Abeta deposition, (67, 68) suggesting that proteolytic activity is decreased rather than increased.

4.3. Is microglial uptake of Abeta₄₂ increased?

An increased microglial uptake of Abeta₄₂ could be hypothesized to lead to the decreased CSF Abeta₄₂ concentration by leaving less Abeta₄₂ available in the ISF for transport to the CSF. Activated microglia are found in the vicinity of compact Abeta plaques, (69) however, the very presence of these plaques shows that microglial uptake of Abeta is insufficient to lead to a lower-than-normal concentration of Abeta in ISF. On the contrary, it has been suggested that with age, microglia become dysfunctional and less able to clear Abeta₄₂. (34, 70, 71) Thus, it is unlikely that microglial ability to degrade Abeta is increased in AD.

4.4. Is Abeta₄₂ clearance affected?

Changes in the clearance of Abeta from either ISF or CSF (the solid arrows in Figure 1) can influence the Abeta concentration. An increased transport of ISF or CSF, containing Abeta, to the lymphatics will not affect the Abeta concentration.

4.4.1. Is clearance of Abeta₄₂ from ISF increased?

An increased clearance of Abeta₄₂ from ISF to blood across the blood-brain barrier would leave a smaller amount of Abeta₄₂ in the ISF that can be transported to the CSF and could thus explain the decreased CSF Abeta₄₂ concentration. The primary transporter of Abeta across the blood-brain barrier is LRP. Even though LRP favors clearance of Abeta₄₀ over Abeta₄₂, it can be hypothesized that in AD, clearance of Abeta₄₂ is upregulated while clearance of Abeta₄₀ remains the same. In support of this hypothesis, expression of LRP by perivascular cells was increased in response to Abeta₄₂ *in vitro*, while Abeta₄₀ had no such effect; however, uptake of Abeta resulted in degeneration of these perivascular cells. (72) It has also been reported that LRP expression was reduced in AD, while RAGE expression (which transports Abeta from blood to brain) was increased. (29) These changes do not support an increased clearance of Abeta₄₂ from ISF over the blood-brain barrier and could actually lead to a reduced transport of Abeta₄₂ out of the brain and an increased uptake of peripheral Abeta₄₂ into the brain.

Another argument against an increased transport of Abeta₄₂ across the blood-brain barrier is the lack of an increase in plasma Abeta₄₂ concentration. Increased transport of Abeta₄₂ to the blood should result in an increase in the plasma concentration of Abeta₄₂ – assuming that the peripheral breakdown and elimination of Abeta₄₂ are not increased in AD. Reports on the plasma Abeta₄₂ concentration in AD patients are contradictory, but a clear increase has not been shown. (73, 74) Finally, enhanced clearance of Abeta₄₂ from the ISF is incompatible with the increased Abeta load that is found in AD brains. In summary, it cannot be concluded that transport across the blood-brain barrier is increased.

4.4.2. Is transport of Abeta₄₂ from ISF to CSF hampered?

4.4.2.1. Hampered flow of ISF to CSF

ISF flows through the perivascular spaces, mixing with CSF in the subarachnoid compartment at the brain surface and at the ventricles. If Abeta₄₂-containing ISF cannot reach the CSF or if Abeta₄₂ is deposited before ISF reaches the CSF, the Abeta₄₂ concentration in CSF will be decreased. Dilated perivascular spaces may be interpreted as a sign that ISF cannot, or has difficulty to, reach the CSF. (75) However, dilated perivascular spaces are often seen in the elderly and are not AD-specific. (76) Their clinical relevance is unclear. The fact that Abeta₄₀ and other proteins can still reach the CSF in AD argues against an obstruction of transport between ISF and CSF.

4.4.2.2. Perivascular deposition of Abeta

Signs that Abeta₄₂ is deposited before it reaches the CSF are present: Abeta deposits are found in the walls of the cerebral vasculature and in the glia limitans in the subarachnoid compartment. (43, 77, 78) The deposits in the cerebral vasculature consist mostly of Abeta₄₀, and to a lesser extent of Abeta₄₂. (56) In clinically diagnosed cerebral amyloid angiopathy, a decreased CSF Abeta₄₀ concentration was found. (79) This suggests that perivascular Abeta deposition can indeed lead to a decreased CSF Abeta concentration. However, cerebral amyloid angiopathy found in AD is mostly mild to moderate, (55) and the amount of Abeta₄₂ that is deposited in the cerebral vasculature is presumably insufficient to be the sole explanation for the decreased CSF Abeta₄₂ concentration.

4.4.2.3. Parenchymal aggregation of Abeta

Aggregation of Abeta₄₂ into insoluble deposits may be a reason why Abeta₄₂ is decreased in CSF: less soluble Abeta₄₂ is left for transport to the CSF, while the insoluble aggregates cannot be transported from ISF to the CSF. Abeta₄₂ is much more prone to aggregation than Abeta₄₀. (56, 80) Both diffuse and classic plaques contain Abeta₄₂, but only the classic plaques, which are fewer in number, contain Abeta₄₀. (54) Thus, much more Abeta₄₂ than Abeta₄₀ is deposited in AD. Considering that Abeta₄₀ is the most abundant Abeta peptide in CSF, (81) the relative amount that is deposited in AD may be too little to induce a clear decrease in the CSF Abeta₄₀ concentration. In contrast, the relative amount of Abeta₄₂ that is deposited is much larger, which may explain why the CSF Abeta₄₂ concentration is decreased.

4.4.3. Is CSF Abeta₄₂ decreased because more Abeta₄₂ is cleared from CSF?

Uptake of Abeta₄₂ from the CSF was inferred from research in cell culture that used Abeta₄₀ as a model compound for all Abeta species. (50) Theoretically, this process could be upregulated in AD, resulting in more Abeta₄₂ to be transported from the CSF into the blood. Since the CSF Abeta₄₀ concentration is unaltered, this upregulation should then selectively affect Abeta₄₂ and not Abeta₄₀. Research to confirm this is currently lacking. In contrast, research in rats suggests that uptake of Abeta from the CSF decreases with aging. (82) Furthermore, the

Decreased CSF Abeta₄₂ in Alzheimer disease

increase in the plasma concentration of Abeta₄₂ that would be expected if upregulated efflux was the case (and peripheral elimination of Abeta was unaltered), has not been shown convincingly. (73, 74) Therefore it seems unlikely that an increased clearance of Abeta₄₂ from the CSF is the cause of the decreased CSF Abeta₄₂ concentration in AD.

5. PERSPECTIVE

We have reviewed a number of possible mechanisms that could theoretically explain the typical observation of a decreased CSF Abeta₄₂ concentration in AD. It appears very difficult to understand the decrease in CSF Abeta₄₂ found in AD. Based on simple but systematic reasoning, a number of theoretical mechanisms can be formulated, however, most of these do not concur with the premise that there is an increased Abeta load in the AD brain, and that the decreased CSF Abeta₄₂ is accompanied by an unaltered CSF Abeta₄₀ concentration. Furthermore, many of these theoretical mechanisms are contradicted by (often fragmented) findings from research. For example, a decreased production of Abeta₄₂ has not been shown: production rates were comparable between controls and AD patients. Increased proteolytic breakdown of Abeta is unlikely, since available evidence suggests a decrease in proteolytic activity rather than an increase. An increased microglial uptake of Abeta₄₂ is not supported, instead, a decrease in activity with aging is found. Increased clearance of Abeta₄₂ from either ISF or CSF has not been reported; rather, decreases in clearance have been suggested.

What, in our opinion, remains as the most plausible cause of the decreased Abeta₄₂ concentration in CSF is Abeta₄₂ aggregation. Abeta₄₂ is known to aggregate easily and once aggregates have been formed, transport of these aggregates from the ISF to the CSF may be seriously impaired. Abeta₄₀ has different properties and is not as prone to aggregation, which may explain why less Abeta₄₀ is found deposited in the AD brain, and why CSF concentrations of Abeta₄₀ are unaltered.

Until novel mechanistic insights with respect to the different steps in Abeta metabolism are obtained, this remains currently the most satisfying explanation for the observed Abeta changes in AD. Our systematic approach may have provided a framework against which the currently available knowledge from research can be laid out, as pieces of a puzzle. In this way, CSF Abeta₄₂ can be much more than a diagnostic test with an empirical cut-off value for use in clinical settings. Better understanding of this biomarker may be the key to a better understanding of the disease.

6. ACKNOWLEDGEMENTS

Marcel Verbeek is supported by grants from the Internationale Stichting Alzheimer Onderzoek (ISAO, no. 07510), the Netherlands Organisation for Scientific Research (NWO/ZonMW, Vidi program, no. 917.46.331) and the Hersenstichting Nederland (no. 14F06.18).

Jurgen Claassen receives grants from Alzheimer Nederland and has received consulting fees from Novartis and lecture fees from Janssen.

7. REFERENCES

1. P. D. Gorevic, F. Goni, B. Pons-Estel, F. Alvarez, N. S. Peress and B. Frangione: Isolation and partial characterization of neurofibrillary tangles and amyloid plaque core in Alzheimer's disease: immunohistological studies. *J Neuropathol Exp Neurol* 45, 647-664 (1986)
2. C. L. Masters, G. Simms, N. A. Weinman, G. Multhaup, B. L. McDonald and K. Beyreuther: Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc Natl Acad Sci U S A* 82, 4245-4249 (1985)
3. A. Roher, D. Wolfe, M. Palutke and D. KuKuruga: Purification, ultrastructure, and chemical analysis of Alzheimer disease amyloid plaque core protein. *Proc Natl Acad Sci U S A* 83, 2662-2666 (1986)
4. J. Kang, H. G. Lemaire, A. Unterbeck, J. M. Salbaum, C. L. Masters, K. H. Grzeschik, G. Multhaup, K. Beyreuther and B. Muller-Hill: The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* 325, 733-736 (1987)
5. M. C. Chartier-Harlin, F. Crawford, H. Houlden, A. Warren, D. Hughes, L. Fidani, A. Goate, M. Rossor, P. Roques and J. Hardy: Early-onset Alzheimer's disease caused by mutations at codon 717 of the beta-amyloid precursor protein gene. *Nature* 353, 844-846 (1991)
6. A. Goate, M. C. Chartier-Harlin, M. Mullan, J. Brown, F. Crawford, L. Fidani, L. Giuffra, A. Haynes, N. Irving and L. James: Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349, 704-706 (1991)
7. M. Mullan, F. Crawford, K. Axelman, H. Houlden, L. Lilius, B. Winblad and L. Lannfelt: A pathogenic mutation for probable Alzheimer's disease in the APP gene at the N-terminus of beta-amyloid. *Nat Genet* 1, 345-347 (1992)
8. J. Murrell, M. Farlow, B. Ghetti and M. D. Benson: A mutation in the amyloid precursor protein associated with hereditary Alzheimer's disease. *Science* 254, 97-99 (1991)
9. P. Seubert, C. Vigo-Pelfrey, F. Esch, M. Lee, H. Dovey, D. Davis, S. Sinha, M. Schlossmacher, J. Whaley and C. Swindlehurst: Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature* 359, 325-327 (1992)
10. M. Shoji, T. E. Golde, J. Ghiso, T. T. Cheung, S. Estus, L. M. Shaffer, X. D. Cai, D. M. McKay, R. Tintner and B. Frangione: Production of the Alzheimer amyloid beta protein by normal proteolytic processing. *Science* 258, 126-129 (1992)
11. D. Galasko, L. Chang, R. Motter, C. M. Clark, J. Kaye, D. Knopman, R. Thomas, D. Kholodenko, D. Schenk, I.

Decreased CSF Abeta42 in Alzheimer disease

- Lieberburg, B. Miller, R. Green, R. Basherad, L. Kertiles, M. A. Boss and P. Seubert: High cerebrospinal fluid tau and low amyloid beta42 levels in the clinical diagnosis of Alzheimer disease and relation to apolipoprotein E genotype. *Arch Neurol* 55, 937-945 (1998)
12. R. Motter, C. Vigo-Pelfrey, D. Kholodenko, R. Barbour, K. Johnson-Wood, D. Galasko, L. Chang, B. Miller, C. Clark and R. Green: Reduction of beta-amyloid peptide42 in the cerebrospinal fluid of patients with Alzheimer's disease. *Ann Neurol* 38, 643-648 (1995)
13. M. Shoji, E. Matsubara, M. Kanai, M. Watanabe, T. Nakamura, Y. Tomidokoro, M. Shizuka, K. Wakabayashi, Y. Igeta, Y. Ikeda, K. Mizushima, M. Amari, K. Ishiguro, T. Kawarabayashi, Y. Harigaya, K. Okamoto and S. Hirai: Combination assay of CSF tau, A beta 1-40 and A beta 1-42 (43) as a biochemical marker of Alzheimer's disease. *J Neurol Sci* 158, 134-140 (1998)
14. A. Tamaoka, N. Sawamura, T. Fukushima, S. Shoji, E. Matsubara, M. Shoji, S. Hirai, Y. Furiya, R. Endoh and H. Mori: Amyloid beta protein 42 (43) in cerebrospinal fluid of patients with Alzheimer's disease. *J Neurol Sci* 148, 41-45 (1997)
15. B. Dubois, H. H. Feldman, C. Jacova, S. T. DeKosky, P. Barberger-Gateau, J. Cummings, A. Delacourte, D. Galasko, S. Gauthier, G. Jicha, K. Meguro, J. O'Brien, F. Pasquier, P. Robert, M. Rossor, S. Salloway, Y. Stern, P. J. Visser and P. Scheltens: Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS-ADRDA criteria. *Lancet Neurol* 6, 734-746 (2007)
16. G. M. McKhann, D. S. Knopman, H. Chertkow, B. T. Hyman, C. R. Jack, Jr., C. H. Kawas, W. E. Klunk, W. J. Koroshetz, J. J. Manly, R. Mayeux, R. C. Mohs, J. C. Morris, M. N. Rossor, P. Scheltens, M. C. Carrillo, B. Thies, S. Weintraub and C. H. Phelps: The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 7, 263-269 (2011)
17. D. Slats, P. E. Spies, M. J. Sjogren, P. J. Visser, M. M. Verbeek, M. G. Olde Rikkert and P. G. Kehoe: CSF biomarker utilisation and ethical considerations of biomarker assisted diagnosis and research in dementia: perspectives from within the European Alzheimer's Disease Consortium (EADC). *J Neurol Neurosurg Psychiatry* 81, 124-125 (2010)
18. P. E. Spies, D. Slats, I. Ramakers, F. Verhey and M. G. Olde Rikkert: Experiences with cerebrospinal fluid analysis in Dutch memory clinics. *Eur J Neurol* DOI: 10.1111/j.1468-1331.2010.03222.x, (2011)
19. M. P. Marzolo, G. Bu: Lipoprotein receptors and cholesterol in APP trafficking and proteolytic processing, implications for Alzheimer's disease. *Semin Cell Dev Biol* 20, 191-200 (2009)
20. G. Thinakaran, E. H. Koo: Amyloid precursor protein trafficking, processing, and function. *J Biol Chem* 283, 29615-29619 (2008)
21. D. M. Skovronsky, D. B. Moore, M. E. Milla, R. W. Doms and V. M. Lee: Protein kinase C-dependent alpha-secretase competes with beta-secretase for cleavage of amyloid-beta precursor protein in the trans-golgi network. *J Biol Chem* 275, 2568-2575 (2000)
22. S. Gandy: The role of cerebral amyloid beta accumulation in common forms of Alzheimer disease. *J Clin Invest* 115, 1121-1129 (2005)
23. S. Kumar-Singh: Cerebral amyloid angiopathy: pathogenetic mechanisms and link to dense amyloid plaques. *Genes Brain Behav* 7 Suppl 1, 67-82 (2008)
24. Q. X. Li, S. J. Fuller, K. Beyreuther and C. L. Masters: The amyloid precursor protein of Alzheimer disease in human brain and blood. *J Leukoc Biol* 66, 567-574 (1999)
25. Y. H. Suh, F. Checler: Amyloid precursor protein, presenilins, and alpha-synuclein: molecular pathogenesis and pharmacological applications in Alzheimer's disease. *Pharmacol Rev* 54, 469-525 (2002)
26. K. S. Vetrivel, G. Thinakaran: Amyloidogenic processing of beta-amyloid precursor protein in intracellular compartments. *Neurology* 66, S69-S73 (2006)
27. F. Kamenetz, T. Tomita, H. Hsieh, G. Seabrook, D. Borchelt, T. Iwatsubo, S. Sisodia and R. Malinow: APP processing and synaptic function. *Neuron* 37, 925-937 (2003)
28. A. Rauk: The chemistry of Alzheimer's disease. *Chem Soc Rev* 38, 2698-2715 (2009)
29. R. Deane, R. D. Bell, A. Sagare and B. V. Zlokovic: Clearance of amyloid-beta peptide across the blood-brain barrier: implication for therapies in Alzheimer's disease. *CNS Neurol Disord Drug Targets* 8, 16-30 (2009)
30. W. Farris, S. Mansourian, Y. Chang, L. Lindsley, E. A. Eckman, M. P. Frosch, C. B. Eckman, R. E. Tanzi, D. J. Selkoe and S. Guenette: Insulin-degrading enzyme regulates the levels of insulin, amyloid beta-protein, and the beta-amyloid precursor protein intracellular domain in vivo. *Proc Natl Acad Sci U S A* 100, 4162-4167 (2003)
31. R. E. Tanzi, R. D. Moir and S. L. Wagner: Clearance of Alzheimer's Abeta peptide: the many roads to perdition. *Neuron* 43, 605-608 (2004)
32. A. J. Turner, L. Fisk and N. N. Nalivaeva: Targeting amyloid-degrading enzymes as therapeutic strategies in neurodegeneration. *Ann N Y Acad Sci* 1035, 1-20 (2004)
33. D. J. Selkoe: Clearing the brain's amyloid cobwebs. *Neuron* 32, 177-180 (2001)

Decreased CSF Abeta42 in Alzheimer disease

34. C. Y. Lee, G. E. Landreth: The role of microglia in amyloid clearance from the AD brain. *J Neural Transm* 117, 949-960 (2010)
35. J. B. El Khoury, K. J. Moore, T. K. Means, J. Leung, K. Terada, M. Toft, M. W. Freeman and A. D. Luster: CD36 mediates the innate host response to beta-amyloid. *J Exp Med* 197, 1657-1666 (2003)
36. S. A. Grathwohl, R. E. Kalin, T. Bolmont, S. Prokop, G. Winkelmann, S. A. Kaeser, J. Odenthal, R. Radde, T. Eldh, S. Gandy, A. Aguzzi, M. Staufenbiel, P. M. Mathews, H. Wolburg, F. L. Heppner and M. Jucker: Formation and maintenance of Alzheimer's disease beta-amyloid plaques in the absence of microglia. *Nat Neurosci* 12, 1361-1363 (2009)
37. H. Akiyama, P. L. McGeer: Specificity of mechanisms for plaque removal after A beta immunotherapy for Alzheimer disease. *Nat Med* 10, 117-118 (2004)
38. A. Majumdar, D. Cruz, N. Asamoah, A. Buxbaum, I. Sohar, P. Lobel and F. R. Maxfield: Activation of microglia acidifies lysosomes and leads to degradation of Alzheimer amyloid fibrils. *Mol Biol Cell* 18, 1490-1496 (2007)
39. J. A. Nicoll, E. Barton, D. Boche, J. W. Neal, I. Ferrer, P. Thompson, C. Vlachouli, D. Wilkinson, A. Bayer, D. Games, P. Seubert, D. Schenk and C. Holmes: Abeta species removal after abeta42 immunization. *J Neuropathol Exp Neurol* 65, 1040-1048 (2006)
40. N. J. Abbott: Evidence for bulk flow of brain interstitial fluid: significance for physiology and pathology. *Neurochem Int* 45, 545-552 (2004)
41. R. Deane, Z. Wu, A. Sagare, J. Davis, Y. S. Du, K. Hamm, F. Xu, M. Parisi, B. LaRue, H. W. Hu, P. Spijkers, H. Guo, X. Song, P. J. Lenting, W. E. Van Nostrand and B. V. Zlokovic: LRP/amyloid beta-peptide interaction mediates differential brain efflux of Abeta isoforms. *Neuron* 43, 333-344 (2004)
42. R. O. Weller, M. Subash, S. D. Preston, I. Mazanti and R. O. Carare: Perivascular drainage of amyloid-beta peptides from the brain and its failure in cerebral amyloid angiopathy and Alzheimer's disease. *Brain Pathol* 18, 253-266 (2008)
43. R. O. Weller, E. Djuanda, H. Y. Yow and R. O. Carare: Lymphatic drainage of the brain and the pathophysiology of neurological disease. *Acta Neuropathol* 117, 1-14 (2009)
44. M. Johnston: The importance of lymphatics in cerebrospinal fluid transport. *Lymphat Res Biol* 1, 41-44 (2003)
45. O. Hansson, H. Zetterberg, P. Buchhave, U. Andreasson, E. Londos, L. Minthon and K. Blennow: Prediction of Alzheimer's disease using the CSF Abeta42/Abeta40 ratio in patients with mild cognitive impairment. *Dement Geriatr Cogn Disord* 23, 316-320 (2007)
46. P. Lewczuk, H. Esselmann, M. Otto, J. M. Maler, A. W. Henkel, M. K. Henkel, O. Eikenberg, C. Antz, W. R. Krause, U. Reulbach, J. Kornhuber and J. Wiltfang: Neurochemical diagnosis of Alzheimer's dementia by CSF Abeta42, Abeta42/Abeta40 ratio and total tau. *Neurobiol Aging* 25, 273-281 (2004)
47. M. Kanai, E. Matsubara, K. Isoe, K. Urakami, K. Nakashima, H. Arai, H. Sasaki, K. Abe, T. Iwatsubo, T. Kosaka, M. Watanabe, Y. Tomidokoro, M. Shizuka, K. Mizushima, T. Nakamura, Y. Igeta, Y. Ikeda, M. Amari, T. Kawarabayashi, K. Ishiguro, Y. Harigaya, K. Wakabayashi, K. Okamoto, S. Hirai and M. Shoji: Longitudinal study of cerebrospinal fluid levels of tau, A beta1-40, and A beta1-42 (43) in Alzheimer's disease: a study in Japan. *Ann Neurol* 44, 17-26 (1998)
48. P. E. Spies, D. Slats, J. M. Sjogren, B. P. Kremer, F. R. Verhey, M. G. Olde Rikkert and M. M. Verbeek: The cerebrospinal fluid amyloid beta42/40 ratio in the differentiation of Alzheimer's disease from non-Alzheimer's dementia. *Curr Alzheimer Res* 7, 470-476 (2010)
49. J. F. Ghersi-Egea, P. D. Gorevic, J. Ghiso, B. Frangione, C. S. Patlak and J. D. Fenstermacher: Fate of cerebrospinal fluid-borne amyloid beta-peptide: rapid clearance into blood and appreciable accumulation by cerebral arteries. *J Neurochem* 67, 880-883 (1996)
50. J. S. Crossgrove, G. J. Li and W. Zheng: The choroid plexus removes beta-amyloid from brain cerebrospinal fluid. *Exp Biol Med* (Maywood) 230, 771-776 (2005)
51. K. G. Kapoor, S. E. Katz, D. M. Grzybowski and M. Lubow: Cerebrospinal fluid outflow: an evolving perspective. *Brain Res Bull* 77, 327-334 (2008)
52. M. Johnston, C. Papaiconomou: Cerebrospinal fluid transport: a lymphatic perspective. *News Physiol Sci* 17, 227-230 (2002)
53. T. Iwatsubo, A. Odaka, N. Suzuki, H. Mizusawa, N. Nukina and Y. Ihara: Visualization of A beta 42 (43) and A beta 40 in senile plaques with end-specific A beta monoclonals: evidence that an initially deposited species is A beta 42 (43). *Neuron* 13, 45-53 (1994)
54. M. M. Verbeek, P. Eikelenboom and R. M. De Waal: Differences between the pathogenesis of senile plaques and congophilic angiopathy in Alzheimer disease. *J Neuropathol Exp Neurol* 56, 751-761 (1997)
55. A. A. Rensink, R. M. De Waal, B. Kremer and M. M. Verbeek: Pathogenesis of cerebral amyloid angiopathy. *Brain Res Brain Res Rev* 43, 207-223 (2003)
56. S. A. Gravina, L. Ho, C. B. Eckman, K. E. Long, L. Otvos, Jr., L. H. Younkin, N. Suzuki and S. G. Younkin:

Decreased CSF Abeta42 in Alzheimer disease

Amyloid beta protein (A beta) in Alzheimer's disease brain. Biochemical and immunocytochemical analysis with antibodies specific for forms ending at A beta 40 or A beta 42 (43). *J Biol Chem* 270, 7013-7016 (1995)

57. H. Funato, M. Yoshimura, K. Kusui, A. Tamaoka, K. Ishikawa, N. Ohkoshi, K. Namekata, R. Okeda and Y. Ihara: Quantitation of amyloid beta-protein (A beta) in the cortex during aging and in Alzheimer's disease. *Am J Pathol* 152, 1633-1640 (1998)

58. Y. Harigaya, M. Shoji, T. Kawarabayashi, M. Kanai, T. Nakamura, T. Iizuka, Y. Igeta, T. C. Saido, N. Sahara and H. Mori: Modified amyloid beta protein ending at 42 or 40 with different solubility accumulates in the brain of Alzheimer's disease. *Biochem Biophys Res Commun* 211, 1015-1022 (1995)

59. M. Moonis, J. M. Swearer, M. P. Dayaw, P. St George-Hyslop, E. Rogaeva, T. Kawarai and D. A. Pollen: Familial Alzheimer disease: decreases in CSF Abeta42 levels precede cognitive decline. *Neurology* 65, 323-325 (2005)

60. E. Portelius, U. Andreasson, J. M. Ringman, K. Buerger, J. Daborg, P. Buchhave, O. Hansson, A. Harmsen, M. K. Gustavsson, E. Hanse, D. Galasko, H. Hampel, K. Blennow and H. Zetterberg: Distinct cerebrospinal fluid amyloid beta peptide signatures in sporadic and PSEN1 A431E-associated familial Alzheimer's disease. *Mol Neurodegener* 5, 2 (2010)

61. A. Tamaoka, Y. Sekijima, S. Matsuno, T. Tokuda, S. Shoji and S. I. Ikeda: Amyloid beta protein species in cerebrospinal fluid and in brain from patients with Down's syndrome. *Ann Neurol* 46, 933 (1999)

62. T. Tapiola, H. Soininen and T. Pirtila: CSF tau and Abeta42 levels in patients with Down's syndrome. *Neurology* 56, 979-980 (2001)

63. F. M. LaFerla, K. N. Green and S. Oddo: Intracellular amyloid-beta in Alzheimer's disease. *Nat Rev Neurosci* 8, 499-509 (2007)

64. K. G. Mawuenyega, W. Sigurdson, V. Ovod, L. Munsell, T. Kasten, J. C. Morris, K. E. Yarasheski and R. J. Bateman: Decreased clearance of CNS beta-amyloid in Alzheimer's disease. *Science* 330, 1774 (2010)

65. M. A. Leissring, W. Farris, A. Y. Chang, D. M. Walsh, X. Wu, X. Sun, M. P. Frosch and D. J. Selkoe: Enhanced proteolysis of beta-amyloid in APP transgenic mice prevents plaque formation, secondary pathology, and premature death. *Neuron* 40, 1087-1093 (2003)

66. J. Apelt, K. Ach and R. Schliebs: Aging-related down-regulation of neprilysin, a putative beta-amyloid-degrading enzyme, in transgenic Tg2576 Alzheimer-like mouse brain is accompanied by an astroglial upregulation in the vicinity of beta-amyloid plaques. *Neurosci Lett* 339, 183-186 (2003)

67. H. Akiyama, H. Kondo, K. Ikeda, M. Kato and P. L. McGeer: Immunohistochemical localization of neprilysin in the human cerebral cortex: inverse association with vulnerability to amyloid beta-protein (Abeta) deposition. *Brain Res* 902, 277-281 (2001)

68. K. Yasojima, H. Akiyama, E. G. McGeer and P. L. McGeer: Reduced neprilysin in high plaque areas of Alzheimer brain: a possible relationship to deficient degradation of beta-amyloid peptide. *Neurosci Lett* 297, 97-100 (2001)

69. T. Ohgami, T. Kitamoto, R. W. Shin, Y. Kaneko, K. Ogomori and J. Tateishi: Increased senile plaques without microglia in Alzheimer's disease. *Acta Neuropathol* 81, 242-247 (1991)

70. W. J. Streit, J. R. Conde, S. E. Fendrick, B. E. Flanary and C. L. Mariani: Role of microglia in the central nervous system's immune response. *Neurol Res* 27, 685-691 (2005)

71. W. J. Streit: Microglial senescence: does the brain's immune system have an expiration date? *Trends Neurosci* 29, 506-510 (2006)

72. M. M. Wilhelmus, I. Otte-Holler, J. J. van Triel, R. Veerhuis, M. L. Maat-Schieman, G. Bu, R. M. De Waal and M. M. Verbeek: Lipoprotein receptor-related protein-1 mediates amyloid-beta-mediated cell death of cerebrovascular cells. *Am J Pathol* 171, 1989-1999 (2007)

73. V. Giedraitis, J. Sundelof, M. C. Irizarry, N. Garevik, B. T. Hyman, L. O. Wahlund, M. Ingelsson and L. Lannfelt: The normal equilibrium between CSF and plasma amyloid beta levels is disrupted in Alzheimer's disease. *Neurosci Lett* 427, 127-131 (2007)

74. T. Kawarabayashi, M. Shoji: Plasma biomarkers of Alzheimer's disease. *Curr Opin Psychiatry* 21, 260-267 (2008)

75. A. E. Roher, Y. M. Kuo, C. Esh, C. Knebel, N. Weiss, W. Kalback, D. C. Luehrs, J. L. Childress, T. G. Beach, R. O. Weller and T. A. Kokjohn: Cortical and leptomeningeal cerebrovascular amyloid and white matter pathology in Alzheimer's disease. *Mol Med* 9, 112-122 (2003)

76. Y. C. Zhu, C. Dufouil, B. Mazoyer, A. Soumare, F. Ricolfi, C. Tzourio and H. Chabriat: Frequency and Location of Dilated Virchow-Robin Spaces in Elderly People: A Population-Based 3D MR Imaging Study. *AJNR Am J Neuroradiol* 32, 709-713 (2011)

77. R. O. Weller: How well does the CSF inform upon pathology in the brain in Creutzfeldt-Jakob and Alzheimer's diseases? *J Pathol* 194, 1-3 (2001)

78. J. Attems, F. Lintner and K. A. Jellinger: Amyloid beta peptide 1-42 highly correlates with capillary cerebral amyloid angiopathy and Alzheimer disease pathology. *Acta Neuropathol* 107, 283-291 (2004)

Decreased CSF Abeta42 in Alzheimer disease

79. M. M. Verbeek, B. P. Kremer, M. O. Rikkert, P. H. Van Domburg, M. E. Skehan and S. M. Greenberg: Cerebrospinal fluid amyloid beta (40) is decreased in cerebral amyloid angiopathy. *Ann Neurol* 66, 245-249 (2009)

80. J. T. Jarrett, E. P. Berger and P. T. Lansbury, Jr.: The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry* 32, 4693-4697 (1993)

81. J. Wiltfang, H. Esselmann, M. Bibl, A. Smirnov, M. Otto, S. Paul, B. Schmidt, H. W. Klafki, M. Maler, T. Dyrks, M. Bienert, M. Beyermann, E. Ruther and J. Kornhuber: Highly conserved and disease-specific patterns of carboxyterminally truncated Abeta peptides 1-37/38/39 in addition to 1-40/42 in Alzheimer's disease and in patients with chronic neuroinflammation. *J Neurochem* 81, 481-496 (2002)

82. J. E. Preston: Ageing choroid plexus-cerebrospinal fluid system. *Microsc Res Tech* 52, 31-37 (2001)

Key words: Dementia, Cerebrospinal fluid, Amyloid-beta peptides, Biomarker, Review

Send correspondence to: Petra E. Spies, Department of Geriatric Medicine, 925, Radboud University Nijmegen Medical Center, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands, Tel.: 31 24 361 6772, Fax: 31 24 361 7408, E-mail: PetraESpies@gmail.com