

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/112687>

Please be advised that this information was generated on 2021-10-20 and may be subject to change.

MATHEMATICAL MODEL FOR CALCIUM OSCILLATIONS IN NON-EXCITABLE CELLS

Joaquin J. Torres, Hilbert J. Kappen, Peter H.G.M. Willems, **Werner J.H. Koopman**:
University of Nijmegen

We introduce a general mathematical model to describe calcium oscillations in non-excitable cells. For this purpose, Ca^{2+} oscillations were recorded in single pancreatic acinar cells of the rat. The model takes into account following Ca^{2+} fluxes: V_{IP_3} , the flow of Ca^{2+} from the ER into the cytosol via the InsP_3 -receptor; V_{CRAC} , the capacitive Ca^{2+} entry pathway; V_{SERCA} and V_{PMCA} , sequestration and extrusion of Ca^{2+} and two leak fluxes (from the ER and across the plasma membrane). After choosing realistic parameters, the dynamics of the model were analyzed by bifurcation analysis and compared with experimental data. We show explicitly that the dynamics of Ca^{2+} oscillations generated by the model agrees qualitatively and quantitatively with those observed in real cells. The model is useful to explain hormone-induced Ca^{2+} patterns recorded in pancreatic acinar cells. These patterns are directly involved in the regulation of subcellular processes that control the functionality of the pancreatic acinar cell. Currently, the one-dimensional model is used to build a 2D model to simulate the spatiotemporal kinetics of Ca^{2+} waves. Numerical simulations reveal the presence of hysteresis and bistability in such a 2D model that explains the presence of traveling and spiral Ca^{2+} waves in the system similar to those in real cells.