

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/125850>

Please be advised that this information was generated on 2019-10-18 and may be subject to change.

RESEARCH

Open Access

IL-1 β processing in mechanical ventilation-induced inflammation is dependent on neutrophil factors rather than caspase-1

Kim Timmermans^{1,3†}, Selina El van der Wal^{1†}, Michiel Vaneker¹, Jeroen AWM van der Laak², Mihai G Netea⁴, Peter Pickkers³, Gert Jan Scheffer¹, Leo AB Joosten⁴ and Matthijs Kox^{1,3*}

* Correspondence: Matthijs.Kox@radboudumc.nl

[†]Equal contributors

¹Department of Anaesthesiology, RUNMC, Nijmegen 6500 HB, The Netherlands

³Department of Intensive Care Medicine, RUNMC, Internal mail 710, P.O. Box 9101, Nijmegen 6500 HB, The Netherlands

Full list of author information is available at the end of the article

Abstract

Purpose: Mechanical ventilation can cause ventilator-induced lung injury, characterized by a sterile inflammatory response in the lungs resulting in tissue damage and respiratory failure. The cytokine interleukin-1 β (IL-1 β) is thought to play an important role in the pathogenesis of ventilator-induced lung injury. Cleavage of the inactive precursor pro-IL-1 β to form bioactive IL-1 β is mediated by several types of proteases, of which caspase-1, activated within the inflammasome, is the most important. Herein, we studied the roles of IL-1 β , caspase-1 and neutrophil factors in the mechanical ventilation-induced inflammatory response in mice.

Methods: Untreated wild-type mice, IL-1 $\alpha\beta$ knockout and caspase-1 knockout mice, pralnacasan (a selective caspase-1 inhibitor)-treated mice, anti-keratinocyte-derived chemokine (KC)-treated mice and cyclophosphamide-treated neutrophil-depleted wild-type mice were ventilated using clinically relevant ventilator settings (tidal volume 8 ml/kg). The lungs and plasma were collected to determine blood gas values, cytokine profiles and neutrophil influx.

Results: Mechanical ventilation resulted in increased pulmonary concentrations of IL-1 β and KC and increased pulmonary neutrophil influx compared with non-ventilated mice. Ventilated IL-1 $\alpha\beta$ knockout mice did not demonstrate this increase in cytokines. No significant differences were observed between wild-type and caspase-1-deficient or pralnacasan-treated mice. In contrast, in anti-KC antibody-treated mice and neutropenic mice, inflammatory parameters decreased in comparison with ventilated non-treated mice.

Conclusions: Our results illustrate that IL-1 is indeed an important cytokine in the inflammatory cascade induced by mechanical ventilation. However, the inflammasome/caspase-1 appears not to be involved in IL-1 β processing in this type of inflammatory response. The attenuated inflammatory response observed in ventilated anti-KC-treated and neutropenic mice suggests that IL-1 β processing in mechanical ventilation-induced inflammation is mainly mediated by neutrophil factors.

Keywords: VILI; Inflammation; Ventilation; IL-1 β

Introduction

Mechanical ventilation is a life-saving therapy, although it can also cause ventilator-induced lung injury (VILI) [1]. VILI is characterized by a sterile inflammatory response in the lungs resulting in tissue damage that may sustain respiratory failure. The mechanical ventilation-induced inflammatory response can also spread systemically, which in severe cases can result in multi-organ dysfunction syndrome (MODS) [2]. Even protective ventilation strategies that do not cause direct mechano-induced tissue damage (baro- or volutrauma) have been shown to elicit the release of pro-inflammatory cytokines, recruitment of leukocytes and subsequent lung injury [3,4]. The mechanisms behind this so-called 'biotrauma' have not yet been completely elucidated.

Previous studies have demonstrated that the TLR4/TRIF pathway is important in the mechanical ventilation-induced inflammatory response [4,5]. Furthermore, it is becoming increasingly clear that the pro-inflammatory cytokine interleukin-1 β (IL-1 β) plays a key role in the pathogenesis of the inflammatory response and VILI by promoting neutrophil recruitment and by increasing epithelial injury and permeability [6-8]. Through recognition by the IL-1 receptor (IL-1R), not only the secreted IL-1 β but also the cell-associated family member IL-1 α may stimulate production of other inflammatory cytokines via IL-1R-associated kinases (IRAKs) and thereby positively amplify the inflammatory response [9]. However, up till now, this has not been studied in the context of mechanical ventilation-induced inflammation.

Upon activation of the innate immune system, e.g. via TLRs, IL-1 β is synthesized as an inactive precursor molecule, pro-IL-1 β , that cannot bind and activate the IL-1R [10]. In order to process pro-IL-1 β and form bioactive IL-1 β , proteolytic cleavage of the N-terminal 116 amino acids from the precursor is required. Caspase-1 is the major protein implicated in cleavage of pro-IL-1 β [10,11].

Caspase-1 exists as an inactive zymogen in cells of myeloid origin (e.g. tissue macrophages, dendritic cells) which needs to be activated to perform its proteolytic tasks [9]. Caspase-1 is also known to be expressed in a wide range of other cell types including lung fibroblasts and epithelial cells [12,13]. The inflammasome is a protein platform that is responsible for the activation of caspase-1 [10,14]. A broad range of infectious and autoimmune diseases that involve IL-1 β have been associated with inappropriate activation of the inflammasome [12,14,15], while in several other disease models in which IL-1 β plays a crucial role, the inflammasome appears not to be involved [16,17]. IL-1 β processing in these models might rely on neutrophil serine proteases, like elastase, granzyme A, cathepsin G or proteinase-3 [10,18-20]. Hitherto, the role of caspase-1 in processing of IL-1 β in the mechanical ventilation-induced inflammatory response is unknown.

We studied the roles of IL-1 β , caspase-1 and neutrophil factors in the mechanical ventilation-induced inflammatory response in mice ventilated with clinically relevant ventilator settings.

Materials and methods

All experiments were approved by the Regional Animal Ethics Committee in Nijmegen and performed under the guidelines of the Dutch Council for Animal Care and the National Institutes of Health. They have therefore been performed in accordance with

the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments.

Animals

Age-matched wild-type C57Bl/6 mice and extensively backcrossed caspase-1 knockout mice or IL-1 $\alpha\beta$ knockout mice (aged 8 to 14 weeks, weight 25 ± 4 g) with C57Bl/6 background were used in this study. The mice were housed in a light- and temperature-controlled room under specific pathogen-free (SPF) conditions. Standard pelleted chow (1.00% Ca, 0.22% Mg, 0.24% Na, 0.70% P, 1.02% K, SSNIFF Spezialdiäten GmbH, Soest, Germany) and drinking water were available *ad libitum*. These conditions are similar to previous studies in which this mouse model was used [4,5,21,22].

Experimental design

IL-1 $\alpha\beta$ knockout experiments

IL-1 can induce inflammation via activation of the IL-1 receptor. To study whether IL-1 is indeed involved in initiation and/or propagation of the inflammatory cascade induced by mechanical ventilation, mechanically ventilated IL-1 $\alpha\beta^{-/-}$ mice ($n = 8$) were compared with ventilated wild-type mice ($n = 8$). As controls, non-ventilated IL-1 $\alpha\beta^{-/-}$ ($n = 8$) and wild-type mice ($n = 8$) were used.

Caspase-1 experiments

Caspase-1 is able to cleave the inactive precursor pro-IL-1 β to form the active cytokine IL-1 β . To study the role of caspase-1 in the mechanical ventilation-induced inflammatory response, mechanically ventilated caspase-1 knockout mice ($n = 8$) and ventilated wild-type mice treated with the selective caspase-1 inhibitor pralnacasan (100 mg/kg) ($n = 8$) were compared with ventilated untreated wild-type mice ($n = 8$) [23,24]. As controls, non-ventilated caspase-1 $^{-/-}$, pralnacasan-treated wild-type and untreated wild-type mice ($n = 8$ per group) were used.

Anti-KC antibody experiments

Apart from caspase-1, neutrophil serine proteases are also able to process IL-1 β [8]. In order to investigate whether the attraction of neutrophils by the chemo-attractant keratinocyte-derived chemokine (KC) is involved in the inflammatory response elicited by mechanical ventilation, mechanically ventilated wild-type mice treated with an intraperitoneal dose of 100 μ g of a neutralizing monoclonal anti-KC antibody (R&D Systems, Minneapolis, MN, USA) 1 h before induction of anaesthesia ($n = 8$) were compared with ventilated untreated wild-type mice ($n = 8$). As controls, non-ventilated untreated wild-type mice ($n = 8$) were used.

Neutrophil depletion experiments

Neutrophil serine proteases are able to process IL-1 β [8]. In order to study the possible role of neutrophil factors in IL-1 β processing in the mechanical ventilation-induced inflammatory response, mechanically ventilated neutrophil-depleted wild-type mice ($n = 8$) were compared with ventilated untreated wild-type mice ($n = 8$). As controls, non-ventilated wild-type mice ($n = 8$) were used. The neutrophil-depleted group was neutrophil-depleted with cyclophosphamide as described previously [25,26].

Experimental procedures

The mice were anaesthetized using an intraperitoneal injection of 7.5 μ l per gram body weight of KMA mix (25.5 mg/ml ketamine, 40 μ g/ml medetomidine, 0.1 mg/ml atropine in saline). Subsequently, the animals were orally intubated, an arterial line was placed in the arteria carotis, and the mice were mechanically ventilated (MiniVent[®], Hugo Sachs Elektronik-Harvard Apparatus, March-Hugstetten, Germany). The ventilation settings used were based on measured tidal volume and respiratory rate during spontaneous ventilation in C57Bl/6 mice [27]: a tidal volume of 8 ml/kg body weight and a frequency of 150/min. All animals received 4 cm H₂O positive end-expiratory pressure (PEEP), and fraction of inspired oxygen was set to 0.4. In order to maintain anaesthesia, boluses of 5.0 μ l per gram body weight maintenance KMA mix (3.6 mg/ml ketamine, 4 μ g/ml medetomidine, 7.5 μ g/ml atropine in saline) were given every 30 min via an intraperitoneally placed catheter. Rectal temperature was monitored continuously and maintained between 36.0°C and 37.5°C using a heating pad. After the 4-h ventilation period, the mice were sacrificed by exsanguination under anaesthesia. The control mice were anaesthetized, but not ventilated, and sacrificed shortly after induction of anaesthesia. Tissue and blood were sampled in order to determine blood gas values (only in ventilated mice), cytokine production and neutrophil influx.

Lipopolysaccharide (LPS) was measured in the mechanical ventilation circuit by Limulus Amebocyte Lysate testing (Cambrex Bio Science, Walkersville, MD, USA; detection limit: 0.06 IU/ml) to rule out contamination and LPS-induced pulmonary inflammation.

Tissue harvesting

Plasma was isolated by centrifugation at 13,000g for 5 min and stored at -80°C. Immediately after exsanguination, the heart and lungs were carefully removed *en block* via midline sternotomy. The right middle lung lobe was fixed in 4% buffered formalin solution overnight at room temperature. The right lung was snap-frozen in liquid nitrogen and stored at -80°C. The left lung was snap-frozen and placed in 500 μ l lysis buffer containing PBS, 0.5% Triton X-100 and protease inhibitor (complete EDTA-free tablets, Roche, Woerden, The Netherlands). Subsequently, the lungs were homogenized using a polytron and subjected to two rapid freeze-thaw cycles using liquid nitrogen. Finally, homogenates were centrifuged (10 min, 16,000g, 4°C), and the supernatant was stored at -80°C until further analysis.

Pulmonary neutrophil influx

After overnight incubation in 4% buffered formalin solution, the right middle lung lobe was dehydrated and embedded in paraplast (Amstelstad, Amsterdam, The Netherlands). Sections of 4- μ m thickness were used. Enzyme histochemistry using chloracetatesterase (LEDER staining) was used to visualize the enzyme activity in the neutrophils. Neutrophils were counted manually (ten fields per mouse), and after automated correction for air/tissue ratio, the average number of neutrophils per square centimetre per mouse was calculated.

Biochemical analysis

KC (murine equivalent of human IL-8) in the lung homogenate was determined by enzyme-linked immunosorbent assay (ELISA; R&D Systems, Minneapolis, MN, USA). The lower detection limit is 160 pg/ml. IL-1 β in the lung homogenate was determined

using a radioimmunoassay (RIA) as described previously [28]. In the samples of the IL-1 $\alpha\beta$ (Figure 1) and caspase (Figure 2) experiments, total protein concentrations in the lung homogenates were determined using a BCA protein assay (Thermo Fisher Scientific, Etten-Leur, The Netherlands), and cytokine concentrations in the homogenates were normalized for protein concentration and therefore expressed as nanogram cytokine per microgram protein. In the anti-KC (Figure 3) and neutrophil depletion (Figure 4) experiments, cytokine concentrations in the lung homogenate were not normalized for total protein content due to insufficient sample volume and therefore expressed as picogram per millilitre.

Statistical analysis


Data were not normally distributed (determined using the Kolmogorov-Smirnov and Shapiro-Wilk tests) and therefore expressed as median and range or median and interquartile range (IQR). Differences between groups were analyzed using the Kruskal-Wallis and Dunn's *post hoc* tests. Statistical analysis was performed using GraphPad Prism 5 software (GraphPad Software, La Jolla, CA, USA). *P* values <0.05 were considered significant.

Results

Mean arterial pressure remained stable and above 65 mmHg in all animals throughout the mechanical ventilation period. Blood gas values that were obtained at the end of the ventilation period did not indicate substantial lung injury (Table 1).

Involvement of IL-1 in the mechanical ventilation-induced inflammatory response

After 4 h of mechanical ventilation, pulmonary levels of pro-inflammatory cytokine KC significantly increased in wild-type mice compared with non-ventilated wild-type mice. In contrast, ventilated IL-1 $\alpha\beta$ knockout mice did not show an increase in pulmonary cytokines compared with non-ventilated IL-1 $\alpha\beta$ knockout mice (Figure 1).







Figure 2 Involvement of caspase-1 in the mechanical ventilation-induced inflammatory response. Pulmonary neutrophil counts expressed as the number of neutrophils per square centimetre tissue and IL-1 β and KC levels in lung homogenates expressed as nanogram cytokine per microgram total protein. Data are expressed as box-and-whiskers plots, with min to max range as whiskers. Results of analysis in the non-ventilated (C) and ventilated (V) wild-type (WT) mice, caspase-1 knockout (-/-) mice and pralnacasan-treated mice are shown. * $p < 0.05$.



Involvement of caspase-1 in the mechanical ventilation-induced inflammatory response

Pulmonary neutrophil influx significantly increased in mechanically ventilated mice compared with non-ventilated wild-type and caspase-1^{-/-} mice, but no differences were observed between wild-type mice, caspase-1^{-/-} mice or pralnacasan-treated mice. Similar to the results described above, 4 h of mechanical ventilation resulted in increased IL-1 β and KC concentrations in lung homogenates in all groups. However, no significant differences in lung cytokine levels were observed between wild-type mice, caspase-1^{-/-} mice or pralnacasan-treated mice. (Figure 2)

Involvement of neutrophil factors in the mechanical ventilation-induced inflammatory response

To determine whether neutrophil factors are involved in the mechanical ventilation-induced inflammatory response and IL-1 β processing, we investigated the effects of treatment with




Table 1 Blood gas values after 4 h of ventilation

	Median	IQR
pH	7.36	7.25 to 7.38
pCO ₂	4.73	4.17 to 5.18
PO ₂	15.3	14.6 to 17.5
BE	-5.5	-7.3 to -4.0
HCO ₃	20.2	18.3 to 20.7
TCO ₂	21.0	19.8 to 21.5
sO ₂ %	99%	98 to 99
Lac	0.98	0.90 to 1.16

Values (median and IQR) from a representative ventilated group (wild-type ventilated mice used as the control group for caspase-1^{-/-} and pralnacasan-treated mice).

an antibody against KC. KC is one of the major factors involved in neutrophil attraction to the site of inflammation (chemo-attractants). Mechanical ventilation resulted in increased levels of pulmonary neutrophils (Figure 3). This increase was abrogated by pre-treatment with an anti-KC antibody. Furthermore, the mechanical ventilation-induced increase in pulmonary IL-1 β levels was less pronounced in anti-KC-treated mice compared with untreated mice, although this did not reach statistical significance (Figure 3).


To further confirm the role of neutrophil factors, we investigated the effects of mechanical ventilation following neutrophil depletion using cyclophosphamide. The effect of cyclophosphamide was visually inspected, and no pulmonary neutrophils were present (data not shown). As depicted in Figure 4, the mechanical ventilation-induced increase in pulmonary IL-1 β and KC concentrations was diminished in neutrophil-depleted mice.

Our hypothesis regarding the role of IL-1 β processing in the inflammatory response following mechanical ventilation is illustrated in Figure 5.

Discussion

Consistent with previous results published by our group [4,5,22] and others [29,30], the present study shows that mechanical ventilation using clinically relevant settings induces a pulmonary inflammatory response in mice. In addition, our data is in support of previous findings that IL-1 plays an important role in initiation and/or propagation of the mechanical ventilation-induced inflammatory response and suggests that processing of IL-1 β in mechanical ventilation-induced inflammation occurs via the release of neutrophil factors and not through caspase-1-dependent mechanisms.

Our finding that caspase-1 does not play a significant role in mechanical ventilation-induced inflammation is in contrast to a recent study where the NLRP3 inflammasome was found to play an important role in the mechanical ventilation-induced inflammatory response and VILI [31]. Several differences between their study and ours might explain the different results. First, in the previous study, ASC and NLRP3 (components of the inflammasome upstream of caspase-1) knockout mice were used, and it was shown that mechanical ventilation activated caspase-1 in a NLRP3-dependent fashion. Nevertheless, it is very well possible that ASC and NLRP3 play other roles in the mechanical ventilation-induced inflammatory cascade than merely activating caspase-1. As abrogation and inhibition of caspase-1 by either a knockout approach or pralnacasan treatment did not have any effect in our



model, the role of caspase-1/the inflammasome appears not to be as crucial as suggested. Second, differences between wild-type and ASC or NLRP3 knockout were only found at a high tidal volume of 15 ml/kg, known to cause extensive lung damage [22], while no effects were found at a low tidal volume of 7.5 ml/kg, which is more representative of the current clinical practice and similar to that used in the present study. This suggests that the inflammasome might play a more important role at higher tidal volumes which lead to apparent lung injury but not in mechanical ventilation-induced inflammation at clinically relevant ventilator settings. Interestingly, a more recent study from the same group showed that pre-treatment with allopurinol or uricase (both degraders of known inflammasome-activating factors [32]) did not decrease mechanical ventilation-induced inflammation, which is in support of a caspase/inflammasome-independent mechanism [33]. As beneficial effects of uricase and allopurinol were observed in terms of alveolar barrier dysfunction, it appears plausible that ASC and NLRP3 are involved in VILI via inflammation-independent mechanisms.

The pronounced influx of neutrophils in the lung observed in our experiments suggests a major role for these inflammatory cells in the inflammatory cascade following mechanical ventilation. Our findings that treatment with an antibody against KC or depletion of neutrophils reduced the mechanical ventilation-induced production of IL-1 β and KC indicate an important role for neutrophils in initiation and/or propagation of the inflammatory response. In this respect, pro-IL-1 β cleavage in our model is probably

achieved through neutrophil factors, such as the serine proteases proteinase-3 (PR-3), elastase or cathepsin G, leading to bioactive IL-1 β and propagation of the inflammatory response through binding of the IL-1-receptor, which in turn leads to production of other inflammatory cytokines such as KC [10,34,35]. Several other IL-1 β -mediated inflammatory responses are described to be partly or completely independent of the inflammasome and caspase-1 and possibly dependent on neutrophil factors, including proteinase-3 and cathepsin G [35]. Future studies should focus on the confirmation of our hypothesis and the identification of these neutrophil factors.

Our study has several limitations. First, we used cyclophosphamide to deplete neutrophils. While this is a widely used method [25,26,36,37], cyclophosphamide treatment may also result in depletion of other cell types that play a role in mechanical ventilation-induced inflammation [38,39]. Nevertheless, our data of mice treated with an anti-KC antibody underline the importance of neutrophils in this process. Second, no histological slides to perform neutrophil counts were collected in the IL-1 $\alpha\beta^{-/-}$ experiments to investigate whether these knockout mice were still able to recruit neutrophils. Finally, we cannot exclude the possibility that next to mechanical ventilation, the procedures related to the instrumentation/ventilation (e.g. intubation, arterial cannulation) also induce inflammation to a certain extent. However, we have previously shown that the inflammatory response is aggravated when mice are ventilated with these parameters for a longer period of time or when higher tidal volumes are used, suggesting that the inflammatory response is mainly ventilation-induced.

Conclusions

In conclusion, our results indicate that IL-1 signalling is important in mechanical ventilation-induced inflammation. We show that following mechanical ventilation, IL-1 β bioactivation is not caspase-1 dependent but appears to be mediated by neutrophil factors, leading to a positive amplification loop and further propagation of the inflammatory response. Further elucidation of the precise mechanism of IL-1 β processing in mechanical ventilation-induced inflammation could provide novel targets for the future treatment of VILI [40].

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KT performed the experiments, carried out the data analysis, interpreted the results and drafted the manuscript. SW wrote the applications for the animal ethics committee, performed the data analysis and corrected the manuscript. MV performed the anti-KC and neutrophil depletion experiments and corrected the manuscript. JL developed the software to automate neutrophil counting and corrected the manuscript. MN and LJ conceived the study design, interpreted the results, were involved in the hypothesis development and corrected the manuscript. PP and GS supervised the experiments, interpreted the results and corrected the manuscript. MK supervised the experiments, interpreted the results, was involved in the hypothesis development and supervised the drafting and correction of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by a Vici Grant of the Netherlands Organization for Scientific Research to Mihai Netea. The authors would like to thank Ilona van den Brink and Francien van de Pol for the assistance with the animal experiments and Ineke Verschueren and Jelle Gerretsen for the technical assistance with the cytokine assays.

Author details

¹Department of Anaesthesiology, RUNMC, Nijmegen 6500 HB, The Netherlands. ²Department of Pathology, RUNMC, Nijmegen 6500 HB, The Netherlands. ³Department of Intensive Care Medicine, RUNMC, Internal mail 710, P.O. Box 9101, Nijmegen 6500 HB, The Netherlands. ⁴Department of Internal Medicine, RUNMC, Nijmegen 6500 HB, The Netherlands.

Received: 9 October 2013 Accepted: 10 October 2013
Published: 29 October 2013

References

1. Lionetti V, Recchia FA, Ranieri VM (2005) Overview of ventilator-induced lung injury mechanisms. *Curr Opin Crit Care* 11:82–86
2. Villar J, Blanco J, Zhang H, Slutsky AS (2011) Ventilator-induced lung injury and sepsis: two sides of the same coin? *Minerva Anesthesiol* 77:647–653
3. Gattinoni L, Protti A, Caironi P, Carlesso E (2010) Ventilator-induced lung injury: the anatomical and physiological framework. *Crit Care Med* 38:S539–548
4. Vaneker M, Joosten LA, Heunks LM, Snijdelaar DG, Halbertsma FJ, van Egmond J, Netea MG, van der Hoeven JG, Scheffer GJ (2008) Low-tidal-volume mechanical ventilation induces a toll-like receptor 4-dependent inflammatory response in healthy mice. *Anesthesiology* 109:465–472
5. Vaneker M, Heunks LM, Joosten LA, van Hees HW, Snijdelaar DG, Halbertsma FJ, van Egmond J, Netea MG, van der Hoeven JG, Scheffer GJ (2009) Mechanical ventilation induces a Toll/interleukin-1 receptor domain-containing adapter-inducing interferon beta-dependent inflammatory response in healthy mice. *Anesthesiology* 111:836–843
6. Frank JA, Pittet JF, Wray C, Matthay MA (2008) Protection from experimental ventilator-induced acute lung injury by IL-1 receptor blockade. *Thorax* 63:147–153
7. Ma SF, Grigoryev DN, Taylor AD, Nonas S, Sammani S, Ye SQ, Garcia JG (2005) Bioinformatic identification of novel early stress response genes in rodent models of lung injury. *Am J Physiol Lung Cell Mol Physiol* 289:L468–477
8. Wallace MJ, Probyn ME, Zahra VA, Crossley K, Cole TJ, Davis PG, Morley CJ, Hooper SB (2009) Early biomarkers and potential mediators of ventilation-induced lung injury in very preterm lambs. *Respir Res* 10:19
9. Dinarello CA (2011) Blocking interleukin-1beta in acute and chronic autoinflammatory diseases. *J Intern Med* 269:16–28
10. Dinarello CA (2011) Interleukin-1 in the pathogenesis and treatment of inflammatory diseases. *Blood* 117:3720–3732
11. Khare S, Luc N, Dorfleutner A, Stehlik C (2010) Inflammasomes and their activation. *Crit Rev Immunol* 30:463–487
12. Ichinohe T, Lee HK, Ogura Y, Flavell R, Iwasaki A (2009) Inflammasome recognition of influenza virus is essential for adaptive immune responses. *J Exp Med* 206:79–87
13. Vats V, Agrawal T, Salhan S, Mittal A (2010) Characterization of apoptotic activities during chlamydia trachomatis infection in primary cervical epithelial cells. *Immunol Invest* 39:674–687
14. Yazdi AS, Guarda G, D'Ombria MC, Drexler SK (2010) Inflammatory caspases in innate immunity and inflammation. *J Innate Immun* 2:228–237
15. Hoffman HM, Wanderer AA (2010) Inflammasome and IL-1beta-mediated disorders. *Curr Allergy Asthma Rep* 10:229–235
16. Joosten LA, Netea MG, Fantuzzi G, Koenders MI, Helsen MM, Sparrer H, Pham CT, van der Meer JW, Dinarello CA, van den Berg WB (2009) Inflammatory arthritis in caspase 1 gene-deficient mice: contribution of proteinase 3 to caspase 1-independent production of bioactive interleukin-1beta. *Arthritis Rheum* 60:3651–3662
17. Mencacci A, Bacci A, Cenci E, Montagnoli C, Fiorucci S, Casagrande A, Flavell RA, Bistoni F, Romani L (2000) Interleukin 18 restores defective Th1 immunity to *Candida albicans* in caspase 1-deficient mice. *Infect Immun* 68:5126–5131
18. Coeshott C, Ohnemus C, Pilyavskaya A, Ross S, Wieczorek M, Kroona H, Leimer AH, Cheronis J (1999) Converting enzyme-independent release of tumor necrosis factor alpha and IL-1beta from a stimulated human monocytic cell line in the presence of activated neutrophils or purified proteinase 3. *Proc Natl Acad Sci U S A* 96:6261–6266
19. Greten FR, Arkan MC, Bollrath J, Hsu LC, Goode J, Miething C, Goktuna SI, Neuenhahn M, Fierer J, Paxian S, Van Rooijen N, Xu Y, O'Cain T, Jaffee BB, Busch DH, Duyster J, Schmid RM, Eckmann L, Karin M (2007) NF-kappaB is a negative regulator of IL-1beta secretion as revealed by genetic and pharmacological inhibition of IKKbeta. *Cell* 130:918–931
20. Fantuzzi G, Dinarello CA (1999) Interleukin-18 and interleukin-1 beta: two cytokine substrates for ICE (caspase-1). *J Clin Immunol* 19:1–11
21. Vaneker M, Santosa JP, Heunks LM, Halbertsma FJ, Snijdelaar DG, van Egmond J, van den Brink IA, van de Pol FM, van der Hoeven JG, Scheffer GJ (2009) Isoflurane attenuates pulmonary interleukin-1beta and systemic tumor necrosis factor-alpha following mechanical ventilation in healthy mice. *Acta Anaesthesiol Scand* 53:742–748
22. Vaneker M, Halbertsma FJ, van Egmond J, Netea MG, Dijkman HB, Snijdelaar DG, Joosten LA, van der Hoeven JG, Scheffer GJ (2007) Mechanical ventilation in healthy mice induces reversible pulmonary and systemic cytokine elevation with preserved alveolar integrity: an in vivo model using clinical relevant ventilation settings. *Anesthesiology* 107:419–426
23. Loher F, Bauer C, Landauer N, Schmall K, Siegmund B, Lehr HA, Dauer M, Schoenharting M, Endres S, Eigler A (2004) The interleukin-1 beta-converting enzyme inhibitor pralnacasan reduces dextran sulfate sodium-induced murine colitis and T helper 1 T-cell activation. *J Pharmacol Exp Ther* 308:583–590
24. Cornelis S, Kersse K, Festjens N, Lamkanfi M, Vandenabeele P (2007) Inflammatory caspases: targets for novel therapies. *Curr Pharm Des* 13:367–385
25. van't Wout JW, Linde I, Leijh PC, van Furth R (1989) Effect of irradiation, cyclophosphamide, and etoposide (VP-16) on number of peripheral blood and peritoneal leukocytes in mice under normal conditions and during acute inflammatory reaction. *Inflammation* 13:1–14
26. Netea MG, Kullberg BJ, Blok WL, Netea RT, van der Meer JW (1997) The role of hyperuricemia in the increased cytokine production after lipopolysaccharide challenge in neutropenic mice. *Blood* 89:577–582
27. Janssen BJ, Smits JF (2002) Autonomic control of blood pressure in mice: basic physiology and effects of genetic modification. *Am J Physiol Regul Integr Comp Physiol* 282:R1545–1564
28. Li LF, Ouyang B, Choukroun G, Matyal R, Mascarenhas M, Jafari B, Bonventre JV, Force T, Quinn DA (2003) Stretch-induced IL-8 depends on c-Jun NH2-terminal and nuclear factor-kappaB-inducing kinases. *Am J Physiol Lung Cell Mol Physiol* 285:L464–475

29. The Acute Respiratory Distress Syndrome Network (2000) Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 342:1301–1308
30. Villar J, Cabrera NE, Casula M, Flores C, Valladares F, Diaz-Flores L, Muros M, Slutsky AS, Kacmarek RM (2010) Mechanical ventilation modulates TLR4 and IRAK-3 in a non-infectious, ventilator-induced lung injury model. *Respir Res* 11:27
31. Kuipers MT, Aslami H, Janczy JR, van der Sluijs KF, Vlaar AP, Wolthuis EK, Choi G, Roelofs JJ, Flavell RA, Sutterwala FS, Bresser P, Leemans JC, van der Poll T, Schultz MJ, Wieland CW (2012) Ventilator-induced lung injury is mediated by the NLRP3 inflammasome. *Anesthesiology* 116:1104–1115
32. Gasse P, Riteau N, Charron S, Girre S, Fick L, Petrilli V, Tschopp J, Lagente V, Quesniaux VF, Ryffel B, Couillin I (2009) Uric acid is a danger signal activating NALP3 inflammasome in lung injury inflammation and fibrosis. *Am J Respir Crit Care Med* 179:903–913
33. Kuipers MT, Aslami H, Vlaar AP, Juffermans NP, Tuip-de Boer AM, Hegeman MA, Jongsma G, Roelofs JJ, van der Poll T, Schultz MJ, Wieland CW (2012) Pre-treatment with allopurinol or uricase attenuates barrier dysfunction but not inflammation during murine ventilator-induced lung injury. *PLoS One* 7:e50559
34. Miller LS, O'Connell RM, Gutierrez MA, Pietras EM, Shahangian A, Gross CE, Thirumala A, Cheung AL, Cheng G, Modlin RL (2006) MyD88 mediates neutrophil recruitment initiated by IL-1R but not TLR2 activation in immunity against *Staphylococcus aureus*. *Immunity* 24:79–91
35. van de Veerdonk FL, Netea MG, Dinarello CA, Joosten LA (2011) Inflammasome activation and IL-1beta and IL-18 processing during infection. *Trends Immunol* 32:110–116
36. Cote CK, Van Rooijen N, Welkos SL (2006) Roles of macrophages and neutrophils in the early host response to *Bacillus anthracis* spores in a mouse model of infection. *Infect Immun* 74:469–480
37. Cirioni O, Ghiselli R, Tomasinsig L, Orlando F, Silvestri C, Skerlavaj B, Riva A, Rocchi M, Saba V, Zanetti M, Scalise G, Giacometti A (2008) Efficacy of LL-37 and granulocyte colony-stimulating factor in a neutropenic murine sepsis due to *Pseudomonas aeruginosa*. *Shock* 30:443–448
38. Becker JC, Schrama D (2013) The dark side of cyclophosphamide: cyclophosphamide-mediated ablation of regulatory T cells. *J Invest Dermatol* 133:1462–1465
39. Venhoff N, Effelsberg NM, Salzer U, Warnatz K, Peter HH, Lebrecht D, Schlesier M, Voll RE, Thiel J (2012) Impact of rituximab on immunoglobulin concentrations and B cell numbers after cyclophosphamide treatment in patients with ANCA-associated vasculitides. *PLoS One* 7:e37626
40. Duranton J, Bieth JG (2003) Inhibition of proteinase 3 by [alpha]1-antitrypsin in vitro predicts very fast inhibition in vivo. *Am J Respir Cell Mol Biol* 29:57–61

doi:10.1186/2197-425X-1-8

Cite this article as: Timmermans *et al.*: IL-1 β processing in mechanical ventilation-induced inflammation is dependent on neutrophil factors rather than caspase-1. *Intensive Care Medicine Experimental* 2013 1:8.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
