

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://repository.ubn.ru.nl/handle/2066/127586>

Please be advised that this information was generated on 2019-12-08 and may be subject to change.



Concise Review: Cell-Based Strategies in Bone Tissue Engineering and Regenerative Medicine

JINLING MA,^{a,b} SANNE K. BOTH,^b FANG YANG,^b FU-ZHAI CUI,^c JULI PAN,^{a,d} GERT J. MEIJER,^e
JOHN A. JANSEN,^b JEROEN J.J.P. VAN DEN BEUCKEN^b

Key Words. Mesenchymal stem cells • Endothelial cells • Bone marrow stromal cells • Adipose stem cells • Vascularization • Tissue regeneration

ABSTRACT

Cellular strategies play an important role in bone tissue engineering and regenerative medicine (BTE/RM). Variability in cell culture procedures (e.g., cell types, cell isolation and expansion, cell seeding methods, and preculture conditions before *in vivo* implantation) may influence experimental outcome. Meanwhile, outcomes from initial clinical trials are far behind those of animal studies, which is suggested to be related to insufficient nutrient and oxygen supply inside the BTE/RM constructs as some complex clinical implementations require bone regeneration in too large a quantity. Coculture strategies, in which angiogenic cells are introduced into osteogenic cell cultures, might provide a solution for improving vascularization and hence increasing bone formation for cell-based constructs. So far, preclinical studies have demonstrated that cell-based tissue-engineered constructs generally induce more bone formation compared with acellular constructs. Further, cocultures have been shown to enhance vascularization and bone formation compared with monocultures. However, translational efficacy from animal studies to clinical use requires improvement, and the role implanted cells play in clinical bone regeneration needs to be further elucidated. In view of this, the present review provides an overview of the critical procedures during *in vitro* and *in vivo* phases for cell-based strategies (both monoculture and coculture) in BTE/RM to achieve more standardized culture conditions for future studies, and hence enhance bone formation. *STEM CELLS TRANSLATIONAL MEDICINE* 2014;3:98–107

INTRODUCTION

Bone is one of the most transplanted tissues, with more than 2.2 million bone graft procedures being performed annually worldwide [1]. Bone tissue engineering/regenerative medicine (BTE/RM) approaches, with the triad principle of applying combinations of the three building blocks: supporting scaffolds, growth factors, and functionally active cells to (re)generate biologically functional tissues, have been suggested as promising strategies to regenerate bone [2].

The potential of BTE/RM constructs becomes especially challenging under compromised conditions, such as in elderly patients with suboptimal medical conditions (e.g., osteoporosis, diabetes, and cancer), or in cases in which the bone defect dimensions are (far) beyond those that can spontaneously heal. Consensus on the difficulty of healing bone defects under such conditions illuminates that the bone regenerative capacity arising from only a scaffold material is often insufficient, and that additional BTE/RM approaches should arise from preseeding the scaffold with cells or incorporating growth factors within the scaffolds. Small successes have been reported for *in vitro* experiments and even animal studies with cell-laden scaffolds, but translation of these results

to the clinic for bone regenerative applications has been insignificant so far [3]. Several issues can be attributed to the lack of this clinical success. First, the quality and quantity of the used cells and the preculture conditions after cell seeding onto the scaffolds are variable and limited, and tiny variations within these procedures may substantially influence the outcome. Second, cells within a construct are subjected to inflammatory conditions and limited nutrient supply on implantation because surgical intervention generates tissue damage and the diffusion of nutrients and oxygen from the nearest capillary is limited to only 150 μm to 200 μm [4]. Researchers have pointed out that rapid vascularization into cell-based BTE/RM-constructs is pivotal to clinical success [3]. From a cellular point of view, the solution for insufficient vascularization is either coculture of osteogenic cells with angiogenic cells [5] or changing the priming differentiation pathway of stem cells (SCs) from osteogenic to chondrogenic, because cartilage is an avascular tissue with less susceptibility to limited vascularization [6].

The aim of this review is to summarize the current state-of-the-art in cell-based BTE/RM in terms of critical procedures and efficacy of monoculture (osteogenic) and coculture approaches.

^aDepartment of VIP Service and ^dDepartment of Oral and Maxillofacial Surgery, Beijing Stomatological Hospital, Capital Medical University, Beijing, China; ^bDepartment of Biomaterials and ^eDepartment of Oral and Maxillofacial Surgery, Radboud University Medical Center, Nijmegen, The Netherlands; ^cState Key Laboratory of New Ceramics and Fine Processing, Department of Materials Science & Engineering, Tsinghua University, Beijing, China

Correspondence: Jeroen J.J.P. van den Beucken, Ph.D., Department of Biomaterials (309), Radboud University Medical Center, Ph v LeijdenIn 25, 6525 EX Nijmegen, The Netherlands. Telephone: 312-436-67305; E-Mail: jeroen.vandenbeucken@radboudumc.nl

Received July 12, 2013; accepted for publication August 30, 2013; first published online in *SCTM EXPRESS* December 3, 2013.

©AlphaMed Press
1066-5099/2013/\$20.00/0

[http://dx.doi.org/
10.5966/sctm.2013-0126](http://dx.doi.org/10.5966/sctm.2013-0126)

Although the authors are aware of the major importance of scaffold properties and the potential of growth factor incorporation and release, the intention is to focus on the cellular component in BTE/RM approaches, and hence critically review the experimental, preclinical, and clinical efforts on this topic.

CRITICAL PROCEDURES

To restore bone defects in clinical applications, some critical issues that are inherently related to the cell quality and quantity (e.g., cell types/sources, cell isolation, and yield), cell seeding efficiency and preculture conditions must be considered, and finally in vivo conditions should be taken into consideration.

Stem Cell Sources

A source of human cells that can be derived in large numbers from a small and easy initial harvest and can differentiate into bone-forming cells is preferable for cell-based BTE/RM constructs [7]. Various cell types have been explored for BTE/RM, each with its own potential and premise.

Nonadult SCs

Nonadult SCs contain two categories: embryonic stem cells (ESCs) and SCs isolated from perinatal tissue, such as aborted fetal tissue and discarded tissue at birth (e.g., umbilical cord and placenta). ESCs are pluripotent, but consistency on bone formation capacity by ESCs progeny has not been achieved [8, 9] and ethical issues exist. The latter category, positioned between embryonic and adult SCs, is multipotent SCs. These cells have similar bone formation capacity compared with adult mesenchymal SCs (MSCs) [10]. Nevertheless, that they can form tumors [11] on in vivo implantation makes the use of these cell types controversial.

Adult SCs

Adult MSCs play a predominant role in the field of BTE/RM. The most common sources are bone marrow (BM), adipose tissue (AT), and dental pulp (DP).

BM-MSCs

BM is the predominant source for adult MSCs. BM-derived MSCs (BM-MSCs) have multipotential differentiation capacity, including osteogenic potential, and can perform pericyte-like functions by secreting both angiogenic and stabilization factors in the process of vessel formation [12]. Although BM-MSCs are widely used in laboratory and preclinical studies, they have several disadvantages, including donor site morbidity and side effects, limited proliferation capacity, and inferior differentiation potential in aged individuals [13]. Thus, alternatives to BM-MSCs have been explored.

AT-MSCs

AT is a popular alternative source for MSCs because of easy and less invasive harvest procedures and larger yield compared with BM [14]. AT-derived MSCs (AT-MSCs) are similar to BM-MSCs regarding gene expression and osteogenic capacity [15] and can also exert pericyte-like functions [16]. Nevertheless, the bone-forming capacity of AT-MSCs needs further confirmation, and whether AT-MSCs exhibit bone-forming capacity similar to that of BM-MSCs is still controversial. Moreover, preclinical safety

and efficacy as well as long-term in vivo studies are required before AT-MSCs can be evaluated clinically.

DP-MSCs

The noninvasive manner of obtaining DP from deciduous/extracted teeth or even nonextracted crown fractured teeth makes it an ideal source for MSCs. DP-derived MSCs (DP-MSCs) have similar gene expression, a faster proliferation rate, and a higher percentage of SCs in the harvested population compared with BM-MSCs [17] and express pericyte markers [18]. Some researchers have reported that DP-MSCs have at least equal bone-forming capacity compared with BM-MSCs [19, 20], and others showed only the formation of connective tissue [21] or dentin pulp-like complex formation [22] from DP-MSCs, suggesting the requirement for in-depth studies on the mechanism of bone-forming capacity by DP-MSCs.

Isolation and Expansion

High-quality cells in relevant quantities are often crucial to satisfy clinical demand (i.e., the successful restoration of a bone defect). To achieve this, effective cell isolation and expansion of harvested cells are of utmost importance. Inadequate isolation methods can lead to polluted isolates and hence inconsistency in cell marker expression [23]. Further, long-term expansion has demonstrated decreased stem cell proliferation and differentiation capacity [24]. Nevertheless, a broadly accepted protocol has not been established for isolation and large-scale expansion of human MSCs, which also makes the comparison among reported results from publicly available databanks difficult.

Isolation of BM-MSCs, AT-MSCs, and DP-MSCs

The isolation of MSCs is a general procedure, meaning tissue harvesting and treatment (mincing and/or enzymatic digestion), washing, filtering and centrifugation, and plating. BM-MSCs can be obtained from either bone chips (cortical or trabecular bone) or BM. The SCs isolated from these two sources have been shown to be identical in terms of phenotype and multilineage differentiation capacity [25]. AT-MSCs can be isolated from a resected adipose block or lipoaspirate. The latter method is easier because lipoaspirate consists of finely minced fat fragments with less volume and a more homogeneous population of cells. Isolation of MSCs from DP basically relies on two methods. The first one is the outgrowth method (i.e., cell migration out of the pulp fragments and adherence to cell culture plates), which is rarely applied because it takes substantially longer to obtain similar cell numbers compared with enzymatic digestion. This latter method is comparable to the earlier mentioned digestion method for BM-MSCs (Fig. 1). Bone chips, resected AT, and DP tissue require mincing, whereas BM and lipoaspirate proceed directly to a washing procedure. After washing, enzymatic digestion is performed for all the tissues except BM aspirates followed by filtering and gradient density centrifugation. The isolated cells are plated for adhesion-dependent selection. To increase the purity of the isolated cells, cell separation methods such as cell sorting using cell surface markers (e.g., CD73, CD105, and Stro-1) are frequently used (Fig. 1).

Expansion of BM-MSCs, AT-MSCs, and DP-MSCs

For expansion, essential culture parameters comprise the type of basic culture medium and the nutritional source (e.g., serum), cell passaging density, and doubling numbers. α -minimum essential




Figure 1. Scheme for isolation of mesenchymal stem cells derived from bone, adipose tissue, and dental pulp. Abbreviations: FACS, fluorescence activated cell sorting; MACS, magnetic activated cell sorting.

medium (α -MEM) is the optimal bare culture medium for isolation and expansion of human BM-MSCs [26], AT-MSCs [27, 28], and DP-MSCs [29, 30]. Human serum [31] or human platelet lysate (PL) [32–34] has been explored as a replacement for fetal bovine serum (FBS) because FBS carries potential hazards related to bovine pathogens and immunological issues. No consensus regarding optimal cell-seeding density for expansion exists, although low-density cell seeding has been recommended (e.g., for BM-MSCs, 50–100, or 1,000 cells per cm^2 for expansion; for AT-MSCs, 100–200 cells per cm^2 , and for DP-MSCs, 800–1,000 cells per cm^2) because this is associated with higher proliferation rates [26, 35, 36].

Small variations within each step of an isolation procedure, including enzyme type and enzymatic digestion time, centrifugation speed and time, or washing liquid (either phosphate-buffered saline or sodium saline), may change the quality and quantity of the isolated SCs and hence influence the experimental outcomes. Additionally, the site of tissue harvest can influence the phenotype of the isolated SCs [37], the percentage of SCs in the whole population [38], and their osteogenic differentiation capacity [39]. However, this review did not reach any significant conclusion about standardized and optimized SC isolation and expansion procedures, which sheds light on the need for relative studies.

Cell-Seeding Techniques and Preculture on Scaffolds

Cell seeding and preculture on scaffolds are essential procedures before *in vivo* implantation. Seeding requirements for cell-based BTE/RM constructs for potential clinical use should allow maximized utilization of donor cells, minimal time for anchorage-dependent and shear-sensitive cells (e.g., osteoblasts) in suspension, and spatially uniform distribution of attached cells [40].

Cell-Seeding Efficiency

Cell-seeding efficiency, which might further correlate with bone-forming capacity, can be increased by either selecting proper scaffolds and modifying the scaffold surface or by optimizing cell-seeding methods. For the former method, the criteria for three-dimensional scaffolds are explored in tandem with properties such as porosity, interconnectivity, biodegradability, and mechanical integrity. As such, diverse forms of scaffolds (e.g., bulk vs. hydrogels; fibrous vs. foam) with different components (e.g., polymers, ceramics, and metal) have been explored. Various scaffold properties affect cell-seeding efficiency; for instance, scaffolds that have more regular and homogeneous pores and are more

accessible for the cell suspension during drop seeding have higher cell-seeding efficiency (e.g., a fiber deposited vs. a foam scaffold) because they avoid cell aggregate entrapment in the small and irregular internal pores [41]. Furthermore, scaffold surfaces can be modified through pretreatment with adhesive proteins (e.g., preimmersion in serum or fibronectin). For the latter method, cell-seeding volume and cell-seeding time affect cell-seeding efficacy as well as cell viability significantly [41]. Moreover, the combination of low pressure (vacuum) with vibration can help remove potential air bubbles around scaffolds and allow more cells to penetrate deeply into the scaffolds to enhance bone formation, especially for porous scaffolds, in comparison with the commonly used static seeding methods [42]. Additionally, dynamic cell seeding (e.g., perfusion bioreactor) has shown to yield higher cell-seeding efficiencies and more homogenous cell distribution compared with static cell seeding [43].

Cell-Seeding Density and Medium Perfusion

Cell proliferation after cell seeding is mainly regulated by contact-inhibition between adjacent cells, which is determined by cell seeding density [44] and nutrient transfer efficiency as a result of the medium perfusion rate, which also determines cell viability [45].

Initial seeding density can alter the expression of osteogenic genes by controlling the distance of paracrine signals among cells. Although no systematic studies on the optimal cell-seeding density are available because of the variety of scaffold properties (Table 1), it has been agreed that a certain threshold of cell density is essential to achieve successful bone regeneration *in vivo*. Low seeding densities may compromise cellular contact and hence influence bone formation, and high seeding densities do not necessarily benefit cell behavior because the overloaded cells may result in limited nutrient transport and insufficient waste removal from the internal structures [46]. In general, below a certain threshold (i.e., the optimal density), osteogenic marker expression and extracellular matrix production (i.e., mineralization) capacities are enhanced with increasing cell-seeding density. In contrast, when the cell-seeding density exceeds this optimal density, a further increase in cell-seeding density can reduce the bone regenerative capacity of cells [46].

Medium perfusion is increased in dynamic cultures using bioreactor systems, leading to an improved nutrient delivery and mechanical stimulation to the cells as well as enhanced osteogenic differentiation compared with static cultures [47]. Three classes of bioreactor systems have been widely used in BTE: spinner flasks, rotating wall, and perfusion systems. Each of these has proven effective to culture cells for BTE. Perfusion systems have been shown to perfuse media effectively throughout the scaffold rather than creating only a homogenous media solution on the exterior of a scaffold [48].

Predifferentiation of Cells

Predifferentiation of the cells into osteogenic [49, 50] or chondrogenic lineage [6, 51] before *in vivo* implantation has been assumed essential for enhancing osteogenesis *in vitro* and *in vivo* in comparison with nonpredifferentiated (i.e., osteogenic or chondrogenic) cells. However, the optimal preculture time for bone regeneration is either controversial (for osteogenic differentiation) or lacking (for chondrogenic differentiation). Short osteogenic induction is probably insufficient for inducing cell

Table 1. In vivo preclinical studies involving human mesenchymal stem cells transplantation (monocultures) in BTE/RM

Cell sources	Scaffold	Seeding density	In vitro preculture time	In vivo time (max)	Species and repair sites	Evaluation method	Efficacy of bone formation (Fold change, cell seeded/cell-free)	Reference
BM-MSCs	Mineralized collagen	20x10 ⁶ /construct (5 cm ³)	1 day PM	24 weeks	Sheep tibia	Histology	Bone percentage (%): 18.0/13.0 = 1.4	[77]
BM-MSCs	Chitosan	1x10 ⁶ /construct (5 mm Ø, 1 mm H)	2 weeks OM	8 weeks	Mouse calvarium	Micro-CT	More bone formation in cell-seeded constructs: (No quantitative analysis)	[78]
BM-MSCs	PCL	3x10 ⁶ /construct (5 mm Ø, 9 mm H)	1 day PM	12 weeks	Rat femur	Micro-CT	Bone volume (mm ³): 25.0/10.0 = 2.5	[10]
BM-MSCs	PCL/β-TCP	1x10 ⁶ /construct (4 mm Ø, 8 mm H)	1 day PM	3 weeks	Rat femur	Micro-CT	Bone volume (mm ³): 1.0/0.3 = 3.3	[79]
BM-MSCs	Hyaluronic acid	1x10 ⁶ /construct	1 day PM	4 weeks	Rat calvarium	Histology	Bone percentage (%): 61.0/16.0 = 3.8	[14]
BM-MSCs	PLGA	0.5x10 ⁶ /construct	9 days in PM +14 days in OM	20 weeks	Rat calvarium	Histology	Bone percentage (%): 54.0/20.0 = 2.7	[80]
BM-MSCs; AT-MSCs	Collagen	Not clear	Not clear	8 weeks	Rat calvarium	X-ray	Bone mineral density percentage (%): 100.0/60.0 = 1.7 (for both BM-MSCs and AT-MSCs)	[81]
AT-MSCs	Collagen; PLGA; HA	1x10 ⁶ /construct (size is unclear)	8-12 hours PM	18 weeks	Rat calvarium	Histology	Bone mineral density (g/cm ³): 4.0 (collagen); 1.6 (PLGA); 2.0 (HA)	[82]
AT-MSCs	PLGA	0.5x10 ⁶ /construct (8 mm Ø, 0.8 mm H)	2 weeks OM	12 weeks	Rat calvarium	Histology	Bone percentage (%): 72.0/38.0 = 1.9	[54]
AT-MSCs	Ti	1x10 ⁶ /construct (5 mm Ø)	1 week OM	8 weeks	Rat calvarium	Histology	Bone percentage (%): 151.0/57.0 = 2.6	[60]
AT-MSCs	PLGA	0.15x10 ⁶ /construct (size is unclear)	1 day PM	8 weeks	Mouse calvarium	Histology	Bone percentage (%): 100.0/42.0 = 2.4	[83]
AT-MSCs	BCP	75x10 ⁶ /ml suspension	1 day PM	16 weeks	Rat femur	Histology	Bone percentage (%): 89.0/40.0 = 2.2	[84]
DP-MSCs	Collagen	1x10 ⁶ /construct (13 mm Ø, 1.5 mm H)	10 days OM	6 weeks	Rat parietal bone	Histology	Bone percentage (%): 69.0/38.0 = 1.8	[85]

Abbreviations: AT, adipose tissue; β-TCP, β-tricalcium phosphate; BCP, biphasic calcium phosphate; BM, bone marrow; BTE, bone tissue engineering; CT, computed tomography; DP, dental pulp; H, height; HA, hydroxyapatite; Ø, diameter; OM, osteogenic medium; PCL, polycaprolactone; PLGA, poly(lactic-co-glycolic acid); PM, proliferation medium; RM, regenerative medicine; Ti, titanium. Search conditions for Table 1:

For human BM-MSCs:

- (i) (human bone marrow mesenchymal stem cell* OR human bone marrow derived mesenchymal stem cell* OR human bone marrow mesenchymal stromal cell* OR human multipotential stromal cell*) AND (bone tissue engineering OR bone regenerat* OR bone form*) AND (spinal OR cranial OR cavariar OR femoral OR radial OR mandible OR maxilla OR tibia), and
- (ii) (human bone marrow mesenchymal stem cell* OR human bone marrow derived mesenchymal stem cell* OR human bone marrow mesenchymal stromal cell* OR human multipotential stromal cell*) AND (bone tissue engineering OR bone regenerat* OR bone form*) AND (animal* OR in vivo).

For human AT-MSCs:

- (i) (human fat derived mesenchymal stem cell* OR human fat tissue derived mesenchymal stem cell* OR human adipose tissue derived mesenchymal stem cell* OR human lipoaspiration) AND (spinal OR cranial OR cavariar OR femoral OR radial OR jaw OR mandible OR maxilla OR tibia), and
- (ii) (human fat derived mesenchymal stem cell* OR human fat tissue derived mesenchymal stem cell* OR human adipose tissue derived mesenchymal stem cell* OR human lipoaspiration) AND (animal* OR in vivo).

For human DP-MSCs:

- (i) (human dental pulp mesenchymal stem cell* OR human dental pulp derived mesenchymal stem cell* OR human dental pulp stem cell*) AND (spinal OR cranial OR cavariar OR femoral OR radial OR jaw OR mandible OR maxilla OR tibia), and
- (ii) (human dental pulp mesenchymal stem cell* OR human dental pulp derived mesenchymal stem cell* OR human dental pulp stem cell*) AND (animal* OR in vivo).

differentiation, whereas long duration osteogenic induction leads to an apoptotic process [52]. Longer preculture time (2 weeks) has been shown to induce more bone formation compared with shorter time (1 week) for both human BM-MSCs [53] and human AT-MSCs [54]. From a clinical point of view, one-step surgical techniques, in which freshly isolated cells (without expansion) are used directly within the operation theater, are recommended for clinical applications of cell-based bone regenerative strategies [55]. This suggests that a balance is needed between the bone regeneration efficacy caused by the relatively longer preculture time and the benefits that patients gain from the one-step surgery.

The aforementioned seeding techniques and preculture conditions (for monocultures) are also used for cocultures. Additional parameters should be considered for cocultures, including choice of cell origin and cell type, culture medium, and cell ratio. The effects of origin of either MSCs (e.g., BM-MSCs or AT-MSCs) or angiogenic cells (e.g., ECs, endothelial progenitor cells [EPCs]) need further investigation. Furthermore, it should be noted that cellular communication in cocultures is cell-type specific. For instance, when ECs were replaced by primary chondrocytes or fibroblasts in osteoblasts/ECs coculture, no increasing effects on alkaline phosphate activity were observed [56]. The culture medium used in cocultures is different depending on the research aims (i.e., angiogenesis or osteogenesis). EC culture medium (EM) has been shown to be more suitable for vessel formation and stabilization, whereas osteogenic medium (OM) has been shown to favor osteogenic differentiation of MSCs [57]. It was reported recently that more bone formation was observed for cells cocultured continuously in OM compared with sequential culture medium variations (e.g., first culturing cells in EM and subsequently in OM) before *in vivo* implantation [58]. Further, it was recently shown that BM-MSCs/ECs cultured in OM achieved higher mineralization while maintaining angiogenic capacity *in vitro* [5]. This indicates that OM may be a more optimal culture medium for cocultures for future research on bone regeneration capacity of such cell-based constructs. A cell ratio of 1:1 is often chosen in cocultures, which was also demonstrated recently as optimal for both BM-MSC/EC [5] and AT-MSC/EC [59] cocultures.

One of the crucial aspects of BTE/RM is the evaluation/prediction of bone healing capacity for the tissue-engineered constructs. To further test such bone-forming efficacy, *in vivo* studies including both preclinical research and clinical trials are necessary.

Preclinical Studies

Monocultures

Because MSCs from humans (rather than other animal cell source) are the cell type essential for future clinical application, and orthotopic implantation (i.e., bony defects) is the preferred model to mimic the clinical situation compared with ectopic implantation sites, a literature review on preclinical studies using BM-MSCs, AT-MSCs, and DP-MSCs of human origin for repair of bony defects was performed in PubMed. A total of 13 such preclinical studies were available, among which 7 included BM-MSCs, 6 included AT-MSCs (including one paper studying both BM-MSCs and AT-MSCs), and 1 included DP-MSCs (Table 1). For human BM-MSCs, three studies used rat cranial defects, one used mouse cranial defects, one used sheep tibial defects, and two used rat femoral defects. For human AT-MSCs, four studies used rat cranial defects, one used mouse cranial defects, and one used rat femoral defects. For human DP-MSCs, the only

available study used rat parietal bone defects. Cell-based constructs demonstrated more efficacy in restoring bone defects in comparison with cell-free constructs, with a fold range of 1.4 to 3.8, 1.6 to 4.0, and 1.8 compared with cell-free control constructs for BM-MSCs, AT-MSCs, and DP-MSCs, respectively.

Cocultures

Although animal studies confirmed the capability of cell-based constructs to form bone and integrate with the host tissues, the repair efficacy could be more promising once the key hurdle of insufficient vascularization can be overcome, which would require clinically sized, fully viable bone grafts to be engineered and implanted. A 50% loss of viable cells in monoculture cell-based constructs, as shown by the presence of apoptotic cells, was observed within two days after implantation *in vivo* [60], most likely because cells encountered an ischemic and inflammatory environment [3]. In view of this, coculture approaches have shown advantages in both vascularization and bone formation compared with monocultures (Table 2). Currently, six preclinical studies have been reported on orthotopic implantation using tissue-engineered constructs seeded with cocultures using human MSCs (i.e., four for BM-MSCs, two for AT-MSCs, and none for DP-MSCs) (Table 2). For human BM-MSCs, one study used rat cranial defects, two used rat femoral defects, and one used rabbit ulna defects. These studies demonstrated a range fold of bone-forming efficacy of 1.2 to 1.4 (cocultures compared with MSCs monocultures) and vessel-forming efficacy of 1.0 to 10.0 (compared with MSCs monocultures) or 2.5 to 2.7 (compared with ECs/EPCs monocultures). For human AT-MSCs, the studies demonstrated a range fold of bone-forming efficacy of 0.3 to 1.3 (cocultures compared with MSCs monocultures). Although cocultures of human DP-MSCs with ECs have been shown to enhance osteogenic and angiogenic potential *in vitro* [61], *in vivo* studies using DP-MSCs in coculture approaches are still lacking.

Fate of the Implanted Cells

After *in vivo* implantation, the fate/contribution of the seeded cells (both MSCs and ECs/EPCs) and the possible mechanisms of their bone-forming (for MSCs) or vessel-forming (for ECs/EPCs) capacity have been explored. The seeded MSCs can proliferate and differentiate into osteogenic lineages to contribute directly to bone formation [62, 63] and have the potential to recruit cells from the host, which can be MSCs [64] or ECs/EPCs [64, 65], to induce/enhance bone formation indirectly. Similarly, implanted EPCs may also act in this way on bone formation by either forming stable vasculature through coculturing with MSCs [66] or by secreting chemotactic factors (e.g., vascular endothelial growth factor) to recruit ECs/EPCs from the host and thereby stimulate vascularization in bone defects [67]. Several studies have shown the synergistic contribution to bone formation [68] as well as vessel formation [69] of both donor and host cells. Most of the studies demonstrated that bone [14, 70] and blood vessel formation [69, 71] were mainly from the implanted donor cells.

Clinical Studies

MSCs have been used in clinical trials in various fields of RM such as treatment of graft versus host disease, diabetes mellitus, amyotrophic lateral sclerosis, and liver failure. However, to our knowledge, human clinical trials addressing cell-based constructs for bone repair are limited to case reports (Table 3), which

Table 2. In vivo preclinical studies involving human MSC transplantation (cocultures) in BTE/RM

Cell sources (cell ratio and medium)	Scaffold	Seeding density	In vitro preculture time	In vivo time (max)	Species and repair sites	Evaluation method	Efficacy of bone and vessel formation (fold change, coculture/monoculture)
BM-MSCs/ECs (not clear, PM)	PLGA	1×10^6 /construct (8.5 mm \varnothing , 2 mm H)	1 hr	12 wk	Rat calvarium	Histology	Bone percentage (%): 33.0/23.0 (MSCs) = 1.4; donor-derived vessels percentage (%): 2.0/0.2 (MSCs) = 10.0 [86]
BM-MSCs/EPCs (1:1, PM)	β -TCP	0.5×10^6 /construct (granule, 0.7–1.4 mm)	0 d	8 wk	Rat femur	Histology	Bone percentage (%): 1.2/1.0 (MSCs) = 1.2; vessel density (/mm ²): 0.4/0.4 (MSCs) = 1.0 0.4/0.15 (EPCs) = 2.7 [87]
BM-MSCs/EPCs (1:1, PM)	β -TCP	0.5×10^6 /construct (granule, 0.7–1.4 mm)	0 d	1 wk	Rat femur	Histology	New bone mass: 7.2/5.4 (MSCs) = 1.3; vessel density (/mm ²): 3.2/0.45 (MSCs) = 7.1 3.2/1.3 (EPCs) = 2.5 [88]
BM-MSCs/ECs (1:1, OM)	β -TCP	1.0×10^6 /construct (3.5 x 2 x 15mm)	2 wk	16 wk	Rabbit ulna	Histology	Bone percentage (%): 37.0/29.0 (MSCs) = 1.3; vessel percentage (%): 5.9/4.9 (MSCs) = 1.2 [89]
AT-MSCs/ECs (2:1, OM)	PCL/PLGA/ β -TCP	0.3×10^6 /construct (8 mm \varnothing , 2 mm H)	2 d	12 wk	Rat calvarium	Micro-CT	Bone percentage (%): 74.0/58.0 (MSCs) = 1.3 [90]
AT-MSCs/ECs (1:1, OM)	Ti	1×10^6 /construct (5 mm \varnothing)	1 wk	8 wk	Rat calvarium	Histology	Bone percentage (%): 44.0/151.0 (MSCs) = 0.3 [60]

Search conditions for Table 2:

Based on the search conditions from Table 1, additional search condition of (coculture* OR co-culture* OR co-seed* OR coseed* OR cotransplant* OR co-transplant* OR co-implant*) OR (endothelial cell* OR endothelial progenitor cell*) was added.

Abbreviations: AT, adipose tissue; BM, bone marrow; BTE, bone tissue engineering; ECs, endothelial cells; EPCs, endothelial progenitor cells; H, height; MSC, mesenchymal stem cells; \varnothing , diameter; OM, osteogenic medium; PCL, polycaprolactone; PLGA, poly(lactic-co-glycolic acid); PM, proliferation medium; RM, regenerative medicine; TCP, tricalcium phosphate; Ti, titanium.

emphasizes the need for randomized control trials and systematic clinical studies. The available case reports cover long bone defects, jaw defects, alveolar cleft regeneration, and sinus augmentation. A search on clinicaltrials.gov using a combination of search terms revealed 24 relevant studies. Eight studies included the terms “bone marrow derived mesenchymal stem cells” AND “bone.” Two studies included the terms “adipose tissue derived mesenchymal stem cells” AND “bone.” No studies included the terms “dental pulp stem cells” AND “bone.” Five studies included the terms “mesenchymal stromal cells” AND “bone.” Two studies included the terms “mesenchymal stromal cells” AND “bone regeneration,” and seven studies included the terms “stem cells” AND “bone regeneration.”

These studies are dedicated to the treatment of bone defects and diseases, such as bone cysts, osteoarthritis, osteonecrosis, spinal fusion, and atrophic nonunion fractures using human MSC monocultures. Some of these studies are still recruiting patients. To date, BM-MSCs are still the main source for clinical trials as well as case reports, and reports on the use of coculture-based constructs for clinical bone regeneration are lacking. Clinical studies conducted so far have demonstrated that it is safe to use human MSCs for bone regeneration, but the outcomes are substantially less promising compared with those of animal studies. In two patients, only 34.5% and 25.6% of bone regeneration (of the original defect volume) was found after four months when BM-MSC-loaded demineralized bone matrix was used to restore human alveolar cleft defects [72]. Meijer et al. observed bone regeneration in 50% (3/6) of patients, and bone formation was induced in only one patient by the tissue-engineered

construct when BM-MSCs seeded β -tricalcium phosphate granules were implanted in jaw defects. A remarkable finding of this study was that despite the low number of construct contributions to clinical bone formation, parallel ectopic implantations with the constructs in nude mice showed bone formation in all specimens [73]. The discrepancy between preclinical results and clinical outcomes is likely explained by several causes. As mentioned, because of the unstandardized cell preparation protocol, the cell populations used are often heterogeneous, which makes comparison among outcomes from different studies difficult. In addition, the long-term survival and proliferation of implanted cells and to what extent these cells can contribute to bone formation remain largely unknown, especially in critical sized bone defects. In view of this, cell tracing such as luciferase labeling enables real time in vivo evaluation of the cells, which is particularly interesting to determine the contribution and distribution of the implanted cells. Finally, the comparison of bone regeneration mechanisms between animals and human beings at a molecular level is necessary to understand the underlying reasons further.

CONCLUSION

Bone regeneration in clinics is compromised in some cases, for example, in patients with typical risk factors like smoking or diabetes, and the incidence of these risk factors is increasing from 5% to 10% (for a healthy population) up to 40% [74]. Cell-based BTE/RM has emerged as an attractive approach for bone regeneration in preclinical studies, although no definite answer about its use has been given for clinical studies. A standardized method

Table 3. Clinical studies using human MSC-based tissue-engineered constructs

Cell sources and treatment	Scaffold	Number of patients	Repair sites	Evaluation method(s)	Reported outcomes
BM-MSCs	HA	3	Long bone defects (1 tibia, 1 ulna, 1 humerus)	X-rays and micro-CT scans	Abundant callus formation along the implants and good integration at the interfaces with the host bones 2 months after surgery [91]
BM-MSCs	HA	4	Long bone (limb)	X-rays and micro-CT scans	No major complications occurred; complete integration between implants and host bone 5–7 months and maintained 6–7 years after surgery [92]
BM-MSCs (with PRP)	Ti	1	Alveolar cleft	Micro-CT scans	Regenerated bone bridged the cleft after 6 months [93]
BM-MSCs	DBM/Calcium sulphate	2	Alveolar cleft	Micro-CT scans	There is 34.5% and 25.6% integrity of the bone defects for the two patients [73]
BM-MSCs (with fibrin glue)	Ti	1	Mandible	Dental CT scans	Successful bone regeneration in large segmental defects of the jaw [94]
BM-MSCs	BCP	7	Sinus augmentation	X-rays and biopsy	The mean percentage of the newly formed bone was 41%; bone height was 11 mm after 1 year compared with the initial height of 2 mm, with a successful rate of 93% (28/30) for dental implants [95]
BM-MSCs	HA (particles)	6	Jaw	X-rays and biopsy	Bone regeneration was observed in 50% (3/6) of the patients, and bone formation was induced by the tissue-engineered construct in only 1 patient [96]
BM-MSCs	Allogenic mandible	3	Mandible	Micro-CT scans	Bone healing was observed in 2 of the 3 patients [97]
AT-MSCs (with BMP-2)	β -TCP	1	Maxilla	X-rays	Regenerated bone resembled mature maxillary bone 2 months after surgery [98]
DP-MSCs	Collagen	17	Mandible	X-rays and biopsy	Bone regeneration in the defects was complete and stable 1 year after surgery [99]

Search conditions for Table 3:

(human mesenchymal stem cell* OR human stem cell* OR human mesenchymal stromal cell* OR human marrow stromal cell*) AND (clinic* OR case report* OR clinical trial*) AND (bone tissue engineering OR bone regenerat* OR bone form*).

Abbreviations: AT, adipose tissue; BCP, biphasic calcium phosphate; BM, bone marrow; BMP, bone morphogenetic protein; CT, computed tomography; DBM, demineralized bone matrix; DP, dental pulp; HA, hydroxyapatite; MSC, mesenchymal stem cells; PRP, platelet-rich-plasma; TCP, tricalcium phosphate; Ti, titanium.

Table 4. Opportunities and threats for cell-based constructs in clinical application for BTE/RM

Opportunities	Threats
Various cell sources available with osteogenic capacity	Limited cell number for clinical cases in which there are large defects, or for one-step surgery
Through expansion, sufficient cell numbers can be obtained	Longer expansion may lead to cell deformation and decreased differentiation capacity
Efficient cell seeding on scaffolds can be achieved before in vivo implantation	Cell survival is compromised after in vivo implantation
Successful bone regeneration using cell-based constructs has been demonstrated in preclinical studies	There have been less promising outcomes of bone regeneration using cell-based constructs in clinics
In some clinical case reports using cell-based strategies for BTE/RM, bone regeneration was observed	Because of the lack of a control group in studies and case reports, the contribution of implanted cells cannot be ascertained

Abbreviations: BTE, bone tissue engineering; RM, regenerative medicine.

of using cell-based tissue engineering constructs in clinical work contains a protocolized cell isolation and culture method; valid characterization of the isolated cells, and efficient cell seeding with an optimized cell source, scaffold type, and cell seeding parameters (e.g., density, time, and volume). A proper predifferentiation method and time are also needed.

Based on the promising outcomes from preclinical studies, in which cell-based constructs (irrespective of monoculture and cocultures) enhanced bone formation compared with acellular constructs, and cocultures showed more pronounced vessel and

bone formation than monocultures, it is logical and straightforward to proceed the translational step from bench to bedside. Because of the limited number of systematic clinical studies and the lack of proven examination methods (e.g., biopsy), very few studies were able to demonstrate the efficacy of cell-based strategies for BTE/RM in a clinical setting. Therefore, a number of areas of ongoing active research are directly relevant to the translation of cell-based constructs for bone regeneration into clinics, including development of bioreactor systems for standardized and scalable cell expansion methods; scale up to clinically sized (large) bone

constructs with potential for further integration with host tissues; evaluation of engineered bone grafts in large animal models; rapid establishment of vascularization through the implanted bone graft; and marketing of off-the-shelf products. For the benefit of patients, intraoperative graft manufacture, also called one-step surgery, in which the graft is assembled during the surgical procedure, may be of interest. A prerequisite for this method is obtaining sufficient cell numbers for cell seeding and bone regeneration without expansion. Moreover, by using cocultures, the needed number of MSCs can be reduced while maintaining equal vascularization and bone formation.

As an overview, the opportunities and threats for cell-based tissue engineered constructs for clinical application in BTE/RM are listed in Table 4. For clinical studies, several aspects should be considered. There is a need for a reduction of the time lapse from cell isolation to in vivo implantation to avoid long waiting times and patient discomfort and ensure the multipotential differentiation capacity of MSCs, which can be lost during (long-term) expansion. Because of the immunomodulatory properties of MSCs [75], standardized off-the-shelf products containing allogeneic MSCs seem feasible. The contribution of implanted MSCs to the regenerative process is still unclear because of the combination of osteoinductive scaffolds and/or growth factors and the lack of control groups (e.g., bare scaffolds) in clinical trials. To determine the regeneration efficacy of cell-based constructs, quantitative measurements should be performed, whereas the current assessment for bone formation is mostly radiological examination (e.g., x-ray and micro-computed tomography). This makes it difficult to distinguish between newly formed bone and the implanted scaffolds, especially when radiopaque scaffolds are used. It is necessary to identify donors with appropriate

osteogenic capacity before in vivo implantation. There is no certain correlation between osteogenic capacity in vitro and bone formation in vivo [76], although researchers are seeking indices to help predict the bone formation capacity. Longer follow-up periods are recommended to evaluate the long-term safety and efficacy of human MSC-based tissue-engineered constructs in bone regeneration.

ACKNOWLEDGMENTS

This work was supported by the Royal Netherlands Academy of Arts and Sciences (KNAW, 08-PSA-M-02) and National 973 Project of China (2011CB606205).

AUTHOR CONTRIBUTIONS

J.M.: conception and design, manuscript writing; S.K.B.: conception and design, administrative support; F.Y.: conception and design, manuscript writing, financial support; F.-Z.C.: financial support, final approval of manuscript; J.P.: administrative support, final approval of manuscript; G.J.M.: conception and design, provision of study material or patients, final approval of manuscript; J.A.J.: conception and design, financial support, final approval of manuscript; J.J.J.P.v.d.B.: conception and design, data analysis and interpretation, manuscript writing, final approval of manuscript.

DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST

The authors indicate no potential conflicts of interest.

REFERENCES

- Giannoudis PV, Dinopoulos H, Tsiridis E. Bone substitutes: An update. *Injury* 2005;36 (suppl 3):S20–S27.
- Cancedda R, Giannoni P, Mastrogiacomo M. A tissue engineering approach to bone repair in large animal models and in clinical practice. *Biomaterials* 2007;28:4240–4250.
- Santos MI, Reis RL. Vascularization in bone tissue engineering: Physiology, current strategies, major hurdles and future challenges. *Macromol Biosci* 2010;10:12–27.
- Jain RK, Au P, Tam J et al. Engineering vascularized tissue. *Nat Biotechnol* 2005;23:821–823.
- Ma J, van den Beucken JJ, Yang F et al. Coculture of osteoblasts and endothelial cells: Optimization of culture medium and cell ratio. *Tissue Eng Part C Methods* 2011;17:349–357.
- Farrell E, Both SK, Odörfer KI et al. In-vivo generation of bone via endochondral ossification by in-vitro chondrogenic priming of adult human and rat mesenchymal stem cells. *BMC Musculoskelet Disord* 2011;12:31.
- Marot D, Knezevic M, Novakovic GV. Bone tissue engineering with human stem cells. *Stem Cell Res Ther* 2010;1:10.
- Both SK, van Apeldoorn AA, Jukes JM et al. Differential bone-forming capacity of osteogenic cells from either embryonic stem cells or bone marrow-derived mesenchymal stem cells. *J Tissue Eng Regen Med* 2010;5:180–190.
- Kuznetsov SA, Cherman N, Robey PG. In vivo bone formation by progeny of human embryonic stem cells. *Stem Cells Dev* 2011;20:269–287.
- Dupont KM, Sharma K, Stevens HY et al. Human stem cell delivery for treatment of large segmental bone defects. *Proc Natl Acad Sci USA* 2010;107:3305–3310.
- Cernansky R. Fetal stem cell therapy causes cancer in teenage boy. Available at <http://blogs.discovermagazine.com/80beats/2009/02/18/embryonic-stem-cell-therapy-causes-cancer-in-teenage-boy/#.UoD2Hvmkqk4> 2009. Accessed February 18, 2009.
- Sorrell JM, Baber MA, Caplan AI. Influence of adult mesenchymal stem cells on in vitro vascular formation. *Tissue Eng Part A* 2009;15:1751–1761.
- Diao Y, Ma Q, Cui F et al. Human umbilical cord mesenchymal stem cells: Osteogenesis in vivo as seed cells for bone tissue engineering. *J Biomed Mater Res A* 2009;91:123–131.
- Cowan CM, Shi YY, Aalami OO et al. Adipose-derived adult stromal cells heal critical-size mouse calvarial defects. *Nat Biotechnol* 2004;22:560–567.
- Niemeyer P, Kornacker M, Mehlhorn A et al. Comparison of immunological properties of bone marrow stromal cells and adipose tissue-derived stem cells before and after osteogenic differentiation in vitro. *Tissue Eng* 2007;13:111–121.
- Verseijden F, Posthumus-van Sluijs SJ, Pavljasevic P et al. Adult human bone marrow and adipose tissue-derived stromal cells support the formation of prevascular-like structures from endothelial cells in vitro. *Tissue Eng Part A* 2010;16:101–114.
- Gronthos S, Mankani M, Brahmi J et al. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci USA* 2000;97:13625–13630.
- Shi S, Gronthos S. Perivascular niche of postnatal mesenchymal stem cells in human bone marrow and dental pulp. *J Bone Miner Res* 2003;18:696–704.
- Ito K, Yamada Y, Nakamura S et al. Osteogenic potential of effective bone engineering using dental pulp stem cells, bone marrow stem cells, and periosteal cells for osseointegration of dental implants. *Int J Oral Maxillofac Implants* 2011;26:947–954.
- Yamada Y, Ito K, Nakamura S et al. Promising cell-based therapy for bone regeneration using stem cells from deciduous teeth, dental pulp, and bone marrow. *Cell Transplant* 2011;20:1003–1013.
- Zhang W, Walboomers XF, van Kuppevelt TH et al. The performance of human dental pulp stem cells on different three-dimensional scaffold materials. *Biomaterials* 2006;27:5658–5668.
- Miura M, Gronthos S, Zhao M et al. SHED: Stem cells from human exfoliated deciduous teeth. *Proc Natl Acad Sci USA* 2003;100:5807–5812.
- Tremp M, Salemi S, Gobet R et al. Adipose-derived stem cells (asc) for tissue engineering. In: Eberli D, ed. *Regenerative Medicine and Tissue Engineering: Cells and Biomaterials*. 2011:179–187.

- 24 Kim J, Kang JW, Park JH et al. Biological characterization of long-term cultured human mesenchymal stem cells. *Arch Pharm Res* 2009;32:117–126.
- 25 Sakaguchi Y, Sekiya I, Yagishita K et al. Suspended cells from trabecular bone by collagenase digestion become virtually identical to mesenchymal stem cells obtained from marrow aspirates. *Blood* 2004;104:2728–2735.
- 26 Chen HH, Decot V, Ouyang JP et al. In vitro initial expansion of mesenchymal stem cells is influenced by the culture parameters used in the isolation process. *Biomed Mater Eng* 2009;19:301–309.
- 27 Fink T, Rasmussen JG, Lund P et al. Isolation and expansion of adipose-derived stem cells for tissue engineering. *Front Biosci (Elite Ed)* 2011;3:256–263 [Elite Ed].
- 28 Lund P, Pilgaard L, Duroux M et al. Effect of growth media and serum replacements on the proliferation and differentiation of adipose-derived stem cells. *Cytotherapy* 2009;11:189–197.
- 29 Patel M, Smith AJ, Sloan AJ et al. Phenotype and behaviour of dental pulp cells during expansion culture. *Arch Oral Biol* 2009;54:898–908.
- 30 Govindasamy V, Ronald VS, Totev S et al. Micromanipulation of culture niche permits long-term expansion of dental pulp stem cells—an economic and commercial angle. *In Vitro Cell Dev Biol Anim* 2010;46:764–773.
- 31 Schallmoser K, Rohde E, Reinisch A et al. Rapid large-scale expansion of functional mesenchymal stem cells from unmanipulated bone marrow without animal serum. *Tissue Eng Part C Methods* 2008;14:185–196.
- 32 Chen B, Sun HH, Wang HG et al. The effects of human platelet lysate on dental pulp stem cells derived from impacted human third molars. *Biomaterials* 2012;33:5023–5035.
- 33 Naaijken BA, Niessen HW, Prins HJ et al. Human platelet lysate as a fetal bovine serum substitute improves human adipose-derived stromal cell culture for future cardiac repair applications. *Cell Tissue Res* 2012;348:119–130.
- 34 Salvadè A, Della Mina P, Gaddi D et al. Characterization of platelet lysate cultured mesenchymal stromal cells and their potential use in tissue-engineered osteogenic devices for the treatment of bone defects. *Tissue Eng Part C Methods* 2010;16:201–214.
- 35 Sotiropoulou PA, Perez SA, Salagianni M et al. Characterization of the optimal culture conditions for clinical scale production of human mesenchymal stem cells. *STEM CELLS* 2006;24:462–471.
- 36 Both SK, van der Muijsenberg AJ, van Blitterswijk CA et al. A rapid and efficient method for expansion of human mesenchymal stem cells. *Tissue Eng* 2007;13:3–9.
- 37 Rada T, Reis RL, Gomes ME. Adipose tissue-derived stem cells and their application in bone and cartilage tissue engineering. *Tissue Eng Part B Rev* 2009;15:113–125.
- 38 Jurgens WJ, Oedayrajsingh-Varma MJ, Helder MN et al. Effect of tissue-harvesting site on yield of stem cells derived from adipose tissue: implications for cell-based therapies. *Cell Tissue Res* 2008;332:415–426.
- 39 Aghaloo TL, Chaichanasakul T, Bezouglia O et al. Osteogenic potential of mandibular vs. long-bone marrow stromal cells. *J Dent Res* 2010;89:1293–1298.
- 40 Vunjak-Novakovic G, Obradovic B, Martin I et al. Dynamic cell seeding of polymer scaffolds for cartilage tissue engineering. *Biotechnol Prog* 1998;14:193–202.
- 41 Chen Y, Bloemen V, Impens S et al. Characterization and optimization of cell seeding in scaffolds by factorial design: quality by design approach for skeletal tissue engineering. *Tissue Eng Part C Methods* 2011;17:1211–1221.
- 42 Hasegawa T, Miwa M, Sakai Y et al. Efficient cell-seeding into scaffolds improves bone formation. *J Dent Res* 2010;89:854–859.
- 43 Zhao F, Ma T. Perfusion bioreactor system for human mesenchymal stem cell tissue engineering: Dynamic cell seeding and construct development. *Biotechnol Bioeng* 2005;91:482–493.
- 44 Kim K, Dean D, Mikos AG et al. Effect of initial cell seeding density on early osteogenic signal expression of rat bone marrow stromal cells cultured on cross-linked poly(propylene fumarate) disks. *Biomacromolecules* 2009;10:1810–1817.
- 45 Grayson WL, Bhumiratana S, Cannizzaro C et al. Effects of initial seeding density and fluid perfusion rate on formation of tissue-engineered bone. *Tissue Eng Part A* 2008;14:1809–1820.
- 46 Zhou H, Weir MD, Xu HH. Effect of cell seeding density on proliferation and osteodifferentiation of umbilical cord stem cells on calcium phosphate cement-fiber scaffold. *Tissue Eng Part A* 2011;17:2603–2613.
- 47 Yeatts AB, Fisher JP. Tubular perfusion system for the long-term dynamic culture of human mesenchymal stem cells. *Tissue Eng Part C Methods* 2011;17:337–348.
- 48 Yeatts AB, Fisher JP. Bone tissue engineering bioreactors: Dynamic culture and the influence of shear stress. *Bone* 2011;48:171–181.
- 49 Thimm BW, Wüst S, Hofmann S et al. Initial cell pre-cultivation can maximize ECM mineralization by human mesenchymal stem cells on silk fibroin scaffolds. *Acta Biomater* 2011;7:2218–2228.
- 50 Song IH, Caplan AI, Dennis JE. In vitro dexamethasone pretreatment enhances bone formation of human mesenchymal stem cells in vivo. *J Orthop Res* 2009;27:916–921.
- 51 Weiss HE, Roberts SJ, Schrooten J et al. A semi-autonomous model of endochondral ossification for developmental tissue engineering. *Tissue Eng Part A* 2012;18:1334–1343.
- 52 Siebers MC, Walboomers XF, Leeuwenburgh SC et al. The influence of the crystallinity of electrostatic spray deposition-derived coatings on osteoblast-like cell behavior, in vitro. *J Biomed Mater Res A* 2006;78:258–267.
- 53 Agata H, Asahina I, Watanabe N et al. Characteristic change and loss of in vivo osteogenic abilities of human bone marrow stromal cells during passage. *Tissue Eng Part A* 2010;16:663–673.
- 54 Yoon E, Dhar S, Chun DE et al. In vivo osteogenic potential of human adipose-derived stem cells/poly lactide-co-glycolic acid constructs for bone regeneration in a rat critical-sized calvarial defect model. *Tissue Eng* 2007;13:619–627.
- 55 Farré-Guasch E, Prins HJ, Overman JR et al. Human maxillary sinus floor elevation as a model for bone regeneration enabling the application of one-step surgical procedures. *Tissue Eng Part B Rev* 2013;19:69–82.
- 56 Hager S, Lampert FM, Orimo H et al. Up-regulation of alkaline phosphatase expression in human primary osteoblasts by cocultivation with primary endothelial cells is mediated by p38 mitogen-activated protein kinase-dependent mRNA stabilization. *Tissue Eng Part A* 2009;15:3437–3447.
- 57 Kolbe M, Xiang Z, Dohle E et al. Paracrine effects influenced by cell culture medium and consequences on microvessel-like structures in cocultures of mesenchymal stem cells and outgrowth endothelial cells. *Tissue Eng Part A* 2011;17:2199–2212.
- 58 Correia C, Grayson WL, Park M et al. In vitro model of vascularized bone: Synergizing vascular development and osteogenesis. *PLoS ONE* 2011;6:e28352.
- 59 Ma J, Both SK, Ji W et al. Adipose tissue-derived mscs as monocultures or cocultures with human umbilical vein endothelial cells: Performance in vitro and in rat cranial defects. *J Biomed Mater Res A* 2013 [Epub ahead of print].
- 60 Giannoni P, Scaglione S, Daga A et al. Short-time survival and engraftment of bone marrow stromal cells in an ectopic model of bone regeneration. *Tissue Eng Part A* 2010;16:489–499.
- 61 Dissanayaka WL, Zhan X, Zhang C et al. Coculture of dental pulp stem cells with endothelial cells enhances osteo-/odontogenic and angiogenic potential in vitro. *J Endod* 2012;38:454–463.
- 62 Ohyabu Y, Kaul Z, Yoshioka T et al. Stable and nondisruptive in vitro/in vivo labeling of mesenchymal stem cells by internalizing quantum dots. *Hum Gene Ther* 2009;20:217–224.
- 63 Xia Z, Locklin RM, Triffitt JT. Fates and osteogenic differentiation potential of human mesenchymal stem cells in immunocompromised mice. *Eur J Cell Biol* 2008;87:353–364.
- 64 Tortelli F, Tasso R, Loiacono F et al. The development of tissue-engineered bone of different origin through endochondral and intramembranous ossification following the implantation of mesenchymal stem cells and osteoblasts in a murine model. *Biomaterials* 2010;31:242–249.
- 65 Tavassol F, Schumann P, Lindhorst D et al. Accelerated angiogenic host tissue response to poly(L-lactide-co-glycolide) scaffolds by vitalization with osteoblast-like cells. *Tissue Eng Part A* 2010;16:2265–2279.
- 66 Aguirre A, Planell JA, Engel E. Dynamics of bone marrow-derived endothelial progenitor cell/mesenchymal stem cell interaction in coculture and its implications in angiogenesis. *Biochem Biophys Res Commun* 2010;400:284–291.
- 67 Seebach C, Henrich D, Wilhelm K et al. Endothelial progenitor cells improve directly and indirectly early vascularization of mesenchymal stem cell-driven bone regeneration in a critical bone defect in rats. *Cell Transplant* 2012;21:1667–1677.
- 68 Tatebe M, Nakamura R, Kagami H et al. Differentiation of transplanted mesenchymal stem cells in a large osteochondral defect in rabbit. *Cytotherapy* 2005;7:520–530.
- 69 Unger RE, Ghanaati S, Orth C et al. The rapid anastomosis between prevascularized networks on silk fibroin scaffolds generated in vitro with cocultures of human microvascular endothelial and osteoblast cells and the host vasculature. *Biomaterials* 2010;31:6959–6967.
- 70 Geuze RE, Prins HJ, Öner FC et al. Luciferase labeling for multipotent stromal cell tracking in spinal fusion versus ectopic bone tissue

engineering in mice and rats. *Tissue Eng Part A* 2010;16:3343–3351.

71 Koob S, Torio-Padron N, Stark GB et al. Bone formation and neovascularization mediated by mesenchymal stem cells and endothelial cells in critical-sized calvarial defects. *Tissue Eng Part A* 2011;17:311–321.

72 Behnia H, Khojasteh A, Soleimani M et al. Secondary repair of alveolar clefts using human mesenchymal stem cells. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;108:e1–e6.

73 Meijer GJ, de Bruijn JD, Koole R et al. Cell-based bone tissue engineering. *PLoS Med* 2007;4:e9.

74 Jakob M, Saxer F, Scotti C et al. Perspective on the evolution of cell-based bone tissue engineering strategies. *Eur Surg Res* 2012;49:1–7.

75 Nauta AJ, Fibbe WE. Immunomodulatory properties of mesenchymal stromal cells. *Blood* 2007;110:3499–3506.

76 Cordonnier T, Langonné A, Corre P et al. Osteoblastic differentiation and potent osteogenicity of three-dimensional hBMSC-BCP particle constructs. *J Tissue Eng Regen Med* 2012 [Epub ahead of print].

77 Niemeyer P, Schönberger TS, Hahn J et al. Xenogenic transplantation of human mesenchymal stem cells in a critical size defect of the sheep tibia for bone regeneration. *Tissue Eng Part A* 2010;16:33–43.

78 Costa-Pinto AR, Correlo VM, Sol PC et al. Chitosan-poly(butylene succinate) scaffolds and human bone marrow stromal cells induce bone repair in a mouse calvaria model. *J Tissue Eng Regen Med* 2012;6:21–28.

79 Rai B, Lin JL, Lim ZX et al. Differences between in vitro viability and differentiation and in vivo bone-forming efficacy of human mesenchymal stem cells cultured on PCL-TCP scaffolds. *Biomaterials* 2010;31:7960–7970.

80 Zong C, Xue D, Yuan W et al. Reconstruction of rat calvarial defects with human mesenchymal stem cells and osteoblast-like cells in poly-lactic-co-glycolic acid scaffolds. *Eur Cell Mater* 2010;20:109–120.

81 Wen Y, Jiang B, Cui J et al. Superior osteogenic capacity of different mesenchymal stem cells for bone tissue engineering. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2013;116:e324–e332.

82 Choi JW, Park EJ, Shin HS et al. In vivo differentiation of undifferentiated human adipose tissue-derived mesenchymal stem cells in critical-sized calvarial bone defects. *Ann Plast Surg* 2012 [Epub ahead of print].

83 Levi B, James AW, Nelson ER et al. Human adipose derived stromal cells heal critical size mouse calvarial defects. *PLoS ONE* 2010;5:e11177.

84 Choi HJ, Kim JM, Kwon E et al. Establishment of efficacy and safety assessment of human adipose tissue-derived mesenchymal stem cells (HATMSCs) in a nude rat femoral segmental defect model. *J Korean Med Sci* 2011;26:482–491.

85 Pisciotta A, Riccio M, Carnevale G et al. Human serum promotes osteogenic differentiation of human dental pulp stem cells in vitro and in vivo. *PLoS ONE* 2012;7:e50542.

86 Kaigler D, Krebsbach PH, Wang Z et al. Transplanted endothelial cells enhance orthotopic bone regeneration. *J Dent Res* 2006;85:633–637.

87 Seebach C, Henrich D, Kähling C et al. Endothelial progenitor cells and mesenchymal stem cells seeded onto beta-TCP granules enhance early vascularization and bone healing in a critical-sized bone defect in rats. *Tissue Eng Part A* 2010;16:1961–1970.

88 Henrich D, Seebach C, Kaehling C et al. Simultaneous cultivation of human endothelial-like differentiated precursor cells and human marrow stromal cells on beta-tricalcium phosphate. *Tissue Eng Part C Methods* 2009;15:551–560.

89 Zhou J, Lin H, Fang T et al. The repair of large segmental bone defects in the rabbit with vascularized tissue engineered bone. *Biomaterials* 2010;31:1171–1179.

90 Kim JY, Jin GZ, Park IS et al. Evaluation of solid free-form fabrication-based scaffolds

seeded with osteoblasts and human umbilical vein endothelial cells for use in vivo osteogenesis. *Tissue Eng Part A* 2010;16:2229–2236.

91 Quarto R, Mastrogiacomo M, Cancedda R et al. Repair of large bone defects with the use of autologous bone marrow stromal cells. *N Engl J Med* 2001;344:385–386.

92 Marcacci M, Kon E, Moukhachev V et al. Stem cells associated with macroporous bioceramics for long bone repair: 6- to 7-year outcome of a pilot clinical study. *Tissue Eng* 2007;13:947–955.

93 Hibi H, Yamada Y, Ueda M et al. Alveolar cleft osteoplasty using tissue-engineered osteogenic material. *Int J Oral Maxillofac Surg* 2006;35:551–555.

94 Lee J, Sung HM, Jang JD et al. Successful reconstruction of 15-cm segmental defects by bone marrow stem cells and resected autogenous bone graft in central hemangioma. *J Oral Maxillofac Surg* 2010;68:188–194.

95 Shayesteh YS, Khojasteh A, Soleimani M et al. Sinus augmentation using human mesenchymal stem cells loaded into a beta-tricalcium phosphate/hydroxyapatite scaffold. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;106:203–209.

96 Meijer GJ, de Bruijn JD, Koole R et al. Cell based bone tissue engineering in jaw defects. *Biomaterials* 2008;29:3053–3061.

97 Zamiri B, Shahidi S, Eslaminejad MB et al. Reconstruction of human mandibular continuity defects with allogenic scaffold and autologous marrow mesenchymal stem cells. *J Craniofac Surg* 2013;24:1292–1297.

98 Mesimäki K, Lindroos B, Törnwall J et al. Novel maxillary reconstruction with ectopic bone formation by GMP adipose stem cells. *Int J Oral Maxillofac Surg* 2009;38:201–209.

99 d'Aquino R, De Rosa A, Lanza V et al. Human mandible bone defect repair by the grafting of dental pulp stem/progenitor cells and collagen sponge biocomplexes. *Eur Cell Mater* 2009;18:75–83.