

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/128742>

Please be advised that this information was generated on 2021-09-16 and may be subject to change.

## Measurement of Time-Dependent $CP$ Asymmetries and the $CP$ -Odd Fraction in the Decay $B^0 \rightarrow D^* + D^{*-}$

B. Aubert,<sup>1</sup> R. Barate,<sup>1</sup> D. Boutigny,<sup>1</sup> F. Couderc,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> V. Poireau,<sup>1</sup> V. Tisserand,<sup>1</sup> A. Zghiche,<sup>1</sup> E. Grauges,<sup>2</sup> A. Palano,<sup>3</sup> M. Pappagallo,<sup>3</sup> A. Pompili,<sup>3</sup> J. C. Chen,<sup>4</sup> N. D. Qi,<sup>4</sup> G. Rong,<sup>4</sup> P. Wang,<sup>4</sup> Y. S. Zhu,<sup>4</sup> G. Eigen,<sup>5</sup> I. Ofte,<sup>5</sup> B. Stugu,<sup>5</sup> G. S. Abrams,<sup>6</sup> M. Battaglia,<sup>6</sup> A. B. Breon,<sup>6</sup> D. N. Brown,<sup>6</sup> J. Button-Shafer,<sup>6</sup> R. N. Cahn,<sup>6</sup> E. Charles,<sup>6</sup> C. T. Day,<sup>6</sup> M. S. Gill,<sup>6</sup> A. V. Gritsan,<sup>6</sup> Y. Groysman,<sup>6</sup> R. G. Jacobsen,<sup>6</sup> R. W. Kadel,<sup>6</sup> J. Kadyk,<sup>6</sup> L. T. Kerth,<sup>6</sup> Yu. G. Kolomoisky,<sup>6</sup> G. Kukartsev,<sup>6</sup> G. Lynch,<sup>6</sup> L. M. Mir,<sup>6</sup> P. J. Oddone,<sup>6</sup> T. J. Orimoto,<sup>6</sup> M. Pripstein,<sup>6</sup> N. A. Roe,<sup>6</sup> M. T. Ronan,<sup>6</sup> W. A. Wenzel,<sup>6</sup> M. Barrett,<sup>7</sup> K. E. Ford,<sup>7</sup> T. J. Harrison,<sup>7</sup> A. J. Hart,<sup>7</sup> C. M. Hawkes,<sup>7</sup> S. E. Morgan,<sup>7</sup> A. T. Watson,<sup>7</sup> M. Fritsch,<sup>8</sup> K. Goetzen,<sup>8</sup> T. Held,<sup>8</sup> H. Koch,<sup>8</sup> B. Lewandowski,<sup>8</sup> M. Pelizaeus,<sup>8</sup> K. Peters,<sup>8</sup> T. Schroeder,<sup>8</sup> M. Steinke,<sup>8</sup> J. T. Boyd,<sup>9</sup> J. P. Burke,<sup>9</sup> N. Chevalier,<sup>9</sup> W. N. Cottingham,<sup>9</sup> M. P. Kelly,<sup>9</sup> T. Cuhadar-Donszelmann,<sup>10</sup> B. G. Fulsom,<sup>10</sup> C. Hearty,<sup>10</sup> N. S. Knecht,<sup>10</sup> T. S. Mattison,<sup>10</sup> J. A. McKenna,<sup>10</sup> A. Khan,<sup>11</sup> P. Kyberd,<sup>11</sup> M. Saleem,<sup>11</sup> L. Teodorescu,<sup>11</sup> A. E. Blinov,<sup>12</sup> V. E. Blinov,<sup>12</sup> A. D. Bukin,<sup>12</sup> V. P. Druzhinin,<sup>12</sup> V. B. Golubev,<sup>12</sup> E. A. Kravchenko,<sup>12</sup> A. P. Onuchin,<sup>12</sup> S. I. Serednyakov,<sup>12</sup> Yu. I. Skovpen,<sup>12</sup> E. P. Solodov,<sup>12</sup> A. N. Yushkov,<sup>12</sup> D. Best,<sup>13</sup> M. Bondioli,<sup>13</sup> M. Bruinsma,<sup>13</sup> M. Chao,<sup>13</sup> I. Eschrich,<sup>13</sup> D. Kirkby,<sup>13</sup> A. J. Lankford,<sup>13</sup> M. Mandelkern,<sup>13</sup> R. K. Mommsen,<sup>13</sup> W. Roethel,<sup>13</sup> D. P. Stoker,<sup>13</sup> C. Buchanan,<sup>14</sup> B. L. Hartfiel,<sup>14</sup> A. J. R. Weinstein,<sup>14</sup> S. D. Foulkes,<sup>15</sup> J. W. Gary,<sup>15</sup> O. Long,<sup>15</sup> B. C. Shen,<sup>15</sup> K. Wang,<sup>15</sup> L. Zhang,<sup>15</sup> D. del Re,<sup>16</sup> H. K. Hadavand,<sup>16</sup> E. J. Hill,<sup>16</sup> D. B. MacFarlane,<sup>16</sup> H. P. Paar,<sup>16</sup> S. Rahatlou,<sup>16</sup> V. Sharma,<sup>16</sup> J. W. Berryhill,<sup>17</sup> C. Campagnari,<sup>17</sup> A. Cunha,<sup>17</sup> B. Dahmes,<sup>17</sup> T. M. Hong,<sup>17</sup> M. A. Mazur,<sup>17</sup> J. D. Richman,<sup>17</sup> W. Verkerke,<sup>17</sup> T. W. Beck,<sup>18</sup> A. M. Eisner,<sup>18</sup> C. J. Flacco,<sup>18</sup> C. A. Heusch,<sup>18</sup> J. Kroseberg,<sup>18</sup> W. S. Lockman,<sup>18</sup> G. Nesom,<sup>18</sup> T. Schalk,<sup>18</sup> B. A. Schumm,<sup>18</sup> A. Seiden,<sup>18</sup> P. Spradlin,<sup>18</sup> D. C. Williams,<sup>18</sup> M. G. Wilson,<sup>18</sup> J. Albert,<sup>19</sup> E. Chen,<sup>19</sup> G. P. Dubois-Felsmann,<sup>19</sup> A. Dvoretzki,<sup>19</sup> D. G. Hitlin,<sup>19</sup> I. Narsky,<sup>19</sup> T. Piatenko,<sup>19</sup> F. C. Porter,<sup>19</sup> A. Ryd,<sup>19</sup> A. Samuel,<sup>19</sup> R. Andreassen,<sup>20</sup> S. Jayatilake,<sup>20</sup> G. Mancinelli,<sup>20</sup> B. T. Meadows,<sup>20</sup> M. D. Sokoloff,<sup>20</sup> F. Blanc,<sup>21</sup> P. Bloom,<sup>21</sup> S. Chen,<sup>21</sup> W. T. Ford,<sup>21</sup> U. Nauenberg,<sup>21</sup> A. Olivas,<sup>21</sup> P. Rankin,<sup>21</sup> W. O. Ruddick,<sup>21</sup> J. G. Smith,<sup>21</sup> K. A. Ulmer,<sup>21</sup> S. R. Wagner,<sup>21</sup> J. Zhang,<sup>21</sup> A. Chen,<sup>22</sup> E. A. Eckhart,<sup>22</sup> A. Soffer,<sup>22</sup> W. H. Toki,<sup>22</sup> R. J. Wilson,<sup>22</sup> Q. Zeng,<sup>22</sup> D. Altenburg,<sup>23</sup> E. Feltresi,<sup>23</sup> A. Hauke,<sup>23</sup> B. Spaan,<sup>23</sup> T. Brandt,<sup>24</sup> J. Brose,<sup>24</sup> M. Dickopp,<sup>24</sup> V. Klose,<sup>24</sup> H. M. Lacker,<sup>24</sup> R. Nogowski,<sup>24</sup> S. Otto,<sup>24</sup> A. Petzold,<sup>24</sup> G. Schott,<sup>24</sup> J. Schubert,<sup>24</sup> K. R. Schubert,<sup>24</sup> R. Schwierz,<sup>24</sup> J. E. Sundermann,<sup>24</sup> D. Bernard,<sup>25</sup> G. R. Bonneaud,<sup>25</sup> P. Grenier,<sup>25</sup> S. Schrenk,<sup>25</sup> Ch. Thiebaut,<sup>25</sup> G. Vasileiadis,<sup>25</sup> M. Verderi,<sup>25</sup> D. J. Bard,<sup>26</sup> P. J. Clark,<sup>26</sup> W. Gradl,<sup>26</sup> F. Muheim,<sup>26</sup> S. Playfer,<sup>26</sup> Y. Xie,<sup>26</sup> M. Andreotti,<sup>27</sup> V. Azzolini,<sup>27</sup> D. Bettoni,<sup>27</sup> C. Bozzi,<sup>27</sup> R. Calabrese,<sup>27</sup> G. Cibinetto,<sup>27</sup> E. Luppi,<sup>27</sup> M. Negrini,<sup>27</sup> L. Piemontese,<sup>27</sup> F. Anulli,<sup>28</sup> R. Baldini-Ferrolli,<sup>28</sup> A. Calcaterra,<sup>28</sup> R. de Sangro,<sup>28</sup> G. Finocchiaro,<sup>28</sup> P. Patteri,<sup>28</sup> I. M. Peruzzi,<sup>28,\*</sup> M. Piccolo,<sup>28</sup> A. Zallo,<sup>28</sup> A. Buzzo,<sup>29</sup> R. Capra,<sup>29</sup> R. Contri,<sup>29</sup> M. Lo Vetere,<sup>29</sup> M. Macri,<sup>29</sup> M. R. Monge,<sup>29</sup> S. Passaggio,<sup>29</sup> C. Patrignani,<sup>29</sup> E. Robutti,<sup>29</sup> A. Santroni,<sup>29</sup> S. Tosi,<sup>29</sup> S. Bailey,<sup>30</sup> G. Brandenburg,<sup>30</sup> K. S. Chaisanguanthum,<sup>30</sup> M. Morii,<sup>30</sup> E. Won,<sup>30</sup> J. Wu,<sup>30</sup> R. S. Dubitzky,<sup>31</sup> U. Langenegger,<sup>31</sup> J. Marks,<sup>31</sup> S. Schenk,<sup>31</sup> U. Uwer,<sup>31</sup> W. Bhimji,<sup>32</sup> D. A. Bowerman,<sup>32</sup> P. D. Dauncey,<sup>32</sup> U. Egede,<sup>32</sup> R. L. Flack,<sup>32</sup> J. R. Gaillard,<sup>32</sup> G. W. Morton,<sup>32</sup> J. A. Nash,<sup>32</sup> M. B. Nikolich,<sup>32</sup> G. P. Taylor,<sup>32</sup> W. P. Vazquez,<sup>32</sup> M. J. Charles,<sup>33</sup> W. F. Mader,<sup>33</sup> U. Mallik,<sup>33</sup> A. K. Mohapatra,<sup>33</sup> J. Cochran,<sup>34</sup> H. B. Crawley,<sup>34</sup> V. Eyges,<sup>34</sup> W. T. Meyer,<sup>34</sup> S. Prell,<sup>34</sup> E. I. Rosenberg,<sup>34</sup> A. E. Rubin,<sup>34</sup> J. Yi,<sup>34</sup> N. Arnaud,<sup>35</sup> M. Davier,<sup>35</sup> X. Giroux,<sup>35</sup> G. Grosdidier,<sup>35</sup> A. Höcker,<sup>35</sup> F. Le Diberder,<sup>35</sup> V. Lepeltier,<sup>35</sup> A. M. Lutz,<sup>35</sup> A. Oyanguren,<sup>35</sup> T. C. Petersen,<sup>35</sup> M. Pierini,<sup>35</sup> S. Plaszczynski,<sup>35</sup> S. Rodier,<sup>35</sup> P. Roudeau,<sup>35</sup> M. H. Schune,<sup>35</sup> A. Stocchi,<sup>35</sup> G. Wormser,<sup>35</sup> C. H. Cheng,<sup>36</sup> D. J. Lange,<sup>36</sup> M. C. Simani,<sup>36</sup> D. M. Wright,<sup>36</sup> A. J. Bevan,<sup>37</sup> C. A. Chavez,<sup>37</sup> J. P. Coleman,<sup>37</sup> I. J. Forster,<sup>37</sup> J. R. Fry,<sup>37</sup> E. Gabathuler,<sup>37</sup> R. Gamet,<sup>37</sup> K. A. George,<sup>37</sup> D. E. Hutchcroft,<sup>37</sup> R. J. Parry,<sup>37</sup> D. J. Payne,<sup>37</sup> K. C. Schofield,<sup>37</sup> C. Touramanis,<sup>37</sup> C. M. Cormack,<sup>38</sup> F. Di Lodovico,<sup>38</sup> R. Sacco,<sup>38</sup> C. L. Brown,<sup>39</sup> G. Cowan,<sup>39</sup> H. U. Flaecher,<sup>39</sup> M. G. Green,<sup>39</sup> D. A. Hopkins,<sup>39</sup> P. S. Jackson,<sup>39</sup> T. R. McMahon,<sup>39</sup> S. Ricciardi,<sup>39</sup> F. Salvatore,<sup>39</sup> D. Brown,<sup>40</sup> C. L. Davis,<sup>40</sup> J. Allison,<sup>41</sup> N. R. Barlow,<sup>41</sup> R. J. Barlow,<sup>41</sup> M. C. Hodgkinson,<sup>41</sup> G. D. Lafferty,<sup>41</sup> M. T. Naisbit,<sup>41</sup> J. C. Williams,<sup>41</sup> C. Chen,<sup>42</sup> A. Farbin,<sup>42</sup> W. D. Hulsbergen,<sup>42</sup> A. Jawahery,<sup>42</sup> D. Kovalskyi,<sup>42</sup> C. K. Lae,<sup>42</sup> V. Lillard,<sup>42</sup> D. A. Roberts,<sup>42</sup> G. Simi,<sup>42</sup> G. Blaylock,<sup>43</sup> C. Dallapiccola,<sup>43</sup> S. S. Hertzbach,<sup>43</sup> R. Kofler,<sup>43</sup> V. B. Koptchev,<sup>43</sup> X. Li,<sup>43</sup> T. B. Moore,<sup>43</sup> S. Saremi,<sup>43</sup> H. Staengle,<sup>43</sup> S. Willocq,<sup>43</sup> R. Cowan,<sup>44</sup> K. Koeneke,<sup>44</sup> G. Sciolla,<sup>44</sup> S. J. Sekula,<sup>44</sup> M. Spitznagel,<sup>44</sup> F. Taylor,<sup>44</sup> R. K. Yamamoto,<sup>44</sup> H. Kim,<sup>45</sup> P. M. Patel,<sup>45</sup> S. H. Robertson,<sup>45</sup> A. Lazzaro,<sup>46</sup> V. Lombardo,<sup>46</sup> F. Palombo,<sup>46</sup> J. M. Bauer,<sup>47</sup> L. Cremaldi,<sup>47</sup> V. Eschenburg,<sup>47</sup> R. Godang,<sup>47</sup> R. Kroeger,<sup>47</sup> J. Reidy,<sup>47</sup> D. A. Sanders,<sup>47</sup> D. J. Summers,<sup>47</sup> H. W. Zhao,<sup>47</sup> S. Brunet,<sup>48</sup> D. Côté,<sup>48</sup> P. Taras,<sup>48</sup> B. Viaud,<sup>48</sup> H. Nicholson,<sup>49</sup> N. Cavallo,<sup>50,†</sup> G. De Nardo,<sup>50</sup> F. Fabozzi,<sup>50,†</sup> C. Gatto,<sup>50</sup> L. Lista,<sup>50</sup> D. Monorchio,<sup>50</sup> P. Paolucci,<sup>50</sup> D. Piccolo,<sup>50</sup> C. Sciacca,<sup>50</sup> M. Baak,<sup>51</sup> H. Bulten,<sup>51</sup> G. Raven,<sup>51</sup> H. L. Snoek,<sup>51</sup> L. Wilden,<sup>51</sup>

C. P. Jessop,<sup>52</sup> J. M. LoSecco,<sup>52</sup> T. Allmendinger,<sup>53</sup> G. Benelli,<sup>53</sup> K. K. Gan,<sup>53</sup> K. Honscheid,<sup>53</sup> D. Hufnagel,<sup>53</sup> P. D. Jackson,<sup>53</sup> H. Kagan,<sup>53</sup> R. Kass,<sup>53</sup> T. Pulliam,<sup>53</sup> A. M. Rahimi,<sup>53</sup> R. Ter-Antonyan,<sup>53</sup> Q. K. Wong,<sup>53</sup> J. Brau,<sup>54</sup> R. Frey,<sup>54</sup> O. Igonkina,<sup>54</sup> M. Lu,<sup>54</sup> C. T. Potter,<sup>54</sup> N. B. Sinev,<sup>54</sup> D. Strom,<sup>54</sup> J. Strube,<sup>54</sup> E. Torrence,<sup>54</sup> A. Dorigo,<sup>55</sup> F. Galeazzi,<sup>55</sup> M. Margoni,<sup>55</sup> M. Morandin,<sup>55</sup> M. Posocco,<sup>55</sup> M. Rotondo,<sup>55</sup> F. Simonetto,<sup>55</sup> R. Stroili,<sup>55</sup> C. Voci,<sup>55</sup> M. Benayoun,<sup>56</sup> H. Briand,<sup>56</sup> J. Chauveau,<sup>56</sup> P. David,<sup>56</sup> L. Del Buono,<sup>56</sup> Ch. de la Vaissière,<sup>56</sup> O. Hamon,<sup>56</sup> M. J. J. John,<sup>56</sup> Ph. Leruste,<sup>56</sup> J. Malclès,<sup>56</sup> J. Ocariz,<sup>56</sup> L. Roos,<sup>56</sup> G. Therin,<sup>56</sup> P. K. Behera,<sup>57</sup> L. Gladney,<sup>57</sup> Q. H. Guo,<sup>57</sup> J. Panetta,<sup>57</sup> M. Biasini,<sup>58</sup> R. Covarelli,<sup>58</sup> S. Pacetti,<sup>58</sup> M. Pioppi,<sup>58</sup> C. Angelini,<sup>59</sup> G. Batignani,<sup>59</sup> S. Bettarini,<sup>59</sup> F. Bucci,<sup>59</sup> G. Calderini,<sup>59</sup> M. Carpinelli,<sup>59</sup> R. Cenci,<sup>59</sup> F. Forti,<sup>59</sup> M. A. Giorgi,<sup>59</sup> A. Lusiani,<sup>59</sup> G. Marchiori,<sup>59</sup> M. Morganti,<sup>59</sup> N. Neri,<sup>59</sup> E. Paoloni,<sup>59</sup> M. Rama,<sup>59</sup> G. Rizzo,<sup>59</sup> J. Walsh,<sup>59</sup> M. Haire,<sup>60</sup> D. Judd,<sup>60</sup> D. E. Wagoner,<sup>60</sup> J. Biesiada,<sup>61</sup> N. Danielson,<sup>61</sup> P. Elmer,<sup>61</sup> Y. P. Lau,<sup>61</sup> C. Lu,<sup>61</sup> J. Olsen,<sup>61</sup> A. J. S. Smith,<sup>61</sup> A. V. Telnov,<sup>61</sup> F. Bellini,<sup>62</sup> G. Cavoto,<sup>62</sup> A. D'Orazio,<sup>62</sup> E. Di Marco,<sup>62</sup> R. Faccini,<sup>62</sup> F. Ferrarotto,<sup>62</sup> F. Ferroni,<sup>62</sup> M. Gaspero,<sup>62</sup> L. Li Gioi,<sup>62</sup> M. A. Mazzoni,<sup>62</sup> S. Morganti,<sup>62</sup> G. Piredda,<sup>62</sup> F. Polci,<sup>62</sup> F. Safai Tehrani,<sup>62</sup> C. Voena,<sup>62</sup> H. Schröder,<sup>63</sup> G. Wagner,<sup>63</sup> R. Waldi,<sup>63</sup> T. Adye,<sup>64</sup> N. De Groot,<sup>64</sup> B. Franek,<sup>64</sup> G. P. Gopal,<sup>64</sup> E. O. Olaiya,<sup>64</sup> F. F. Wilson,<sup>64</sup> R. Aleksan,<sup>65</sup> S. Emery,<sup>65</sup> A. Gaidot,<sup>65</sup> S. F. Ganzhur,<sup>65</sup> P.-F. Giraud,<sup>65</sup> G. Graziani,<sup>65</sup> G. Hamel de Monchenault,<sup>65</sup> W. Kozanecki,<sup>65</sup> M. Legendre,<sup>65</sup> G. W. London,<sup>65</sup> B. Mayer,<sup>65</sup> G. Vasseur,<sup>65</sup> Ch. Yèche,<sup>65</sup> M. Zito,<sup>65</sup> M. V. Purohit,<sup>66</sup> A. W. Weidemann,<sup>66</sup> J. R. Wilson,<sup>66</sup> F. X. Yumiceva,<sup>66</sup> T. Abe,<sup>67</sup> M. T. Allen,<sup>67</sup> D. Aston,<sup>67</sup> R. Bartoldus,<sup>67</sup> N. Berger,<sup>67</sup> A. M. Boyarski,<sup>67</sup> O. L. Buchmueller,<sup>67</sup> R. Claus,<sup>67</sup> M. R. Convery,<sup>67</sup> M. Cristinziani,<sup>67</sup> J. C. Dingfelder,<sup>67</sup> D. Dong,<sup>67</sup> J. Dorfan,<sup>67</sup> D. Dujmic,<sup>67</sup> W. Dunwoodie,<sup>67</sup> S. Fan,<sup>67</sup> R. C. Field,<sup>67</sup> T. Glanzman,<sup>67</sup> S. J. Gowdy,<sup>67</sup> T. Hadig,<sup>67</sup> V. Halyo,<sup>67</sup> C. Hast,<sup>67</sup> T. Hryn'ova,<sup>67</sup> W. R. Innes,<sup>67</sup> M. H. Kelsey,<sup>67</sup> P. Kim,<sup>67</sup> M. L. Kocian,<sup>67</sup> D. W. G. S. Leith,<sup>67</sup> J. Libby,<sup>67</sup> S. Luitz,<sup>67</sup> V. Luth,<sup>67</sup> H. L. Lynch,<sup>67</sup> H. Marsiske,<sup>67</sup> R. Messner,<sup>67</sup> D. R. Muller,<sup>67</sup> C. P. O'Grady,<sup>67</sup> V. E. Ozcan,<sup>67</sup> A. Perazzo,<sup>67</sup> M. Perl,<sup>67</sup> B. N. Ratcliff,<sup>67</sup> A. Roodman,<sup>67</sup> A. A. Salnikov,<sup>67</sup> R. H. Schindler,<sup>67</sup> J. Schwiening,<sup>67</sup> A. Snyder,<sup>67</sup> J. Stelzer,<sup>67</sup> D. Su,<sup>67</sup> M. K. Sullivan,<sup>67</sup> K. Suzuki,<sup>67</sup> S. Swain,<sup>67</sup> J. M. Thompson,<sup>67</sup> J. Va'vra,<sup>67</sup> M. Weaver,<sup>67</sup> W. J. Wisniewski,<sup>67</sup> M. Wittgen,<sup>67</sup> D. H. Wright,<sup>67</sup> A. K. Yarritu,<sup>67</sup> K. Yi,<sup>67</sup> C. C. Young,<sup>67</sup> P. R. Burchat,<sup>68</sup> A. J. Edwards,<sup>68</sup> S. A. Majewski,<sup>68</sup> B. A. Petersen,<sup>68</sup> C. Roat,<sup>68</sup> M. Ahmed,<sup>69</sup> S. Ahmed,<sup>69</sup> M. S. Alam,<sup>69</sup> J. A. Ernst,<sup>69</sup> M. A. Saeed,<sup>69</sup> F. R. Wappler,<sup>69</sup> S. B. Zain,<sup>69</sup> W. Bugg,<sup>70</sup> M. Krishnamurthy,<sup>70</sup> S. M. Spanier,<sup>70</sup> R. Eckmann,<sup>71</sup> J. L. Ritchie,<sup>71</sup> A. Satpathy,<sup>71</sup> R. F. Schwitters,<sup>71</sup> J. M. Izen,<sup>72</sup> I. Kitayama,<sup>72</sup> X. C. Lou,<sup>72</sup> S. Ye,<sup>72</sup> F. Bianchi,<sup>73</sup> M. Bona,<sup>73</sup> F. Gallo,<sup>73</sup> D. Gamba,<sup>73</sup> M. Bomben,<sup>74</sup> L. Bosisio,<sup>74</sup> C. Cartaro,<sup>74</sup> F. Cossutti,<sup>74</sup> G. Della Ricca,<sup>74</sup> S. Dittongo,<sup>74</sup> S. Grancagnolo,<sup>74</sup> L. Lanceri,<sup>74</sup> L. Vitale,<sup>74</sup> F. Martinez-Vidal,<sup>75</sup> R. S. Panvini,<sup>76,‡</sup> Sw. Banerjee,<sup>77</sup> B. Bhuyan,<sup>77</sup> C. M. Brown,<sup>77</sup> D. Fortin,<sup>77</sup> K. Hamano,<sup>77</sup> R. Kowalewski,<sup>77</sup> J. M. Roney,<sup>77</sup> R. J. Sobie,<sup>77</sup> J. J. Back,<sup>78</sup> P. F. Harrison,<sup>78</sup> T. E. Latham,<sup>78</sup> G. B. Mohanty,<sup>78</sup> H. R. Band,<sup>79</sup> X. Chen,<sup>79</sup> B. Cheng,<sup>79</sup> S. Dasu,<sup>79</sup> M. Datta,<sup>79</sup> A. M. Eichenbaum,<sup>79</sup> K. T. Flood,<sup>79</sup> M. Graham,<sup>79</sup> J. J. Hollar,<sup>79</sup> J. R. Johnson,<sup>79</sup> P. E. Kutter,<sup>79</sup> H. Li,<sup>79</sup> R. Liu,<sup>79</sup> B. Mellado,<sup>79</sup> A. Mihalyi,<sup>79</sup> Y. Pan,<sup>79</sup> R. Prepost,<sup>79</sup> P. Tan,<sup>79</sup> J. H. von Wimmersperg-Toeller,<sup>79</sup> S. L. Wu,<sup>79</sup> Z. Yu,<sup>79</sup> and H. Neal<sup>80</sup>

(BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

<sup>2</sup>IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

<sup>3</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

<sup>4</sup>Institute of High Energy Physics, Beijing 100039, China

<sup>5</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway

<sup>6</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

<sup>7</sup>University of Birmingham, Birmingham B15 2TT, United Kingdom

<sup>8</sup>Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany

<sup>9</sup>University of Bristol, Bristol BS8 1TL, United Kingdom

<sup>10</sup>University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

<sup>11</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>12</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>13</sup>University of California at Irvine, Irvine, California 92697, USA

<sup>14</sup>University of California at Los Angeles, Los Angeles, California 90024, USA

<sup>15</sup>University of California at Riverside, Riverside, California 92521, USA

<sup>16</sup>University of California at San Diego, La Jolla, California 92093, USA

<sup>17</sup>University of California at Santa Barbara, Santa Barbara, California 93106, USA

<sup>18</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

<sup>19</sup>California Institute of Technology, Pasadena, California 91125, USA

- <sup>20</sup>University of Cincinnati, Cincinnati, Ohio 45221, USA  
<sup>21</sup>University of Colorado, Boulder, Colorado 80309, USA  
<sup>22</sup>Colorado State University, Fort Collins, Colorado 80523, USA  
<sup>23</sup>Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany  
<sup>24</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany  
<sup>25</sup>Ecole Polytechnique, LLR, F-91128 Palaiseau, France  
<sup>26</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom  
<sup>27</sup>Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy  
<sup>28</sup>Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy  
<sup>29</sup>Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy  
<sup>30</sup>Harvard University, Cambridge, Massachusetts 02138, USA  
<sup>31</sup>Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany  
<sup>32</sup>Imperial College London, London SW7 2AZ, United Kingdom  
<sup>33</sup>University of Iowa, Iowa City, Iowa 52242, USA  
<sup>34</sup>Iowa State University, Ames, Iowa 50011-3160, USA  
<sup>35</sup>Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France  
<sup>36</sup>Lawrence Livermore National Laboratory, Livermore, California 94550, USA  
<sup>37</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom  
<sup>38</sup>Queen Mary, University of London, London E1 4NS, United Kingdom  
<sup>39</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom  
<sup>40</sup>University of Louisville, Louisville, Kentucky 40292, USA  
<sup>41</sup>University of Manchester, Manchester M13 9PL, United Kingdom  
<sup>42</sup>University of Maryland, College Park, Maryland 20742, USA  
<sup>43</sup>University of Massachusetts, Amherst, Massachusetts 01003, USA  
<sup>44</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA  
<sup>45</sup>McGill University, Montréal, Québec, Canada H3A 2T8  
<sup>46</sup>Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy  
<sup>47</sup>University of Mississippi, University, Mississippi 38677, USA  
<sup>48</sup>Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Québec, Canada H3C 3J7  
<sup>49</sup>Mount Holyoke College, South Hadley, Massachusetts 01075, USA  
<sup>50</sup>Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy  
<sup>51</sup>NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands  
<sup>52</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA  
<sup>53</sup>The Ohio State University, Columbus, Ohio 43210, USA  
<sup>54</sup>University of Oregon, Eugene, Oregon 97403, USA  
<sup>55</sup>Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy  
<sup>56</sup>Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France  
<sup>57</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA  
<sup>58</sup>Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy  
<sup>59</sup>Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy  
<sup>60</sup>Prairie View A&M University, Prairie View, Texas 77446, USA  
<sup>61</sup>Princeton University, Princeton, New Jersey 08544, USA  
<sup>62</sup>Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy  
<sup>63</sup>Universität Rostock, D-18051 Rostock, Germany  
<sup>64</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, United Kingdom  
<sup>65</sup>DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France  
<sup>66</sup>University of South Carolina, Columbia, South Carolina 29208, USA  
<sup>67</sup>Stanford Linear Accelerator Center, Stanford, California 94309, USA  
<sup>68</sup>Stanford University, Stanford, California 94305-4060, USA  
<sup>69</sup>State University of New York, Albany, New York 12222, USA  
<sup>70</sup>University of Tennessee, Knoxville, Tennessee 37996, USA  
<sup>71</sup>University of Texas at Austin, Austin, Texas 78712, USA  
<sup>72</sup>University of Texas at Dallas, Richardson, Texas 75083, USA  
<sup>73</sup>Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy  
<sup>74</sup>Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy  
<sup>75</sup>IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain  
<sup>76</sup>Vanderbilt University, Nashville, Tennessee 37235, USA  
<sup>77</sup>University of Victoria, Victoria, British Columbia, Canada V8W 3P6  
<sup>78</sup>Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom  
<sup>79</sup>University of Wisconsin, Madison, Wisconsin 53706, USA  
<sup>80</sup>Yale University, New Haven, Connecticut 06511, USA

(Received 30 June 2005; published 5 October 2005)

We present an updated measurement of time-dependent  $CP$  asymmetries and the  $CP$ -odd fraction in the decay  $B^0 \rightarrow D^{*+}D^{*-}$  using  $232 \times 10^6 B\bar{B}$  pairs collected by the *BABAR* detector at the SLAC PEP-II  $B$  factory. We determine the  $CP$ -odd fraction to be  $0.125 \pm 0.044(\text{stat}) \pm 0.007(\text{syst})$ . The time-dependent  $CP$  asymmetry parameters  $C_+$  and  $S_+$  are determined to be  $0.06 \pm 0.17(\text{stat}) \pm 0.03(\text{syst})$  and  $-0.75 \pm 0.25(\text{stat}) \pm 0.03(\text{syst})$ , respectively. The standard model predicts these parameters to be 0 and  $-\sin 2\beta$ , respectively, in the absence of penguin amplitude contributions.

DOI: 10.1103/PhysRevLett.95.151804

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

The time-dependent  $CP$  asymmetry measurement in  $B^0 \rightarrow D^{*+}D^{*-}$  decay provides an important test of the standard model (SM). In the SM,  $CP$  violation arises from a complex phase in the Cabibbo-Kobayashi-Maskawa quark-mixing matrix [1]. Measurements of  $CP$  asymmetries by the *BABAR* [2] and Belle [3] Collaborations have firmly established this effect in the  $B^0 \rightarrow J\psi K_S^0$  decay [4] and related modes that are governed by the  $b \rightarrow c\bar{c}s$  transition. The  $B^0 \rightarrow D^{*+}D^{*-}$  decay is dominated by the  $b \rightarrow c\bar{c}d$  transition. Within the framework of the SM, the  $CP$  asymmetry of  $B^0 \rightarrow D^{*+}D^{*-}$  is related to  $\sin 2\beta$  when the correction due to penguin diagram contributions are neglected. The penguin-induced correction has been estimated in models based on the factorization approximation and heavy quark symmetry and was predicted to be about 2% [5]. A significant deviation of the measured  $\sin 2\beta$  from the one observed in  $b \rightarrow c\bar{c}s$  decays would be evidence for a new  $CP$ -violating interaction. The enhanced sensitivity of  $B^0 \rightarrow D^{*+}D^{*-}$  to such a process arises from its much smaller SM amplitude compared with that of the  $b \rightarrow c\bar{c}s$  transition.

The  $B^0 \rightarrow D^{*+}D^{*-}$  decay proceeds through the  $CP$ -even  $S$  and  $D$  waves and through the  $CP$ -odd  $P$  wave. In this Letter, we present an improved measurement of the  $CP$ -odd fraction [6,7]  $R_\perp$  based on a time-integrated one-dimensional angular analysis. We also present an improved measurement of the time-dependent  $CP$  asymmetry [6,7], obtained from a combined analysis of time-dependent flavor-tagged decays and the one-dimensional angular distribution of the decay products.

The data used in this analysis comprise  $232 \times 10^6$   $\Upsilon(4S) \rightarrow B\bar{B}$  decays collected by the *BABAR* detector at the SLAC PEP-II storage ring. The *BABAR* detector is described in detail elsewhere [8]. We use a Monte Carlo (MC) simulation based on GEANT4 [9] to validate the analysis procedure and to study the relevant backgrounds.

We select  $B^0 \rightarrow D^{*+}D^{*-}$  decay by combining two charged  $D^*$  candidates reconstructed in the modes  $D^{*+} \rightarrow D^0\pi^+$  and  $D^{*+} \rightarrow D^+\pi^0$ . We include the  $D^{*+}D^{*-}$  combinations ( $D^0\pi^+, \bar{D}^0\pi^-$ ) and ( $D^0\pi^+, D^-\pi^0$ ), but not ( $D^+\pi^0, D^-\pi^0$ ) because of the smaller branching fraction and larger backgrounds. To suppress the  $e^+e^- \rightarrow q\bar{q}$  ( $q = u, d, s, \text{ and } c$ ) continuum background, we require the ratio of the second and zeroth order Fox-Wolfram moments [10] to be less than 0.6.

Candidates for  $D^0$  and  $D^+$  mesons are reconstructed in the modes  $D^0 \rightarrow K^-\pi^+, K^-\pi^+\pi^0, K^-\pi^+\pi^+\pi^-, K_S^0\pi^+\pi^-$  and  $D^+ \rightarrow K^-\pi^+\pi^+, K_S^0\pi^+, K^-K^+\pi^+$ . The reconstructed mass of the  $D^0$  ( $D^+$ ) candidate is required to be within 20 MeV/ $c^2$  of its nominal mass [11], except for the  $D^0 \rightarrow K^-\pi^+\pi^0$  candidate, where a looser requirement of 40 MeV/ $c^2$  is applied.

The  $K_S^0$  candidates are reconstructed from two oppositely charged tracks with an invariant mass within 20 MeV/ $c^2$  of the nominal  $K_S^0$  mass. The  $\chi^2$  probability of the  $\pi^+\pi^-$  vertex fit must be greater than 0.1%. Charged kaon candidates are required to be inconsistent with the pion hypothesis, as inferred from the Cherenkov angle measured by the Cherenkov detector and the ionization energy loss measured by the charged-particle tracking system [8]. Neutral pion candidates are formed from two photons detected in the electromagnetic calorimeter [8], each with energy above 30 MeV. The mass of the pair must be within 30 MeV/ $c^2$  of the nominal  $\pi^0$  mass, and their summed energy is required to be greater than 200 MeV. In addition, a mass-constrained fit is applied to the  $\pi^0$  candidates for further analysis.

The  $D^0$  and  $D^+$  candidates are subject to a mass-constrained fit prior to the formation of the  $D^{*+}$  candidates. A slow  $\pi^+$  from  $D^{*+}$  decay is required to have a momentum in the  $\Upsilon(4S)$  center-of-mass (c.m.) frame less than 450 MeV/ $c$ . A slow  $\pi^0$  from  $D^{*+}$  must have a momentum between 70 and 450 MeV/ $c$  in the c.m. frame. No requirement on the photon-energy sum is applied to the  $\pi^0$  candidates from the  $D^{*+}$  decays.

For each  $B^0 \rightarrow D^{*+}D^{*-}$  candidate, we construct a likelihood function [12]  $\mathcal{L}_{\text{mass}}$  from the masses and mass uncertainties of the  $D$  and  $D^*$  candidates. The likelihood  $\mathcal{L}_{\text{mass}}$  is calculated as the product of the likelihoods for the  $D$  and  $D^*$  candidates. The  $D$  mass resolution is modeled by a Gaussian whose variance is determined on a candidate-by-candidate basis. The  $D^* - D$  mass difference resolution is modeled by a double-Gaussian distribution whose parameters are determined from simulated events. The values of  $\mathcal{L}_{\text{mass}}$  and the difference of the  $B^0$  candidate energy  $E_B$  from the beam energy  $E_{\text{Beam}}$ ,  $\Delta E \equiv E_B - E_{\text{Beam}}$ , in the  $\Upsilon(4S)$  c.m. frame are used to reduce the combinatoric background further. From the simulated events, the maximum allowed values of  $-\ln \mathcal{L}_{\text{mass}}$  and  $|\Delta E|$  are optimized for each individual

final state to obtain the highest expected signal significance using the previously measured  $B^0 \rightarrow D^{*+} D^{*-}$  branching fraction [6].

The energy-substituted mass,  $m_{ES} \equiv \sqrt{E_{\text{Beam}}^2 - p_B^{*2}}$ , where  $p_B^*$  is the  $B^0$  candidate momentum in the  $Y(4S)$  c.m. frame, is used to extract the signal yield from the events satisfying the aforementioned selection. We select the  $B^0$  candidates that have  $m_{ES} \geq 5.23 \text{ GeV}/c^2$ . In cases where more than one  $B^0$  candidate is reconstructed in an event, the candidate with the smallest value of  $-\ln \mathcal{L}_{\text{mass}}$  is chosen. A fit to the  $m_{ES}$  distribution with a probability density function (PDF) given by the sum of a Gaussian shape for the signal and an ARGUS [13] function for the background yields  $391 \pm 28(\text{stat})$  signal events. In

the region of  $m_{ES} > 5.27 \text{ GeV}/c^2$ , the signal purity is approximately 70%.

In the transversity basis [14], we define the following three angles: the angle  $\theta_1$  between the momentum of the slow pion from the  $D^{*-}$  and the opposite direction of flight of the  $D^{*+}$  in the  $D^{*-}$  rest frame; the polar angle  $\theta_{\text{tr}}$  and azimuthal angle  $\phi_{\text{tr}}$  of the slow pion from the  $D^{*+}$  defined in the  $D^{*+}$  rest frame, where the opposite direction of flight of the  $D^{*-}$  is chosen as the  $x$  axis, and the  $z$  axis is defined as the normal to the  $D^{*-}$  decay plane.

The time-dependent angular distribution of the decay products is given in Ref. [15]. Taking into account the detector angular acceptance efficiency and integrating over the decay time and the angles  $\theta_1$  and  $\phi_{\text{tr}}$ , we obtain a one-dimensional differential decay rate:

$$\frac{1}{\Gamma} \frac{d\Gamma}{d \cos \theta_{\text{tr}}} = \frac{9}{32\pi} \left[ (1 - R_{\perp}) \sin^2 \theta_{\text{tr}} \left\{ \frac{1 + \alpha}{2} I_0(\cos \theta_{\text{tr}}) + \frac{1 - \alpha}{2} I_{\parallel}(\cos \theta_{\text{tr}}) \right\} + 2R_{\perp} \cos^2 \theta_{\text{tr}} \times I_{\perp}(\cos \theta_{\text{tr}}) \right], \quad (1)$$

where  $R_{\perp} = |A_{\perp}|^2 / (|A_0|^2 + |A_{\parallel}|^2 + |A_{\perp}|^2)$ ,  $\alpha = (|A_0|^2 - |A_{\parallel}|^2) / (|A_0|^2 + |A_{\parallel}|^2)$ ,  $A_0$  is the amplitude for longitudinally polarized  $D^*$  mesons,  $A_{\parallel}$  and  $A_{\perp}$  are the amplitudes for parallel and perpendicular transversely polarized  $D^*$  mesons. The three efficiency moments,  $I_k$  ( $k = 0, \parallel, \perp$ ), are defined as

$$I_k(\cos \theta_{\text{tr}}) = \int d \cos \theta_1 d \phi_{\text{tr}} g_k(\theta_1, \phi_{\text{tr}}) \varepsilon(\theta_1, \theta_{\text{tr}}, \phi_{\text{tr}}), \quad (2)$$

where  $g_0 = 4 \cos^2 \theta_1 \cos^2 \phi_{\text{tr}}$ ,  $g_{\parallel} = 2 \sin^2 \theta_1 \sin^2 \phi_{\text{tr}}$ ,  $g_{\perp} = \sin^2 \theta_1$ , and  $\varepsilon$  is the detector efficiency. The efficiency moments are parametrized as second-order even polynomials of  $\cos \theta_{\text{tr}}$ . Their parameter values are determined from the MC calculation and are subsequently fixed in the likelihood fit to the differential decay distribution of  $\cos \theta_{\text{tr}}$ . In fact, the three  $I_k$  functions deviate only slightly from a constant, making the distribution, Eq. (1), nearly independent of the amplitude ratio  $\alpha$ .

The  $CP$ -odd fraction  $R_{\perp}$  is measured in a simultaneous unbinned maximum likelihood fit to the  $\cos \theta_{\text{tr}}$  and the  $m_{ES}$  distribution. The background shape is modeled as an even second-order polynomial in  $\cos \theta_{\text{tr}}$ , while the signal PDF is given by Eq. (1). The finite detector resolution of the  $\theta_{\text{tr}}$  measurement is modeled as a double Gaussian plus a small tail component that accounts for misreconstructed events. The parametrization of the  $\theta_{\text{tr}}$  resolution function is fixed from the MC simulation and subsequently used to convolve the signal PDF in the maximum likelihood fit. Since the angle  $\theta_{\text{tr}}$  is calculated with the slow pion from the  $D^{*+}$ , we categorize events into three types:  $D^{*+} D^{*-} \rightarrow (D^0 \pi^+, \bar{D}^0 \pi^-)$ ,  $(D^0 \pi^+, D^- \pi^0)$ , and  $(D^+ \pi^0, \bar{D}^0 \pi^-)$ , each with different signal-fraction parameters in the likelihood fit. Their angular efficiency moments and  $\cos \theta_{\text{tr}}$  resolutions are also separately determined from the MC simulation. The other parameters determined in the likelihood fit

are the  $\cos \theta_{\text{tr}}$  background-shape parameter, three  $m_{ES}$  parameters ( $\sigma$  and mean of the signal Gaussian, and the ARGUS shape parameter  $\kappa$ ), as well as  $R_{\perp}$ . The fit to the data yields

$$R_{\perp} = 0.125 \pm 0.044(\text{stat}) \pm 0.007(\text{syst}). \quad (3)$$

The projections of the fitted result onto  $m_{ES}$  and  $\cos \theta_{\text{tr}}$  are shown in Fig. 1.

In the fit described above, the value of  $\alpha$  is fixed to zero. We estimate the corresponding systematic uncertainty by varying its value from  $-1$  to  $+1$  and find negligible change (less than 0.002) in the fitted value of  $R_{\perp}$ . Other systematic uncertainties arise from the parametrization of the angular resolution, the determination of the efficiency moments, and the background parametrization. The total systematic uncertainty on  $R_{\perp}$  is 0.007, significantly smaller than the statistical error.

We subsequently perform a combined analysis of the  $\cos \theta_{\text{tr}}$  distribution and the time dependence to extract the




FIG. 1. Measured distribution of  $m_{ES}$  (left) and of  $\cos \theta_{\text{tr}}$  in the region  $m_{ES} > 5.27 \text{ GeV}/c^2$  (right). The solid line is the projection of the fit result. The dotted line represents the background component.

time-dependent  $CP$  asymmetry, using the event sample described previously. We use information from the other  $B$  meson in the event to tag the initial flavor of the fully reconstructed  $B^0 \rightarrow D^{*+}D^{*-}$  candidate.

The decay rate  $f_+(f_-)$  for a neutral  $B$  meson accompanied by a  $B^0(\bar{B}^0)$  tag is given by

$$f_{\pm}(\Delta t, \cos\theta_{\text{tr}}) \propto e^{-|\Delta t|/\tau_{B^0}} \{G(1 \mp \Delta\omega) \mp (1 - 2\omega) \\ \times [F \sin(\Delta m_d \Delta t) + H \cos(\Delta m_d \Delta t)]\}, \quad (4)$$

where  $\Delta t = t_{\text{rec}} - t_{\text{tag}}$  is the difference between the proper decay time of the reconstructed  $B$  meson ( $B_{\text{rec}}$ ) and that of the tagging  $B$  meson ( $B_{\text{tag}}$ ),  $\tau_{B^0}$  is the  $B^0$  lifetime, and  $\Delta m_d$  is the mass difference determined from the  $B^0 - \bar{B}^0$  oscillation frequency [11]. The average mistag probability  $\omega$  describes the effect of incorrect tags, and  $\Delta\omega$  is the difference between the mistag rate for  $B^0$  and  $\bar{B}^0$ . The  $G$ ,  $F$ , and  $H$  coefficients are defined as

$$\begin{aligned} G &= (1 - R_{\perp})\sin^2\theta_{\text{tr}} + 2R_{\perp}\cos^2\theta_{\text{tr}}, \\ F &= (1 - R_{\perp})S_{+}\sin^2\theta_{\text{tr}} - 2R_{\perp}S_{\perp}\cos^2\theta_{\text{tr}}, \\ H &= (1 - R_{\perp})C_{+}\sin^2\theta_{\text{tr}} + 2R_{\perp}C_{\perp}\cos^2\theta_{\text{tr}}, \end{aligned} \quad (5)$$

where we allow the three transversity amplitudes to have different  $\lambda_k = (q/p)(\bar{A}_k/A_k)$  ( $k = 0, \parallel, \perp$ ) [15] due to possibly different penguin-to-tree amplitude ratios, and define the  $CP$  asymmetry  $C_k = 1 - |\lambda_k|^2/1 + |\lambda_k|^2$ ,  $S_k = 2\text{Im}(\lambda_k)/1 + |\lambda_k|^2$ . Here we also have




FIG. 2. From top to bottom: the distribution of  $\Delta t$  in the region  $m_{\text{ES}} > 5.27 \text{ GeV}/c^2$  for  $B^0$  ( $\bar{B}^0$ ) tag candidates, and the raw asymmetry  $(N_{B^0} - N_{\bar{B}^0}) / (N_{B^0} + N_{\bar{B}^0})$ , as functions of  $\Delta t$ . In the upper plot the solid (dashed) curves represent the fit projections in  $\Delta t$  for  $B^0$  ( $\bar{B}^0$ ) tags.

$$C_{+} = \frac{C_{\parallel}|A_{\parallel}|^2 + C_0|A_0|^2}{|A_{\parallel}|^2 + |A_0|^2}, \quad S_{+} = \frac{S_{\parallel}|A_{\parallel}|^2 + S_0|A_0|^2}{|A_{\parallel}|^2 + |A_0|^2}. \quad (6)$$

In the absence of penguin contributions, we expect that  $C_0 = C_{\parallel} = C_{\perp} = 0$ , and  $S_0 = S_{\parallel} = S_{\perp} = -\sin 2\beta$ .

In Eq. (4), the small angular acceptance effects are not incorporated, but absorbed into the “effective” value of  $R_{\perp}$ , which is left free to vary in the final fit. No bias is seen in the resulting values of  $C_{+}$ ,  $C_{\perp}$ ,  $S_{+}$ , and  $S_{\perp}$  in MC simulation.

The technique used to measure the  $CP$  asymmetry is analogous to previous  $BABAR$  measurements as described in Ref. [16]. Only events with a  $\Delta t$  uncertainty less than 2.5 ps and a measured  $|\Delta t|$  less than 20 ps are accepted. We performed a simultaneous unbinned maximum likelihood fit to the  $\cos\theta_{\text{tr}}$ ,  $\Delta t$ , and  $m_{\text{ES}}$  distributions to extract the  $CP$  asymmetry. The signal PDF in  $\theta_{\text{tr}}$  and  $\Delta t$  is given by Eq. (4). The signal mistag probability is determined from a sample of neutral  $B$  decays to flavor eigenstates,  $B_{\text{flav}}$ . In the likelihood fit, the expression in Eq. (4) is convolved with an empirical  $\Delta t$  resolution function determined from the  $B_{\text{flav}}$  sample. The  $\theta_{\text{tr}}$  resolution is accounted for in the same way as described previously.

The background  $\Delta t$  distributions are parametrized with an empirical description that includes prompt and non-prompt components. We allow the nonprompt background to have two free parameters,  $C_{\text{eff}}$  and  $S_{\text{eff}}$ , the effective  $CP$  asymmetries, in the likelihood fit. The background shape in  $\theta_{\text{tr}}$  is modeled as an even second-order polynomial in  $\cos\theta_{\text{tr}}$ , much as it is in the time-integrated angular analysis.

The fit to the data yields

$$\begin{aligned} C_{+} &= 0.06 \pm 0.17(\text{stat}) \pm 0.03(\text{syst}), \\ C_{\perp} &= -0.20 \pm 0.96(\text{stat}) \pm 0.11(\text{syst}), \\ S_{+} &= -0.75 \pm 0.25(\text{stat}) \pm 0.03(\text{syst}), \\ S_{\perp} &= -1.75 \pm 1.78(\text{stat}) \pm 0.22(\text{syst}). \end{aligned} \quad (7)$$

Figure 2 shows the  $\Delta t$  distributions and asymmetries in yields between  $B^0$  and  $\bar{B}^0$  tags, overlaid with the projection of the likelihood fit result. Because the  $CP$ -odd fraction is small, we have rather large statistical uncertainties for the measured  $C_{\perp}$  and  $S_{\perp}$  values. For comparison, we repeat the fit with the assumption that both  $CP$ -even and  $CP$ -odd states have the same  $CP$  asymmetry. We find that  $C_{+} = C_{\perp} = 0.03 \pm 0.13(\text{stat}) \pm 0.02(\text{syst})$ , and  $S_{+} = S_{\perp} = -0.69 \pm 0.23(\text{stat}) \pm 0.03(\text{syst})$ . In both cases, the effective  $CP$  asymmetries in the background are found to be consistent with zero within the statistical uncertainties.

The systematic uncertainties on  $C_{+}$ ,  $C_{\perp}$ ,  $S_{+}$ , and  $S_{\perp}$  arise from the amount of possible backgrounds that tend to peak under the signal and their  $CP$  asymmetry, the assumed parametrization of the  $\Delta t$  resolution function, the possible differences between the  $B_{\text{flav}}$  and  $B_{CP}$  mistag fractions, knowledge of the event-by-event beam-spot

position, and the possible interference between the suppressed  $\bar{b} \rightarrow \bar{u}c\bar{d}$  amplitude and the favored  $b \rightarrow c\bar{u}d$  amplitude for some tagside decays [17]. It also includes the systematic uncertainties from the finite MC calculation sample used to verify the fitting method. In general, all of the systematic uncertainties are found to be much smaller than the statistical uncertainties.

In summary, we have reported measurements of the  $CP$ -odd fraction and time-dependent  $CP$  asymmetries for the decay  $B^0 \rightarrow D^{*+}D^{*-}$ . The measurement supersedes the previous *BABAR* result [6], with more than 50% reduction in the statistical uncertainty, and indicates that  $B^0 \rightarrow D^{*+}D^{*-}$  is mostly  $CP$  even. The time-dependent asymmetries are found to be consistent with the SM predictions within the statistical uncertainty.

We are grateful for the excellent luminosity and machine conditions provided by our SLAC PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

---

\*Also at Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

<sup>†</sup>Also at Università della Basilicata, Potenza, Italy.

<sup>‡</sup>Deceased.

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **66**, 032003 (2002).
- [3] K. Abe *et al.* (Belle Collaboration), Phys. Rev. D **66**, 071102 (2002).
- [4] We imply charged conjugate modes throughout the Letter.
- [5] X. Y. Pham and Z. Z. Xing, Phys. Lett. B **458**, 375 (1999).
- [6] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **91**, 131801 (2003).
- [7] H. Miyake *et al.* (Belle Collaboration), Phys. Lett. B **618**, 34 (2005).
- [8] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [9] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [10] G. C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
- [11] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
- [12] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **89**, 061801 (2002).
- [13] H. Albrecht *et al.* (ARGUS Collaboration), Z. Phys. C **48**, 543 (1990).
- [14] I. Dunietz, H. R. Quinn, A. Snyder, W. Toki, and H. J. Lipkin, Phys. Rev. D **43**, 2193 (1991).
- [15] *The BABAR Physics Book*, edited by P. F. Harrison and H. R. Quinn, SLAC-R504 (Stanford Linear Acceleration Center, Stanford, CA, 1998).
- [16] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **94**, 161803 (2005).
- [17] O. Long, M. Baak, R. N. Cahn, and D. Kirkby, Phys. Rev. D **68**, 034010 (2003).