

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/129963>

Please be advised that this information was generated on 2019-02-16 and may be subject to change.

Corrigendum

Antidepressant efficacy of high-frequency transcranial magnetic stimulation over the left dorsolateral prefrontal cortex in double-blind sham-controlled designs: a meta-analysis – Corrigendum

D. J. L. G. Schutter

doi: 10.1017/S0033291708003462. Published online by Cambridge University Press, 30 April 2008 (volume 39, 2009, pages 65–75).

There were a number of errors in the above paper for which the author apologizes.

Where the total 1164 appears *it should read* 1120:

Page 65, Methods section, line 3.

Page 67, Results section, line 1.

Page 70, Table 2, column 5.

Page 72, right column, line 9.

Where the total 606 appears *it should read* 593:

Page 67, Results section, line 3.

Page 70, Table 2, column 3.

Where the total 558 appears *it should read* 527:

Page 67, Results section, line 4.

Page 70, Table 2, column 4.

Page 69, Table 2, Study no. 24 (Rossini *et al.* 2005):

The number of Active patients (column 4) should read 36 *not* 49.

The number of Sham patients (column 4) should read 16 *not* 47.

Reference

Schutter DJLG (2008). Antidepressant efficacy of high-frequency transcranial magnetic stimulation over the left dorsolateral prefrontal cortex in double-blind sham-controlled designs: a meta-analysis. *Psychological Medicine*. doi:10.1017/S0033291708003462. Published online by Cambridge University Press, 30 April 2008.