

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/131460>

Please be advised that this information was generated on 2019-03-24 and may be subject to change.



Open access – Invited review

THIS ARTICLE IS PART OF A SPECIAL ISSUE ENTITLED
'TRACE GAS BIOLOGY'

Nitric oxide in plants: an assessment of the current state of knowledge

Luis A. J. Mur^{1*}, Julien Mandon², Stefan Persijn², Simona M. Cristescu², Igor E. Moshkov³, Galina V. Novikova³, Michael A. Hall¹, Frans J. M. Harren², Kim H. Hebelstrup⁴ and Kapuganti J. Gupta⁵¹ Institute of Environmental and Rural Science, Aberystwyth University, Edward Llwyd Building, Aberystwyth SY23 3DA, UK² Life Science Trace Gas Facility, Molecular and Laser Physics, Institute for Molecules and Materials, Radboud University, PO Box 9010, 6500 GL Nijmegen, The Netherlands³ Timiryazev Institute of Plant Physiology, Russian Academy of Sciences, ul. Botanicheskaya 35, Moscow 127276, Russia⁴ Department of Molecular Biology and Genetics, Section of Crop Genetics and Biotechnology, Aarhus University, Forsøgsvej 1, DK-4200 Slagelse, Denmark⁵ Department of Plant Sciences, University of Oxford, South Parks Road, Oxford OX1 3RB, UK**Received:** 31 August 2012; **Revised:** 11 December 2012; **Accepted:** 12 December 2012; **Published:** 20 December 2012**Citation details:** Mur LAJ, Mandon J, Persijn S, Cristescu SM, Moshkov IE, Novikova GV, Hall MA, Harren FJM, Hebelstrup KH, Gupta KJ. 2013. Nitric oxide in plants: an assessment of the current state of knowledge. *AoB PLANTS* 5: pls052; doi:10.1093/aobpla/pls052

Abstract

Background and aims

After a series of seminal works during the last decade of the 20th century, nitric oxide (NO) is now firmly placed in the pantheon of plant signals. Nitric oxide acts in plant–microbe interactions, responses to abiotic stress, stomatal regulation and a range of developmental processes. By considering the recent advances in plant NO biology, this review will highlight certain key aspects that require further attention.

Scope and conclusions

The following questions will be considered. While cytosolic nitrate reductase is an important source of NO, the contributions of other mechanisms, including a poorly defined arginine oxidizing activity, need to be characterized at the molecular level. Other oxidative pathways utilizing polyamine and hydroxylamine also need further attention. Nitric oxide action is dependent on its concentration and spatial generation patterns. However, no single technology currently available is able to provide accurate *in planta* measurements of spatio-temporal patterns of NO production. It is also the case that pharmaceutical NO donors are used in studies, sometimes with little consideration of the kinetics of NO production. We here include *in planta* assessments of NO production from diethylamine nitric oxide, S-nitrosoglutathione and sodium nitroprusside following infiltration of tobacco leaves, which could aid workers in their experiments. Further, based on current data it is difficult to define a bespoke plant NO signalling pathway, but rather NO appears to act as a modifier of other signalling pathways. Thus, early reports that NO signalling involves cGMP—as in animal systems—require revisiting. Finally, as plants are exposed to NO from a number of external sources, investigations into the control of NO scavenging by such as non-symbiotic haemoglobins and other sinks for NO should feature more highly. By crystallizing these questions the authors encourage their resolution through the concerted efforts of the plant NO community.

* Corresponding author's e-mail address: lum@aber.ac.uk

Published by Oxford University Press on behalf of the Annals of Botany Company. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Nitric oxide (NO) emerged as a signal in plants during the last decade of the 20th century and has since come to be associated with a large number of phenomena. These have been extensively reviewed in a recent series of special reviews (Wendehenne and Hancock 2011). However, although impressive progress has been made, some persistent questions remain unanswered. Most fundamentally, various means of NO generation have been described, but the contexts of when and where they are deployed need a fuller description. In particular, the underlying basis of frequently reported arginine-oxidizing ‘nitric oxide synthase’-like activity requires final resolution. It must also be admitted that NO detection methods are often inadequate and that the accuracy and tissue specificity of methods used for measuring *in planta* NO production can be questioned. We also suggest that more attention should be paid to understanding how NO metabolism in plants interferes with and contributes to the larger scale nitrogen cycle. The purposes of this review are to highlight the progress made in plant NO research to date and to point out areas where further work is required.

Nitric oxide: some major themes from animal research

Nitric oxide has long been a major research topic in animals and some important aspects of this work merit a brief outline, thereby allowing a comparison with the current understanding in plants. The NO story in mammals started when it was noted that treatment of cultured macrophages with bacterial lipopolysaccharides (LPS) resulted in the production of NO (Hauschildt *et al.* 1990). Nitric oxide has emerged as an important component of innate resistance mechanisms elicited by pathogens or endotoxins such as LPS. In the innate resistance, NO generated by phagocytes may be directly antimicrobial via DNA damage or disruption of iron-sulfur (4Fe–4S) enzymes (Nathan 1995). Another important driver for NO research is its role in the regulation of smooth muscle contraction as its signalling pathways are pharmaceutical targets to mitigate the effects of cardiac infarctions (Murad 2004). This important role for NO grew out of observations that the vasodilatory effects of neurotransmitters such as acetylcholine were only observed when the endothelium cells covering the smooth muscle of the vessel wall were retained. Screens for soluble endothelium-derived relaxing factors led to the identification of NO as well as prostacyclin and endothelium-derived hyperpolarizing factor. To promote vasodilation, NO diffuses into smooth

muscle cells to bind to iron centres within the active site of a soluble guanylate cyclase complex which generates cGMP from GTP. The soluble forms of guanylate cyclase are complexes of α and β subunits, each unit ranging between 79 and 80 kDa, and each with a haem site and catalytic domains. cGMP activates protein kinase G, which in smooth muscle suppresses calcium influxes to reduce calcium-dependent muscle contraction. The effects of this NO-initiated pathway can be reversed through the activity of phosphodiesterases, which convert cGMP to GMP (Derbyshire and Marletta 2012). These phosphodiesterases are targets for Viagra (sildenafil) and their inhibition leads to a more persistent NO effect (Moreland *et al.* 1999).

In animals, an important mechanism of NO generation is the deamination of arginine by nitric oxide synthase (NOS) to form citrulline and NO. Nitric oxide synthase is homologous to P450 cytochrome c reductases, its activity being dependent on the reductant NADPH, flavin mononucleotide, flavin adenine dinucleotide (FAD) and tetrahydrobiopterin. The NOS group of enzymes is usually sub-classified as Ca^{2+} /calmodulin-activated brain NOS (nNOS), endothelial NOS (eNOS) and inducible NOS (iNOS). These NOS are associated, respectively, with the neuronal, smooth muscle relaxation and induction following immunological challenge (Forstermann and Sessa 2012). Besides NOS, mammalian tissues can generate NO through the reduction of NO_2^- in the mitochondrion via reduction at complex III or cytochrome c oxidase (complex IV) (Shiva 2010) or enzymes with nitrate reductase (NR) activity that may be xanthine oxidoreductases (XOR) (Jansson *et al.* 2008).

Plants join the NO party

Nitric oxide first came to prominence within the context of regulating plant defence during plant–pathogen interactions (Delledonne *et al.* 1998; Durner *et al.* 1998). Nitric oxide has been implicated in defence against *Pseudomonas syringae* pathogens (Delledonne *et al.* 1998; Clarke *et al.* 2000; Mur *et al.* 2005), in barley infected with powdery mildew and downy mildew on pearl millet (Prats *et al.* 2005; Manjunatha *et al.* 2009) or *Botrytis cinerea*-challenged *Arabidopsis* (Mur *et al.* 2012). As with mammalian systems, bacterial LPS, a contributor to pathogen-associated molecular patterns triggered immunity (PTI), proved to be a highly effective initiator of NO (Zeidler *et al.* 2004). Given these plant responses, it is unsurprising that many pathogens have evolved genes that could suppress NO-associated event(s). For example, *Erwinia chrysanthemi* expresses the flavohaemoglobin (fHb) *HmpX*,

which oxidizes NO to NO₃⁻ (Boccarda et al. 2005). In other cases, the pathogen may actively elicit host NO to aid in the infection process. For example, the virulence factor cryptogein produced by the oomycete *Phytophthora cryptogea* aids pathogenesis by promoting host cell death via NO generation (Foissner et al. 2000; Lamotte et al. 2004). In addition, pathogen-generated NO can promote the formation of key fungal infection structures (Prats et al. 2008; Turrion-Gomez and Benito 2011). Thus, depending on the pathogenic lifestyle, NO can act as either a pathogen virulence or a host defence factor (Perchepped et al. 2010; Turrion-Gomez and Benito 2011; Mur et al. 2012; Rasul et al. 2012).

Nitric oxide also plays an important role in symbiotic organisms, particularly between legumes and *Sinorhizobium* (Baudouin et al. 2006). Nitric oxide was first detected complexed to leghaemoglobin (within nitrogen-fixing nodules of cowpea and pea; Kanayama and Yamamoto 1991). Transcriptional analyses of legumes suggested that the NO played an early role in nodule development, being observed as early as during infection thread development within root hairs, through which *Sinorhizobium* colonizes the host (del Giudice et al. 2011). Indeed, NO may induce the expression of flavonoids which are essential in establishing bacterial *nod* gene expression, which initiates root hair deformations and represents one of the earliest stages of the symbiotic interaction (Dénarié et al. 1992). Categorical proof of an important role for NO in *Sinorhizobium* interactions with its host was obtained from *Medicago truncatula* plants expressing an NO-oxidizing fHb gene regulated by a nodule-specific promoter. Following inoculation with *Sinorhizobium meliloti* strain, nodule formation was significantly delayed. Transcriptional analysis indicated that the host gene *MtCRE1*, which encodes a cytokinin receptor, and the cell cycle-switching gene *MtCCS52A* were suppressed in expressing nodules (del Giudice et al. 2011). *MtCCS52A* triggers selected cells within the root primordium to switch from mitotic cycles to endoreplicating cycles (where genomes duplicate without cell division), which is essential for nodular symbiotic cell differentiation in *M. truncatula* (Vinardell et al. 2003). Moving beyond nitrogen fixation, NO generation also plays a role in symbiotic interactions involving arbuscular mycorrhizal fungi (Calcagno et al. 2012).

It is immediately obvious, even in this brief overview, that NO seems to be involved in a series of apparently incongruous events; it is involved in host defence (both cell death and PTI), pathogen virulence and also many forms of symbiotic interaction. This poses a question that is not only relevant to plant pathology: How exactly could NO fulfil these very different roles? It seems probable that the relative concentration of NO is important (Beligni

and Lamattina 1999; Turrion-Gomez and Benito 2011) but equally, cell-, tissue- or organelle-specific roles are vital, reflecting interactions with differing components and signalling pathways.

Spatio-temporal subtlety, as well as key roles for interactions with hormone signalling, is also a feature of the developmental actions of NO. Excellent examples of the role of NO in root development have been described in a series of papers produced by the Lamattina group. Thus, NO is required for root organogenesis (Pagnussat et al. 2002), the formation of adventitious roots (Pagnussat et al. 2003), lateral root development (Correa-Aragunde et al. 2004) and root hair formation (Lombardo et al. 2006). The role of auxin is very well established in various features of root development (Kramer and Bennett 2006) so that an important advance was made when NO and cGMP were implicated as downstream effectors of at least some auxin effects (Pagnussat et al. 2003). In adventitious root formation, auxin (indole acetic acid, IAA) was suggested to act through NO to activate mitogen-activated protein kinase (MAPK) signalling (Pagnussat et al. 2004) and through a modification of the auxin receptor TIR1 (Terrile et al. 2012). Ultimately, NO appears to influence root development through the initiation of cell cycle genes and patterns of cellulose synthesis (Correa-Aragunde et al. 2006, 2008), and influencing vesicle trafficking in root hair formation (Lombardo and Lamattina 2012). At the root apices, NO has been shown to influence the arrangement of the actin cytoskeleton (Yemets et al. 2011).

At the cross-roads between developmental and abiotic stress tolerance lies the regulation of the stomatal aperture by NO (Hancock et al. 2011). Early work showed that NO was produced in stomata and was an output of well-characterized abscisic acid (ABA) signalling pathways (Neill et al. 2002). Thus, an ABA-induced increase in cytoplasmic pH acts together with H₂O₂ to initiate NO generation. In *Vicia faba* guard cells, NO regulates Ca²⁺ release from intercellular Ca²⁺ stores, which regulates inward-rectifying K⁺ channels to close stomata (Garcia-Mata et al. 2003; Bright et al. 2006; Wilson et al. 2009). However, NO can appear to be a redundant element in stomatal regulation as in, for instance, conditions of rapid dehydration (Ribeiro et al. 2009). It is apparent that H₂O₂ effects can stimulate and at least partially parallel the effects of NO in *Arabidopsis* (Bright et al. 2006). This redox-sensitive step appears to involve the ethylene receptor ETR1, adding yet another level of NO hormone interactions at the stomatal level (Desikan et al. 2006). Additionally, NO concentration appears to be important in its effects in plants, with high amounts opening stomata (Sakihama et al. 2003). These effects were linked to NO effects on

outward-rectifying K^+ channels which are Ca^{2+} insensitive, possibly by direct modification of the K^+ channel by NO (Sokolovski and Blatt 2004).

A finer level of spatial effects is demanded when considering the differential intracellular role of NO. Various cellular compartments such as mitochondria (Gupta *et al.* 2009), peroxisomes (Corpas *et al.* 2009) and chloroplasts (Jasid *et al.* 2006) have been shown to produce NO. It is very probable that NO has a specific role in each compartment, possibly interacting with local signal events. For example, NO has recently been shown to modulate mitochondrial alternative oxidase activity to influence the generation of reactive oxygen species (ROS), net NO production and shift primary metabolism towards amino acid biosynthesis via inhibition of aconitase (Cvetkovska and Vanlerberghe 2012; Gupta *et al.* 2012). Another role for NO was indicated by the work of Palmieri *et al.* (2008), who showed that NO produced during plant–pathogen interactions can inhibit the P protein of glycine decarboxylase (GDC) activity by S-nitrosylation (see below) to promote the hypersensitive response (HR). In this case GDC inhibition could limit NADH to the electron transport chain and leads to a change in redox of the electron transport chain of mitochondria. In the peroxisome, Ortega-Galisteo *et al.* (2012) recently showed that the NO that is produced in peroxisomes nitrosylates proteins such as catalase and glyoxylate oxidase, which are involved in photorespiration, β -oxidation and the detoxification of ROS.

Taking some lessons from these papers reveals the need for a careful experimental design that needs to be considered by plant NO scientists. In particular, as far as possible, strategies should be followed where the subtlety of NO effects is not lost. An example of this could be through the use of pharmaceutical NO donors, which represent an easy method to apply NO exogenously. If the NO concentration hypothesis is correct, these should be coupled to a better means of visualizing and/or measuring *in situ* NO generation (see below). Experimentally, it is preferable to utilize approaches where the spatial, temporal and biochemical features of NO action in the biological phenomena under investigation are preserved. Admittedly, this is easier for such as stomata and root hairs but should be a major driver in experimental design for all NO biologists.

How NO is generated in plants?

As NO is now firmly established as an important signal in plant science, a remaining task is to describe the various mechanisms of NO generation in plants. After some wrong turns this is now much less of a contentious issue. Guo *et al.* (2003) identified a novel NOS in

Arabidopsis which had the same co-factor requirements as NOS but exhibited no significant sequence homology to the mammalian form. The derived mutant *atnos1* has proven to be a useful tool in NO research as it does display reduced NO production, and has been used to show the roles of NO in floral development (He *et al.* 2004) or the interaction of NO with ROS (Zhao *et al.* 2007). However, subsequently AtNOS1 was found not to possess NOS activity and was renamed AtNOA1 (Nitric Oxide Associated1; Zemojtel *et al.* 2006) and indeed has been established to be a GTPase (Moreau *et al.* 2008).

Currently, a seemingly bewildering number of sources for NO (at least seven) have been identified (Gupta *et al.* 2011a). Any discomfort that arises from this is most likely due to a comparison with mammalian cells, where NOS represents a bespoke NO-generating system with different forms having well-defined roles and expression patterns. In the apparent absence of a true plant NOS, it may be better to consider the varieties of ROS generation as a better paradigm for understanding NO production. Thus, although ROS generation is most often linked to NADPH oxidases, other sources, peroxidases, polyamine oxidases and non-enzymatically from photosynthetic and respiratory electron transport chains, have important roles (Wojtaszek 1997).

A series of reductive pathways for NO generation have been described (Gupta *et al.* 2011a), including a peroxisomally located XOR which reduces nitrite to NO at the expense of NADH under anaerobic conditions (Corpas *et al.* 2008) or a plasma membrane-bound nitrite: NO reductase (Ni-NOR) (Stöhr *et al.* 2001). However, it is cytosolic NR that is rapidly emerging as the main source of NO in plants under aerobic conditions. Nitrate reductase is implicated in the NO production during bacterially induced defence (Modolo *et al.* 2005), disease development in certain pathogenic interactions (Shi and Li 2008), drought (Freschi *et al.* 2010), cold (Zhao *et al.* 2009), stomatal regulation (Srivastava *et al.* 2009) and many developmental processes, for example, the initiation of flowering (Seligman *et al.* 2008). Nitrate reductase is a cytosolic enzyme that undergoes a regulatory switch from its preferential high-affinity substrate NO_3^- ($K_m \text{ nitrate} = <40 \mu\text{M}$) to NO_2^- (low affinity; $K_m \text{ nitrite} = 100 \mu\text{M}$) and producing NO. An important question to consider is how this switch comes about (Gupta *et al.* 2011a). Currently, regulation seems to be at the level of substrate competition so that high nitrite levels are required to competitively inhibit NO_3^- reduction. This could come about through either increased NO_3^- influx into the vacuole or efflux from the cell. In *Arabidopsis*, accumulation into the vacuole involves AtCLCa (*Arabidopsis thaliana* Chloride Channel a) which acts as a proton antiporter (Geelen *et al.* 2000). Thus, it is possible that

NO_3^- vacuolar import is promoted by cytoplasmic pH changes, possibly driven by tonoplastic H^+ -ATPases (De Angeli et al. 2007). This could enable the build-up of a 50-fold excess of nitrate in the vacuole compared with the cytoplasm (De Angeli et al. 2007). An alternative/additional mechanism could be to promote NO_3^- efflux from the cells. In the case of the HR elicited by cryptogin in tobacco cells, nitrate efflux was shown to be vital to cell death and defence. This resulted in a rapid 60 % drop in the concentration of internal NO_3^- (Wendehenne et al. 2002). Concomitant with NO_3^- efflux is an influx of calcium, on which NO production was completely dependent (Lamotte et al. 2004) and acts via calmodulin/calmodulin-like proteins (Ma et al. 2008). The longer-term generation of calcium is influenced by NO (Lamotte et al. 2006) and this integrates Ca^{2+} -NO effects into the web of positive and negative feedback loops that typify plant-pathogen interactions (Robert-Seilaniantz et al. 2011). In terms of triggering events, it seems likely that NO_3^- efflux is one of a series of calcium ion-mediated events which contribute to plasma membrane depolarization and also causing the extrusion of K^+ ions and water loss, all of which are early features of HR-type cell death (Garcia-Brugger et al. 2006). Clearly, more work is needed on the regulation of NO_3^- fluxes in disease responses and more widely, and especially the possible perturbation that could occur when plants are supplied with high levels of nitrogen fertilizer must be assessed.

As well as modulation of nitrogen flux within cells, direct regulation of NR itself should be considered. Nitrate reductase is a relatively labile protein, the levels of which are based on its relative expression and degradation. Thus, NR may be phosphorylated at a serine residue to interact with 14-3-3 proteins to inactivate the protein and, mostly likely, promote proteolysis (Kaiser and Huber 2001). However, the phosphorylation of NR is Ca^{2+} dependent for at least three types of kinase, and the formation of the inactive form is influenced by divalent cations (Weiner and Kaiser 1999, 2000). Thus, we need to understand why Ca^{2+} influxes which activate NO_3^- efflux and NO production do not also encourage NR degradation. This could be via the action of Type 2A phosphatases (Deruere et al. 1999) or, interestingly, hexose sugars via unknown mechanisms (Cotelle et al. 2000).

One other reductive mechanism of NO generation merits mention as it allows NO production under very small partial pressures of oxygen (Planchet et al. 2005). This involves a mitochondrial-based NR activity where NO_2^- acts as a terminal electron acceptor for cytochrome c oxidase/reductase (Castello et al. 2006). This can maintain some ATP generation under hypoxic conditions and will also reduce NO_2^- to generate NO which can be scavenged by non-symbiotic Hbs (Stoimenova et al. 2007).

The nitrite reductase activity of cytochrome oxidase/reductase is well established under hypoxia (Gupta et al. 2011a) but may be increasingly important as partial pressures of oxygen are reduced from ambient. The importance of this generational route has been demonstrated by our recent work where we studied NO production under various degrees of hypoxia and identified a threshold at 0.5 % O_2 , below which substantial NO is produced (Hebelstrup et al. 2012). This led to the production of ethylene which promotes petiole hypernasty and/or vertical elongation, an important plant mechanism for avoiding the negative effects of submergence in *Rumex palustris* and a wide variety of other species (Voesenek et al. 2006).

Moving on to consider oxidative mechanisms of NO generation in plants, there are frequent reports of NOS-like activity in plants, notwithstanding the absence of an isolated enzyme or gene (Corpas et al. 2009). Over the last decade the authors have described the NOS activities in peroxisomes (Barroso et al. 1999) and chloroplasts (Jasid et al. 2006), and in isolated root mitochondria (Gupta and Kaiser 2010). Further, this NOS-like activity shares several of the co-factor requirements for mammalian NOS (Corpas et al. 2009). There are also many examples of the use of arginine-based analogues, which inhibit mammalian NOS, to suppress plant NO production, including from our work (Mur et al. 2005, 2008). Independent confirmation of this arginine pathway has come from *Arabidopsis* arginase mutants which exhibited increased levels of NO production (Flores et al. 2008). Taking all of this work together, we are left with strong data suggesting that an arginine-utilizing pathway can generate NO, although this need not be one that leads to the co-production of citrulline, an important characteristic for true NOS (Tischner et al. 2007). The vagueness of this situation begs the question of how specific really is this arginine oxidizing capacity. For example, another oxidative pathway leading to NO generation based on polyamines has been described (Tun et al. 2006). The mechanism through which it acts is also unknown, but although very different, mammalian NOS could be a target for mammalian NOS inhibitors (Gupta et al. 2011a). Given the chequered history of NOS studies in plants, the sources of NOS-like activity in plants and why arginine-based inhibitors are effective in plants need resolving.

Recently, it was discovered that NR-free plant cells are able to oxidize externally supplied hydroxylamine (HA) to NO (Rumer et al. 2009a, b), a pathway that is well characterized in bacteria and animal systems (Vetrovsky et al. 1996). Conditions that increase ROS are able to increase the NO production from HA, for instance increasing ROS by mitochondrial complex III inhibitor

myxothiazol enhanced NO production from HA (Rumer et al. 2009b). Interestingly, it was recently shown that isolated non-rice non-symbiotic haemoglobin 1 enzyme could reduce NO_2^- to NO with a rate constant that was far in excess of that reported for Hbs (Sturms et al. 2011). Given such an observation, the endogenous production of HA by plants needs to be characterized and the possible enzymatic basis of NO generation from this substrate should also be a target for further research.

How much NO is made and where?

If the sources of NO generation are now coming to be well characterized, the next problem that needs addressing is the patterns of NO generation. We have already highlighted how several authors are suggesting that the concentration of NO is important for its activity (for example, Beligni and Lamattina 1999; Garcia-Mata et al. 2003); therefore, actual measures of *in planta* NO should be a major feature of NO studies. However, as outlined in our recent review (Mur et al. 2011), nearly all the approaches currently employed have associated problems.

The most superficially attractive approach is to use diaminofluoresceins (DAF)—fluorescent dyes that are available from many manufacturers. Diaminofluorescein dyes react with N_2O_3 , a by-product of NO oxidation, with a resulting dramatic increase in fluorescence to allow an approximate quantification of NO content (Kojima et al. 1998a, b). Diaminofluorescein dyes are sensitive NO sensors with detection limits in the nanomolar range and specific to NO as no increased fluorescence was observed with NO_2^- , NO_3^- , H_2O_2 and peroxynitrite (ONOO^-) (Kojima et al. 1998b). When used in conjunction with confocal microscopy, DAF dyes offer the possibility of exactly defining the site of NO generation. As suggested above, such a precise tissue resolution is essential in order to begin to assign roles for NO in plant development and in plant–microbe interactions. However, the specificity of DAF dyes has recently been questioned and it is also possible that a differential take-up of the dye by some tissue types or organelles could give artefactual results (Mur et al. 2011). For example, Rumer et al. (2012) have shown that reactions involving horseradish peroxidase and H_2O_2 were sufficient to generate DAF fluorescence *in vitro*. This stated, there is currently no well-established alternative approach to reveal high tissue-specific patterns of NO generation. We therefore expect DAF dyes to be continued to be used. We have therefore recommended that users employing DAF dyes should follow the following steps (Mur et al. 2011). First, the background fluorescence of tissues in the absence of the

DAF dye should be assessed. Then, most importantly, a fluorescence in the presence of DAF (not the absence of) dyes should be suppressed by co-application of NO scavengers (for example, 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazoline-l-oxyl-3-oxide (cPTIO); $\text{NO} + \text{cPTIO} \rightarrow \text{NO}_2^- + \text{cPTI}$) or inhibitors of mammalian NOS, although this last approach may be questioned. Additionally, we note with interest that newer fluorescent probes have been developed, although these have yet to be widely used by the plant NO community. These include rhodamine B (2-amino-3',6'-bis(diethylamino)-2,3-dihydrospiro [iso-indole-1,9'-xanthene]-3-one)-based dyes which can act to detect NO_2^- and ONOO^- (Rieth and Sasamoto 1998; Yang et al. 2002). In a comprehensive assessment of the specificity of rhodamine B dyes, the fluorescence response was greatest with NO and ONOO^- . However, fluorescence was also significant with hydroxyl radicals (Wu et al. 2011), which are likely to be generated in stressed plants (Apel and Hirt 2004). Europium(III) chelate has recently been advanced as an NO fluorescent probe but its wider specificities are yet to be fully assessed (Chen et al. 2011).

To be confident in the NO measurements obtained, it is preferable that alternative approaches are employed in parallel. The Griess assay is an indirect colorimetric assay for NO that is available as a kit. In this assay, NO is oxidized to NO_2^- which reacts with sulfanilic acid and α -naphthylamine under acidic conditions to yield an azo dye. The sensitivity of the Griess assay can be improved by employing a flow-through system where NO emitted from a plant or cell culture is passed via a flow into a receiving vessel where the Griess assay is carried out; the dye accumulates as more NO enters the vessels (Vitecek et al. 2008). Another highly popular commercially available means of NO detection from the gas phase is chemiluminescence. The detection system is based on the reactivity of NO with O_3 , which produces excited-state nitrogen dioxide (NO_2^*), which emits a photon upon relaxation to the ground state. Therefore each photon is related to a single NO molecule; the NO concentration is then determined by measuring the light intensity. The chemiluminescent approach has been utilized by the present authors and has proved to be a highly sensitive and accurate measure of NO emission from plants (Planchet et al. 2005; Mur et al. 2011). Its main drawback concerns its lack of selectivity, as molecules such as water can dampen the chemical reaction and lead to erroneous NO measurements. Two additional platforms that we have extensively used to measure NO in the gas phase are laser photoacoustic detection (LPAD) and tunable diode laser absorption spectroscopy (TDLAS) (Mur et al. 2005, 2011, 2012), neither of which is currently available in a commercial form.

Both target the strongest absorption band of NO centred at 5.3 μm (1876 cm^{-1}) (Rothman et al. 2005), and use different laser sources such as CO gas lasers, quantum cascade lasers (QCL) or interband cascade lasers. Compared with other methods, laser-based systems directly measure NO molecules and are extremely selective. In LPAD the absorption of NO in bursts of laser light results in pressure variations that generate sound, which is detected by a microphone located within a photoacoustic cell. Traditional LPAD systems require high-power sources and are not user friendly, while compact QCLs are still unable to reach the sub-ppb detection limit (Kosterev et al. 2002; Spagnolo et al. 2010). However, TDLAS systems are offering the better potential for miniaturization and commercialization. In combination with a thermoelectrically cooled QCL, the key part of TDLAS systems is a multipass absorption cell where the light undergoes multiple reflections between two mirrors. This represents an interaction path length with the NO gas sample of 76 m in a compact design. It allows the detection of NO at and below 1 ppb by volume with a second measuring time (Cristescu et al. 2012). Each of these gaseous detection platforms offers the ability to obtain multiple real-time measures of NO production from plants, but there are some important deficiencies. The relationship between the concentration of *in planta* NO and that lost by the plant to the atmosphere is difficult to establish; also it is difficult to measure tissue-specific patterns of NO generation and it is impossible to measure organellar production.

The requirement for accurate measurements of NO production is also important due to the widespread use of chemical NO donors as surrogates for biologically generated NO production in experiments. Many NO donors have been developed, often for pharmaceutical use (Napoli and Ignarro 2003), and are readily obtained commercially. Nitric oxide donors include NONOates (spermidine- or diethylamine-NONOate) (Keefer et al. 1996), S-nitroso-N-acetylpenicillamine (SNAP), S-nitrosoglutathione (GSNO) and sodium nitroprusside (SNP). In plant biology NONOate has been used in, for example, the analysis of pathogen-associated cell death (Lamotte et al. 2004) and mitochondrial function (Fu et al. 2010). Example studies using SNAP to show NO effects have focused on wound healing (Orozco-Cardenas and Ryan 2002), thermotolerance (Xuan et al. 2010) and root meristem development by influencing the key auxin effector PIN1 (Fernandez-Marcos et al. 2011).

Sodium nitroprusside is frequently used as an NO donor, including by ourselves to validate the LPAD approach (Mur et al. 2005) or to investigate the bolting time in *Arabidopsis* (Hebelstrup and Jensen 2008). Others have used SNP to show a role in

photomorphogenesis by regulating phytochrome and gibberellin signalling (Lozano-Juste and Leon 2011). Sodium nitroprusside is in fact an NO^+ donor, which is a highly stable electrophile but can be stabilized by coordination with metals (Roncaroli et al. 2007). NO^+ readily forms nitrosothiolate adducts which can be reduced (for example, $\text{GSNO} + \text{H}^+ \rightarrow \text{GSH} + \text{NO}^+$) to generate NO gas (Wang et al. 2002). An additional problem with SNP is the activity of the 'spent' donor (Bethke et al. 2006). When assessing the ability of NO to break *Arabidopsis* seed dormancy, it was found that SNP and the 'spent donor' products potassium ferricyanide (Fe (III) CN) and potassium ferrocyanide (Fe (II) CN) were all acting via the generation of cyanide (CN^-). To counter this problem, experimenters must include controls involving the NO 'spent' controls (i.e. the remaining products when all NO has been produced).

S-nitrosoglutathione is often used as a 'clean' NO donor in nitrosylation studies (see below) as it presents no known problem with spent products. This undergoes spontaneous homolytic cleavage of the Cys-based S-NO bond to release NO (Ederli et al. 2009). The S-nitrosylation process involves initial reaction with O_2 to form substances such as N_2O_3 , which dissociates to form the nitrosonium ion, NO^+ . The electrophile NO^+ will attack thiolate to form S-NO adducts.

Nitric oxide concentration is clearly important for action but there have been very few attempts to assess the *in planta* kinetics of NO generation by NO donors. Ederli et al. (2009) investigated the *in planta* production of NO from SNP, GSNO and injections of mammalian NOS using the oxyhaemoglobin NO assay method. Similarly, we have examined the kinetics of NO production from donors following infiltration of tobacco leaves using our QCL-based approach (Fig. 1). The NO donors diethylamine nitric oxide (DEANO) (Fig. 1A) and SNAP (data not shown) proved to give rise to a short burst of NO. S-nitrosoglutathione increased NO for a longer period (Fig. 1B) but SNP gave rise to a more persistent pattern of NO generation (Fig. 1C). This pattern of NO production was similar to that observed during the HR in tobacco elicited by the bacterial pathogen *P. syringae* pv. *phaseolicola*. Taking the above points together, we experimenters need to be circumspect when using NO donors and give thought to *in planta* patterns and amounts of NO production as well as the possible effects of spent donor products.

How far is there a discrete NO signalling module?

As the mechanisms of NO generation come to be established, the mechanisms of NO signalling are increasingly




Fig. 1 Nitric oxide production from tobacco leaves infiltrated with NO donor chemicals or *P. syringae pv. phaseolicola*. Nitric oxide production was measured using a QCL from tobacco leaves removed from 5-week-old plants and infiltrated with various concentrations of (A) diethylamine nitric oxide (DEANO); (B) *S*-nitrosoglutathione (GSNO) or (C) sodium nitroprusside and compared with (D) infiltrations with suspensions of 10^6 cells mL^{-1} 10 mM MgCl_2 of the HR-eliciting *P. syringae pv. phaseolicola* (*PspH*). Results are given as mean NO production: $\text{nmol g}^{-1} \text{h}^{-1}$ ($n = 3$) \pm SE. Plant cultivation conditions and bacterial culture are described in Mur et al. (2005). Quantum cascade laser measurement protocols are described in Mur et al. (2012).

being resolved. Initial work focused on transcriptomic assessments of NO effects, which revealed the importance of NO-regulated antioxidant responses (Krause and Durner 2004) or the possible roles of iron-deficiency genes in NO-mediated responses to Cd^{2+} (Besson et al. 2007). Currently, many groups concentrate on two protein modifications which are specific to NO signalling: thiol protein *S*-nitrosylation and tyrosine nitration, to reveal target signalling components and modes of action. Excellent reviews are available describing the current status of our understanding of the roles of *S*-nitrosylation and nitration (Astier et al. 2011; Malik et al. 2011; Vandelle and Delledonne 2011) and so we will confine ourselves to a general overview and some general points that emerge.

The regulation of proteins by *S*-nitrosylation is attracting a lot of interest as an important reversible post-transcriptional modification. The following represents some elegant regulatory mechanisms which have recently been reported in the literature. *Nonexpressor of pathogenesis related proteins 1* (NPR1) is a key regulator within the signalling cascade of the key defence hormone salicylic acid. In its non-activated state, NPR1 exists as a cytoplasmically located oligomer with each monomer linked by redox-active disulfide bridges. When activated, cytoplasmic changes lead to reduction

of the thiol groups to release monomeric NPR1 forms which are translocated to the nucleus (Mou et al. 2003). Apparently, gratuitous translocation of NPR1 into the nucleus leads to proteasome-mediated digestion (Spoel et al. 2009). In the nucleus, NPR1 interacts with a range of TGA-class transcription factors, some of which are involved in binding to *as-1* elements to activate *pathogenesis-related protein 1* gene expression. Thus, the oligomeric status of NPR1 is essential to its action and *S*-nitrosylation of cysteine-156 has been shown to facilitate oligomerization (Tada et al. 2008). The de-*S*-nitrosylation process is also receiving considerable attention, with particular focus being the roles of thioredoxins (TR). With NPR1 it is the *S*-nitrosylated cysteine-156 that is reduced by TR to promote monomer formation (Tada et al. 2008). These observations neatly link oxidative and salicylic acid signalling modules with NO events. In stomata, NO and oxidative events are well understood to work in tandem (Wilson et al. 2009) and this is clearly also the case during plant defence. Interestingly, NO is well known to influence the biosynthesis of salicylic acid (SA) so that this could promote NPR1 signalling, while NO, presumably by promoting GSNO, could also encourage oligomerization. These could be seen to be contradictory events but GSNO also promotes the NPR1–TGA1 interaction to

strengthen binding to cognate promoter sequences, thereby elevating the expression of *pathogenesis-related protein* gene expression (Lindermayr et al. 2010).

Other subtle interactions between NO/GSNO, SA and oxidative stress are also revealed by the recent work of Yun et al. (2011). SA is known to drive the speed of the HR by augmenting the generation of oxidative stresses, but these authors suggested that at lower GSNO concentrations cell death was augmented by other as yet unknown mechanisms. At higher concentrations of S-nitrosothiol, the activity of the ROS-generating enzyme NADPH oxidase, AtRbohD, was suppressed. This was shown to arise from S-nitrosylation of a cysteine-890 to affect enzyme binding to its key cofactor FAD. Thus, S-nitrosylation of NADPH oxidase appears to be a mechanism through which the plant can regulate oxidatively driven cell death. However, other members of the NADPH gene family have other roles, for example AtRbohC in root development and root hair formation (Foreman et al. 2003), so this regulatory mechanism could have much wider physiological roles.

Tyrosine nitration is dependent on peroxynitrite, which is generated via superoxide and NO ($O_2^- + NO \rightarrow ONOO^-$). This is a highly toxic molecule that can generate hydroxyl radicals ($ONOO^- + H^+ \rightarrow NO_2^- + OH$) and cause considerable macromolecular damage via proton abstraction, and could lead ultimately to cell death as in mammalian systems (Pacher et al. 2007). Alternatively, NO_2^- may be added at the *ortho* position on the tyrosine aromatic ring, which could exert steric effects to alter the configuration of a protein. Compared with studies on S-nitrosylation, there have been relatively few studies on nitration, possibly because the latter's irreversible nature made it a less attractive regulatory switch. However, one of the most elegant studies has focused on the role of nitration during the HR in *Arabidopsis*, which revealed a novel regulation of peroxiredoxins (Prx). Peroxiredoxins can detoxify the highly reactive peroxynitrite ion and this has been demonstrated by two plastid-located Prx: PrxIIIE and 2-Cys-Prx (Sakamoto et al. 2003; Romero-Puertas et al. 2007). If $ONOO^-$ contributes to HR-mediated cell death, then Prx would suppress this, but PrxIIIE is inhibited by S-nitrosylation, thereby allowing the propagation of $ONOO^-$ (Romero-Puertas et al. 2007; Cecconi et al. 2009). Clearly, the identification of nitrated proteins is an ongoing process, but it also highlights the need to develop an organelle-specific understanding of NO effects as, due to its reactivity, it is unlikely that $ONOO^-$ will move out of the plastids to nitrate cytoplasmic proteins.

Taking the S-nitrosylation/nitration studies together, some tentative observations can be made. Invariably, it has been shown that NO modifies the activity of

enzymes and some key signalling components, and many have suggested an integration between NO/GSNO-SA-ROS-associated events. Although this may simply reflect the research interest of the main workers in the field, it may suggest that NO effects could be exerted to a large extent by influencing the redox status of the cell. Another feature is that S-nitrosylation/nitration studies seem to be suggesting that NO modifies signalling pathways which have been characterized as part of the action of another signal. Thus, there are redox-associated proteins, PrxIIIE (Romero-Puertas et al. 2007), AtRbohD (Malik et al. 2011), SA signalling pathway proteins salicylic acid-binding protein 3 (Wang et al. 2009), NPR1 (see above) (Tada 2009), TGA1 (Lindermayr et al. 2010) and auxin signalling TIR1 (Terrile et al. 2012). Other pathways such as jasmonates are also being suggested (Manjunatha et al. 2012; Mur et al. 2012). Thus, NO could also mainly act as a modifier of other signalling cascades. This was also suggested from a bioinformatic analysis of 'NO-responsive' promoters where salicylate- and jasmonate-responsive *cis*-elements were prominent (Palmieri et al. 2008). This stated, it is possible that further characterization of S-nitrosylative control of the R2R3-MYB class of transcription factors may identify an NO-specific transcriptional output.

Modulation of guanylate cyclase represents an NO-specific mechanism of signalling and there are many reports showing that cGMP is a facet of NO effects in plants (Klessig et al. 2000; Pagnussat et al. 2003; Szmjdt-Jaworska et al. 2004, 2008, 2009; Suita et al. 2009; Li and Xue 2010; Wang et al. 2010; Dubovskaya et al. 2011). Indeed, even Viagra has been shown to have effects in plants (Siegel-Itzkovich 1999). Paradoxically, until recently, higher plants were thought to lack the required soluble form of guanylate cyclase (Schaap 2005), although it is present in single-cell algae (de Montaigu et al. 2010). Indeed, in *Chlamydomonas*, a soluble guanylate cyclase influenced NR expression (de Montaigu et al. 2010). Recently, a novel guanylate cyclase that generates cGMP and binds NO has been described (Mulaudzi et al. 2011). It is imperative that this be extensively characterized and integrated into our existing knowledge of NO networks.

How does plant-generated NO fit into the nitrogen cycle economy?

An important point frequently disregarded by plant NO scientists is that plants are being continually exposed to NO from the soil. Nitric oxide production is a feature of the oxido-reductive steps ranging from NH_4^+ to NO_3^- that form the nitrogen cycle. Various factors also influence NO production in soil such as temperature,

oxygen availability, humidity, soil pH and nitrogen status. These influence the activities of nitrifying and denitrifying bacteria which under different conditions can produce NO at differing rates (Anderson and Levine 1986).

If the plant is being continuously exposed to NO, how can NO function as a discrete endogenous signalling molecule, particularly in the root? We suggest that these questions highlight the importance of *in planta* NO scavenging mechanisms as key in NO biology. Some simple chemicals such as urate have been shown to prevent NO toxicity (Alamillo and Garcia-Olmedo 2001), but more selectivity is offered by enzymatic regulation by GSNO reductase (GSNOR) (Malik et al. 2011) and nsHb (Gupta et al. 2011b).

Pools of reduced glutathione (GSH) are readily available within plants and GSH can be oxidized by NO to form GSNO. This represents a stable reservoir of potential NO signal so that the regulation of GSNO content represents an important regulatory step in NO regulation. The key enzyme regulating GSNO pools is GSNOR; GSNOR will reduce GSNO to ultimately produce glutathione disulfide (GSSG) and ammonia (NH₃), and GSSG can be reduced by glutathione reductase to re-enter the GSH pool. In *Arabidopsis*, GSNOR exists as a single gene (AtGSNOR1; At5g43940), resulting in an increase in nitrosothiolates such as GSNO and thus also increases in S-nitrosylated proteins (Feechan et al. 2005). Thus, GSNOR represents an important enzyme regulating indirect NO effects via S-nitrosylation.

Our interest has focused on the role of the direct oxidation of NO by nsHb and its role in regulating NO content within plants (Gupta et al. 2011b; Hebelstrup et al. 2012; Mur et al. 2012). Plant Hbs are able to regulate several of the effects of NO, as recently reviewed by Hill (2012). Plant Hbs may be sub-divided into three classes: I, II and III. Most Hbs found in association with nitrogen-fixing bacteria in root nodules of plants appear to have evolved from class II Hb, which has a relatively low affinity for O₂ (K_m ~ 150 nM), so that this is readily released under low partial pressures of O₂. As such, the functions of most of those Hbs called 'symbiotic haemoglobins' are in facilitating oxygen supply to tissues within nitrogen-fixing nodules. However, this requires a high concentration of Hb. Other class II 'non-symbiotic' Hbs (nsHbs) are found in other tissues at low concentration where the contribution to facilitated oxygen diffusion is negligible (Heckmann et al. 2006). We have previously shown that such class II nsHbs do contribute to NO removal when over-expressed (Hebelstrup et al. 2006, 2012). Class III Hbs are truncated Hbs with a very low affinity for O₂ (K_m ~ 1500 nM) and, given their closer homology to bacterial

Hbs, they may have been acquired by horizontal gene transfer (Watts et al. 2001). The function of these truncated Hbs is obscure and no observable phenotype was noted in *gfb3*, an *Arabidopsis* class III mutant (Mur et al. 2012). Class I nsHbs possess very high affinity for O₂ (2 nM) so that they represent poor oxygen carriers (Smagghe et al. 2009). Given this it appears that NO oxidation is an important role. During hypoxic/anoxic conditions, the oxidation of NO to NO₃⁻ by oxyhaemoglobin [Hb(Fe²⁺)O₂] is coupled to the reduction of NO₃⁻ and NO₂⁻, resulting in an Hb/NO cycle (Dordas et al. 2004). In this Hb/NO cycle, excess NAD(P)H is oxidized to maintain electron flow and ATP production under hypoxic conditions (Dordas et al. 2003; Stioménova et al. 2007). Nitric oxide oxidation by Hb(Fe²⁺)O₂ results in the formation of oxidized ferric metHb [Hb(Fe³⁺)] and so the reaction is $\text{Hb(Fe}^{2+}\text{)O}_2 + \text{NO}^+ \rightarrow \text{Hb(Fe}^{3+}\text{)} + \text{NO}_3^-$. Haemoglobin can be reduced back to its ferrous form [Hb(Fe²⁺)] by an associated reductase $2\text{Hb(Fe}^{3+}\text{)} + \text{NAD(P)H} \rightarrow 2\text{Hb(Fe}^{2+}\text{)} + \text{NAD(P)}^+ + \text{H}^+$. The NO₃⁻ produced is reduced to NO₂⁻ by NR ($\text{NO}_3^- + \text{NAD(P)H} \rightarrow \text{NO}_2^- + \text{NAD(P)}^+ + \text{OH}^-$) and subsequently by mitochondrial nitrite NO-reductase activity (Mt NINOR) at complex III and cytochrome c oxidase NO₂⁻ is reduced back to NO to restart the cycle $2\text{NO}_2^- + \text{H}^+ + \text{NAD(P)H} \rightarrow 2\text{NO} + \text{NAD(P)}^+ + 2\text{OH}^-$. Following this reasoning, hypoxically generated NO could improve the plant's energy status by adding to the Hb/NO cycle (Igamberdiev and Hill 2009); and possibly not only during hypoxia but also in bulky tissues where low internal oxygen is present in the centre of tissues. This, again, should be actively explored in future. Besides the Hb/NO cycle, the role of nsHb in NO removal has also attracted considerable interest. Early work used over-expressed bacterial Hb *hmpX* in transgenic lines as a useful method to reduce NO production and show the roles for NO in the HR, responses to UV-B, symbiotic interactions and senescence (Zeier et al. 2004; Boccara et al. 2005; Mishina et al. 2007; del Giudice et al. 2011; Tossi et al. 2011). The logical inference of this work is that nsHbs reduce endogenous NO production from plants and possibly the environment. We have provided evidence for this in our recent work where unstressed *Arabidopsis* lines with reduced expression of nsHbs [*GLB1* (At2g16060) and *GLB2* (At3g10520)] exhibited increased NO production (Hebelstrup et al. 2012; Mur et al. 2012). When NO-generating systems are deployed, it may be assumed that the NO generated by (for example) NR will swamp the scavenging capacity of nsHbs. However, some of our data have revealed an interesting regulatory mechanism whereby Hb expression (*GLB1*) is reduced in a manner which is apparently inversely correlated with the patterns of NO generation (Mur et al. 2012). Although we focused on responses to pathogens,

interrogation of transcriptome data in the Genevestigator database (Zimmermann et al. 2004) has suggested that *GLB1* (the major Hb) is also suppressed in response to heat, Fe deficiency, salt and drought stress (Fig. 2). Interestingly, in stresses linked to low oxygen caused by flooding (both root and shoot expression), hypoxia or anoxia where the Hb would be expected to contribute to plant fitness via the Hb/NO cycle, *GLB1* is induced. Similarly, in response to external nitrate when both NO-generating NR and Hb would be required, the expression of genes encoding these proteins appeared to be co-regulated in maize (Trevisan et al. 2011). There are several indications that plant Hbs can control development and physiological reactions by modulating cellular NO levels, which interfere with the actions of various hormones (Hill 2012). For example, *Arabidopsis* plants with silencing of *GLB1* (class I nsHbs) exhibit increased cellular NO levels, resulting in modified development with stunted organs, loss of apical dominance and late flowering (Hebelstrup et al. 2006; Hebelstrup and Jensen 2008). Thus, it would seem that coupled to characterization of the NO generation mechanism, the means through which Hb expression is regulated is as important and this is actively being investigated by ourselves and others. Comparisons of Genevestigator expression data for *GLB1* with those of the SA

marker genes *PR1* and the jasmonate marker gene *PDF1.2* suggested that these defence signals were unlikely to be playing a key regulatory role (Fig. 2). Another aspect that needs considering is how far Hb could help to preserve plant nitrogen. Our recent results using *glb1* suppressed lines have suggested that in the absence of Hb, nitrogen loss during hypoxia via NO generation is substantial: $\sim 0.2 \text{ mmol NO}_3^- \text{ g FW}^{-1}$ lost over a 24-h period (Hebelstrup et al. 2012).

Conclusions and forward look

To conclude, in the course of this review we have highlighted only some basic and persistent questions. The sources of NO generation have now been extensively defined but NO generation from polyamines, HA and especially arginine has resisted elucidation. This may be because no appropriate mutants, genes or proteins have been identified. Thus, plant biologists have been lucky that *NIA1* has proven to be a major source of NO despite some functional redundancy with *NIA2*. Thus, the *nia1* mutant exhibits reduced NO production even when *NIA2* is still functional (Wilson et al. 2009). However, for other NO generation mechanisms, problems with lethality, functional redundancy or their activation only under precise conditions (for example, normoxia and hypoxia) may be the reason that no generation mutants have been isolated. Thus, it may be that the plant ROS field offers a salutary lesson, as here generation mechanisms have often been characterized via biochemical means. This also highlights another theme of our review, the necessity to develop a better means of measuring NO, both to assay NO generation and the site of its generation. Currently, no technique fully meets all these requirements but we have noted ongoing developments in fluorescent dyes that could ultimately provide NO scientists with a key resource.

Moving to consider NO signalling, currently a major focus is on S-nitrosylation and nitration events. We hope that our review of the early work suggesting that NO acts with cGMP (for example, Durner et al. 1998) will serve to inspire a revisiting of this possibility and may, incidentally, reveal a signalling pathway that is similar to that found in animals. Our last theme is one that is, understandably, often not considered by laboratory-based plant scientists, namely how do plant signalling pathways function in an open environment. This is particularly apposite for NO as plants are being exposed to this signal from many exogenous sources. We therefore suggest that NO scavenging, e.g. by endogenous Hb, should be considered to be as important as NO generation in understanding *in planta* NO signalling.




Fig. 2 Transcriptional regulation of *Arabidopsis* non-symbiotic Hb genes in response to abiotic stress. Transcriptomic data for the non-symbiotic Hbs *GLB1* (At2g16060), *GLB2* (At3g10520) and *GLB3* (At4g32690) and for comparative purposes the salicylic acid marker gene *PR1* (At2g14610) and jasmonic acid marker gene *PDF1.2* (At5g44420) were extracted from the Genevestigator database (Zimmermann et al. 2004). The data were grouped by hierarchical cluster analysis and displayed using a heat map using Eplust software. Green represents down-regulation compared with controls and red up-regulation. Black squares indicate no net change in expression. The fold up/down regulation range is indicated.

Sources of funding

The work of L.A.J.M. is supported by grants from the BBRSC and the Centre for Integrated Research in the Rural Environment (CIRRE; <http://www.cirre.ac.uk/>), UK; that of K.H.H. by the Danish Council for Independent Research Technology and Production Sciences; that of K.J.G. by a Marie Curie Intra European Fellowship for Career Development within the 7th European Community Framework Programme; and that of J.M., S.M.C. and F.J.M.H. by the EU-FP6-Infrastructures-5 programme, project FP6-026183 Life Science Trace Gas Facility.

Contributions by the authors

Most of the text of this review was written by L.A.J.M., K.H.H. and K.J.G. Comments and editorial changes on approaches to NO measurement were provided by J.M., S.M.C. and F.J.M.H., while S.P. aided L.A.J.M. in measuring *in planta* NO production following tobacco leaf infiltration with NO donors. I.E.M. and G.V.N. provided comments and edited the section on NO-associated signalling events. Overall editorial comments and changes were provided by M.A.H.

Acknowledgements

The authors appreciate the comments made by the anonymous reviewers, which improved the manuscript.

Conflict of interest statement

None declared.

Literature cited

- Alamillo JM, Garcia-Olmedo F. 2001. Effects of urate, a natural inhibitor of peroxynitrite-mediated toxicity, in the response of *Arabidopsis thaliana* to the bacterial pathogen *Pseudomonas syringae*. *The Plant Journal* **25**: 529–540.
- Anderson IC, Levine JS. 1986. Relative rates of nitric oxide and nitrous oxide production by nitrifiers, denitrifiers, and nitrate respirers. *Applied and Environmental Microbiology* **51**: 938–945.
- Apel K, Hirt H. 2004. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annual Review of Plant Biology* **55**: 373–399.
- Astier J, Rasul S, Koen E, Manzoor H, Besson-Bard A, Lamotte O, Jeandroz S, Durner J, Lindermayr C, Wendehenne D. 2011. S-nitrosylation: an emerging post-translational protein modification in plants. *Plant Science* **181**: 527–533.
- Barroso JB, Corpas FJ, Carreras A, Sandalio LM, Valderrama R, Palma JM, Lupianez JA, del Rio LA. 1999. Localization of nitric-oxide synthase in plant peroxisomes. *Journal of Biological Chemistry* **274**: 36729–36733.
- Baudouin E, Pieuchot L, Engler G, Pauly N, Puppo A. 2006. Nitric oxide is formed in *Medicago truncatula*–*Sinorhizobium meliloti* functional nodules. *Molecular Plant-Microbe Interactions* **19**: 970–975.
- Beligni MV, Lamattina L. 1999. Is nitric oxide toxic or protective? *Trends in Plant Science* **4**: 299–300.
- Besson A, Gravot A, Pugin A, Wendehenne D. 2007. NO contributes to cadmium toxicity in *Arabidopsis thaliana*. *Comparative Biochemistry and Physiology A—Molecular & Integrative Physiology* **146**: S262–S263.
- Bethke PC, Libourel IGL, Jones RL. 2006. Nitric oxide reduces seed dormancy in *Arabidopsis*. *Journal of Experimental Botany* **57**: 517–526.
- Boccarda M, Mills CE, Zeier J, Anzi C, Lamb C, Poole RK, Delledonne M. 2005. Flavohaemoglobin HmpX from *Erwinia chrysanthemi* confers nitrosative stress tolerance and affects the plant hypersensitive reaction by intercepting nitric oxide produced by the host. *The Plant Journal* **43**: 226–237.
- Bright J, Desikan R, Hancock JT, Weir IS, Neill SJ. 2006. ABA-induced NO generation and stomatal closure in *Arabidopsis* are dependent on H₂O₂ synthesis. *The Plant Journal* **45**: 113–122.
- Calcagno C, Novero M, Genre A, Bonfante P, Lanfranco L. 2012. The exudate from an arbuscular mycorrhizal fungus induces nitric oxide accumulation in *Medicago truncatula* roots. *Mycorrhiza* **22**: 259–269.
- Castello PR, David PS, McClure T, Crook Z, Poyton RO. 2006. Mitochondrial cytochrome oxidase produces nitric oxide under hypoxic conditions: implications for oxygen sensing and hypoxic signaling in eukaryotes. *Cell Metabolism* **3**: 277–287.
- Cecconi D, Orzetti S, Vandelle E, Rinalducci S, Zolla L, Delledonne M. 2009. Protein nitration during defense response in *Arabidopsis thaliana*. *Electrophoresis* **30**: 2460–2468.
- Chen YG, Guo WH, Ye ZQ, Wang GL, Yuan JL. 2011. A europium(III) chelate as an efficient time-gated luminescent probe for nitric oxide. *Chemical Communications* **47**: 6266–6268.
- Clarke A, Desikan R, Hurst RD, Hancock JT, Neill SJ. 2000. NO way back: nitric oxide and programmed cell death in *Arabidopsis thaliana* suspension cultures. *The Plant Journal* **24**: 667–677.
- Corpas FJ, Carreras A, Esteban FJ, Chaki M, Valderrama R, del Rio LA, Barroso JB. 2008. Localization of S-nitrosothiols and assay of nitric oxide synthase and S-nitrosoglutathione reductase activity in plants. *Methods in Enzymology* **437**: 561–574.
- Corpas FJ, Palma JM, del Rio LA, Barroso JB. 2009. Evidence supporting the existence of L-arginine-dependent nitric oxide synthase activity in plants. *New Phytologist* **184**: 9–14.
- Correa-Aragunde N, Graziano M, Lamattina L. 2004. Nitric oxide plays a central role in determining lateral root development in tomato. *Planta* **218**: 900–905.
- Correa-Aragunde N, Graziano M, Chevalier C, Lamattina L. 2006. Nitric oxide modulates the expression of cell cycle regulatory genes during lateral root formation in tomato. *Journal of Experimental Botany* **57**: 581–588.
- Correa-Aragunde N, Lombardo C, Lamattina L. 2008. Nitric oxide: an active nitrogen molecule that modulates cellulose synthesis in tomato roots. *New Phytologist* **179**: 386–396.
- Cotelle V, Meek SEM, Provan F, Milne FC, Morrice N, MacKintosh C. 2000. 14–3–3s regulate global cleavage of their diverse binding partners in sugar-starved *Arabidopsis* cells. *EMBO Journal* **19**: 2869–2876.
- Cristescu SM, Marchenko D, Mandon J, Hebelstrup K, Griffith GW, Mur LAJ, Harren FJM. 2012. Spectroscopic monitoring of NO traces in plants and human breath: applications and perspectives. *Applied Physics B*: 1–9.

- Cvetkovska M, Vanlerberghe GC. 2012. Alternative oxidase modulates leaf mitochondrial concentrations of superoxide and nitric oxide. *New Phytologist* **195**: 32–39.
- De Angeli A, Thomine S, Frachisse JM, Ephritikhine G, Gambale F, Barbier-Brygoo H. 2007. Anion channels and transporters in plant cell membranes. *FEBS Letters* **581**: 2367–2374.
- de Montaigu A, Sanz-Luque E, Galvan A, Fernandez E. 2010. A soluble guanylate cyclase mediates negative signaling by ammonium on expression of nitrate reductase in *Chlamydomonas*. *The Plant Cell* **22**: 1532–1548.
- del Giudice J, Cam Y, Damiani I, Fung-Chat F, Meilhoc E, Bruand C, Brouquisse R, Puppo A, Boscari A. 2011. Nitric oxide is required for an optimal establishment of the *Medicago truncatula*–*Sinorhizobium meliloti* symbiosis. *New Phytologist* **191**: 405–417.
- Delledonne M, Xia Y, Dixon RA, Lamb C. 1998. Nitric oxide functions as a signal in plant disease resistance. *Nature* **394**: 585–588.
- Dénarié J, Debelle F, Rosenberg C. 1992. Signaling and host range variation in nodulation. *Annual Review of Microbiology* **46**: 497–531.
- Derbyshire ER, Marletta MA. 2012. Structure and regulation of soluble guanylate cyclase. *Annual Review of Biochemistry* **81**: 533–559.
- Deruere J, Jackson K, Garbers C, Soll D, DeLong A. 1999. The RCN1-encoded A subunit of protein phosphatase 2A increases phosphatase activity *in vivo*. *The Plant Journal* **20**: 389–399.
- Desikan R, Last K, Harrett-Williams R, Tagliavia C, Harter K, Hooley R, Hancock JT, Neill SJ. 2006. Ethylene-induced stomatal closure in *Arabidopsis* occurs via AtrbohF-mediated hydrogen peroxide synthesis. *The Plant Journal* **47**: 907–916.
- Dordas C, Hasinoff BB, Igamberdiev AU, Manac'h N, Rivoal J, Hill RD. 2003. Expression of a stress-induced hemoglobin affects NO levels produced by alfalfa root cultures under hypoxic stress. *The Plant Journal* **35**: 763–770.
- Dordas C, Hasinoff BB, Rivoal J, Hill RD. 2004. Class-1 hemoglobins, nitrate and NO levels in anoxic maize cell-suspension cultures. *Planta* **219**: 66–72.
- Dubovskaya LV, Bakakina YS, Kolesneva EV, Sodel DL, McAinsh MR, Hetherington AM, Volotovskii ID. 2011. cGMP-dependent ABA-induced stomatal closure in the ABA-insensitive *Arabidopsis* mutant *abi1-1*. *New Phytologist* **191**: 57–69.
- Durner J, Wendehenne D, Klessig DF. 1998. Defense gene induction in tobacco by nitric oxide, cyclic GMP, and cyclic ADP-ribose. *Proceedings of the National Academy of Sciences of the USA* **95**: 10328–10333.
- Ederli L, Reale L, Madeo L, Ferranti F, Gehring C, Fornaciari M, Romano B, Pasqualini S. 2009. NO release by nitric oxide donors *in vitro* and *in planta*. *Plant Physiology and Biochemistry* **47**: 42–48.
- Feechan A, Kwon E, Yun BW, Wang YQ, Pallas JA, Loake GJ. 2005. A central role for S-nitrosothiols in plant disease resistance. *Proceedings of the National Academy of Sciences of the USA* **102**: 8054–8059.
- Fernandez-Marcos M, Sanz L, Lewis DR, Muday GK, Lorenzo O. 2011. Nitric oxide causes root apical meristem defects and growth inhibition while reducing PIN-FORMED 1 (PIN1)-dependent acropetal auxin transport. *Proceedings of the National Academy of Sciences of the USA* **108**: 18506–18511.
- Flores T, Todd CD, Tovar-Mendez A, Dhanoa PK, Correa-Aragunde N, Hoyos ME, Brownfield DM, Mullen RT, Lamattina L, Polacco JC. 2008. Arginase-negative mutants of *Arabidopsis* exhibit increased nitric oxide signaling in root development. *Plant Physiology* **147**: 1936–1946.
- Foissner I, Wendehenne D, Langebartels C, Durner J. 2000. *In vivo* imaging of an elicitor-induced nitric oxide burst in tobacco. *The Plant Journal* **23**: 817–824.
- Foreman J, Demidchik V, Bothwell JH, Mylona P, Miedema H, Torres MA, Linstead P, Costa S, Brownlee C, Jones JD, Davies JM, Dolan L. 2003. Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* **422**: 442–446.
- Forstermann U, Sessa WC. 2012. Nitric oxide synthases: regulation and function. *European Heart Journal* **33**: 829–837.
- Freschi L, Rodrigues MA, Domingues DS, Purgatto E, Van Sluys MA, Magalhaes JR, Kaiser WM, Mercier H. 2010. Nitric oxide mediates the hormonal control of Crassulacean acid metabolism expression in young pineapple plants. *Plant Physiology* **152**: 1971–1985.
- Fu LJ, Shi K, Gu M, Zhou YH, Dong DK, Liang WS, Song FM, Yu JQ. 2010. Systemic induction and role of mitochondrial alternative oxidase and nitric oxide in a compatible tomato-tobacco mosaic virus interaction. *Molecular Plant-Microbe Interactions* **23**: 39–48.
- Garcia-Brugger A, Lamotte O, Vandelle E, Bourque S, Lecourieux D, Poinssot B, Wendehenne D, Pugin A. 2006. Early signaling events induced by elicitors of plant defenses. *Molecular Plant-Microbe Interactions* **19**: 711–724.
- Garcia-Mata C, Gay R, Sokolovski S, Hills A, Lamattina L, Blatt MR. 2003. Nitric oxide regulates K⁺ and Cl⁻ channels in guard cells through a subset of abscisic acid-evoked signaling pathways. *Proceedings of the National Academy of Sciences of the USA* **100**: 11116–11121.
- Geelen D, Lurin C, Bouchez D, Frachisse JM, Lelievre F, Courtial B, Barbier-Brygoo H, Maurel C. 2000. Disruption of putative anion channel gene *AtCLC-a* in *Arabidopsis* suggests a role in the regulation of nitrate content. *The Plant Journal* **21**: 259–267.
- Guo FQ, Okamoto M, Crawford NM. 2003. Identification of a plant nitric oxide synthase gene involved in hormonal signaling. *Science* **302**: 100–103.
- Gupta KJ, Kaiser WM. 2010. Production and scavenging of nitric oxide by barley root mitochondria. *Plant and Cell Physiology* **51**: 576–584.
- Gupta KJ, Zabalza A, van Dongen JT. 2009. Regulation of respiration when the oxygen availability changes. *Physiologia Plantarum* **137**: 383–391.
- Gupta KJ, Fernie AR, Kaiser WM, van Dongen JT. 2011a. On the origins of nitric oxide. *Trends in Plant Science* **16**: 160–168.
- Gupta KJ, Hebelstrup KH, Mur LA, Igamberdiev AU. 2011b. Plant hemoglobins: important players at the crossroads between oxygen and nitric oxide. *FEBS Letters* **585**: 3843–3849.
- Gupta KJ, Shah JK, Brotman Y, Jahnke K, Willmitzer L, Kaiser WM, Bauwe H, Igamberdiev AU. 2012. Inhibition of aconitase by nitric oxide leads to induction of the alternative oxidase and to a shift of metabolism towards biosynthesis of amino acids. *Journal of Experimental Botany* **63**: 1773–1784.
- Hancock JT, Neill SJ, Wilson ID. 2011. Nitric oxide and ABA in the control of plant function. *Plant Science* **181**: 555–559.
- Hauschildt S, Bassenge E, Bessler W, Busse R, Mulsch A. 1990. L-Arginine-dependent nitric-oxide formation and nitrite

- release in bone marrow-derived macrophages stimulated with bacterial lipopeptide and lipopolysaccharide. *Immunology* **70**: 332–337.
- He Y, Tang RH, Hao Y, Stevens RD, Cook CW, Ahn SM, Jing L, Yang Z, Chen L, Guo F, Fiorani F, Jackson RB, Crawford NM, Pei ZM. 2004. Nitric oxide represses the *Arabidopsis* floral transition. *Science* **305**: 1968–1971.
- Hebelstrup KH, Jensen EO. 2008. Expression of NO scavenging hemoglobin is involved in the timing of bolting in *Arabidopsis thaliana*. *Planta* **227**: 917–927.
- Hebelstrup KH, Hunt P, Dennis E, Jensen SB, Jensen EO. 2006. Hemoglobin is essential for normal growth of *Arabidopsis* organs. *Physiologia Plantarum* **127**: 157–166.
- Hebelstrup KH, van Zanten M, Mandon J, Voesenek LA, Harren FJ, Cristescu SM, Moller IM, Mur LA. 2012. Haemoglobin modulates NO emission and hyponasty under hypoxia-related stress in *Arabidopsis thaliana*. *Journal of Experimental Botany* **3**: 5581–5591.
- Heckmann AB, Hebelstrup KH, Larsen K, Micaelo NM, Jensen EO. 2006. A single hemoglobin gene in *Myrica gale* retains both symbiotic and non-symbiotic specificity. *Plant Molecular Biology* **61**: 769–779.
- Hill RD. 2012. Non-symbiotic haemoglobins—What's happening beyond nitric oxide scavenging? *AoB PLANTS* **2012**: pls004; doi:10.1093/aobpla/pls004.
- Igamberdiev AU, Hill RD. 2009. Plant mitochondrial function during anaerobiosis. *Annals of Botany* **103**: 259–268.
- Jansson EA, Huang L, Malkey R, Govoni M, Nihlen C, Olsson A, Stensdotter M, Petersson J, Holm L, Weitzberg E, Lundberg JO. 2008. A mammalian functional nitrate reductase that regulates nitrite and nitric oxide homeostasis. *Nature Chemical Biology* **4**: 411–417.
- Jasid S, Simontacchi M, Bartoli CG, Puntarulo S. 2006. Chloroplasts as a nitric oxide cellular source. Effect of reactive nitrogen species on chloroplastic lipids and proteins. *Plant Physiology* **142**: 1246–1255.
- Kaiser WM, Huber SC. 2001. Post-translational regulation of nitrate reductase: mechanism, physiological relevance and environmental triggers. *Journal of Experimental Botany* **52**: 1981–1989.
- Kanayama Y, Yamamoto Y. 1991. Formation of nitrosylhemoglobin in nodules of nitrate-treated cowpea and pea-plants. *Plant and Cell Physiology* **32**: 19–24.
- Keefer LK, Nims RW, Davies KM, Wink DA. 1996. 'NONOates' (1-substituted diazen-1-ium-1,2-diolates) as nitric oxide donors: convenient nitric oxide dosage forms. *Methods in Enzymology* **268**: 281–293.
- Klessig DF, Durner J, Noad R, Navarre DA, Wendehenne D, Kumar D, Zhou JM, Shah J, Zhang SQ, Kachroo P, Trifa Y, Pontier D, Lam E, Silva H. 2000. Nitric oxide and salicylic acid signaling in plant defense. *Proceedings of the National Academy of Sciences of the USA* **97**: 8849–8855.
- Kojima H, Sakurai K, Kikuchi K, Kawahara S, Kirino Y, Nagoshi H, Hirata Y, Nagano T. 1998a. Development of a fluorescent indicator for nitric oxide based on the fluorescein chromophore. *Chemical & Pharmaceutical Bulletin* **46**: 373–375.
- Kojima H, Nakatsubo N, Kikuchi K, Kawahara S, Kirino Y, Nagoshi H, Hirata Y, Nagano T. 1998b. Detection and imaging of nitric oxide with novel fluorescent indicators: diaminofluoresceins. *Analytical Chemistry* **70**: 2446–2453.
- Kosterev AA, Tittel FK, Kohler R, Gmachl C, Capasso F, Sivco DL, Cho AY, Wehe S, Allen MG. 2002. Thermoelectrically cooled quantum-cascade-laser-based sensor for the continuous monitoring of ambient atmospheric carbon monoxide. *Applied Optics* **41**: 1169–1173.
- Kramer EM, Bennett MJ. 2006. Auxin transport: a field in flux. *Trends in Plant Science* **11**: 382–386.
- Krause M, Durner J. 2004. Harpin inactivates mitochondria in *Arabidopsis* suspension cells. *Molecular Plant-Microbe Interactions* **17**: 131–139.
- Lamotte O, Gould K, Lecourieux D, Sequeira-Legrand A, Lebrun-Garcia A, Durner J, Pugin A, Wendehenne D. 2004. Analysis of nitric oxide signaling functions in tobacco cells challenged by the elicitor cryptogein. *Plant Physiology* **135**: 516–529.
- Lamotte O, Courtois C, Dobrowolska G, Besson A, Pugin A, Wendehenne D. 2006. Mechanisms of nitric-oxide-induced increase of free cytosolic Ca²⁺ concentration in *Nicotiana glauca* cells. *Free Radical Biology and Medicine* **40**: 1369–1376.
- Li SW, Xue LG. 2010. The interaction between H₂O₂ and NO, Ca²⁺, cGMP, and MAPKs during adventitious rooting in mung bean seedlings. *In Vitro Cellular & Developmental Biology-Plant* **46**: 142–148.
- Lindermayr C, Sell S, Muller B, Leister D, Durner J. 2010. Redox regulation of the NPR1-TGA1 system of *Arabidopsis thaliana* by nitric oxide. *The Plant Cell* **22**: 2894–2907.
- Lombardo MC, Lamattina L. 2012. Nitric oxide is essential for vesicle formation and trafficking in *Arabidopsis* root hair growth. *Journal of Experimental Botany* **63**: 4875–4885.
- Lombardo MC, Graziano M, Polacco JC, Lamattina L. 2006. Nitric oxide functions as a positive regulator of root hair development. *Plant Signaling and Behaviour* **1**: 28–33.
- Lozano-Juste J, Leon J. 2011. Nitric oxide regulates DELLA content and PIF expression to promote photomorphogenesis in *Arabidopsis*. *Plant Physiology* **156**: 1410–1423.
- Ma W, Smigel A, Tsai YC, Braam J, Berkowitz GA. 2008. Innate immunity signaling: cytosolic Ca²⁺ elevation is linked to downstream nitric oxide generation through the action of calmodulin or a calmodulin-like protein. *Plant Physiology* **148**: 818–828.
- Malik SI, Hussain A, Yun BW, Spoel SH, Loake GJ. 2011. GSNOR-mediated de-nitrosylation in the plant defence response. *Plant Science* **181**: 540–544.
- Manjunatha G, Niranjan-Raj S, Prashanth GN, Deepak S, Amruthesh KN, Shetty HS. 2009. Nitric oxide is involved in chitosan-induced systemic resistance in pearl millet against downy mildew disease. *Pest Management Science* **65**: 737–743.
- Manjunatha G, Gupta KJ, Lokesh V, Mur LA, Neelwarne B. 2012. Nitric oxide counters ethylene effects on ripening fruits. *Plant Signaling and Behaviour* **7**: 476–483.
- Mishina TE, Lamb C, Zeier J. 2007. Expression of a nitric oxide degrading enzyme induces a senescence programme in *Arabidopsis*. *Plant, Cell and Environment* **30**: 39–52.
- Modolo LV, Augusto O, Almeida IMG, Magalhaes JR, Salgado I. 2005. Nitrite as the major source of nitric oxide production by *Arabidopsis thaliana* in response to *Pseudomonas syringae*. *FEBS Letters* **579**: 3814–3820.
- Moreau M, Lee GI, Wang Y, Crane BR, Klessig DF. 2008. AtNOS/AtNOA1 is a functional *Arabidopsis thaliana* cGTPase and not a nitric-oxide synthase. *Journal of Biological Chemistry* **283**: 32957–32967.

- Moreland RB, Goldstein I, Kim NN, Traish A. 1999. Sildenafil citrate, a selective phosphodiesterase type 5 inhibitor: research and clinical implications in erectile dysfunction. *Trends in Endocrinology and Metabolism* **10**: 97–104.
- Mou Z, Fan W, Dong X. 2003. Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell* **113**: 935–944.
- Mulaudzi T, Ludidi N, Ruzvidzo O, Morse M, Hendricks N, Iwuoha E, Gehring C. 2011. Identification of a novel *Arabidopsis thaliana* nitric oxide-binding molecule with guanylate cyclase activity *in vitro*. *FEBS Letters* **585**: 2693–2697.
- Mur LAJ, Santosa IE, Laarhoven LJJ, Holton NJ, Harren FJM, Smith AR. 2005. Laser photoacoustic detection allows *in planta* detection of nitric oxide in tobacco following challenge with avirulent and virulent *Pseudomonas syringae* pathovars. *Plant Physiology* **138**: 1247–1258.
- Mur LAJ, Laarhoven LJJ, Harren FJM, Hall MA, Smith AR. 2008. Nitric oxide interacts with salicylate to regulate biphasic ethylene production during the hypersensitive response. *Plant Physiology* **148**: 1537–1546.
- Mur LAJ, Mandon J, Cristescu SM, Harren FJM, Prats E. 2011. Methods of nitric oxide detection in plants: a commentary. *Plant Science* **181**: 509–519.
- Mur LAJ, Sivakumaran A, Mandon J, Cristescu SM, Harren FJ, Hebelstrup KH. 2012. Haemoglobin modulates salicylate and jasmonate/ethylene-mediated resistance mechanisms against pathogens. *Journal of Experimental Botany* **63**: 4375–4387.
- Murad F. 2004. Discovery of some of the biological effects of nitric oxide and its role in cell signaling. *Bioscience Reports* **24**: 452–474.
- Napoli C, Ignarro LJ. 2003. Nitric oxide-releasing drugs. *Annual Reviews in Pharmacology and Toxicology* **43**: 97–123.
- Nathan C. 1995. Natural resistance and nitric oxide. *Cell* **82**: 873–876.
- Neill SJ, Desikan R, Clarke A, Hancock JT. 2002. Nitric oxide is a novel component of abscisic acid signaling in stomatal guard cells. *Plant Physiology* **128**: 13–16.
- Orozco-Cardenas ML, Ryan CA. 2002. Nitric oxide negatively modulates wound signaling in tomato plants. *Plant Physiology* **130**: 487–493.
- Ortega-Galisteo AP, Rodriguez-Serrano M, Pazmino DM, Gupta DK, Sandalio LM, Romero-Puertas MC. 2012. S-nitrosylated proteins in pea (*Pisum sativum* L.) leaf peroxisomes: changes under abiotic stress. *Journal of Experimental Botany* **63**: 2089–2103.
- Pacher P, Beckman JS, Liaudet L. 2007. Nitric oxide and peroxynitrite in health and disease. *Physiological Reviews* **87**: 315–424.
- Pagnussat GC, Simontacchi M, Puntarulo S, Lamattina L. 2002. Nitric oxide is required for root organogenesis. *Plant Physiology* **129**: 954–956.
- Pagnussat GC, Lanteri ML, Lamattina L. 2003. Nitric oxide and cyclic GMP are messengers in the indole acetic acid-induced adventitious rooting process. *Plant Physiology* **132**: 1241–1248.
- Pagnussat GC, Lanteri ML, Lombardo MC, Lamattina L. 2004. Nitric oxide mediates the indole acetic acid induction activation of a mitogen-activated protein kinase cascade involved in adventitious root development. *Plant Physiology* **135**: 279–286.
- Palmieri MC, Sell S, Huang X, Scherf M, Werner T, Durner J, Lindermayr C. 2008. Nitric oxide-responsive genes and promoters in *Arabidopsis thaliana*: a bioinformatics approach. *Journal of Experimental Botany* **59**: 177–186.
- Percheviel L, Balague C, Riou C, Claudel-Renard C, Riviere N, Grezes-Besset B, Roby D. 2010. Nitric oxide participates in the complex interplay of defense-related signaling pathways controlling disease resistance to *Sclerotinia sclerotiorum* in *Arabidopsis thaliana*. *Molecular Plant-Microbe Interactions* **23**: 846–860.
- Planchet E, Jagadis Gupta K, Sonoda M, Kaiser WM. 2005. Nitric oxide emission from tobacco leaves and cell suspensions: rate limiting factors and evidence for the involvement of mitochondrial electron transport. *The Plant Journal* **41**: 732–743.
- Prats E, Mur LAJ, Sanderson R, Carver TLW. 2005. Nitric oxide contributes both to papilla-based resistance and the hypersensitive response in barley attacked by *Blumeria graminis* f. sp. *hordei*. *Molecular Plant Pathology* **6**: 65–78.
- Prats E, Carver TL, Mur LA. 2008. Pathogen-derived nitric oxide influences formation of the appressorium infection structure in the phytopathogenic fungus *Blumeria graminis*. *Research in Microbiology* **159**: 476–480.
- Rasul S, Dubreuil-Maurizi C, Lamotte O, Koen E, Poinssot B, Alcaraz G, Wendehenne D, Jeandroz S. 2012. Nitric oxide production mediates oligogalacturonide-triggered immunity and resistance to *Botrytis cinerea* in *Arabidopsis thaliana*. *Plant, Cell and Environment* **35**: 1483–1499.
- Ribeiro DM, Desikan R, Bright J, Confraria A, Harrison J, Hancock JT, Barros RS, Neill SJ, Wilson ID. 2009. Differential requirement for NO during ABA-induced stomatal closure in turgid and wilted leaves. *Plant, Cell and Environment* **32**: 46–57.
- Rieth T, Sasamoto K. 1998. Detection of nitric oxide and nitrite by using a rhodamine-type fluorescent indicator. *Analytical Communications* **35**: 195–197.
- Robert-Seilaniantz A, Grant M, Jones JDG. 2011. Hormone crosstalk in plant disease and defense: more than just JASMONATE-SALICYLATE antagonism. *Annual Review of Phytopathology* **49**: 317–343.
- Romero-Puertas MC, Laxa M, Matte A, Zaninotto F, Finkemeier I, Jones AME, Perazzoli M, Vandelle E, Dietz KJ, Delledonne M. 2007. S-nitrosylation of peroxiredoxin II E promotes peroxynitrite-mediated tyrosine nitration. *The Plant Cell* **19**: 4120–4130.
- Roncaroli F, Videla M, Slep LD, Olabe JA. 2007. New features in the redox coordination chemistry of metal nitrosyls (M-NO⁺; M-NO; M-NO- (HNO)). *Coordination Chemistry Reviews* **251**: 1903–1930.
- Rothman LS, Jacquemart D, Barbe A, Benner DC, Birk M, Brown LR, Carleer MR, Chackerian C, Chance K, Coudert LH, Dana V, Devi VM, Flaud JM, Gamache RR, Goldman A, Hartmann JM, Jucks KW, Maki AG, Mandin JY, Massie ST, Orphal J, Perrin A, Rinsland CP, Smith MAH, Tennyson J, Tolchenov RN, Toth RA, Vander Auwera J, Varanasi P, Wagner G. 2005. The HITRAN 2004 molecular spectroscopic database. *Journal of Quantitative Spectroscopy & Radiative Transfer* **96**: 139–204.
- Rumer S, Gupta KJ, Kaiser WM. 2009a. Plant cells oxidize hydroxylamines to NO. *Journal of Experimental Botany* **60**: 2065–2072.
- Rumer S, Kapuganti JG, Kaiser WM. 2009b. Oxidation of hydroxylamines to NO by plant cells. *Plant Signaling and Behavior* **4**: 853–855.
- Rumer S, Kruschke M, Fekete A, Mueller MJ, Kaiser WM. 2012. DAF-fluorescence without NO: elicitor treated tobacco cells produce fluorescing DAF-derivatives not related to DAF-2 triazol. *Nitric Oxide* **27**: 123–135.
- Sakamoto A, Tsukamoto S, Yamamoto H, Ueda-Hashimoto M, Takahashi M, Suzuki H, Morikawa H. 2003. Functional

- complementation in yeast reveals a protective role of chloroplast 2-Cys peroxidoredoxin against reactive nitrogen species. *The Plant Journal* **33**: 841–851.
- Sakihama Y, Murakami S, Yamasaki H. 2003.** Involvement of nitric oxide in the mechanism for stomatal opening in *Vicia faba* leaves. *Biologia Plantarum* **46**: 117–119.
- Schaap P. 2005.** Guanylyl cyclases across the tree of life. *Frontiers in Bioscience-Landmark* **10**: 1485–1498.
- Seligman K, Saviani EE, Oliveira HC, Pinto-Maglio CAF, Salgado I. 2008.** Floral transition and nitric oxide emission during flower development in *Arabidopsis thaliana* is affected in nitrate reductase-deficient plants. *Plant and Cell Physiology* **49**: 1112–1121.
- Shi FM, Li YZ. 2008.** *Verticillium dahliae* toxins-induced nitric oxide production in *Arabidopsis* is major dependent on nitrate reductase. *BMB Reports* **41**: 79–85.
- Shiva S. 2010.** Mitochondria as metabolizers and targets of nitrite. *Nitric Oxide-Biology and Chemistry* **22**: 64–74.
- Siegel-Itzkovich J. 1999.** Viagra makes flowers stand up straight. *British Medical Journal* **319**: 274A.
- Smaghe BJ, Hoy JA, Percifield R, Kundu S, Hargrove MS, Sarath G, Hilbert JL, Watts RA, Dennis ES, Peacock WJ, Dewilde S, Moens L, Blouin GC, Olson JS, Appleby CA. 2009.** Correlations between oxygen affinity and sequence classifications of plant hemoglobins. *Biopolymers* **91**: 1083–1096.
- Sokolovski S, Blatt MR. 2004.** Nitric oxide block of outward-rectifying K⁺ channels indicates direct control by protein nitrosylation in guard cells. *Plant Physiology* **136**: 4275–4284.
- Spagnolo V, Kosterev AA, Dong L, Lewicki R, Tittel FK. 2010.** NO trace gas sensor based on quartz-enhanced photoacoustic spectroscopy and external cavity quantum cascade laser. *Applied Physics B—Lasers and Optics* **100**: 125–130.
- Spoel SH, Mou Z, Tada Y, Spivey NW, Genschik P, Dong X. 2009.** Proteasome-mediated turnover of the transcription coactivator NPR1 plays dual roles in regulating plant immunity. *Cell* **137**: 860–872.
- Srivastava N, Gonugunta VK, Puli MR, Raghavendra AS. 2009.** Nitric oxide production occurs downstream of reactive oxygen species in guard cells during stomatal closure induced by chitosan in abaxial epidermis of *Pisum sativum*. *Planta* **229**: 757–765.
- Stöhr C, Strube F, Marx G, Ullrich WR, Rockel P. 2001.** A plasma membrane-bound enzyme of tobacco roots catalyses the formation of nitric oxide from nitrite. *Planta* **212**: 835–841.
- Stoimenova M, Igamberdiev AU, Gupta KJ, Hill RD. 2007.** Nitrite-driven anaerobic ATP synthesis in barley and rice root mitochondria. *Planta* **226**: 465–474.
- Sturms R, DiSpirito AA, Hargrove MS. 2011.** Plant and cyanobacterial hemoglobins reduce nitrite to nitric oxide under anoxic conditions. *Biochemistry* **50**: 3873–3878.
- Suita K, Kiryu T, Sawada M, Mitsui M, Nakagawa M, Kanamaru K, Yamagata H. 2009.** Cyclic GMP acts as a common regulator for the transcriptional activation of the flavonoid biosynthetic pathway in soybean. *Planta* **229**: 403–413.
- Szmidt-Jaworska A, Jaworski K, Tretyn A, Kopcewicz J. 2004.** The involvement of cyclic GMP in the photoperiodic flower induction of *Pharbitis nil*. *Journal of Plant Physiology* **161**: 277–284.
- Szmidt-Jaworska A, Jaworski K, Kopcewicz J. 2008.** Involvement of cyclic GMP in phytochrome-controlled flowering of *Pharbitis nil*. *Journal of Plant Physiology* **165**: 858–867.
- Szmidt-Jaworska A, Jaworski K, Kopcewicz J. 2009.** Cyclic GMP stimulates flower induction of *Pharbitis nil* via its influence on cGMP regulated protein kinase. *Plant Growth Regulation* **57**: 115–126.
- Tada Y. 2009.** Plant immunity requires conformational changes of NPR1 via S-nitrosylation and thioredoxins. *Science* **325**: 1072–1072.
- Tada Y, Spoel SH, Pajerowska-Mukhtar K, Mou Z, Song J, Wang C, Zuo J, Dong X. 2008.** Plant immunity requires conformational changes [corrected] of NPR1 via S-nitrosylation and thioredoxins. *Science* **321**: 952–956.
- Terrile MC, Paris R, Calderon-Villalobos LIA, Iglesias MJ, Lamattina L, Estelle M, Casalogue CA. 2012.** Nitric oxide influences auxin signaling through S-nitrosylation of the *Arabidopsis* TRANSPORT INHIBITOR RESPONSE 1 auxin receptor. *The Plant Journal* **70**: 492–500.
- Tischner R, Galli M, Heimer YM, Bielefeld S, Okamoto M, Mack A, Crawford NM. 2007.** Interference with the citrulline-based nitric oxide synthase assay by argininosuccinate lyase activity in *Arabidopsis* extracts. *FEBS Journal* **274**: 4238–4245.
- Tossi V, Amenta M, Lamattina L, Cassia R. 2011.** Nitric oxide enhances plant ultraviolet-B protection up-regulating gene expression of the phenylpropanoid biosynthetic pathway. *Plant, Cell and Environment* **34**: 909–921.
- Trevisan S, Manoli A, Begheldo M, Nonis A, Enna M, Vaccaro S, Caporale G, Ruperti B, Quaggiotti S. 2011.** Transcriptome analysis reveals coordinated spatiotemporal regulation of hemoglobin and nitrate reductase in response to nitrate in maize roots. *New Phytologist* **192**: 338–352.
- Tun NN, Santa-Catarina C, Begum T, Silveira V, Handro W, Floh EIS, Scherer GFE. 2006.** Polyamines induce rapid biosynthesis of nitric oxide (NO) in *Arabidopsis thaliana* seedlings. *Plant and Cell Physiology* **47**: 346–354.
- Turrion-Gomez JL, Benito EP. 2011.** Flux of nitric oxide between the necrotrophic pathogen *Botrytis cinerea* and the host plant. *Molecular Plant Pathology* **12**: 606–616.
- Vandelle E, Delledonne M. 2011.** Peroxynitrite formation and function in plants. *Plant Science* **181**: 534–539.
- Vetrovsky P, Stoclet JC, Entlicher G. 1996.** Possible mechanism of nitric oxide production from N(G)-hydroxy-L-arginine or hydroxylamine by superoxide ion. *International Journal of Biochemistry & Cell Biology* **28**: 1311–1318.
- Vinardell JM, Fedorova E, Cebolla A, Kevei Z, Horvath G, Kelemen Z, Tarayre S, Roudier F, Mergaert P, Kondorosi A, Kondorosi E. 2003.** Endoreduplication mediated by the anaphase-promoting complex activator CCS52A is required for symbiotic cell differentiation in *Medicago truncatula* nodules. *The Plant Cell* **15**: 2093–2105.
- Víteček J, Reinohl V, Jones RL. 2008.** Measuring NO production by plant tissues and suspension cultured cells. *Molecular Plant* **1**: 270–284.
- Voesenek LACJ, Colmer TD, Pierik R, Millenaar FF, Peeters AJM. 2006.** How plants cope with complete submergence. *New Phytologist* **170**: 213–226.

- Wang PG, Xian M, Tang X, Wu X, Wen Z, Cai T, Janczuk AJ. 2002.** Nitric oxide donors: chemical activities and biological applications. *Chemical Reviews* **102**: 1091–1134.
- Wang XM, Li JS, Liu J, He WL, Bi YR. 2010.** Nitric oxide increases mitochondrial respiration in a cGMP-dependent manner in the callus from *Arabidopsis thaliana*. *Nitric Oxide—Biology and Chemistry* **23**: 242–250.
- Wang YQ, Feechan A, Yun BW, Shafiei R, Hofmann A, Taylor P, Xue P, Yang FQ, Xie ZS, Pallas JA, Chu CC, Loake GJ. 2009.** S-nitrosylation of AtSABP3 antagonizes the expression of plant immunity. *Journal of Biological Chemistry* **284**: 2131–2137.
- Watts RA, Hunt PW, Hvitved AN, Hargrove MS, Peacock WJ, Dennis ES. 2001.** A hemoglobin from plants homologous to truncated hemoglobins of microorganisms. *Proceedings of the National Academy of Sciences of the USA* **98**: 10119–10124.
- Weiner H, Kaiser WM. 1999.** 14-3-3 proteins control proteolysis of nitrate reductase in spinach leaves. *FEBS Letters* **455**: 75–78.
- Weiner H, Kaiser WM. 2000.** Binding to 14-3-3 proteins is not sufficient to inhibit nitrate reductase in spinach leaves. *FEBS Letters* **480**: 217–220.
- Wendehenne D, Hancock JT. 2011.** New frontiers in nitric oxide biology in plant. *Plant Science* **181**: 507–508.
- Wendehenne D, Lamotte O, Frachisse JM, Barbier-Brygoo H, Pugin A. 2002.** Nitrate efflux is an essential component of the cryptogein signaling pathway leading to defense responses and hypersensitive cell death in tobacco. *The Plant Cell* **14**: 1937–1951.
- Wilson ID, Ribeiro DM, Bright J, Confraria A, Harrison J, Barros RS, Desikan R, Neill SJ, Hancock JT. 2009.** Role of nitric oxide in regulating stomatal apertures. *Plant Signaling and Behavior* **4**: 467–469.
- Wojtaszek P. 1997.** Oxidative burst: an early plant response to pathogen infection. *Biochemical Journal* **322**: 681–692.
- Wu CM, Chen YH, Dayananda K, Shiue TW, Hung CH, Liaw WF, Chen PY, Wang YM. 2011.** Sensitivity evaluation of rhodamine B hydrazide towards nitric oxide and its application for macrophage cells imaging. *Analytica Chimica Acta* **708**: 141–148.
- Xuan Y, Zhou S, Wang L, Cheng YD, Zhao LQ. 2010.** Nitric oxide functions as a signal and acts upstream of AtCaM3 in thermotolerance in *Arabidopsis* seedlings. *Plant Physiology* **153**: 1895–1906.
- Yang XF, Guo XQ, Zhao YB. 2002.** Development of a novel rhodamine-type fluorescent probe to determine peroxynitrite. *Talanta* **57**: 883–890.
- Yemets AI, Krasylenko YA, Lytvyn DI, Sheremet YA, Blume YB. 2011.** Nitric oxide signalling via cytoskeleton in plants. *Plant Science* **181**: 545–554.
- Yun BW, Feechan A, Yin M, Saidi NB, Le Bihan T, Yu M, Moore JW, Kang JG, Kwon E, Spoel SH, Pallas JA, Loake GJ. 2011.** S-nitrosylation of NADPH oxidase regulates cell death in plant immunity. *Nature* **478**: 264–268.
- Zeidler D, Zahringer U, Gerber I, Dubery I, Hartung T, Bors W, Hutzler P, Durner J. 2004.** Innate immunity in *Arabidopsis thaliana*: lipopolysaccharides activate nitric oxide synthase (NOS) and induce defense genes. *Proceedings of the National Academy of Sciences of the USA* **101**: 15811–15816.
- Zeier J, Delledonne M, Mishina T, Severi E, Sonoda M, Lamb C. 2004.** Genetic elucidation of nitric oxide signaling in incompatible plant-pathogen interactions. *Plant Physiology* **136**: 2875–2886.
- Zemojtel T, Frohlich A, Palmieri MC, Kolanczyk M, Mikula I, Wyrwicz LS, Wanker EE, Mundlos S, Vingron M, Martasek P, Durner J. 2006.** Plant nitric oxide synthase: a never-ending story? *Trends in Plant Science* **11**: 524–525.
- Zhao MG, Zhao X, Wu YX, Zhang LX. 2007.** Enhanced sensitivity to oxidative stress in an *Arabidopsis* nitric oxide synthase mutant. *Journal of Plant Physiology* **164**: 737–745.