

tic dynamics discipline, passing by the most successful predictions and models, finishing with an integral vision of what is known from the Milky Way structure from its dynamics and the prospects with the new large scale surveys to understand it in the next decades.

¹ Instituto de Astronomía, Universidad Nacional Autónoma de México (UNAM).

USING MASSCLEAN TO DESCRIBE STELLAR CLUSTERS FOUND IN THE VISTA VARIABLES IN THE VIA LACTEA (VVV) SURVEY

B. Popescu¹, M. M. Hanson¹, J. Borissova², R. Kurtev², V. D. Ivanov³, M. Catelan⁴, S. S. Larsen⁵, D. Minniti⁴, and P. Lucas⁶

The important parameters: age, mass and distance of resolved or partially resolved stellar clusters are better accurately determined by using color-magnitude diagrams (CMD). However, when the main sequence turnoff is not available or clearly identifiable, large errors in all parameters result when using simple isochrone fitting, particularly when observations are limited to near-infrared bands. We used the MASSCLEAN package to perform 5 million Monte Carlo simulations of stochastically sampled stellar clusters in order to generate CMD templates for a variety of cluster masses and ages and which mimic the observational photometric errors. This results in the creation of tens of thousands of n-dimensional stellar density maps (templates) in numerous color planes as a function of age and mass. We use these MASSCLEAN CMD templates to refine and sharpen traditional isochrone fitting to analyze the newly discovered stellar clusters/cluster candidates from the Vista Variables in the Via Lactea (VVV) Survey. Our MASSCLEAN templates are also being used to design and optimize search algorithms for stellar clusters in broad-band surveys.

¹ Department of Physics, University of Cincinnati, PO Box 210011, Cincinnati, OH 45221-0011, USA (bogdan.popescu@uc.edu, margaret.hanson@uc.edu).

² Departamento de Física y Astronomía, Facultad de Ciencias, Universidad de Valparaíso, Ave. Gran Bretaña 1111, Playa Ancha, Valparaíso, Chile.

³ European Southern Observatory, Ave. Alonso de Córdoba 3107, Casilla 19001, Santiago, Chile.

⁴ Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile.

⁵ Department of Astrophysics/ IMAPP, Radboud University Nijmegen, PO Box 9010, 6500 GL Nijmegen, The Netherlands.

⁶ Centre for Astrophysics Research, Science and Technology Research Inst., University of Hertfordshire, Hatfield, AL10 9AB, UK.

MASS EXTINCTIONS, GALACTIC ORBITS IN THE SOLAR NEIGHBORHOOD AND THE SUN: A CONNECTION?

G. F. Porto de Mello¹, W. S. Dias², J. Lépine³, D. Lorenzo-Oliveira¹, and R. K. Siqueira⁴

The orbits of the stars in the disk of the Galaxy, and their passages through the Galactic spiral arms, are a rarely mentioned factor of biosphere stability which might be important for long-term planetary climate evolution, with a possible bearing on mass extinctions. The Sun lies very near the co-rotation radius, where stars revolve around the Galaxy in the same period as the density wave perturbations of the spiral arms. Conventional wisdom generally considers that this status makes for few passages through the spiral arms. Controversy still surrounds whether time spent inside or around spiral arms is dangerous to biospheres and conducive to mass extinctions. Possible threats include giant molecular clouds disturbing the Oort comet cloud and provoking heavy bombardment; a higher exposure to cosmic rays near star forming regions triggering increased cloudiness in Earth's atmosphere and ice ages; and the destruction of Earth's ozone layer posed by supernova explosions. We present detailed calculations of the history of spiral arm passages for all 212 solar-type stars nearer than 20 parsecs, including the total time spent inside the spiral arms in the last 500 Myr, when the spiral arm position can be traced with good accuracy. We found that there is a large diversity of stellar orbits in the solar neighborhood, and the time fraction spent inside spiral arms can vary from a few percent to nearly half the time. The Sun, despite its proximity to the galactic co-rotation radius, has exceptionally low eccentricity and a low vertical velocity component, and therefore spends 30% of its lifetime crossing the spiral arms, more than most nearby stars. We discuss the possible implications of this fact to the long-term habitability of the Earth, and possible correlations of the Sun's passage through the spiral arms with the five great mass extinctions of the Earth's biosphere from the Late Ordovician to the Cretaceous-Tertiary.

¹ Observatório do Valongo, Universidade Federal do Rio de Janeiro.