

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/137965>

Please be advised that this information was generated on 2019-04-25 and may be subject to change.

Original Paper

Evaluation of an eHealth Intervention in Chronic Care for Frail Older People: Why Adherence is the First Target

Peter Makai¹, MSc, PhD; Marieke Perry¹, MD, PhD; Sarah HM Robben², MD, PhD; Henk J Schers³, MD, PhD; Maud M Heinen⁴, RN, PhD; Marcel GM Olde Rikkert¹, MD, PhD; René F Melis¹, MD, PhD

¹Radboud University Medical Center, Department of Geriatrics, Nijmegen, Netherlands

²Jeroen Bosch Hospital, Department of Internal Medicine, Hertogenbosch, Netherlands

³Radboud University Medical Center, Department of Primary and Community Care, Nijmegen, Netherlands

⁴Radboud University Medical Center, Scientific Institute for Quality in Healthcare, Nijmegen, Netherlands

Corresponding Author:

Peter Makai, MSc, PhD

Radboud University Medical Center

Department of Geriatrics

Renier Postlaan 4

Nijmegen, 6500 HB

Netherlands

Phone: 31 24 361 98 07

Fax: 31 24 361 74 08

Email: Peter.Makai@radboudumc.nl

Abstract

Background: Older people suffering from frailty often receive fragmented chronic care from multiple professionals. According to the literature, there is an urgent need for coordination of care.

Objective: The objective of this study was to investigate the effectiveness of an online health community (OHC) intervention for older people with frailty aimed at facilitating multidisciplinary communication.

Methods: The design was a controlled before-after study with 12 months follow-up in 11 family practices in the eastern part of the Netherlands. Participants consisted of frail older people living in the community requiring multidisciplinary (long-term) care. The intervention used was the health and welfare portal (ZWIP): an OHC for frail elderly patients, their informal caregivers and professionals. ZWIP contains a secure messaging system supplemented by a shared electronic health record. Primary outcomes were scores on the Instrumental Activities of Daily Living scale (IADL), mental health, and social activity limitations.

Results: There were 290 patients in the intervention group and 392 in the control group. Of these, 76/290 (26.2%) in the intervention group actively used ZWIP. After 12 months follow-up, we observed no significant improvement on primary patient outcomes. ADL improved in the intervention group with a standardized score of 0.21 ($P=.27$); IADL improved with 0.50 points, $P=.64$.

Conclusions: Only a small percentage of frail elderly people in the study intensively used ZWIP, our newly developed and innovative eHealth tool. The use of this OHC did not significantly improve patient outcomes. This was most likely due to the limited use of the OHC, and a relatively short follow-up time. Increasing actual use of eHealth intervention seems a precondition for large-scale evaluation, and earlier adoption before frailty develops may improve later use and effectiveness of ZWIP.

(*J Med Internet Res* 2014;16(6):e156) doi:[10.2196/jmir.3057](https://doi.org/10.2196/jmir.3057)

KEYWORDS

eHealth; frail elderly; care coordination; chronic care

Introduction

Chronic care for frail older people is fragmented, with involvement from a large and constantly changing group of professionals who are frequently unaware that they provide care

to the same patient [1]. Such professionals include home care professionals, general practitioners (GPs), clinicians, physiotherapists, and case managers dedicated to long-term care of the patients in the community. Frail elderly often suffer from comorbidities, which results in care by multiple health care


professionals [2]. Therefore lack of communication between professionals leads to a fragmented and ineffective health care delivery for frail elderly [3]. To reduce fragmentation and promote continuity of care, better coordination and communication between professionals and with patients is necessary. Online health communities (OHCs) have been recognized as an effective mechanism for supporting continuous care for frail older people [4], allowing better coordination and more efficient communication with patients and among professionals. OHCs consist of Internet-based platforms that unite groups of individuals with a shared goal or similar interest, including both professionals and patients [5]. The main strength of OHCs is that they allow communication between people who would not have met each other otherwise [5]. Thus, OHCs are particularly suited for improving the coordination of care for frail elderly who have multiple professional caregivers. For this purpose, we developed and evaluated the Health and Welfare Information Portal (Zorg en Welzijns Informatie Portaal, ZWIP, in Dutch) [1,6] on its effectiveness.

Methods

Intervention

ZWIP is an OHC [5] that aims to facilitate communication for patients, their informal caregivers, and their professionals. ZWIP contains a secure messaging system supplemented by a shared electronic health record. All messages shared in a patient's ZWIP are visible for all users, thus stimulating involvement of and discussion between patients and a team of health professionals. All informal caregivers and health care professionals have access to the electronic health record. To ensure confidentiality, professionals can participate in a patient's personal care network in ZWIP only at the invitation of the patient. Patients who were not able to manage their own ZWIP account could appoint an informal caregiver to act on their behalf. Figure 1 demonstrates the conceptual model underlying ZWIP, and the video in Multimedia Appendix 1 illustrates the use of ZWIP by a patient and an informal caregiver.

Figure 1. A conceptual model of the ZWIP.



Development and Implementation

The development of ZWIP and the process of implementation have been described elsewhere [1,6]. In brief, ZWIP was developed using intervention mapping [7], a stepwise approach for the systematic development of interventions informed by both evidence and theory [1]. Main steps of intervention mapping for ZWIP were (1) needs assessment in frail elderly, (2) developing program objectives, (3) selecting theory informed intervention methods and strategies, (4) creating and pilot testing

program components, (5) planning program adoption and implementation, and (6) planning for evaluation [1]. Theoretically, ZWIP was based on social cognitive theory [8], with special attention paid to improving self-efficacy, the belief people have in their ability to complete tasks and achieve specific goals [9]. Following the steps of intervention mapping and as suggested in the guideline on development and evaluation of complex interventions [8], the ZWIP was piloted by 2 frail elderly and 7 professionals, including one GP. Furthermore,

newly developed elements of ZWIP were regularly piloted by similar user panels.

To enhance implementation of the ZWIP, we used several strategies for professionals such as a continuing medical education (CME) accredited education program based on active learning theory [10], direct experience, and modeling [1]. Additionally, drawing from organization theory, we installed a telephonic helpdesk and provided e-coaching and financial compensation to support the uptake of ZWIP by professionals [7,11]. To facilitate the use of ZWIP among elderly patients, a number of approaches were used: flyers were distributed in the primary care centers, a hard-copy version of ZWIP was provided in order for patients to familiarize themselves with ZWIP, coaching on the use of ZWIP was made available, involvement of informal caregivers was encouraged, and the GPs actively advocated the use of ZWIP, thus drawing on modeling, guided practice, and tailoring support for use of the intervention [1,6]. During the implementation phase, we designated one key person in each family practice who coordinated implementation activities and helped colleagues with questions [6].

Inclusion and Design

Between July 2010 and July 2011, frail older patients were included in an observational, controlled before-after study with 12 months follow-up to investigate ZWIP's effects on patient outcomes. Participating primary care centers were recruited from the university primary care network around the city of Nijmegen, the Netherlands. These centers identified their frail older people using the EASYcare Two-step Older person Screening (TOS) instrument [12]. Therefore, both intervention and control practices had to implement an identification scheme and redesign care for their frail elderly. Intervention centers were selected based on willingness to participate in ZWIP, whereas control practices were selected from a separate project: the EASYcare-TOS validation study [13] {van Kempen, 2013 #7718}. Frail status as determined by the EASYcare-TOS was the only inclusion criterion for included patients. Patients in the intervention group patients needed to agree to the creation of a ZWIP account. No exclusion criteria were specified.

All measurements were performed by trained nurses in the patients' homes, using a face-to-face questionnaire at baseline and at follow-up. The study was exempt from ethics review by the local ethics committee because of its observational nature and nonintrusive data collection. Nevertheless, oral informed consent was obtained to analyze the data during data collection.

Outcomes

Primary outcomes were Activities of Daily Living (ADL) as measured by the Katz index [14], combined ADL and Instrumental Activities of Daily Living (IADL) as measured by the Katz-15, a combined measure of the ADL and Lawton-index [14,15], SF-36 mental health and social activity limitation dimensions [16]. The Katz index consists of yes or no responses on ADL items such as bathing or dressing. ADL scores range from 0-6 with higher scores indicating higher dependency. The Katz-15 consists of yes or no responses on ADL and additional IADL items such as using the telephone and managing money [14]. The Katz-15 scores range from 0-15 with higher scores

indicating more limitations. Both scales are established in the literature and have adequate reliability and validity [17]. The SF-36 mental health dimension, consisting of the following subscales: happy, calm, blue, down, nervous scoring from 0-5 with higher numbers indicating a higher score. The scores were summed into a summary score ranging from 0-100, with 100 indicating full mental health, and 0 low mental health [16]. To assess differences in social activity limitations, the social activity limitation item from the SF-36 was used [16]. This item measures the frequency in which respondents experienced social activity limitations due to health. The item used in this current study is scored from 0 (none of the time) to 5 (all of the time). The various SF-36 subscales have excellent reliability and validity [17]. Secondary outcomes were several self-developed scales of patient satisfaction and GPs' subjective experience with care coordination. Patient satisfaction items were scored on a 5-point Likert scale ranging from 1 (way too little or way too much) to 5 (optimal), similar to this article [18]. GP experience with coordination of care was scored between 1 (uncoordinated) to 10 (optimal coordination). Important covariates were measured including a frailty index based on the accumulation of deficits concept [19,20]. The frailty index is the number of deficits present divided by a total possible number of deficits [2]. As such, the frailty index can account for all kinds of health-related imbalances between the intervention and control group and provides an accurate measure of individuals' frailty.

Analysis

For comparing baseline characteristics, chi-square tests were used to compare nominal variables, and *t* tests were used for normally distributed continuous variables. Effects were determined using linear mixed models within a highly efficient analysis of covariance (ANCOVA) framework [21] to allow for clustering within a primary care center. Adjustments were made for frailty status and centered baseline status of the outcome variable and additional covariates with baseline imbalance. All analyses were performed with SAS 9.2.

Results

Overall, 290/622 (46.6%) of all frail persons identified within 11 practices participated in the intervention group. From 6 practices 392 frail older people participated in the control group. At 12-month follow-up, in the intervention group 179/290 (61.7% of original) patients provided data at follow-up, versus 270/392 (68.8% of original) patients in the control group. At baseline, participants in the intervention group were more likely to have completed primary education only, have more informal caregivers, and have higher complexity of care compared to the participants in the control group. Further, participants in the intervention groups also had a higher average frailty index score, and GPs had lower experience with coordination of care (Table 1).

One quarter 45/117 (25%) of all patients in the intervention group used ZWIP at least once a month during a period of 12 months. Controlling for frailty and other unbalanced baseline characteristics, we found no significant differences in primary

patient outcomes (Table 2). Change in coordination of care as reported by GPs improved in the control group.

Table 1. Demographic and care-related characteristics in the intervention and control group.

Demographic and care-related characteristics	Category	Total	Intervention ZWIP, n=179, n (%)	Control regular care, n=270, n (%)	P value ^a
Sex, n (%)	Female	284	117 (65.4)	167 (61.8)	.45
Age, mean (SD)		449	81.69 (5.38)	81.32 (5.72)	.49
Education, n (%)					
	Primary or less than primary education	67	30 (19.1)	37 (14.8)	<.01
	Secondary education	339	123 (76.5)	216 (82.1)	
	University/tertiary education	15	7 (4.4)	8 (3.1)	
Marital status, n (%)					
	Married	196	80 (45.2)	116 (44.3)	.43
	Divorced	28	7 (3.4)	21 (7.4)	
	Widow/widower/partner deceased	199	79 (44.3)	120 (44.1)	
	Unmarried	24	12 (6.7)	12 (4.2)	
Informal caregiver, n (%)	Available	253	147 (82.6)	106 (39.7)	<.01 ^a
Living independently, n (%)	Yes	230	86 (50.3)	144 (53.0)	.46
Complexity of care, n (%)					
	One professional	63	12 (6.8)	51 (18.7)	<.01 ^a
	2 or 3 professionals	311	128 (71.9)	183 (67.4)	
	>3 professionals	74	38 (21.3)	36 (13.9)	
Frailty index, mean (SD)		447	0.29 (0.07)	0.27 (0.07)	.02 ^a
Multimorbidity, mean (SD)		447	1.70 (1.22)	1.73 (1.35)	.78
GP experience with coordination of care around the patient, mean (SD)		449	5.92 (2.36)	6.76 (3.45)	<.01 ^a

^a2-sided chi-square for discrete and *t* tests for continuous variables.

Table 2. Change in outcomes by 12 months application of the ZWIP Web-based tool for patient-professional and interprofessional communication.

Variable	Total	Intervention ZWIP baseline	Intervention ZWIP follow-up	Control baseline	Control follow-up	Standardized difference between study groups	<i>P</i> value (mixed model)
	N	mean (95% CI)	mean (95% CI)	mean (95% CI)	mean (95% CI)	mean, (95% CI)	
Katz ADL	442	1.09 (0.91-1.27)	1.35 (1.14-1.56)	0.85 (0.72-0.98)	1.02 (0.86-1.18)	0.21 (-0.17-0.59)	.27
Katz-15	442	5.08 (4.73-5.44)	5.76 (5.32-6.21)	4.24 (3.92-4.57)	4.93 (4.58-5.28)	0.50 (-1.59-2.60)	.64
SF-36 mental health	440	76.30 (74.32-78.28)	74.59 (72.83-76.36)	76.27 (74.77-77.77)	79.06 (77.36-80.75)	-8.34 (-17.02-0.34)	.06
SF-36 social	436	1.44 (1.23-1.64)	1.20 (1.01-1.39)	0.87 (0.73-1.01)	0.93 (0.79-1.08)	0.84 (-0.78-2.45)	.31
Patient experience with coordination of care	303	4.66 (4.50-4.83)	4.78 (4.65-4.91)	4.59 (4.44-4.75)	4.77 (4.67-4.88)	-0.25 (-0.99-0.49)	.58
Patient experience with co-decision making	399	3.51 (3.41-3.61)	4.86 (4.77-4.95)	3.63 (3.54-3.72)	4.73 (4.62-4.83)	0.16 (-0.40-0.71)	.64
Patient preferences for influence	414	3.59 (3.42-3.75)	3.59 (3.42-3.76)	3.22 (3.08-3.36)	3.33 (3.20-3.45)	0.36 (-1.28-1.99)	.08
Patient knowledge of providers (health and social)	432	3.51 (3.41-3.61)	3.57 (3.46-3.68)	3.63 (3.54-3.72)	3.68 (3.59-3.77)	-0.68 (-1.44-0.08)	.08
Patient experience with self-management	386	4.85 (4.76-4.93)	4.87 (4.80-4.95)	4.78 (4.67-4.88)	4.72 (4.61-4.83)	0.38 (-0.29-1.06)	.26
GP experience with coordination of care around the patient	432	5.92 (5.58-6.27)	7.11 (6.81-7.42)	6.76 (6.34-7.17)	8.18 (7.94-8.42)	-5.28 (-10.64-0.07)	.04

Discussion

Summary of Results

There were 290 patients who participated in the intervention group and 392 in the control group. In the intervention group 76/290 (26.2%) of the patients actively used ZWIP. After a follow-up of 12 months, we observed no significant improvement on primary patient outcomes, ADL, IADL, and mental health.

Strengths and Limitations

The online ZWIP platform was specifically developed for reducing fragmentation of care delivery in older people. Almost half of a frail elderly population without exclusion criteria could be included in the intervention group for using the online ZWIP tool [6]. This is modestly higher than what can be expected in the Dutch context, where 39% persons older than 75 years report having Internet access [22]. This study has two important limitations that can impact results. First, due to the observational nature of the study, comparability between the intervention and the control groups was limited. Despite adjusting for a range of covariates, there may be residual confounding.

Observational, controlled before-after designs are common for complex interventions, where randomized controlled trials (RCTs) are often not appropriate or feasible for evaluation [23].

In the case of ZWIP, contamination between patients would have made individual-level randomization inappropriate. Cluster randomization was not feasible because the level of commitment required from a number of local stakeholders could not be sustained in the control group.

A second limitation was the fact that actual usage of ZWIP was low, even though the implementation of ZWIP was prepared systematically during the development of ZWIP, as this is a structural part of intervention mapping [1,6,7,24,25]. Additionally, implementation strategies were added or adapted when needed during the actual implementation phase. A wide range of implementation strategies were used to encourage uptake; for example, a training program was developed for professionals and an active recruitment phase led to a high participation of older persons. Therefore, low levels of use were attained not because of the lack of, but despite using state of the art implementation techniques. Failure to integrate eHealth interventions in health care is widespread [26], and therefore the low levels of use of these frail older subjects is not surprising. This is especially true for sustained usage of an eHealth intervention [27]. As in other studies [26], further efforts should be focused on improving usability of the intervention, in terms of compatibility for frail older people in chronic disease trajectories [28].

Future Directions

In addition to further refinement, it is essential to identify those who benefit most from ZWIP and eHealth applications in general. The use of eHealth applications in frail populations could be increased by first identifying frail people with a high likelihood of early adoption of the eHealth intervention, such as people with high computer literacy. Which frail elderly are likely adopters requires further research [26]. Therefore, we plan to perform a quantitative and qualitative evaluation of ZWIP usage as well, going beyond the scope of this paper. We must recognize that in the early stages of evaluation, we take more of an efficacy approach to the evaluation, rather than a pragmatic trial approach. Although the efficacy approach limits generalizability, it allows a thorough investigation of the intervention's working mechanisms under more controlled, laboratory-like conditions. Such work may also reveal ideal levels of use of ZWIP, as it is possible that communication was already adequate in the case of some patients, making ZWIP usage superfluous. Using both quantitative and qualitative methods in this development phase may elicit remaining barriers

and reveal more effective implementation strategies. Only after adapting to this group and proven efficacy is large-scale implementation warranted. Successful wide-scale implementation is a precondition for investigating the effectiveness of eHealth interventions. Otherwise finding no differences between treatment arms cannot be interpreted as a lack of effectiveness. These arguments show that, sufficient time and resources are required to develop, test, and retest new eHealth interventions before finally evaluating their effectiveness in pragmatic trials [29,30].

Conclusions

Overall, the study confirmed that introducing eHealth interventions in the elderly is a difficult task. Despite using a theory-driven intervention design and state of the art implementation techniques, usage remained low and effectiveness was not observed. Performing a thorough proof of principle study in early adopters may be crucial to improving the use of eHealth interventions in the elderly before evaluating effects on a larger scale.

Acknowledgments

This study was supported by the Netherlands Organization for Health Service and Development (ZonMw) project #311050201. The funder had no role in conducting the study, analysis of the data, or publication of the manuscript.

We would like to thank Wilma Derksen-Driessen, Emile ter Horst Mark Kuster, Charlotte Neger, Jean Nielen, Joep Scheltinga, and Leontien van Nieuwenhuijzen for their assistance in the development and implementation of the program. We would also like to thank Jennifer Lutomski for English editing.

Authors' Contributions

PM drafted the manuscript, conducted the analysis, and interpreted the data. MP, SHMR, and HJS contributed to the design, data collection, and revised the manuscript for important content. MMH and MGMOR contributed to the design and revised the manuscript for important intellectual content. RJFM contributed to the design, interpretation of the data, and revised the manuscript for important intellectual content. All authors approved the final manuscript.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Short video of ZWIP in use.

[[WMV File \(Windows Media Video\), 1MB - jmir_v16i6e156_app1.wmv](#)]

References

1. Robben SH, Huisjes M, van Achterberg T, Zuidema SU, Olde Rikkert MG, Schers HJ, ZOWEL NN Study Group. Filling the Gaps in a Fragmented Health Care System: Development of the Health and Welfare Information Portal (ZWIP). *JMIR Res Protoc* 2012;1(2):e10 [[FREE Full text](#)] [doi: [10.2196/resprot.1945](#)] [Medline: [23611877](#)]
2. Mitnitski AB, Mogilner AJ, Rockwood K. Accumulation of deficits as a proxy measure of aging. *ScientificWorldJournal* 2001 Aug 8;1:323-336 [[FREE Full text](#)] [doi: [10.1100/tsw.2001.58](#)] [Medline: [12806071](#)]
3. Stange KC. The problem of fragmentation and the need for integrative solutions. *Ann Fam Med* 2009;7(2):100-103 [[FREE Full text](#)] [doi: [10.1370/afm.971](#)] [Medline: [19273863](#)]
4. Weiner M, Callahan CM, Tierney WM, Overhage JM, Mamlin B, Dexter PR, et al. Using information technology to improve the health care of older adults. *Ann Intern Med* 2003 Sep 2;139(5 Pt 2):430-436. [Medline: [12965971](#)]
5. van der Eijk M, Faber MJ, Aarts JW, Kremer JA, Munneke M, Bloem BR. Using online health communities to deliver patient-centered care to people with chronic conditions. *J Med Internet Res* 2013;15(6):e115 [[FREE Full text](#)] [doi: [10.2196/jmir.2476](#)] [Medline: [23803284](#)]

6. Robben SH, Perry M, Huisjes M, van Nieuwenhuijzen L, Schers HJ, van Weel C, et al. Implementation of an innovative web-based conference table for community-dwelling frail older people, their informal caregivers and professionals: a process evaluation. *BMC Health Serv Res* 2012;12:251 [FREE Full text] [doi: [10.1186/1472-6963-12-251](https://doi.org/10.1186/1472-6963-12-251)] [Medline: [22894654](https://pubmed.ncbi.nlm.nih.gov/22894654/)]
7. Bartholomew LK, Parcel GS, Kok G, Gottlieb NH, Fernandez ME. Planning health promotion programs: An intervention mapping approach. New York: John Wiley & Sons; Jan 25, 2011:-768.
8. Bandura A. Health promotion by social cognitive means. *Health Educ Behav* 2004 Apr;31(2):143-164. [doi: [10.1177/1090198104263660](https://doi.org/10.1177/1090198104263660)] [Medline: [15090118](https://pubmed.ncbi.nlm.nih.gov/15090118/)]
9. Bandura A. Self-efficacy. In: Ramachandran VS, editor. *Encyclopedia of Human Behavior, Second Edition*. CA: Academic Press; 2012.
10. Prince M. Does Active Learning Work? A Review of the Research. *Journal of Engineering Education* 2004 Jul;93(3):223-231. [doi: [10.1002/j.2168-9830.2004.tb00809.x](https://doi.org/10.1002/j.2168-9830.2004.tb00809.x)]
11. Butterfoss F, Kegler M, Francisco V. Mobilizing organizations for health promotion: theories of organizational change. In: Glanz K, editor. *Health behavior and health education: theory, research and practice*. San Francisco, CA: Jossey-Bass; 2008:335-361.
12. van Kempen JA, Schers HJ, Jacobs A, Zuidema SU, Ruikes F, Robben SH, et al. Development of an instrument for the identification of frail older people as a target population for integrated care. *Br J Gen Pract* 2013 Mar;63(608):e225-e231 [FREE Full text] [doi: [10.3399/bjgp13X664289](https://doi.org/10.3399/bjgp13X664289)] [Medline: [23561790](https://pubmed.ncbi.nlm.nih.gov/23561790/)]
13. van Kempen JA, Schers HJ, Jacobs A, Zuidema SU, Ruikes F, Robben SH, et al. Development of an instrument for the identification of frail older people as a target population for integrated care. *Br J Gen Pract* 2013 Mar;63(608):e225-e231 [FREE Full text] [doi: [10.3399/bjgp13X664289](https://doi.org/10.3399/bjgp13X664289)] [Medline: [23561790](https://pubmed.ncbi.nlm.nih.gov/23561790/)]
14. Katz S. Assessing self-maintenance: activities of daily living, mobility, and instrumental activities of daily living. *J Am Geriatr Soc* 1983 Dec;31(12):721-727. [Medline: [6418786](https://pubmed.ncbi.nlm.nih.gov/6418786/)]
15. Weinberger M, Samsa GP, Schmader K, Greenberg SM, Carr DB, Wildman DS. Comparing proxy and patients' perceptions of patients' functional status: results from an outpatient geriatric clinic. *J Am Geriatr Soc* 1992 Jun;40(6):585-588. [Medline: [1587975](https://pubmed.ncbi.nlm.nih.gov/1587975/)]
16. Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992 Jun;30(6):473-483. [Medline: [1593914](https://pubmed.ncbi.nlm.nih.gov/1593914/)]
17. McDowell I, Newell C. *Measuring health: a guide to rating scales and questionnaires*. New York: Oxford University Press; 2006:-748.
18. Goldberg SE, Bradshaw LE, Kearney FC, Russell C, Whittamore KH, Foster PE, Medical Crises in Older People Study Group. Care in specialist medical and mental health unit compared with standard care for older people with cognitive impairment admitted to general hospital: randomised controlled trial (NIHR TEAM trial). *BMJ* 2013;347:f4132 [FREE Full text] [Medline: [23819964](https://pubmed.ncbi.nlm.nih.gov/23819964/)]
19. Lutomski JE, Baars MA, van Kempen JA, Buurman BM, den Elzen WP, Jansen AP, et al. Validation of a frailty index from the older persons and informal caregivers survey minimum data set. *J Am Geriatr Soc* 2013 Sep;61(9):1625-1627. [doi: [10.1111/jgs.12430](https://doi.org/10.1111/jgs.12430)] [Medline: [24028364](https://pubmed.ncbi.nlm.nih.gov/24028364/)]
20. Rockwood K, Mitnitski A. Frailty in relation to the accumulation of deficits. *J Gerontol A Biol Sci Med Sci* 2007 Jul;62(7):722-727. [Medline: [17634318](https://pubmed.ncbi.nlm.nih.gov/17634318/)]
21. Teerenstra S, Eldridge S, Graff M, de Hoop E, Borm GF. A simple sample size formula for analysis of covariance in cluster randomized trials. *Stat Med* 2012 Sep 10;31(20):2169-2178. [doi: [10.1002/sim.5352](https://doi.org/10.1002/sim.5352)] [Medline: [22495809](https://pubmed.ncbi.nlm.nih.gov/22495809/)]
22. Statistics Netherlands. Web Magazine. 2013. Rapid increase internet usage among older people URL: <http://www.cbs.nl/en-GB/menu/themas/vrije-tijd-cultuur/publicaties/artikelen/archief/2013/2013-4005-wm.htm?Languageswitch=on> [accessed 2014-06-10] [WebCite Cache ID 6QE0xa1Q7]
23. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M, Medical Research Council Guidance. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008;337:a1655 [FREE Full text] [Medline: [18824488](https://pubmed.ncbi.nlm.nih.gov/18824488/)]
24. Brendryen H, Johansen A, Nesvåg S, Kok G, Duckert F. Constructing a Theory- and Evidence-Based Treatment Rationale for Complex eHealth Interventions: Development of an Online Alcohol Intervention Using an Intervention Mapping Approach. *JMIR Res Protoc* 2013;2(1):e6 [FREE Full text] [doi: [10.2196/resprot.2371](https://doi.org/10.2196/resprot.2371)] [Medline: [23612478](https://pubmed.ncbi.nlm.nih.gov/23612478/)]
25. Vonk Noordegraaf A, Huirne JA, Pittens CA, van Mechelen W, Broerse JE, Brölmann HA, et al. eHealth program to empower patients in returning to normal activities and work after gynecological surgery: intervention mapping as a useful method for development. *J Med Internet Res* 2012;14(5):e124 [FREE Full text] [doi: [10.2196/jmir.1915](https://doi.org/10.2196/jmir.1915)] [Medline: [23086834](https://pubmed.ncbi.nlm.nih.gov/23086834/)]
26. Vedel I, Akhlaghpour S, Vaghefi I, Bergman H, Lapointe L. Health information technologies in geriatrics and gerontology: a mixed systematic review. *J Am Med Inform Assoc* 2013;20(6):1109-1119. [doi: [10.1136/amiajnl-2013-001705](https://doi.org/10.1136/amiajnl-2013-001705)] [Medline: [23666776](https://pubmed.ncbi.nlm.nih.gov/23666776/)]
27. Eysenbach G. The law of attrition. *J Med Internet Res* 2005;7(1):e11 [FREE Full text] [doi: [10.2196/jmir.7.1.e11](https://doi.org/10.2196/jmir.7.1.e11)] [Medline: [15829473](https://pubmed.ncbi.nlm.nih.gov/15829473/)]
28. Scandurra I, Sjölander M. Participatory Design With Seniors: Design of Future Services and Iterative Refinements of Interactive eHealth Services for Old Citizens. *Med 2.0* 2013 Oct 08;2(2):e12 [FREE Full text] [doi: [10.2196/med20.2729](https://doi.org/10.2196/med20.2729)]

29. van Gemert-Pijnen JE, Nijland N, van Limburg M, Ossebaard HC, Kelders SM, Eysenbach G, et al. A holistic framework to improve the uptake and impact of eHealth technologies. *J Med Internet Res* 2011;13(4):e111 [[FREE Full text](#)] [doi: [10.2196/jmir.1672](https://doi.org/10.2196/jmir.1672)] [Medline: [22155738](https://pubmed.ncbi.nlm.nih.gov/22155738/)]
30. Makai P, Melis RJ, Olde-Rikkert MG. Technical difficulties and evaluating e-health interventions. *JAMA Intern Med* 2014 Feb 1;174(2):304-305. [doi: [10.1001/jamainternmed.2013.12753](https://doi.org/10.1001/jamainternmed.2013.12753)] [Medline: [24493611](https://pubmed.ncbi.nlm.nih.gov/24493611/)]

Abbreviations

ADL: Activities of Daily Living
ANCOVA: analysis of covariance
EASYcare-TOS: EASYcare Two-step Older person Screening
GP: general practitioner
IADL: Instrumental Activities of Daily Living
OHC: Online Health Community
ZWIP: Health and Welfare portal

Edited by G Eysenbach; submitted 29.10.13; peer-reviewed by M Wise, S Langrial; comments to author 09.01.14; revised version received 10.02.14; accepted 31.05.14; published 23.06.14

Please cite as:

Makai P, Perry M, Robben SHM, Schers HJ, Heinen MM, Olde Rikkert MGM, Melis RF
Evaluation of an eHealth Intervention in Chronic Care for Frail Older People: Why Adherence is the First Target
J Med Internet Res 2014;16(6):e156
URL: <http://www.jmir.org/2014/6/e156/>
doi: [10.2196/jmir.3057](https://doi.org/10.2196/jmir.3057)
PMID: [24966146](https://pubmed.ncbi.nlm.nih.gov/24966146/)

©Peter Makai, Marieke Perry, Sarah HM Robben, Henk J Schers, Maud M Heinen, Marcel GM Olde Rikkert, René F Melis. Originally published in the *Journal of Medical Internet Research* (<http://www.jmir.org>), 23.06.2014. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the *Journal of Medical Internet Research*, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.