

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/142256>

Please be advised that this information was generated on 2021-10-26 and may be subject to change.

SYNTHESIS OF DNA COMPLEMENTARY TO 14S CALF LENS
CRYSTALLIN MESSENGER RNA BY REVERSE TRANSCRIPTASE

Anton J.M. Berns and Hans Bloemendaal*

Laboratorium voor Biochemie, Universiteit van Nijmegen, Nijmegen, Nederland

Stephen J. Kaufman and Inder M. Verma

Department of Biology, Massachusetts Institute of Technology, Cambridge,
Massachusetts 02139 U.S.A.

Received April 13, 1973

SUMMARY

The 14S messenger RNA (mRNA), coding for the A₂ chain of α -crystallin was isolated from calf lens polyribosomes by zonal centrifugation and further purified by oligo(dT)-cellulose chromatography. The purified mRNA was used as a template to synthesize complementary DNA by reverse transcriptase. The DNA product appeared to be smaller in size than the RNA template but a faithful transcript of base sequences in the 14S α -crystallin mRNA.

INTRODUCTION

In vitro synthesis of DNA complementary to several eukaryotic mRNAs by the avian myeloblastosis virus (AMV) DNA polymerase has been recently reported (1-6). Most eukaryotic mRNAs have been shown to contain regions rich in adenylic acid (poly(A)) (7) as an integral part of their structure. If these poly(A) regions are located at or near the 3' end of the RNA, then an oligomer of (dT) can hydrogen bond to it and serve as primer to support the synthesis of complementary DNA by the reverse transcriptase. The 14S lens mRNA has been shown to contain poly(A) containing regions (8) and can thus serve as a template for the reverse transcriptase using oligomer of (dT) as primer.

In this communication we describe the synthesis of DNA complementary to the 14 S calf lens mRNA that codes for the A₂ chain of α -crystallin (9,10,11,12).

MATERIALS AND METHODS

Calf lens polyribosomes were isolated as described earlier (13). RNA was prepared by a chloroform-phenol extraction procedure described by Perry et al (14). The lens mRNA was purified by zonal centrifugation followed by oligo(dT)-cellulose chromatography (15) which resulted in a further 3 to 4 fold purification. The ability of the RNA to direct the synthesis of α -crystallin polypeptides in vitro was tested as described elsewhere (12). The reverse transcriptase from avian myelo-

*to whom requests for reprints should be addressed




Figure 1. Identification of 14S lens mRNA directed translational products, ³⁵S-methionine as described in ref. 12. After incubation an acidic acetone precipitation was carried out. The *in vitro* products were analyzed by SDS gel electrophoresis according to Laemmli (19). To each gel about 200 μg of protein was applied. 20 μg of crystallin polypeptides were added in order to compare the radioactive pattern with the stained bands. After staining and destaining longitudinal slices were dried down on filter paper under vacuo and autoradiographed for 3 weeks. The details of the procedure are described elsewhere (12).
 1) without 14S mRNA 2) with 14S mRNA 3) stained gel

blastosis virus was purified according to Verma and Baltimore (5). The single strand specific nuclease from *Aspergillus oryzae* was a generous gift of Drs. D. Housman and A. Skoultchi.

The reaction mixture consisted of the following components: 50 mM Tris-HCl, pH 8.3, 20 mM dithiothreitol, 6 mM magnesium acetate, 60 mM NaCl, 600 μM of the three unlabeled deoxyribonucleoside triphosphates, 80 μM of the ³H-labeled deoxyribonucleoside triphosphate, 10 μg/ml oligo(dT)₁₂₋₁₈, 100 μg/ml Actinomycin D, 0.2 to 0.5 μg of reverse transcriptase and 6 μg/ml of 14S mRNA. Incubations were carried out at 37°C for the specified times. The amount of incorporation was determined by measuring the acid-precipitable radioactivity as described previously (16).

TABLE 1

REQUIREMENTS FOR DNA SYNTHESIS USING 14S CALF LENS mRNA AS TEMPLATE

Exp. I	pmoles dGMP incorporated
Complete	15
-Actinomycin	25
-dTTP	2.3
-dCTP	2.5
-dATP	2.5
+RNAase	1.7
Exp. II	
Complete	43
-oligo(dT)+oligo(dG)	29
-oligo(dT)+oligo(dA)	3.4
-oligo(dT)-oligo(dC)	4.5
-primer	4.1

Except where indicated, the complete reaction mixture described in the experimental section was used. The amount of RNA in exp. I and exp. II was 850 pmoles nucleotides of RNA. Oligomers of dT, dG, dA and dC have chain length of approximately 12-18 nucleotides long and were obtained from Collaborative Res. Waltham, Mass. The ribonuclease reagent contained 400 μ g of Pancreatic ribonuclease A/ml (Worthington Biochemical), 80 μ g of ribonuclease T₁/ml (Calbiochem, 5000 U/mg) and 1 mg bovine serum albumin/ml in 0.01 M Tris-Cl, pH 7.6, and 0.01 M NaCl. The RNAase was added together with the other reaction constituents. Acid precipitable radioactivity was determined as described before (16).

RESULTS AND DISCUSSIONA) Purity of mRNA

Calf lens 14S mRNA used in these experiments was purified by zonal centrifugation and oligo(dT)-cellulose chromatography. Identification of the translational products suggest that the 14S mRNA preparation contained only a minor contamination of other lens mRNAs (Fig. 1).

synthesis of complementary DNA. The maximal synthesis was achieved between 120 to 180 min. of incubation. The amount of incorporation was proportional to the amount of RNA added. However at elevated RNA concentrations the incorporation is no longer linear (above 15,000 pmoles of nucleotide/ml).

C) Size of the complementary DNA

Figure 3 depicts the sedimentation pattern of the DNA on an alkaline sucrose gradient. The complementary 8S rabbit globin DNA was included as marker. The DNA complementary to the 14S lens mRNA appears slightly smaller and more heterogenous than globin DNA. Since the 14S mRNA appears larger than the 10S globin mRNA (although it directs the synthesis of a polypeptide with a molecular weight of only 20,000) this indicates that the mRNA is only partially transcribed or has been transcribed in several pieces.




Figure 4. Hybridization of the DNA product with the 14S mRNA.

High specific activity labeled DNA transcript (2400 cpm of ^3H dGMP/pmole) was made using the complete reaction mixture described in the Method section. The DNA was purified as described before (20). DNA-RNA hybridization was performed in 10 μl capillary tubes in a medium containing 0.2 M phosphate buffer, pH 7.0 and 0.5% SDS. After heating the mixture for 5 min. at 95°C the capillary tubes were transferred immediately to a 65°C waterbath. After hybridization the capillary tubes were cooled in ice, broken and the solution expelled into 1.0 ml of digestion buffer (1mM ZnSO_4 , 0.1 M sodium acetate, pH 4.5 and 10 $\mu\text{g}/\text{ml}$ calf thymus DNA). 6 μl of *Aspergillus* single strand nuclease was added and the solution was incubated at 45°C for 30 min. Acid-precipitable radioactivity was determined. The hybridization mixture in each capillary contained 1 pmole of complementary DNA and 100 pmol 14S mRNA.

The same situation has been encountered with the transcription of the AMV RNA where only DNA pieces of 5 to 7 were synthesized (17). It is possible that the secondary structure of the 14S crystallin mRNA and the AMV-RNA prevent their complete transcription.

(D) Nature of the product

To determine if a faithful transcript of DNA complementary to the 14S mRNA was synthesized we studied the ability of the DNA to hybridize with the RNA

template. The kinetics of hybridization are shown in Figure 4. About 90% of the DNA annealed to the RNA, indicating that the DNA product is a faithful transcript of the lens mRNA. Some self annealing could be detected in the absence of RNA suggesting that either the synthesis of double stranded DNA is not completely inhibited by actinomycin D or the single stranded DNA formed "hairpins". The extent of self annealing was subtracted from the RNA-DNA annealing values. When the DNA was hybridized to tRNA or 10S globin mRNA it remained completely sensitive to single strand specific nuclease.

CONCLUSION

The 14S calf lens mRNA can be transcribed into complementary DNA using purified DNA polymerase from AMV. The product though smaller in size than the template is a faithful transcript of the template. Attempts are being made to determine if there are any homologies in the base sequences of various fractions of calf lens mRNAs using the labeled complementary DNA as a probe. We are also trying to transcribe the complementary DNA into RNA by E. coli RNA polymerase in an attempt to synthesize labeled RNA for nucleotide sequences analysis. The availability of labeled complementary DNA will also enable us to determine the number of gene copies of α -crystallin 14S mRNA in the calf lens.

ACKNOWLEDGEMENT

The present investigations have partly been carried out under the auspices of the Netherlands Foundation for Chemical Research (SON) with financial aid of the Netherlands Organization for the Advancement of Pure Research (ZWO) and a contract to D. Baltimore from the Special Virus Cancer Program of the National Cancer Institute. Stephen J. Kaufman and Inder M. Verma were fellows of the Jane Coffin Child Memorial Fund for Medical Research.

We are thankful to Drs. A. Rich and D. Baltimore in whose laboratories this work was performed.

REFERENCES

1. Verma, I.M., Temple, G.F., Fan, H. and Baltimore, D. (1972). *Nature New Biology* 235, 163-167.
2. Kacian, D.L., Spiegelman, S., Bank, A., Terada, M., Metafora, S., Dow, L. and Marks, P.A. (1972). *Nature New Biology* 235, 167-169.
3. Ross, J., Aviv, H., Scolnick, E. and Leder, P. (1972). *Proc. Natl. Acad. Sci. U.S.A.* 69, 264-268.
4. Zassenhous, P. and Kates, J. (1972). *Nature New Biology* 238, 139-141.
5. Verma, I.M. and Baltimore, D. (1973). *Methods in Enzymology*, in press.
6. Verma, I.M. and Firtel, R.A., manuscript in preparation.
7. Slater, D.W., Slater, I. and Gillespie, D. (1972). *Nature*, 240, 333-337.
8. Piperno, G., Bertazzoni, U., Berns, A. and Bloemendal, H., manuscript in preparation.
9. Berns, A.J.M., Strous, G.J.A.M. and Bloemendal, H. (1972). *Nature New Biology* 236, 7-9.

10. Mathews, M.B., Osborn, M., Berns, A.J.M. and Bloemendal, H. (1972). *Nature New Biology* 236, 5-7.
11. Berns, A.J.M., van Kraaikamp, M., Bloemendal, H. and Lane, C.D. (1972). *Proc.Natl. Acad.Sci. U.S.A.* 69, 1606-1609.
12. Berns, A.J.M., Schreurs, V.V.A.M., van Kraaikamp, M. and Bloemendal, H. (1973). *Eur. J. Biochem.* 33, 551-557.
13. Bloemendal, H., Schoenmakers, J.G.G., Zweers, A., Matze, R. and Benedetti, E.L. (1966). *Biochim.Biophys.Acta* 123, 217-220.
14. Perry, R.P., La Torre, J., Kelley, D.E. and Greenberg, J.R.(1972). *Biochim. Biophys. Acta*, 262, 220-226.
15. Aviv, H. and Leder, P. (1972). *Proc.Natl.Acad.Sci. U.S.A.*, 69, 1408-1412.
16. Baltimore, D., Huang, A.S. and Stampfer, M. (1970). *Proc.Natl.Acad.Sci. U.S.A.* 66, 572-576.
17. Temin, H. and Baltimore, D. (1972). *Advances in Virus Research* 17, 129, Academic Press, Inc., New York and London.
18. Verma, I.M., Temple, G.F., Fan, H. and Baltimore, D. (1973). "Symposium on Control of Transcription" held at Calcutta, India Feb. 12-15, to be published.
19. Laemmli, U.K. (1970). *Nature* 227, 680-685.
20. Verma, I.M., Meuth, N.L., Bromfield, E., Manly, K.F. and Baltimore, D. (1971). *Nature New Biology* 233, 131-134.

