

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/143773>

Please be advised that this information was generated on 2019-11-19 and may be subject to change.

The Degree of Squares is an Atom (Extended Version^{*})

Jörg Endrullis¹, Clemens Grabmayer¹, Dimitri Hendriks¹, and
Hans Zantema^{2,3}

¹ Department of Computer Science, VU University Amsterdam

² Department of Computer Science, Eindhoven University of Technology

³ Institute for Computing and Information Science, Radboud University Nijmegen

Abstract. We answer an open question in the theory of degrees of infinite sequences with respect to transducibility by finite-state transducers. An initial study of this partial order of degrees was carried out in [5], but many basic questions remain unanswered. One of the central questions concerns the existence of atom degrees, other than the degree of the ‘identity sequence’ $10^0 10^1 10^2 10^3 \dots$. A degree is called an ‘atom’ if below it there is only the bottom degree $\mathbf{0}$, which consists of the ultimately periodic sequences. We show that also the degree of the ‘squares sequence’ $10^0 10^1 10^4 10^9 10^{16} \dots$ is an atom.

As the main tool for this result we characterise the transducts of ‘spiralling’ sequences and their degrees. We use this to show that every transduct of a ‘polynomial sequence’ either is in $\mathbf{0}$ or can be transduced back to a polynomial sequence for a polynomial of the same order.

1 Introduction

Finite-state transducers are ubiquitous in computer science, but little is known about the transducibility relation they induce on infinite sequences. A finite-state transducer (FST) is a deterministic finite automaton which reads the input sequence letter by letter, in each step producing an output word and changing its state. An example of an FST is depicted in Figure 1, where we write ‘ $a|w$ ’ along the transitions to indicate that the input letter is a and the output word is w . For example, it transduces the Thue-Morse sequence $T = 0110100110010110\dots$ to the period doubling sequence $P = 1011101010111011\dots$.

We are interested in transductions of infinite sequences. We say that a sequence σ is *transducible* to a sequence τ , $\sigma \geq \tau$, if there exists an FST that transforms σ into τ . The relation \geq is a preorder on the set $\Sigma^{\mathbb{N}}$ of infinite sequences, which induces an equivalence relation \equiv on $\Sigma^{\mathbb{N}}$, and a partial order on the set of (FST) *degrees*, that is, the equivalence classes with respect to \equiv .

So we have $T \geq P$. Also the back transformation can be realised by an FST, $P \geq T$. Hence the sequences are equivalent, $T \equiv P$, and are in the same degree.

^{*} This is an extended version of [3].




Fig. 1. A finite-state transducer realising the sum of consecutive bits modulo 2.

The bottom degree $\mathbf{0}$ is formed by the ultimately periodic sequences, that is, all sequences of the form $uvvv\dots$ for finite words u, v with v non-empty. Every infinite sequence can be transduced to any ultimately periodic sequence.

There is a clear analogy between degrees induced by transducibility and the recursion-theoretic degrees of unsolvability (Turing degrees). Hence many of the problems settled for Turing degrees, predominantly in the 1940s, 50s and 60s, can be asked again for FST degrees.




Fig. 2. Possible structures in the hierarchy (no intermediate points on the arrows).

Some initial results on FST degrees have been obtained in [5]: the partial order of degrees is not dense, not well-founded, there exist no maximal degrees, and a set of degrees has an upper bound if and only if the set is countable. The morphic degrees, and the computable degrees form subhierarchies. Also shown in [5] is the existence of an ‘atom’ degree, namely the degree of $10^010^110^210^3\dots$ ¹. A degree $D \neq \mathbf{0}$ is an atom if there exists no degree strictly between D and the bottom degree $\mathbf{0}$. Thus every transduct of a sequence in an atom degree D is either in $\mathbf{0}$ or still in D . The following questions, which have been answered for Turing degrees, remain open for the FST degrees:

- (i) How many atom degrees exist?
- (ii) When do two degrees have a supremum (or infimum)? In particular, are there pairs of degrees without a supremum (infimum)?

¹ In [5] atom degrees were called ‘prime degrees’. We prefer the more general notion of ‘atom’ because prime factorisation does not hold, see Theorem 4.27.

- (iii) Do the configurations displayed in Figure 1 exist?
- (iv) Can every finite distributive lattice be embedded in the hierarchy?

At the British Colloquium for Theoretical Computer Science 2014, Jeffrey Shallit offered £20 for each of the following questions, see [10]:

- (a) Is the degree of the Thue-Morse sequence an atom?
- (b) Are there are atom degrees other than that of $10^010^110^210^3\dots$?

We answer (b) by showing that the degree of the ‘squares’ $10^010^110^410^9\dots$ is an atom. The main tool that we use in the proof is a characterisation of transducts of ‘spiralling’ sequences (Theorem 4.25), which has the following consequence (Proposition 5.1): For all $k \geq 1$ it holds that every transduct $\sigma \notin \mathbf{0}$ of the sequence $\langle n^k \rangle = 10^{0^k}10^{1^k}10^{2^k}10^{3^k}\dots$ has, in its turn, a transduct of the form $\langle p(n) \rangle = 10^{p(0)}10^{p(1)}10^{p(2)}10^{p(3)}\dots$ where $p(n)$ is a polynomial also of degree k . This fact not only enables us to show that the degree of the squares sequence is an atom, by using it for $k = 2$, but it also suggests that the analogous result for the degree of $\langle n^k \rangle$, for arbitrary $k \geq 1$, has come within reach. For this it would namely suffice to show, for polynomials $p(n)$ of degree k , that $\langle p(n) \rangle$ can be transduced back to $\langle n^k \rangle$.

We obtain that there is a pair of non-zero degrees whose infimum is $\mathbf{0}$, namely the pair of atom degrees of $\langle n \rangle$ and $\langle n^2 \rangle$. We moreover use Theorem 4.25 to show that there is a non-atom degree that has no atom degree below it (Theorem 4.27).

2 Preliminaries

We use standard terminology and notation, see, e.g., [8]. Let Σ be an alphabet, i.e., a finite non-empty set of symbols. We denote by Σ^* the set of all finite words over Σ , and by ε the empty word. We let $\Sigma^+ = \Sigma^* \setminus \{\varepsilon\}$. The set of infinite sequences over Σ is $\Sigma^{\mathbb{N}} = \{\sigma \mid \sigma : \mathbb{N} \rightarrow \Sigma\}$ with $\mathbb{N} = \{0, 1, 2, \dots\}$, the set of natural numbers. For $u \in \Sigma^*$, we let $u^\omega = uuu\dots$. We define $\mathbb{N}_{<k} = \{0, 1, \dots, k-1\}$. We let $\Sigma^\infty = \Sigma^* \cup \Sigma^{\mathbb{N}}$ denote the set of all words over Σ . For $u \in \Sigma^*$ and $v \in \Sigma^\infty$ we write $u \sqsubseteq v$ to denote that u is a prefix of v , that is, when $v = uv'$ for some $v' \in \Sigma^\infty$. For $f : \mathbb{N} \rightarrow A$ and $k \in \mathbb{N}$, the k -th shift of f is defined by $\mathcal{S}^k(f)(n) = f(n+k)$, for all $n \in \mathbb{N}$.

We use the notation \mathbf{a} to denote a tuple $\mathbf{a} = \langle a_0, a_1, \dots, a_{k-1} \rangle$ where the a_i are elements from some set A ; the length of \mathbf{a} is k . Given a tuple \mathbf{a} we use a_i for the element indexed i , that is, we start indexing from 0 onward. We use the notation \mathbf{a}' to denote the rotated tuple $\mathbf{a}' = \langle a_1, \dots, a_{k-1}, a_0 \rangle$.

3 Finite-State Transducers and Degrees

For a thorough introduction to finite-state transducers, we refer to [8]. Here we only consider *complete pure sequential* finite-state transducers, where for every state q and input letter a there is precisely one successor state $\delta(q, a)$, and the functions realised by these transducers preserve prefixes. We use $\mathbf{2} = \{0, 1\}$ both for the input and the output alphabet.

Definition 3.1. A *finite-state transducer (FST)* is a tuple $T = \langle Q, q_0, \delta, \lambda \rangle$ where Q is a finite set of states, $q_0 \in Q$ is the initial state, $\delta : Q \times \mathbf{2} \rightarrow Q$ is the transition function, and $\lambda : Q \times \mathbf{2} \rightarrow \mathbf{2}^*$ is the output function.

We homomorphically extend the transition function δ to $Q \times \mathbf{2}^* \rightarrow Q$ and the output function λ to $Q \times \mathbf{2}^\infty \rightarrow \mathbf{2}^\infty$ as follows:

$$\begin{aligned} \delta(q, \varepsilon) &= q & \delta(q, au) &= \delta(\delta(q, a), u) & (q \in Q, a \in \mathbf{2}, u \in \mathbf{2}^*) \\ \lambda(q, \varepsilon) &= \varepsilon & \lambda(q, au) &= \lambda(q, a) \cdot \lambda(\delta(q, a), u) & (q \in Q, a \in \mathbf{2}, u \in \mathbf{2}^\infty). \end{aligned}$$

The function $T : \mathbf{2}^\infty \rightarrow \mathbf{2}^\infty$ realised by the FST T is defined by $T(u) = \lambda(q_0, u)$, for all $u \in \mathbf{2}^\infty$.

We note that finite state transducers and transduction of infinite sequences can be implemented as infinitary rewrite systems [2,6,4].

Definition 3.2. Let $T = \langle Q, q_0, \delta, \lambda \rangle$ be an FST. A *zero-loop* in T is a sequence of states q_1, \dots, q_n with $n > 1$ such that $q_1 = q_n$ and $q_i \neq q_j$ for all i, j with $1 \leq i < j < n$ and $q_{i+1} = \delta(q_i, 0)$ for all $1 \leq i < n$. The *length* of the zero-loop is $n - 1$.

Note that there can only be finitely many zero-loops in an FST.

Definition 3.3. Let $T = \langle Q, q_0, \delta, \lambda \rangle$ be an FST. We define $Z(T)$ as the least common multiple of the lengths of all zero-loops of T .

Let T be an FST with states Q . From any state $q \in Q$, after reading the word $0^{|Q|}$, the automaton must have entered a zero-loop (by the pigeonhole principle there must be a state repetition). By definition of $Z(T)$, the length ℓ of this loop divides $Z(T)$; say $Z(T) = d\ell$ for some $d \geq 1$. As a consequence, reading $0^{|Q|+i \cdot Z(T)}$ yields the output $\lambda(q, 0^{|Q|})$ followed by di copies of the output of the zero-loop. This yields a pumping lemma for FSTs, see also [5, Lemma 29].

Lemma 3.4. Let $T = \langle Q, q_0, \delta, \lambda \rangle$ be an FST. For every $q \in Q$ and $n \geq |Q|$ there exist words $p, c \in \mathbf{2}^*$ such that for all $i \in \mathbb{N}$, $\delta(q, 10^{n+i \cdot Z(T)}) = \delta(q, 10^n)$ and $\lambda(q, 10^{n+i \cdot Z(T)}) = pc^i$.

Proof. Let $q \in Q$ and $n \geq |Q|$. Then there exist $k, \ell \in \mathbb{N}$ such that $k + \ell \leq n$ and $\delta(q, 10^k) = \delta(q, 10^{k+\ell})$. Let k, ℓ be minimal with these properties. Then ℓ is the length of a zero-loop in T , and so $Z(T) = d\ell$ for some $d \geq 1$. We have $\delta(q, 10^{k'}) = \delta(q, 10^{k'+\ell})$ for all $k' \geq k$, and so $\delta(q, 10^n) = \delta(q, 10^{n+i \cdot d\ell}) = \delta(q, 10^{n+i \cdot Z(T)})$, for all $i \in \mathbb{N}$. Let $q' = \delta(q, 10^n)$, $p = \lambda(q, 10^n)$ and $c = \lambda(q', 0^{Z(T)})$. Then we have $\lambda(q, 10^{n+i \cdot Z(T)}) = \lambda(q, 10^n) \cdot \lambda(q', 0^{i \cdot Z(T)}) = pc^i$, for all $i \in \mathbb{N}$. \square

Definition 3.5. Let T be an FST, and let $\sigma, \tau \in \mathbf{2}^\mathbb{N}$ be infinite sequences. We say that T *transduces* σ to τ , and that τ is the *T -transduct* of σ , which we denote by $\sigma \geq_T \tau$, whenever $T(\sigma) = \tau$. We write $\sigma \geq \tau$, and call τ a *transduct* of σ , if there exists an FST T such that $\sigma \geq_T \tau$.

Clearly, the relation \geq is reflexive. By composition of FSTs (the so-called ‘wreath product’), the relation \geq is also transitive, see [5, Remark 9]. We write $\sigma > \tau$ when $\sigma \geq \tau$ but not $\tau \geq \sigma$. Whenever we have $\sigma \geq \tau$ as well as a back-transduction $\tau \geq \sigma$, we consider σ and τ to be equivalent.

Definition 3.6. We define the relation $\equiv \subseteq \mathbf{2}^{\mathbb{N}} \times \mathbf{2}^{\mathbb{N}}$ by $\equiv = \geq \cap \geq^{-1}$. For a sequence $\sigma \in \mathbf{2}^{\mathbb{N}}$ the equivalence class $[\sigma] = \{\tau \mid \sigma \equiv \tau\}$ is called the *degree* of σ .

A degree $[\sigma]$ is an *atom* if $[\sigma] \neq \mathbf{0}$ and there is no degree $[\tau]$ such that $[\sigma] > [\tau] > \mathbf{0}$.

4 Characterising Transducts of Spiralling Sequences

In this section we characterize the transducts of ‘spiralling’ sequences. Proofs omitted in the text can be found in the appendix.

Definition 4.1. For a function $f : \mathbb{N} \rightarrow \mathbb{N}$ we define the sequence $\langle f \rangle \in \mathbf{2}^{\mathbb{N}}$ by

$$\langle f \rangle = \prod_{i=0}^{\infty} 10^{f(i)} = 10^{f(0)} 10^{f(1)} 10^{f(2)} \dots$$

For a sequence $\langle f \rangle$, we often speak of the n -th *block* of $\langle f \rangle$ to refer to the occurrence of the word $10^{f(n)}$ in $\langle f \rangle$.

In the sequel we often write $\langle f(n) \rangle$ to denote the sequence $\langle n \mapsto f(n) \rangle$. We note that there is a one-to-one correspondence between functions $f : \mathbb{N} \rightarrow \mathbb{N}$, and infinite sequences over the alphabet $\mathbf{2}$ that start with the letter 1 and that contain infinitely many occurrences of the letter 1. Every degree is of the form $[\langle f \rangle]$ for some $f : \mathbb{N} \rightarrow \mathbb{N}$.

The following lemma is concerned with some basic operations on functions that have no effect on the degree of $\langle f \rangle$ (multiplication with a constant, and x - and y -shifts), and others by which we may go to a lower degree (taking subsequences, merging blocks).

Lemma 4.2. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$, and $a, b \in \mathbb{N}$. It holds that:*

- (i) $\langle af(n) \rangle \equiv \langle f(n) \rangle$, for $a > 0$,
- (ii) $\langle f(n+a) \rangle \equiv \langle f(n) \rangle$,
- (iii) $\langle f(n) + a \rangle \equiv \langle f(n) \rangle$,
- (iv) $\langle f(n) \rangle \geq \langle f(an) \rangle$, for $a > 0$,
- (v) $\langle f(n) \rangle \geq \langle af(2n) + bf(2n+1) \rangle$.

Proof. Recall that, for any $g : \mathbb{N} \rightarrow \mathbb{N}$, with the n -th *block* of $\langle g \rangle$ we mean the factor $10^{g(n)}$ of $\langle g \rangle$.

- (i) We have $\langle af(n) \rangle \geq_T \langle f(n) \rangle$ where T is an FST that replaces every factor 0^a by 0. Clearly, the inverse operation is also FST-realizable, so $\langle f(n) \rangle \geq \langle af(n) \rangle$.

- (ii) An FST that prepends $\prod_{i=0}^{a-1} 10^{f(i)}$ to the incoming word witnesses $\langle f(n+a) \rangle \geq \langle f(n) \rangle$. For the converse direction, let the FST remove the first a blocks from the input.
- (iii) To see that $\langle f(n+a) \rangle \geq \langle f(n) \rangle$, consider an FST that removes a factor 0^a from every block in $\langle f(n+a) \rangle$. Conversely, $\langle f(n) \rangle \geq \langle f(n+a) \rangle$ is witnessed by an FST that appends 0^a to every block.
- (iv) It is clear that the operation of selecting every a -th block is realisable by an FST.
- (v) By Theorem 4.25. □

Definition 4.3. Let A be a set. A function $f : \mathbb{N} \rightarrow A$ is *ultimately periodic* if for some integers $n_0 \geq 0$, $p > 0$ we have $f(n+p) = f(n)$ for all $n \geq n_0$.

Definition 4.4. A function $f : \mathbb{N} \rightarrow \mathbb{N}$ is called *spiralling* if

- (i) $\lim_{n \rightarrow \infty} f(n) = \infty$, and
- (ii) for every $m \geq 1$, the function $n \mapsto f(n) \bmod m$ is ultimately periodic.

Functions with the property (ii) in Definition 4.4 have been called ‘ultimately periodic reducible’ by Siefkes [11] (quotation from [9]). Note that the identity function is spiralling. Furthermore scalar products, and pointwise sums and products of spiralling functions are again spiralling. As a consequence also polynomials $a_k n^k + a_{k-1} n^{k-1} + \dots + a_0$ are spiralling.

In the remainder of this section, we will characterise the transducts σ of $\langle f \rangle$ for spiralling f . We will show that if such a transduct σ is not ultimately periodic, then it is equivalent to a sequence $\langle g \rangle$ for a spiralling function g , and moreover, g can be obtained from f by a ‘weighted product’.

Lemma 4.5. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a spiralling function. We have $\langle f \rangle \geq \sigma$ if and only if σ is of the form*

$$\sigma = w \cdot \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_j c_j^{\varphi(i,j)} \quad \text{where} \quad \varphi(i,j) = \frac{f(n_0 + mi + j) - a_j}{z}, \quad (1)$$

for some integers $n_0, m, a_j \geq 0$ and $z > 0$, and finite words $w, p_j, c_j \in \mathbf{2}^*$ ($0 \leq j < m$) such that $\varphi(i,j) \in \mathbb{N}$ for all $i \in \mathbb{N}$ and $j \in \mathbb{N}_{< m}$.

Proof. Assume $\langle f \rangle \geq \sigma$ and let $T = \langle Q, q_0, \delta, \lambda \rangle$ be an FST that transduces $\langle f \rangle$ to σ . As f is spiralling, there exist $\ell_0, p \in \mathbb{N}$, $p > 0$ such that $f(n) \equiv f(n+p) \pmod{Z(T)}$ for every $n \geq \ell_0$. Moreover, as a consequence of $\lim_{n \rightarrow \infty} f(n) = \infty$, there exists $\ell_1 \in \mathbb{N}$ such that $f(n) \geq |Q|$ for every $n \geq \ell_1$.

For $n \in \mathbb{N}$, let $q_n \in Q$ be the state that the automaton T is in, before reading the n -th occurrence of 1 in the sequence σ (i.e., the start of the block $10^{f(n)}$). By the pigeonhole principle there exist $n_0, m \in \mathbb{N}$ with $\max\{\ell_0, \ell_1\} < n_0$ and $m > 0$ such that $m \equiv 0 \pmod{p}$ and $q_{n_0} = q_{n_0+m}$. Then, for every $i \in \mathbb{N}$ and $j \in \mathbb{N}_{< m}$, we have $f(n_0 + mi + j) \geq |Q|$ and $f(n_0 + mi + j) \equiv f(n_0 + j) \pmod{Z(t)}$. For $j \in \mathbb{N}_{< m}$, we define $a_j = \min\{f(n_0 + mi + j) \mid i \in \mathbb{N}\}$. Note that $a_j \geq |Q|$.




Fig. 3. An FST that transduces $\prod_{i=0}^{\infty} \prod_{j=0}^{m-1} 10^{\varphi(i,j)}$ into $\prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_j c_j^{\varphi(i,j)}$.

Then for all $i \in \mathbb{N}$ and $j \in \mathbb{N}_{< m}$ we have: $f(n_0 + mi + j) = a_j + \varphi(i, j) \cdot z$ where $\varphi(i, j) = (f(n_0 + mi + j) - a_j)/z$ and $z = Z(T)$. Using Lemma 3.4 it follows by induction that $q_{n_0+n} = q_{n_0+m+n}$ for every $n \in \mathbb{N}$. Hence $q_{n_0+j} = q_{n_0+mi+j}$ for every $i \in \mathbb{N}$ and $j \in \mathbb{N}_{< m}$. Then, again by Lemma 3.4, for every $j \in \mathbb{N}_{< m}$ there exist words $p_j, c_j \in \mathbf{2}^*$ such that $\lambda(q_{n_0+mi+j}, 10^{f(n_0+mi+j)}) = \lambda(q_{n_0+j}, 10^{a_j + \varphi(i,j) \cdot Z(T)}) = p_j c_j^{\varphi(i,j)}$ for all $i \in \mathbb{N}$. We conclude that

$$\sigma = \prod_{n=0}^{\infty} \lambda(q_n, 10^{f(n)}) = w \cdot \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_j c_j^{\varphi(i,j)},$$

where $w = \prod_{n=0}^{n_0-1} \lambda(q_n, 10^{f(n)})$.

For the other direction, assume σ is of the form (1) for some $n_0, m, a_j, z \in \mathbb{N}$ with $z > 0$, and $w, p_j, c_j \in \mathbf{2}^*$ ($0 \leq j < m$) such that $\varphi(i, j) = (f(n_0 + mi + j) - a_j)/z \in \mathbb{N}$ for all $i \in \mathbb{N}$ and $j \in \mathbb{N}_{< m}$. We define an FST T such that $\langle f \rangle \geq_T \sigma'$, where


$$\sigma' = \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_j c_j^{\varphi(i,j)}$$

Define $T = \langle Q, q_{m-1}^0, \delta, \lambda \rangle$ where $Q = \{q_j^h \mid j \in \mathbb{N}_{< m}, -a_j \leq h < z\}$, and

$$\begin{aligned} \langle \delta, \lambda \rangle (q_j^h, 0) &= \langle q_{j'}^{h'}, 0^e \rangle && \text{where } e = 1 \text{ and } h' = 0 \text{ if } h + 1 = z, \\ &&& \text{and } e = 0 \text{ and } h' = h + 1, \text{ otherwise,} \\ \langle \delta, \lambda \rangle (q_j^h, 1) &= \langle q_{j'}^{-a_{j'}}, 1 \rangle && (j' = j + 1 \bmod m). \end{aligned}$$

For the verification of $\langle f \rangle \geq_T \sigma'$ we refer to the proof of Lemma 4.19 that contains a more general construction. The transduction $\sigma' \geq \mathcal{S}^{|\omega|}(\sigma)$ is realised by the FST in Figure 3. Clearly also $\mathcal{S}^{|\omega|}(\sigma) \geq \sigma$ holds, and we conclude $\langle f \rangle \geq \sigma$. \square

Example 4.6. We illustrate the influence of both the zero-loops of the automaton as well as the growth of the block lengths of the input sequence, on the size m of the inner product of Lemma 4.5. Consider the FST $T = \langle \{q_0, q_1, q_2\}, q_0, \delta, \lambda \rangle$ given by



and the sequence

$$\langle \lfloor \frac{n}{2} \rfloor \rangle = 11101010^210^210^310^310^410^4 \dots ,$$

where $\lfloor x \rfloor = \max\{n \in \mathbb{N} \mid n \leq x\}$. We investigate the sequence $r_0 a_0 r_1 a_1 r_2 \dots$ of states r of T alternated with letters a from the input sequence, in such a way that T is in state r_n after having read the word $a_0 a_1 \dots a_{n-1}$,

$$\underline{q_0} 1 q_2 1 q_1 1 \underline{q_0} 0 q_1 1 q_0 0 q_1 1 q_0 0 q_1 0 q_2 1 q_1 0 q_2 0 \underline{q_0} 1 q_2 0 q_0 0 q_1 0 q_2 1 q_1 0 q_2 0 q_0 0 \dots$$

The two underlined occurrences of q_0 indicate a repetition of states in combination with a repetition of the block size modulo $Z(T) = 3$. The number of blocks between these occurrences, forming the repetition, is $m = 6$. Actually, following the proof of Lemma 4.5 precisely, the algorithm would instead select the repetition starting from the second underlined occurrence of q_0 . The reason is that in general, when reading a block $100 \dots 0$, only after reading $|Q|$ zeros, we are guaranteed to be in a zero-loop. For this FST T , all states are on a zero-loop, and so we enter the loop immediately.


From Lemma 4.5 it follows that FSTs can only transform the length of blocks by linear functions (and merge blocks). As a consequence, FSTs typically cannot slow down the growth rate of blocks in spiralling sequences by more than a linear factor. This yields the following simple criterion for non-reducibility.

Lemma 4.7. *Let $f, g : \mathbb{N} \rightarrow \mathbb{N}$ be such that f is spiralling, g is not ultimately periodic and $g \in o(f)$, i.e., for every $a \in \mathbb{N}$ there is $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$ it holds that $f(n) \geq ag(n)$. Then $\langle f \rangle \not\geq \langle g \rangle$.*

Note that the reverse $\langle g \rangle \not\geq \langle f \rangle$ does *not* follow. For example, we have $\langle 4^n \rangle \not\geq \langle 2^n \rangle$, but $\langle 2^n \rangle \geq \langle 4^n \rangle$.

There can be several ways to factor the transduct σ in the statement of Lemma 4.5, as shown by the following example.

Example 4.8. Consider the following FST $T = \langle \{q_0, q_1\}, q_0, \delta, \lambda \rangle$



together with the sequence $\langle n \rangle = 11010^210^310^4 \dots$. The T -transduct of $\langle n \rangle$ is

$$T(\langle n \rangle) = 11(01)1(10)^21(01)^31(10)^4 \dots$$

Using the double product (1) of Lemma 4.5 we have $w = \varepsilon$, $n_0 = 0$, $m = 2$, $p_0 = p_1 = 1$, $c_0 = 01$, $c_1 = 10$, $a_0 = a_1 = 0$, and $z = Z(T) = 1$. Thus, $\varphi(i, j) = 2i + j$. Note that $c_1^\omega = p_0 c_0^\omega$, and so we can merge the factors of the inner product, and decrease its size to $m = 1$. Then $T(\langle n \rangle) = 11011101(0101)1101(0101)^21101(0101)^3 \dots$, that is, in the double product (1), $T(\langle n \rangle)$ can be factored by choosing $m = 1$, $p_0 = 1101$, and $c_0 = 0101$.

Whenever we have the situation $c_j^\omega = p_{j+1}c_{j+1}^\omega$ in the representation (1) for some $j \in \mathbb{N}_{< m}$ (addition is modulo m), we speak of a ‘transition ambiguity’. We will eliminate such ambiguities by merging factors of the inner product, as in Lemma 4.10. In general, this merging may involve weighted sums in the exponents. The following lemma is folklore, see for instance [1].

Lemma 4.9. *If a sequence σ is periodic with period lengths k and ℓ , then it is periodic with period length $\gcd(k, \ell)$.*

Lemma 4.10. *Let $u, v, w \in \mathbf{2}^*$ be finite words with $u, w \neq \varepsilon$ such that $u^\omega = vw^\omega$. Then there exists a word $x \in \mathbf{2}^*$ and $a, b \in \mathbb{N}$ such that $u^m v w^n = vx^{am+bn}$ for all $m, n \in \mathbb{N}$.*

Proof. Let $d = \gcd(|u|, |w|)$, $a = |u|/d$ and $b = |w|/d$. Let x be the suffix of w of length d . We argue that this choice has the desired property. The sequence $\sigma = u^\omega = vw^\omega$ is periodic with period lengths $|u|$ and $|w|$. By Lemma 4.9 it follows that σ is also periodic with period length d . Hence $w = x^b$ and for all $m, n \in \mathbb{N}$, $u^m v w^n$ is a prefix of σ . Likewise vx^{am+bn} is a prefix of $\sigma = vx^\omega$, and both prefixes have the same length: $|u^m v w^n| = m|u| + |v| + n|w| = |v| + am|x| + bn|x| = |vx^{am+bn}|$. We conclude $u^m v w^n = vx^{am+bn}$. \square

Our goal is to obtain a simple characterisation of the degrees of the transducts σ of spiralling sequences $\langle f \rangle$. To this end, we transform the transduct σ into a sequence σ' by replacing in the double product of Lemma 4.5 the displayed occurrence of p_j by 1 and of c_j by 0 for every $j \in \mathbb{N}_{< m}$. To guarantee that this transformation does not change the degree, that is $\sigma' \equiv \sigma$, we first have to resolve transition ambiguities.

For the back transformation blocks $10^{\varphi(i,j)}$ have to be replaced by $p_j c_j^{\varphi(i,j)}$, an operation that is easily realised by an FST, see Figure 3.

If the product does not contain transition ambiguities, then also the transformation from σ into σ' can be realised by an FST and thus does not change the degree of the sequence, hence $\sigma \equiv \sigma'$. If there exists $j \in \mathbb{N}_{< m}$ with $c_j^\omega = p_{j+1}c_{j+1}^\omega$, then, by this transformation of σ into σ' , one possibly leaves the degree of σ , i.e., $\sigma' > \sigma$. because for large enough $i \in \mathbb{N}$, an FST cannot recognise where a block $p_j c_j^{\varphi(i,j)}$ ends and where the next block $p_{j+1} c_{j+1}^{\varphi(i,j+1)}$ starts. This might make it impossible to realise the transformation by an FST, as then the FST cannot replace p_{j+1} by 1.

Definition 4.11. A *weight* is a tuple $\langle a_0, \dots, a_{k-1}, b \rangle \in \mathbb{Q}^{k+1}$ of rational numbers such that $a_0, \dots, a_{k-1} \geq 0$. Given a weight $\alpha = \langle a_0, \dots, a_{k-1}, b \rangle$ and a function $f : \mathbb{N} \rightarrow \mathbb{N}$ we define $\alpha \cdot f \in \mathbb{Q}$ by

$$\alpha \cdot f = a_0 f(0) + a_1 f(1) + \dots + a_{k-1} f(k-1) + b.$$

The weight α is called *constant* when $a_j = 0$ for all $j \in \mathbb{N}_{<k}$. For a tuple of weights $\boldsymbol{\alpha} = \langle \alpha_0, \dots, \alpha_{m-1} \rangle$ we define its *rotation* by $\boldsymbol{\alpha}' = \langle \alpha_1, \dots, \alpha_{m-1}, \alpha_0 \rangle$.

For functions $f : \mathbb{N} \rightarrow \mathbb{N}$, and tuples $\boldsymbol{\alpha} = \langle \alpha_0, \alpha_1, \dots, \alpha_{m-1} \rangle$ of weights, the *weighted product* of $\boldsymbol{\alpha}$ and f is a function $\boldsymbol{\alpha} \otimes f : \mathbb{N} \rightarrow \mathbb{Q}$ that is defined by induction on n through the following scheme of equations:

$$\begin{aligned} (\boldsymbol{\alpha} \otimes f)(0) &= \alpha_0 \cdot f \\ (\boldsymbol{\alpha} \otimes f)(n+1) &= (\boldsymbol{\alpha}' \otimes \mathcal{S}^{|\alpha_0|-1}(f))(n) \quad (n \in \mathbb{N}) \end{aligned}$$

where $|\alpha_i|$ is the length of the tuple α_i , and $\mathcal{S}^k(f)$ is the k -th shift of f . A weighted product $\boldsymbol{\alpha} \otimes f$ is called *natural* if $(\boldsymbol{\alpha} \otimes f)(n) \in \mathbb{N}$ for all $n \in \mathbb{N}$.

In what follows, all weighted products $\boldsymbol{\alpha} \otimes f$ that we consider are assumed to be natural.

Example 4.12. Let $f(n) = n$ for all $n \in \mathbb{N}$, and $\boldsymbol{\alpha} = \langle \alpha_1, \alpha_2 \rangle$ with $\alpha_1 = \langle 1, 2, 3, 4 \rangle$, $\alpha_2 = \langle 0, 1, 1 \rangle$. Interpreting the functions f and $\boldsymbol{\alpha} \otimes f$ as sequences, the computation of $\boldsymbol{\alpha} \otimes f$ can be visualised as follows:

$$\begin{array}{cccccccccccc} f & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & \dots \\ & & \swarrow \times 1 & \downarrow \times 2 & \swarrow \times 3 & & \swarrow \times 0 & \downarrow \times 1 & \swarrow \times 1 & \downarrow \times 2 & \swarrow \times 3 & & \swarrow \times 0 & \downarrow \times 1 & \swarrow \times 1 & & \dots \\ \boldsymbol{\alpha} \otimes f & & & 12 & & 5 & & 42 & & 10 & & \dots \end{array}$$

Thus, for $n = 0, 1, 2, 3, \dots$, $(\boldsymbol{\alpha} \otimes f)(n)$ takes the values 12, 5, 42, 10, \dots .

Lemma 4.13. Let $\boldsymbol{\alpha}$ be an m -tuple of weights ($m > 0$), and let $f : \mathbb{N} \rightarrow \mathbb{N}$. For all $n \in \mathbb{N}$ we have $(\boldsymbol{\alpha} \otimes f)(n) = \alpha_r \cdot \mathcal{S}^t(f)$ where $q, r \in \mathbb{N}$ with $r < m$ are such that $n = qm + r$, and $t = q \cdot \sum_{j=0}^{m-1} (|\alpha_j| - 1) + \sum_{j=0}^{r-1} (|\alpha_j| - 1)$.

Proof. By induction on $n \in \mathbb{N}$ we show that, for all $f : \mathbb{N} \rightarrow \mathbb{N}$, and all m -tuples of weights $\boldsymbol{\alpha}$, we have

$$\begin{aligned} (\boldsymbol{\alpha} \otimes f)(n) &= \alpha_r \cdot \mathcal{S}^t(f) \quad \text{where } n = qm + r, \text{ and } q, r \in \mathbb{N} \text{ with } r < m, \\ &\text{and } t = q \cdot \sum_{i=0}^{m-1} (|\alpha_i| - 1) + \sum_{i=0}^{r-1} (|\alpha_i| - 1). \end{aligned}$$

The base case, $n = 0$ follows directly from the definition. For the induction step let $\boldsymbol{\alpha}$ and $f : \mathbb{N} \rightarrow \mathbb{N}$ be arbitrary, let $n = n' + 1$, $q' = \lfloor n'/m \rfloor$, and $r' = n' - q'm$. We abbreviate $k_i = |\alpha_i| - 1$ ($i \in \mathbb{N}_{<m}$) and $s_\ell = \sum_{i=0}^{\ell-1} k_i$ ($\ell \in \mathbb{N}_{\leq m}$). By Definition 4.11 and the induction hypothesis we obtain

$$(\boldsymbol{\alpha} \otimes f)(n) = (\boldsymbol{\alpha}' \otimes \mathcal{S}^{k_0}(f))(n') = \alpha_{r'+1} \cdot \mathcal{S}^{s_{q'+1}}(\mathcal{S}^{k_0}(f))$$

with $t' = q' \cdot s_m + s_{r'+1} - k_0$, and where addition in the subscript of α is modulo m . We conclude $(\alpha \otimes f)(n) = \alpha_r \cdot \mathcal{S}^t(f)$ where $n = (q' + 1)m$, $r = 0$ and $t = (q' + 1) \cdot s_m$ if $r' = m - 1$, and $n = q'm + r$, $r = r' + 1$ and $t = q' \cdot s_m + r$ if $r' < m - 1$.

Lemma 4.14. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$. If $\langle \alpha \otimes f \rangle \notin \mathbf{0}$, then there exists $i \in \mathbb{N}_{<|\alpha|}$ such that α_i is a non-constant weight.*

Proof. Assume α_i is a constant weight for all $i \in \mathbb{N}_{<|\alpha|}$; let $\alpha_i = \langle 0, 0, \dots, 0, b_i \rangle$. Then $\alpha \otimes f : \mathbb{N} \rightarrow \mathbb{N}$ is periodic: for all $n \in \mathbb{N}$, we have $(\alpha \otimes f)(n) = b_i$ where $i \equiv n \pmod{|\alpha|}$ and $i \in \mathbb{N}_{<|\alpha|}$. Hence $\langle \alpha \otimes f \rangle \in \mathbf{0}$.

We define the operation zip [7] also known as ‘perfect shuffle’.

Definition 4.15. Let $k \in \mathbb{N}$. For $i \in \mathbb{N}_{<k}$, let $f_i : \mathbb{N} \rightarrow \mathbb{N}$ be a function. We define the function $\text{zip}_k(f_0, f_1, \dots, f_{k-1}) : \mathbb{N} \rightarrow \mathbb{N}$ by

$$\text{zip}_k(f_0, f_1, \dots, f_{k-1})(kn + i) = f_i(n) \quad (n \in \mathbb{N}, i \in \mathbb{N}_{<k}).$$

Lemma 4.16. *Let $f_0, f_1, \dots, f_{k-1} : \mathbb{N} \rightarrow \mathbb{N}$ be such that f_i ($i \in \mathbb{N}_{<k}$) is ultimately periodic modulo every $m \geq 1$ (see Definition 4.4 (ii)). Then the function $\text{zip}_k(f_0, f_1, \dots, f_{k-1})$ is ultimately periodic modulo every $m \geq 1$.*

Proof. Fix an arbitrary $m \geq 1$. From the assumption we obtain, for every $i \in \mathbb{N}_{<k}$, the existence of $n_i, p_i \in \mathbb{N}$ with $p_i \geq 1$ such that $f_i(n+p_i) \equiv f_i(n) \pmod{m}$ for all $n \geq n_0$. Define $n' = \max\{n_i \mid i \in \mathbb{N}_{<k}\}$ and $p = \text{lcm}\{p_i \mid i \in \mathbb{N}_{<k}\}$. Let $f = \text{zip}_k(f_0, f_1, \dots, f_{k-1})$. Let $n \geq n'$ be arbitrary, and let $q, i \in \mathbb{N}$ with $i < k$ be such that $n = kq + i$. Then we have $f(n+kp) = f(k(q+p) + i) = f_i(q+p) \equiv f_i(q) \pmod{m}$, and $f(n) = f(kq + i) = f_i(q)$, as desired. \square

Lemma 4.17. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a function, and let α be an m -tuple of weights. Then there is a m -tuple of weights β such that*

$$\alpha \otimes f = \text{zip}_m(g_0, g_1, \dots, g_{m-1}) \quad \text{where } g_i = \langle \beta_i \rangle \otimes f \text{ for } i \in \mathbb{N}_{<m}.$$

Proof. Let $k_i = |\alpha_i| - 1$ ($i \in \mathbb{N}_{<m}$), and $\alpha_i = \langle a_{i,0}, a_{i,1}, \dots, a_{i,k_i-1}, b_i \rangle$. For $\ell \in \mathbb{N}_{\leq m}$ we define $s_\ell = \sum_{i=0}^{\ell-1} k_i$. For $i \in \mathbb{N}_{<m}$, we define the weight β_i of length $s_m + 1$ by

$$(\beta_i)_{s_m} = b_i \quad (\beta_i)_{s_j+h} = \begin{cases} a_{i,h} & \text{if } j = i, \\ 0 & \text{if } j \neq i \end{cases} \quad (j \in \mathbb{N}_{<m}, h \in \mathbb{N}_{<k_i}).$$

Let $g = \text{zip}_m(g_0, g_1, \dots, g_{m-1})$. Now it is just a matter of unfolding definitions to derive that $\alpha \otimes f = g$. Fix an arbitrary $n \in \mathbb{N}$, and let $k, i \in \mathbb{N}$ with $i < m$ be such that $n = km + i$. Then we have, using Lemma 4.13 twice, $(\alpha \otimes f)(n) = (\alpha_i \cdot \mathcal{S}^{ks_m+s_i}(f))(n) = a_{i,0} \cdot f(ks_m + s_i) + a_{i,1} \cdot f(ks_m + s_i + 1) + \dots + a_{i,k_i-1} \cdot f(ks_m + s_i + k_i - 1) + b_i = (\beta_i)_0 \cdot f(ks_m) + \dots + (\beta_i)_{s_m-1} \cdot f(ks_m + s_m - 1) + (\beta_i)_{s_m} = (\beta_i \cdot \mathcal{S}^{ks_m}(f))(k) = (\langle \beta_i \rangle \otimes f)(k) = g_i(k) = g(km + i) = g(n)$. \square

Lemma 4.18. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be spiralling, and let α be a tuple of non-constant weights. Then $\alpha \otimes f$ is spiralling.*

Proof. Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be spiralling, and let α be an r -tuple of non-constant weights. It is easy to see that $\lim_{n \rightarrow \infty} (\alpha \otimes f)(n) = \infty$.

To see that $g = \alpha \otimes f$ is ultimately periodic modulo every $m \geq 1$, i.e., $\forall m. \exists n_0, p. \forall n \geq n_0. g(n) \equiv g(n+p) \pmod{m}$, we first prove the claim for $r = 1$. So $\alpha = \langle \alpha \rangle$ is a tuple consisting of one non-constant weight. Without loss of generality, let $\alpha = \langle a_0/d, a_1/d, \dots, a_{k-1}/d, b/d \rangle$, for some integer $d > 0$. Let $g = \langle \alpha \rangle \otimes f$. By Lemma 4.13 we have $g(n) = \sum_{j=0}^{k-1} ((a_j/d) \cdot f(kn+j)) + b/d$, for all $n \in \mathbb{N}$. Fix an arbitrary integer $m > 0$. By f being spiralling there exist $n_0, p \in \mathbb{N}$ with $p > 0$ such that $f(n+p) \equiv f(n) \pmod{dm}$ for all $n \geq n_0$. This implies that $(a_j/d) \cdot f(k(n+p)+j) \equiv (a_j/d) \cdot f(kn+j) \pmod{m}$ for all $n \geq n_0$. Hence we obtain $g(n+p) \equiv g(n) \pmod{m}$.

For the case $r > 1$ we reason as follows. By Lemma 4.17 there exists an r -tuple β such that $g = \alpha \otimes f = \text{zip}_r(g_0, g_1, \dots, g_{m-1})$, where $g_i = \langle \beta_i \rangle \otimes f$ ($i \in \mathbb{N}_{<r}$). By the above argument (for $r = 1$), we have that g_i is ultimately periodic modulo every $m \geq 1$. We conclude by Lemma 4.16. \square

We will show that weighted products give rise to a characterisation, up to equivalence \equiv , of functions realised by FSTs on the set of spiralling sequences, see Theorem 4.25.

Lemma 4.19. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$, and α a tuple of weights. If $\alpha \otimes f$ is a natural weighted product, then we have $\langle f \rangle \geq \langle \alpha \otimes f \rangle$.*

Proof. Let $m = |\alpha|$, $k_i = |\alpha_i| - 1$, and $\alpha_i = \langle a_{i,0}/d_i, a_{i,1}/d_i, \dots, a_{i,k_i-1}/d_i, b_i/d_i \rangle$ with $a_{i,j}, d_i \in \mathbb{N}$ and $b_i \in \mathbb{Z}$ for all $i \in \mathbb{N}_{<m}$ and $j \in \mathbb{N}_{<k_i}$ (clearly weights can always be brought into this form).

We define $T = \langle Q, q_{m-1, k_{m-1}-1}^0, \delta, \lambda \rangle$ consisting of states $q_{i,j}^h$ for every $i \in \mathbb{N}_{<m}$, $j \in \mathbb{N}_{<k_i}$, and h such that $\min(0, b_i) \leq h < d_i$. The superscript h ($\min(0, b_i) \leq h < d_i$) in a state $q_{i,j}^h$ indicates the amount h/d_i of zeros that still has to be consumed/produced. The transition and output functions $\langle \delta, \lambda \rangle : Q \times \mathbf{2} \rightarrow Q \times \mathbf{2}^*$ of T are defined as follows; let $\text{id}_{\geq 0} : \mathbb{Z} \rightarrow \mathbb{N}$ be defined by $\text{id}_{\geq 0}(z) = z$ if $z \geq 0$ and $\text{id}_{\geq 0}(z) = 0$ otherwise.

$$\begin{aligned} \langle \delta, \lambda \rangle(q_{i,j}^h, 0) &= \langle q_{i,j}^{h'}, 0^e \rangle \quad \text{where } e = \lfloor \text{id}_{\geq 0}(h + a_{i,j})/d_i \rfloor, h' = h + a_{i,j} - ed_i \\ \langle \delta, \lambda \rangle(q_{i,j}^h, 1) &= \langle q_{i,j+1}^h, \varepsilon \rangle \quad (j < k_i - 1) \\ \langle \delta, \lambda \rangle(q_{i,k_i-1}^h, 1) &= \langle q_{i',0}^{h'}, 10^e \rangle \quad \text{where } e = \lfloor \text{id}_{\geq 0}(b_{i'})/d_{i'} \rfloor, h' = b_{i'} - ed_{i'}, \end{aligned}$$

where $i' = i + 1 \pmod{m}$.

We now show $\langle f \rangle \geq_T \langle \alpha \otimes f \rangle$. For $i \in \mathbb{N}_{<m}$, define $s_i = k_0 + k_1 + \dots + k_{i-1}$. We fix arbitrary $i \in \mathbb{N}_{<m}$ and $n \in \mathbb{N}$ with $n \equiv s_i \pmod{s_m}$. After reading the prefix $10^{f(0)} \dots 10^{f(n-1)}1$ of $\langle f \rangle$, T is in state $q_{i,0}^{b_i \bmod d_i}$. Let u denote the prefix of $\mathcal{S}(\langle \mathcal{S}^n(f) \rangle)$ of the form $u = 0^{f(n)}10^{f(n+1)} \dots 10^{f(n+k_i-1)}1$. We show that

$$\langle \delta, \lambda \rangle(q_{i,0}^{b_i - ed_i}, u) = \langle q_{i',0}^{b_{i'} - e'd_{i'}}, 0^{\alpha_i \cdot \mathcal{S}^n(f) - e} 10^{e'} \rangle \quad (2)$$

where $i' = i + 1 \bmod m$, $e = \lfloor \text{id}_{\geq 0}(b_i)/d_i \rfloor$, and $e' = \lfloor \text{id}_{\geq 0}(b_{i'})/d_{i'} \rfloor$. By the (implicit) assumption that $(\alpha \otimes f)(n) \in \mathbb{N}$ for all $n \in \mathbb{N}$, we have $\alpha_i \cdot \mathcal{S}^n(f) \in \mathbb{N}$. Also note that $\alpha_i \cdot \mathcal{S}^n(f) = (b_i + \sum_{j=0}^{k_i-1} a_{i,j} \cdot f(n+j))/d_i$.

For $j \in \mathbb{N}_{<k_i}$, we define $h_j, e_j \in \mathbb{N}$ with $\min(0, b_i) \leq h_j < d_i$ as follows:

$$\begin{aligned} h_0 &= b_i - ed_i \\ h_{j+1} &= h_j + a_{i,j} \cdot f(n+j) - e_j \cdot d_i & (j < k_i - 1) \\ e_j &= \left\lfloor \frac{h_j + a_{i,j} \cdot f(n+j)}{d_i} \right\rfloor. \end{aligned}$$

Then we have

$$\begin{aligned} \langle \delta, \lambda \rangle (q_{i,j}^{h_j}, 0^{f(n+j)} 1) &= \langle q_{i,j+1}^{h_{j+1}}, 0^{e_j} \rangle & \text{for } j < k_i - 1, \text{ and} \\ \langle \delta, \lambda \rangle (q_{i,k_i-1}^{h_{k_i-1}}, 0^{f(n+k_i-1)} 1) &= \langle q_{i',0}^{b_{i'}}, 0^{e_{k_i-1}} 1 \rangle & \text{where } i' = i + 1 \bmod m. \end{aligned}$$

Moreover, as we have

$$\sum_{j=0}^{k_i-1} e_j = \left\lfloor \frac{h_0 + \sum_{j=0}^{k_i-1} a_{i,j} \cdot f(n+j)}{d_i} \right\rfloor = \alpha \cdot \mathcal{S}^n(f) - e,$$

we conclude that (2) holds. \square

Definition 4.20. Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a function, and, for some $m > 0$, let α be an m -tuple of weights such that $\alpha \otimes f$ is a natural weighted product. Let \mathbf{p} and \mathbf{c} be m -tuples of finite words. We define the sequence $\Phi(f, \alpha, \mathbf{p}, \mathbf{c}) \in \mathbf{2}^{\mathbb{N}}$ by

$$\Phi(f, \alpha, \mathbf{p}, \mathbf{c}) = \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_j c_j^{\varphi(i,j)} \quad \text{where} \quad \varphi(i, j) = (\alpha \otimes f)(mi + j).$$

Note that $\langle f \rangle$ can also be cast into this notation: $\langle f \rangle = \Phi(f, \langle \langle 1, 0 \rangle \rangle, \langle 1 \rangle, \langle 0 \rangle)$. For the following lemma we recall that for a tuple $\mathbf{a} = \langle a_0, a_1, \dots, a_{k-1} \rangle$, we write \mathbf{a}' for the rotation $\langle a_1, \dots, a_{k-1}, a_0 \rangle$.

Lemma 4.21. Let $f, \alpha, \mathbf{p}, \mathbf{c}$ be as in Definition 4.20. We have $\Phi(f, \alpha, \mathbf{p}, \mathbf{c}) = p_0 c_0^{\alpha \cdot f} \cdot \Phi(\mathcal{S}^{|\alpha_0|-1}(f), \alpha', \mathbf{p}', \mathbf{c}')$.

Proof. This is a straightforward calculation. Let $|\alpha| = |\mathbf{p}| = |\mathbf{c}| = m$, and let $\varphi(i, j) = (\alpha \otimes f)(mi + j)$. For $i \in \mathbb{N}$ and $j \in \mathbb{N}_{<m}$ we define $\varphi'(i, j)$ and $\psi(i, j)$ by

$$\begin{aligned} \varphi'(i, j) &= \begin{cases} \varphi(i, j+1) & \text{if } j < m-1, \\ \varphi(i+1, 0) & \text{if } j = m-1, \end{cases} \\ \psi(i, j) &= (\alpha' \otimes \mathcal{S}^{|\alpha_0|-1}(f))(mi + j) \end{aligned}$$

First we note that $\varphi'(i, j) = \psi(i, j)$ for all $i \in \mathbb{N}$ and $j \in \mathbb{N}_{<m}$: Let $i \in \mathbb{N}$ and $j \in \mathbb{N}_{<m}$ be arbitrary. If $j < m-1$, then $\varphi'(i, j) = \varphi(i, j+1) = (\alpha \otimes$

$f)(mi + j + 1) = (\boldsymbol{\alpha}' \otimes \mathcal{S}^{|\alpha_0|-1}(f))(mi + j) = \psi(i, j)$ by Definition 4.11. For the case $j = m - 1$, we find $\varphi'(i, m - 1) = \varphi(i + 1, 0) = (\boldsymbol{\alpha} \otimes f)(m(i + 1)) = (\boldsymbol{\alpha}' \otimes \mathcal{S}^{|\alpha_0|-1}(f))(mi + m - 1) = \psi(i, m - 1)$, again by definition of weighted products. Then we have

$$\begin{aligned} \Phi(f, \boldsymbol{\alpha}, \mathbf{p}, \mathbf{c}) &= \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_j c_j^{\varphi(i,j)} \\ &= p_0 c_0^{\varphi(0,0)} \cdot \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_{j+1} c_{j+1}^{\varphi'(i,j)} \\ &= p_0 c_0^{\alpha_0 \cdot f} \cdot \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_{j+1} c_{j+1}^{\psi(i,j)} \\ &= p_0 c_0^{\alpha_0 \cdot f} \cdot \Phi(\mathcal{S}^{|\alpha_0|-1}(f), \boldsymbol{\alpha}', \mathbf{p}', \mathbf{c}') \end{aligned}$$

where addition in the subscripts is computed modulo m . \square

Lemma 4.22. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a spiralling function, and let $\sigma \in \mathbf{2}^{\mathbb{N}}$ be such that $\langle f \rangle \geq \sigma$ and $\sigma \notin \mathbf{0}$. Then there exist $n_0, m \in \mathbb{N}$, a word $w \in \mathbf{2}^*$, a tuple of weights $\boldsymbol{\alpha}$, and tuples of words \mathbf{p} and \mathbf{c} with $|\boldsymbol{\alpha}| = |\mathbf{p}| = |\mathbf{c}| = m > 0$ such that:*

- (i) $\sigma = w \cdot \Phi(\mathcal{S}^{n_0}(f), \boldsymbol{\alpha}, \mathbf{p}, \mathbf{c})$,
- (ii) $c_j^\omega \neq p_{j+1} c_{j+1}^\omega$ for every j with $0 \leq j < m - 1$, and $c_{m-1}^\omega \neq p_0 c_0^\omega$, and
- (iii) $c_j \neq \varepsilon$, and α_j is non-constant, for all $j \in \mathbb{N}_{< m}$.

Proof. By Lemma 4.5, there exist $n_0, m, a_j, z \in \mathbb{N}$ ($j \in \mathbb{N}_{< m}$), $w \in \mathbf{2}^*$, and $\mathbf{p}, \mathbf{c} \in (\mathbf{2}^*)^m$ such that $\sigma = w \cdot \Phi(\mathcal{S}^{n_0}(f), \boldsymbol{\alpha}, \mathbf{p}, \mathbf{c})$, where, for $j \in \mathbb{N}_{< m}$, α_j is defined by $\alpha_j = \langle \frac{1}{z}, -\frac{a_j}{z} \rangle$.

We now repeatedly alter the tuples $\boldsymbol{\alpha}, \mathbf{p}, \mathbf{c}$ until conditions (ii) and (iii) are fulfilled while condition (i) is upheld. For this we let $n_0 \in \mathbb{N}$, $w \in \mathbf{2}^*$, $m \in \mathbb{N}$, $\boldsymbol{\alpha}, \mathbf{p}$, and \mathbf{c} with $|\boldsymbol{\alpha}| = |\mathbf{p}| = |\mathbf{c}| = m$ be arbitrary such that (i) holds.

First note that, if $m = 1$ and condition (ii) or (iii) are violated, then $\sigma \in \mathbf{0}$, contradicting the assumption.

In case (ii) does not hold, consider the smallest $h \in \mathbb{N}_{< m}$ such that $c_h^\omega = p_{h+1} c_{h+1}^\omega$ where addition in the subscripts is computed modulo m . We assume $h < m - 1$; the case $h = m - 1$ proceeds analogously, using Lemma 4.21. For $i \in \mathbb{N}$ and $j \in \mathbb{N}_{< m}$, we let $\varphi(i, j) = (\boldsymbol{\alpha} \otimes \mathcal{S}^{n_0}(f))(mi + j)$. By Lemma 4.10 there are integers $a, b \geq 0$ and a word $x \in \mathbf{2}^*$ such that $c_h^{\varphi(i,h)} p_{h+1} c_{h+1}^{\varphi(i,h+1)} = p_{h+1} x^{a\varphi(i,h)+b\varphi(i,h+1)}$ (\star). We now define a tuple of weights $\boldsymbol{\beta}$, and tuples of words \mathbf{q} and \mathbf{d} , with $|\boldsymbol{\beta}| = |\mathbf{q}| = |\mathbf{d}| = m - 1$, as follows: Let $j \in \mathbb{N}_{< m-1}$. If $j < h$, then we define $q_j = p_j$, $d_j = c_j$, and $\beta_j = \alpha_j$. If $j > h$, we define $q_j = p_{j+1}$, $d_j = c_{j+1}$, and $\beta_j = \alpha_{j+1}$. If $j = h$, we define $q_j = p_h p_{h+1}$, $d_j = x$, and we let the weight β_j be defined as follows: For $\alpha_h = \langle r_0, r_1, \dots, r_{k-1}, e \rangle$ and $\alpha_{h+1} = \langle r'_0, r'_1, \dots, r'_{\ell-1}, e' \rangle$, let $\beta_h = \langle ar_0, ar_1, \dots, ar_{k-1}, br'_0, br'_1, \dots, br'_{\ell-1}, ae + be' \rangle$. By definition of \mathbf{q}, \mathbf{d} , and $\boldsymbol{\beta}$, to verify $\Phi(\mathcal{S}^{n_0}(f), \boldsymbol{\alpha}, \mathbf{p}, \mathbf{c}) = \Phi(\mathcal{S}^{n_0}(f), \boldsymbol{\beta}, \mathbf{q}, \mathbf{d})$, it suffices to check, for all $i \in \mathbb{N}$, $p_h c_h^{\varphi(i,h)} p_{h+1} c_{h+1}^{\varphi(i,h+1)} = q_h d_h^{\varphi(i,h)}$; here, for $i \in \mathbb{N}$

and $j \in \{0, \dots, m-2\}$, $\varphi'(i, j)$ is defined by $\varphi'(i, j) = (\beta \otimes \mathcal{S}^{n_0}(f))((m-1)i+j)$. Fix $i \in \mathbb{N}$. By definition of weighted products we have $\varphi(i, h) = \alpha_h \cdot \mathcal{S}^t(f)$ and $\varphi(i, h+1) = \alpha_{h+1} \cdot \mathcal{S}^{t+k}(f)$, for some $t \in \mathbb{N}$ (see Lemma 4.13). Also we have $\varphi'(i, h) = \beta_h \cdot \mathcal{S}^{t'}(f)$ for some $t' \in \mathbb{N}$. By definition of β we obtain $t' = t$. It follows that $\varphi'(i, h) = a \cdot \varphi(i, h) + b \cdot \varphi(i, h+1)$, and we conclude by (\star) . Repeat the procedure with $\beta, \mathbf{q}, \mathbf{d}$.

In case (iii) does not hold because of $c_h = \varepsilon$ for some $h \in \mathbb{N}_{< m}$, we change α_h into a constant weight $\langle 0, 0, \dots, 0 \rangle$ of length $|\alpha_h|$. This clearly does not change σ . Now consider the case that (iii) does not hold because α_h is constant. Then let $h \in \mathbb{N}_{< m}$ be minimal with this property. We assume $h < m-1$. The case $h = m-1$ proceeds analogously, again by Lemma 4.21. We now define a tuple of weights β , and tuples of words \mathbf{q} and \mathbf{d} , with $|\beta| = |\mathbf{q}| = |\mathbf{d}| = m-1$, as follows: For $j < h$, we define $\beta_j = \alpha_j$, $q_j = p_j$, and $d_j = c_j$. For $j > h$, we define $\beta_j = \alpha_{j+1}$, $q_j = p_{j+1}$, and $d_j = c_{j+1}$. For the case $j = h$, let $\alpha_h = \langle 0, 0, \dots, 0, e \rangle$ and $\alpha_{h+1} = \langle r_0, r_1, \dots, r_{\ell-1}, e' \rangle$. We define $q_j = p_h c_h^e p_{h+1}$, $d_j = c_{h+1}$, and $\beta_j = \langle 0, 0, \dots, 0, r_0, r_1, \dots, r_{\ell-1}, e' \rangle$ of length $|\alpha_h| + |\alpha_{h+1}| - 1$. The verification of $\Phi(\mathcal{S}^{n_0}(f), \alpha, \mathbf{p}, \mathbf{c}) = \Phi(\mathcal{S}^{n_0}(f), \beta, \mathbf{q}, \mathbf{d})$ is similar as above. Repeat the procedure with $\beta, \mathbf{q}, \mathbf{d}$. \square

For the proof of the following theorem we allow for a more liberal version of transducers. Instead of input letters along the edges we now allow input *words*. Transitions of these transducers are of the form $q \xrightarrow{\langle u, v \rangle^w} q'$. The idea is that this transition is taken if the automaton is in state q and the input word is of the form $uv\tau$. Then the automaton produces output w and switches to state q' , consuming u and continuing with $v\tau$.

Definition 4.23. An FST with *look-ahead* (FST *) is a tuple $T = \langle Q, q_0, D, \delta, \lambda \rangle$ where Q is a finite set of states, $q_0 \in Q$ is the initial state, the finite set $D \subseteq Q \times \mathbf{2}^+ \times \mathbf{2}^*$ is the input domain of the transition function $\delta : D \rightarrow Q$, and the output function $\lambda : D \rightarrow \mathbf{2}^*$, satisfying the following condition: for all $q \in Q$, $u_1, u_2, v_1, v_2 \in \mathbf{2}^*$ if $u_1 u_2$ is a prefix of $v_1 v_2$ and $\langle q, u_1, u_2 \rangle \in D$ and $\langle q, v_1, v_2 \rangle \in D$, then $u_1 = v_1$ and $u_2 = v_2$.

We lift δ to a partial function $\delta^* : Q \times \mathbf{2}^* \rightarrow Q$ by $\delta^*(q, \varepsilon) = q$ and

$$\delta^*(q, u_1 u_2 v) = \delta^*(\delta(q, u_1, u_2), u_2 v) \quad (\langle q, u_1, u_2 \rangle \in D, v \in \mathbf{2}^*).$$

Similarly, we lift λ to a partial function $\lambda^* : Q \times \mathbf{2}^\infty \rightarrow \mathbf{2}^\infty$ by $\lambda^*(q, \varepsilon) = \varepsilon$ and

$$\lambda^*(q, u_1 u_2 v) = \lambda(q, u_1, u_2) \cdot \lambda^*(\delta(q, u_1, u_2), u_2 v) \quad (\langle q, u_1, u_2 \rangle \in D, v \in \mathbf{2}^\infty).$$

The partial function $T : \mathbf{2}^\infty \rightarrow \mathbf{2}^\infty$ realised by the FST * T is defined by $T(u) = \lambda^*(q_0, u)$, for all $u \in \mathbf{2}^\infty$.

These transducers can be simulated by FSTs.

Lemma 4.24. For every FST * T there is an FST T' such that for all $u \in \mathbf{2}^\infty$, $T'(u) = T(u)$ whenever $T(u)$ is defined.

Proof. Let $T = \langle Q, q_0, D, \delta, \lambda \rangle$ be an FST^{*}. We define an FST $T' = \langle Q', q'_0, \delta', \lambda' \rangle$ as follows. Define $\ell = \max\{|u_1 u_2| \mid \exists q. \langle q, u_1, u_2 \rangle \in D\}$. We choose $Q' = Q \times \mathbf{2}^{\leq \ell}$ and $q'_0 = \langle q_0, \varepsilon \rangle$ where $\mathbf{2}^{\leq \ell}$ are all words of length at most ℓ . For every $q \in Q$ and $a \in \mathbf{2}$ we define:

$$\begin{aligned} \langle \delta', \lambda' \rangle(\langle q, v \rangle, a) &= \langle \langle q, va \rangle, \varepsilon \rangle & v \in \mathbf{2}^* \text{ with } |v| < \ell \\ \langle \delta', \lambda' \rangle(\langle q, u_1 u_2 v \rangle, a) &= \langle \langle q, u_2 va \rangle, \lambda(q, u_1, u_2) \rangle & u_1, u_2, v \in \mathbf{2}^*, \langle q, u_1, u_2 \rangle \in D \\ & & \text{with } |u_1 u_2 v| = \ell \end{aligned}$$

To make the FST T' complete, we let $\langle \delta', \lambda' \rangle(\langle q, v \rangle, a) = \langle \langle q, v \rangle, \varepsilon \rangle$ whenever $v \in \mathbf{2}^*$ and $|v| = \ell$ and neither of the two clauses above applies. It is straightforward to verify that for all $u \in \mathbf{2}^\infty$, $T'(u) = T(u)$ whenever $T(u)$ is defined. \square

Theorem 4.25. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be spiralling, and $\sigma \in \mathbf{2}^\mathbb{N}$. Then $\langle f \rangle \geq \sigma$ if and only if $\sigma \equiv \langle \alpha \otimes \mathcal{S}^{n_0}(f) \rangle$ for some integer $n_0 \geq 0$, and a tuple of weights α .*

Proof. One direction is by Lemma 4.19. For the other, assume $\langle f \rangle \geq \sigma$. If $\sigma \in \mathbf{0}$, then $\sigma \equiv \langle \langle \langle 0, 0 \rangle \rangle \otimes \mathcal{S}^0(f) \rangle = \langle n \mapsto 0 \rangle = 1^\omega$. Thus let $\sigma \notin \mathbf{0}$. By Lemma 4.22 there exist $n_1, m \in \mathbb{N}$, $w \in \mathbf{2}^*$, α, \mathbf{p} and \mathbf{c} with $|\alpha| = |\mathbf{p}| = |\mathbf{c}| = m > 0$ such that $\sigma = w \cdot \Phi(\mathcal{S}^{n_1}(f), \alpha, \mathbf{p}, \mathbf{c})$, and fulfilling the conditions (ii) and (iii) of Lemma 4.22. We abbreviate $g = \alpha \otimes \mathcal{S}^{n_1}(f)$. We will show that $\sigma \equiv \langle g \rangle$. By Lemma 4.18 we have that the function g is spiralling too.

By conditions (ii) and (iii), for every $j \in \mathbb{N}_{< m}$, there exists $t_j \in \mathbb{N}$ such that $c_j^\omega(t_j) \neq (p_{j+1} c_{j+1}^\omega)(t_j)$ (where addition is modulo m); let t_j be minimal with this property. For $j \in \mathbb{N}_{< m}$, let $\ell_j, \ell'_j \in \mathbb{N}$ be minimal such that $|c_j c_j^{\ell'_j}| > t_j$ and $|p_{j+1} c_{j+1}^{\ell'_j}| > t_j$. Then by minimality of t_j and ℓ_j , we obtain

- (i) $c_j c_j^{\ell_j - 1} \sqsubseteq p_{j+1} c_{j+1}^{\ell'_j} \tau$ for every $\tau \in \mathbf{2}^\mathbb{N}$, and
- (ii) $c_j c_j^{\ell_j} \not\sqsubseteq p_{j+1} c_{j+1}^{\ell'_j} \tau$ for every $\tau \in \mathbf{2}^\mathbb{N}$

(with again addition computed modulo m). From (i) we moreover obtain

- (iii) $c_j c_j^{\ell_j} \sqsubseteq c_j^n p_{j+1} c_{j+1}^{\ell'_j} \tau$ for every $n > 0$ and $\tau \in \mathbf{2}^\mathbb{N}$.

Next, we take a suffix σ' of σ such that every occurrence of a block $p_{j+1} c_{j+1}^{\varphi(i,j)}$ has as a prefix $p_{j+1} c_{j+1}^{\ell'_j}$. Let $n_2 \in \mathbb{N}$ be such that for $g' = \mathcal{S}^{n_2 \cdot m}(g)$ we have that $g'(n) > \max\{t_j \mid j \in \mathbb{N}_{< m}\}$ for all $n \in \mathbb{N}$; the existence of such an n_2 follows from g being spiralling. To prove $\sigma \equiv \langle g \rangle$ it suffices to show $\sigma' \equiv \langle g' \rangle$ where

$$\sigma' = \prod_{i=n_2}^{\infty} \prod_{j=0}^{m-1} p_j c_j^{\varphi(i,j)} = \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} p_j c_j^{\varphi'(i,j)} \quad \langle g' \rangle = \prod_{i=0}^{\infty} \prod_{j=0}^{m-1} 10^{\varphi'(i,j)}$$

where $\varphi(i, j) = g(mi + j)$, and $\varphi'(i, j) = g'(mi + j)$. Note that by the choice of n_2 , we have $\varphi'(i, j + 1) \geq \ell'_j$ for all $i \in \mathbb{N}$ and $j \in \mathbb{N}_{< m}$.

It is clear how to construct an FST that transduces $\langle g' \rangle$ to σ' , see Figure 3. For $\sigma' \geq \langle g' \rangle$, we define a FST^{*} $T = \langle Q, q_{m-1}, D, \delta, \lambda \rangle$, as follows, and apply

Lemma 4.24. Let $Q = \{q_j \mid j \in \mathbb{N}_{< m}\}$ and $D = \{\langle q_j, c_j, c_j^{\ell_j} \rangle \mid j \in \mathbb{N}_{< m}\} \cup \{\langle q_j, p_{j+1}, c_{j+1}^{\ell_j} \rangle \mid j \in \mathbb{N}_{< m}\}$, and define δ, λ by

$$\langle \delta, \lambda \rangle(q_j, c_j, c_j^{\ell_j}) = \langle q_j, 0 \rangle \quad \langle \delta, \lambda \rangle(q_j, p_{j+1}, c_{j+1}^{\ell_j}) = \langle q_{j+1}, 1 \rangle.$$

We now argue that $\sigma' \geq_T \langle g' \rangle$. This follows from the following facts:

- (a) $\lambda^*(q_j, p_{j+1} c^{\varphi'(i, j+1)} \tau) = 1 \cdot \lambda^*(q_{j+1}, c^{\varphi'(i, j+1)} \tau)$,
- (b) $\lambda^*(q_j, c_j^n p_{j+1} c^{\varphi'(i, j+1)} \tau) = 0 \cdot \lambda^*(q_{j+1}, c^{\varphi'(i, j+1)} \tau)$ for all $n > 0$, since by item (iii) we have $c_j c_j^{\ell_j} \sqsubseteq c_j^n p_{j+1} c^{\varphi'(i, j+1)} \tau$. \square

Lemma 4.26. *Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be spiralling, and $\sigma \notin \mathbf{0}$ with $\langle f \rangle \geq \sigma$. Then we have $\sigma \geq \langle \langle \beta \rangle \otimes \mathcal{S}^{n_0}(f) \rangle$ for some integer $n_0 \geq 0$, and a non-constant weight β .*

Proof. Let $\sigma \notin \mathbf{0}$ be a transduct of $\langle f \rangle$. By Theorem 4.25 we have $\sigma \equiv \langle \alpha \otimes \mathcal{S}^{n_0}(f) \rangle$ for some $n_0 \in \mathbb{N}$, and an m -tuple of weights α . By Lemma 4.14, there exists $i \in \mathbb{N}_{< m}$ such that $\alpha_i = \langle a_0, a_1, \dots, a_{k_i-1}, b_i \rangle$ is not constant. Let $k_j = |\alpha_j| - 1$ ($j \in \mathbb{N}_{< m}$), and for $\ell \in \mathbb{N}_{\leq m}$ let $s_\ell = \sum_{j=0}^{\ell-1} k_j$. We define a weight β of length $s_m + 1$ where, for $j \in \mathbb{N}_{< m}$ and $h \in \mathbb{N}_{< k_j}$, $\beta_{s_j+h} = 0$ if $j \neq i$, and $\beta_{s_j+h} = a_h$ if $j = i$, and $\beta_{s_m} = b_i$. Then we have $\sigma \geq \langle n \mapsto (\alpha \otimes \mathcal{S}^{n_0}(f))(nm + i) \rangle = \langle \langle \beta \rangle \otimes \mathcal{S}^{n_0}(f) \rangle$.

Theorem 4.27. *There is a non-atom, non-zero degree $[\sigma]$ that has no atom degree below it. Hence, non-zero transducts of σ start an infinite descending chain.*

Proof. We define the function $f : \mathbb{N} \rightarrow \mathbb{N}$ by $f(n) = 2^n$. We show that the degree $[\langle f \rangle]$ has no atom degree below it. Let $\sigma \notin \mathbf{0}$ with $\langle f \rangle \geq \sigma$. By Lemma 4.26 there is a non-constant weight $\beta = \langle a_0, a_1, \dots, a_{k-1}, b \rangle$ such that $\sigma \geq \langle g \rangle$ where $g = \langle \beta \rangle \otimes \mathcal{S}^{n_0}(f)$. Since $f(n) = 2^n$ it follows that

$$g(n) = b + \sum_{i=0}^{k-1} a_i 2^{n_0+nk+i} = b + 2^{nk} \sum_{i=0}^{k-1} a_i 2^{n_0+i} = (g(0) - b) \cdot 2^{nk} + b.$$

By Lemma 4.19 we have that $\langle g \rangle = \langle (g(0) - b) \cdot 2^{nk} + b \rangle \equiv \langle 2^{nk} \rangle$. Thus we have $\sigma \geq \langle 2^{nk} \rangle$. Also $\langle 2^{nk} \rangle \geq \langle 2^{2nk} \rangle$ holds by Lemma 4.2 (iv), and by Lemma 4.7 we conclude $\langle 2^{2nk} \rangle \not\geq \langle 2^{nk} \rangle$. \square

5 Squares

In [5] it is shown that $[\langle n \rangle]$ is an atom degree. One of the main questions of [5] is whether there exist other atom degrees. Here we show that also $[\langle n^2 \rangle]$ is an atom degree. The main tool is Theorem 4.25, the characterisation of transducts of spiralling sequences, which implies the following proposition.

Proposition 5.1. *Let $p(n)$ be a polynomial of degree k with non-negative integer coefficients, and let σ be a transduct of $\langle p(n) \rangle$ with $\sigma \notin \mathbf{0}$. Then $\sigma \geq \langle q(n) \rangle$ for some polynomial $q(n)$ of degree k with non-negative integer coefficients.*

Proof. By Lemma 4.26 it follows that $\sigma \geq \langle \langle \alpha \rangle \otimes \mathcal{S}^{n_0}(p) \rangle$ for some integer $n_0 \geq 0$, and a non-constant weight $\alpha = \langle a_0, \dots, a_{k-1}, b \rangle$. Let $h \in \mathbb{N}_{<k}$ be such that $a_h \neq 0$. Then we find $(\langle \alpha \rangle \otimes \mathcal{S}^{n_0}(p))(n) = b + \sum_{j=0}^{k-1} a_j \cdot p(n_0 + nk + j)$, which can easily be recognised to be a polynomial $q(n)$ of degree k . \square

Theorem 5.2. *The degree $\langle n^2 \rangle$ is an atom.*

Proof. Let $\sigma \in \mathbf{2}^{\mathbb{N}}$ be a transduct of $\langle n^2 \rangle$ such that σ is not ultimately periodic. By Proposition 5.1 there are integers $a > 0, b, c \geq 0$ such that $\sigma \geq \langle an^2 + bn + c \rangle$. We first assume $2a \geq b$. Abbreviate $f(n) = an^2 + (2a + b)n$. The roman numbers below refer to Lemma 4.2. We derive

$$\begin{aligned}
\langle an^2 + bn + c \rangle &\equiv \langle an^2 + bn \rangle && \text{by (iii)} \\
&\equiv \langle a(n+1)^2 + b(n+1) \rangle && \text{by (ii)} \\
&\equiv \langle f(n) \rangle && \text{by (iii)} \\
&\geq \langle b(f(2n)) + (2a-b)(f(2n+1)) \rangle && \text{by (v)} \\
&\equiv \langle 8a^2n^2 + 16a^2n \rangle \equiv \langle 8a^2(n+1)^2 \rangle && \text{by (iii)} \\
&\equiv \langle (n+1)^2 \rangle && \text{by (i)} \\
&\equiv \langle n^2 \rangle && \text{by (ii)}.
\end{aligned}$$

If $2a < b$, choose d such that $2ad \geq b$. Then we have $\langle an^2 + bn \rangle \geq \langle ad^2n^2 + bdn \rangle$ by (iv), and we reason as above for $\langle a'n^2 + b'n \rangle$ with $a' = ad^2$ and $b' = bd$.

This shows that every non-ultimately periodic transduct of $\langle n^2 \rangle$ can be transduced back to $\langle n^2 \rangle$. Hence, the degree of $\langle n^2 \rangle$ is an atom. \square

References

1. S. Cautis, F. Mignosi, J. Shallit, M. Wang, and S. Yazdani. Periodicity, morphisms, and matrices. *Theoretical Computer Science*, 295:107–121, 2003.
2. J. Endrullis, C. Grabmayer, and D. Hendriks. Data-Oblivious Stream Productivity. In *Logic for Programming, Artificial Intelligence, and Reasoning, 15th International Conference, LPAR 2008, Qatar*, pages 79–96, 2008.
3. J. Endrullis, C. Grabmayer, D. Hendriks, and H. Zantema. The Degree of Squares is an Atom. In *Proc. Conf. on Combinatorics on Words (WORDS 2015)*, Lecture Notes in Computer Science. Springer, 2015. To appear.
4. J. Endrullis, H. H. Hansen, D. Hendriks, A. Polonsky, and A. Silva. A Coinductive Framework for Infinitary Rewriting and Equational Reasoning. In *Proc. Conf. on Rewriting Techniques and Applications (RTA 2015)*, 2015. To appear.
5. J. Endrullis, D. Hendriks, and J.W. Klop. Degrees of Streams. *Journal of Integers*, 11B(A6):1–40, 2011. Proceedings of the Leiden Numeration Conference 2010.
6. J. Endrullis, D. Hendriks, and J.W. Klop. Highlights in Infinitary Rewriting and Lambda Calculus. *Theoretical Computer Science*, 464:48–71, 2012.
7. C. Grabmayer, J. Endrullis, D. Hendriks, J.W. Klop, and L. S. Moss. Automatic Sequences and Zip-Specifications. In *Proc. Symp. on Logic in Computer Science (LICS 2012)*, pages 335–344. IEEE Computer Society, 2012.
8. J. Sakarovitch. *Elements Of Automata Theory*. Cambridge, 2003.

9. J.I. Seiferas and R. McNaughton. Regularity-preserving relations. *Theoretical Computer Science*, 2(2):147–154, 1976.
10. J. Shallit. Open problems in automata theory: an idiosyncratic view. <https://cs.uwaterloo.ca/~shallit/Talks/bc4.pdf>, 2014. LMS Keynote Talk in Discrete Mathematics, BCTCS.
11. D. Siefkes. Undecidable extensions of monadic second order successor arithmetic. *Mathematical Logic Quarterly*, 17(1):385–394, 1971.