

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/144068>


Please be advised that this information was generated on 2019-02-19 and may be subject to change.

CORRECTION

# Correction: New Insights into Phosphorus Mobilisation from Sulphur-Rich Sediments: Time-Dependent Effects of Salinisation

The PLOS ONE Staff

There is an error in Fig. 4, “Oxygen ( $O_2$ ) concentration ( $mg\ L^{-1}$ ) profile per mm of both sediments (A and B), at the sediment-water interface (indicated by vertical dotted line) during aerobic and anaerobic conditions.” The title of the y-axis should read: Depth in sediment (mm). Please see the corrected Fig. 4 here.



**Fig 4. Oxygen ( $O_2$ ) concentration ( $mg\ L^{-1}$ ) profile per mm of both sediments (A and B), at the sediment-water interface (indicated by vertical dotted line) during aerobic and anaerobic conditions.**

doi:10.1371/journal.pone.0118387.g001



**OPEN ACCESS**

**Citation:** The PLOS ONE Staff (2015) Correction: New Insights into Phosphorus Mobilisation from Sulphur-Rich Sediments: Time-Dependent Effects of Salinisation. PLoS ONE 10(3): e0118387. doi:10.1371/journal.pone.0118387

**Published:** March 30, 2015

**Copyright:** © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Reference**

1. van Diggelen JMH, Lamers LPM, van Dijk G, Schaafsma MJ, Roelofs JGM, et al. (2014) New Insights into Phosphorus Mobilisation from Sulphur-Rich Sediments: Time-Dependent Effects of Salinisation. PLoS ONE 9(11): e1111106. doi:10.1371/journal.pone.0111106 PMID: 25369128