

RESEARCH ARTICLE | DECEMBER 01 2015

On the Colbeck–Renner theorem

Klaas Landsman



J. Math. Phys. 56, 122103 (2015)

<https://doi.org/10.1063/1.4936556>



Articles You May Be Interested In

A most compendious and facile quantum de Finetti theorem


J. Math. Phys. (January 2009)

Finite de Finetti theorem for conditional probability distributions describing physical theories

J. Math. Phys. (April 2009)

Renner-Teller nonadiabatic coupling terms: An *ab-initio* study of the HNH molecule

J. Chem. Phys. (February 2006)



Journal of Mathematical Physics

Young Researcher Award:
Recognizing the Outstanding Work
of Early Career Researchers

[Learn More!](#)

On the Colbeck–Renner theorem

Klaas Landsman^{a)}

Institute for Mathematics, Astrophysics, and Particle Physics, Faculty of Science, Radboud University Nijmegen, Nijmegen, Netherlands

(Received 28 September 2015; accepted 10 November 2015; published online 1 December 2015)

In three papers, Colbeck and Renner [Nat. Commun. **2**, 411 (2011); Phys. Rev. Lett. **108**, 150402 (2012); e-print [arXiv:1208.4123](https://arxiv.org/abs/1208.4123)] argued that “no alternative theory compatible with quantum theory and satisfying the freedom of choice assumption can give improved predictions.” We give a more precise version of the formulation and proof of this remarkable claim. Our proof broadly follows theirs, which relies on physically well motivated axioms, but to fill in some crucial details, certain technical assumptions have had to be added, whose physical status seems somewhat obscure. © 2015 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4936556>]

I. INTRODUCTION

The claim by Colbeck and Renner that “no alternative theory compatible with quantum theory and satisfying the freedom of choice assumption can give improved predictions”^{4–6} has attracted considerable attention (see, e.g., the review of Ref. 10), some of which has been rather critical.^{7–9} The aim of this paper is to give a watertight proof of their theorem, including a statement of precise, mathematically formulated assumptions.

Our proof broadly follows the dazzling reasoning of Colbeck and Renner, except that some of their theoretical physics style heuristic arguments have been replaced by rigorous mathematics. However, if this had been a routine exercise in mathematical physics, we would not have taken the effort. The point of our analysis is to show that additional assumptions are necessary to make the proof work, so that the theorem is weaker than it may appear to be at first sight: it does not show that quantum mechanics is complete, but that (informative) extensions are subject to (possibly undesirable) constraints.

Indeed, apart from three physically natural (and unavoidable) assumptions, namely, *Compatibility with Quantum Mechanics*, *Parameter Independence* (the latter being a well-known hidden variable version of the no-signaling axiom), and what we call *Product Extension*, we also need three assumptions that are satisfied by quantum mechanics itself but might seem somewhat unnatural if imposed on a hidden variable theory, viz., *Continuity of Probabilities*, *Unitary Invariance*, and what we call *Schmidt Extension*. We also replaced the original probabilistic setting in which almost everything (including even the quantum state) was treated as a random variable, by a more conventional hidden variable theory perspective (which circumvents some unnecessary controversies^{7,11}). Our approach differs significantly from interesting recent work of Leegwater,⁹ which has a similar goal.

II. NOTATION

A hidden variable theory \mathcal{T} underlying quantum mechanics yields probabilities

$$P(Z_1 = z_1, \dots, Z_n = z_n | \lambda) \equiv P(\vec{Z} = \vec{z} | \lambda)$$

for the possible outcomes $\vec{z} = (z_1, \dots, z_n)$ of a measurement of any family $\vec{Z} = (Z_1, \dots, Z_n)$ of commuting Hermitian operators on any Hilbert space H (here assumed to be finite dimensional for simplicity), given an arbitrary parameter $\lambda \in \Lambda$ (i.e., the “hidden variable”), where Λ is some Borel space.¹⁹ Being “classical” probabilities, these numbers are *a priori* only supposed to satisfy

^{a)}landsman@math.ru.nl

$0 \leq P(\vec{Z} = \vec{z}|\lambda) \leq 1$ and $\sum_{\vec{z}} P(\vec{Z} = \vec{z}|\lambda) = 1$, where the sum is over all possible outcomes. It will follow from the assumptions below that necessarily $z_i \in \sigma(Z_i)$ (i.e., the spectrum of Z_i) for each $i = 1, \dots, n$, in the sense that $P(\vec{Z} = \vec{z}|\lambda) = 0$ if this is not the case. Families of operators \vec{Z}_c (all defined on the same H) are indexed by some parameter $c \in C$, called the “setting” of the experiment.¹⁵

An important special case will be the bipartite setting $H = H_1 \otimes H_2$, where Alice and Bob measure Hermitian operators X and Y on H_1 and H_2 , respectively, so that $n = 2$, $Z_1 = X \otimes 1_{H_2}$, and $Z_2 = 1_{H_1} \otimes Y$. We then write $z_1 = x$, $z_2 = y$, and $c = (a, b)$, so that we typically look at expressions like $P(X_a = x, Y_b = y|\lambda)$. The other case of interest will simply be $n = 1$ with $Z_1 \equiv Z$, $z_1 \equiv z$; indeed, this will be the case in the statement of the theorem (the bipartite case playing a role only in the proof, though a crucial one!).

In this paper, quantum-mechanical states will just be unit vectors $\psi \in H$. The corresponding prediction for the above probabilities, i.e., the “Born rule,” is given by Ref. 13,

$$P_\psi(\vec{Z} = \vec{z}) = \langle \psi, E_{\vec{Z}}(\vec{z})\psi \rangle, \tag{2.1}$$

where $E_{\vec{Z}}(\vec{z}) = \prod_{i=1}^n E_{Z_i}(z_i)$, in which $E_{Z_i}(z_i)$ is the spectral projection on the eigenspace $H_{z_i} \subset H$ of Z_i (i.e., $Z_i\psi = z_i\psi$ iff $\psi \in H_{z_i}$). As detailed in Sec. III, \mathcal{T} assigns a probability measure μ_ψ on Λ to each state ψ . The following notation occurs throughout the paper:

$$P_\psi(\vec{Z} = \vec{z}|\lambda) = \alpha(\lambda), \tag{2.2}$$

with $\alpha : \Lambda \rightarrow [0, 1]$ an explicitly given measurable function (often constant). This means¹⁶

$$P(\vec{Z} = \vec{z}|\lambda) = \alpha(\lambda) \text{ for almost every } \lambda \text{ with respect to the measure } \mu_\psi. \tag{17}$$

Since this notation renders equalities like

$$P_\psi(\vec{Z} = \vec{z}|\lambda) = P_\varphi(\vec{Z}' = \vec{z}'|\lambda), \tag{2.3}$$

ambiguous (where ψ, φ are states in H), we explicitly define (2.3) as the double implication

$$P_\psi(\vec{Z} = \vec{z}|\lambda) = \alpha(\lambda) \Leftrightarrow P_\varphi(\vec{Z}' = \vec{z}'|\lambda) = \alpha(\lambda).$$

This notation also appears in our final pair of conventions: for $\varepsilon \rightarrow 0$, we write

$$\psi \stackrel{\varepsilon}{\approx} \varphi \Leftrightarrow (1 - \varepsilon) \leq |\langle \psi, \varphi \rangle| \leq 1, \tag{2.4}$$

$$P_\psi(\vec{Z} = \vec{z}|\lambda) \stackrel{\varepsilon}{\approx} P_\varphi(\vec{Z}' = \vec{z}'|\lambda) \Leftrightarrow P_\psi(\vec{Z} = \vec{z}|\lambda) = P_\varphi(\vec{Z}' = \vec{z}'|\lambda) + O(\sqrt{\varepsilon}). \tag{2.5}$$

III. ASSUMPTIONS

The assumptions in our reformulation of the Colbeck–Renner theorem are as follows.

CQ *Compatibility with Quantum Mechanics:* for any unit vector $\psi \in H$, the theory \mathcal{T} yields a state μ_ψ (i.e., a probability measure on Λ),¹⁴ such that (cf. (2.1))

$$\int_{\Lambda} d\mu_\psi(\lambda) P(\vec{Z} = \vec{z}|\lambda) = P_\psi(\vec{Z} = \vec{z}). \tag{3.1}$$

UI *Unitary Invariance:* for any unit vector $\psi \in H$ and unitary operator U on H ,²⁰

$$P_{U\psi}(\vec{Z} = \vec{z}|\lambda) = P_\psi(U^{-1}\vec{Z}U = \vec{z}|\lambda). \tag{3.2}$$

CP *Continuity of Probabilities:* if $\psi \stackrel{\varepsilon}{\approx} \varphi$, then $P_\psi(\vec{Z} = \vec{z}|\lambda) \stackrel{\varepsilon}{\approx} P_\varphi(\vec{Z} = \vec{z}|\lambda)$.

In the remaining three axioms, $H = H_1 \otimes H_2$, and X and Y are Hermitian operators on H_1 and H_2 , respectively (identified with operators $X \otimes 1_{H_2}$ and $1_{H_1} \otimes Y$ on H as appropriate).

PI *Parameter Independence*,¹⁸

$$\sum_{y \in \sigma(Y)} P(X = x, Y = y | \lambda) = P(X = x | \lambda), \tag{3.3}$$

$$\sum_{x \in \sigma(X)} P(X = x, Y = y | \lambda) = P(Y = y | \lambda). \tag{3.4}$$

PE *Product Extension*: for any pair of states $\psi_1 \in H_1, \psi_2 \in H_2$,

$$P_{\psi_1}(X = x | \lambda) = P_{\psi_1 \otimes \psi_2}(X = x | \lambda). \tag{3.5}$$

SE *Schmidt Extension*: if $e_i \in H_1$ ($i = 1, \dots, \dim(H)$) are eigenstates of X , then for arbitrary orthogonal states $u_i \in H_2$ and arbitrary coefficients $c_i > 0$ with $\sum_i c_i^2 = 1$,

$$P_{\sum_i c_i \cdot e_i}(X = x | \lambda) = P_{\sum_i c_i \cdot e_i \otimes u_i}(X = x | \lambda). \tag{3.6}$$

Comments. All assumptions are satisfied by quantum mechanics itself (seen as a “hidden” variable theory, with the state ψ as the “hidden” variable λ^1). In the broader context of hidden variable theories,³ **CQ** seems unavoidable in any such discussion, and also **PI** and **PE** have convincing physical plausibility. Unfortunately, the other assumptions are purely technical and have solely been invented to carry out certain steps in the proof.

In particular, although **UI**, **CP**, and **SE** represent the essence of quantum mechanics itself, these assumptions are far from self-evident for a hidden variable theory. Moreover, the former two are quite unsatisfactory in that they do not merely constrain the probabilities $P(\vec{Z} = \vec{z} | \lambda)$ of \mathcal{T} : they rather involve an interplay between these probabilities and the supports of the measures μ_ψ and $\mu_{U\psi}$. We challenge the reader to economize this!

IV. THEOREM AND PROOF

Our reformulation of the Colbeck–Renner theorem, then, is as follows.

Theorem 4.1. *If some hidden variable-theory \mathcal{T} satisfies **CQ**, **UI**, **CP**, **PI**, **PE**, and **SE**, then for any (finite-dimensional) Hilbert space H , state $\psi \in H$, and observable Z on H ,*

$$P_\psi(Z = z | \lambda) = P_\psi(Z = z). \tag{4.1}$$

We first assume (without loss of generality) that Z is nondegenerate as a Hermitian matrix in that it has distinct eigenvalues $(z_1, \dots, z_{\dim(H)})$. This assumption will be justified at the end of the proof. The proof consists of three steps as follows:

1. The theorem holds for $H = \mathbb{C}^2$ and any pair (Z, ψ) for which

$$P_\psi(Z = z_1) = P_\psi(Z = z_2) = 1/2. \tag{4.2}$$

This only requires assumptions **CQ**, **PI**, and **SE**.

2. The theorem holds for $H = \mathbb{C}^l, l < \infty$ arbitrary, and any pair (Z, ψ) for which

$$P_\psi(Z = z_1) = \dots = P_\psi(Z = z_l) = 1/l. \tag{4.3}$$

This is just a slight extension of step 1 and uses the same three assumptions.

3. The theorem holds in general. This requires all assumptions (as well as step 2).

The first step is mathematically straightforward but physically quite deep, depending on *chained Bell inequalities*,² and is due to Ref. 6 (we will give a slightly simplified proof below). The second step is easy. The third step, relying on the technique of *embezzlement*,¹² is highly nontrivial. This is the step that our analysis mainly attempts to clarify.

Step 1

Let $H = \mathbb{C}^2$, with basis (e_1, e_2) of eigenvectors of Z , so that $\psi \in \mathbb{C}^2$ may be written as

$$\psi = (e_1 + e_2)/\sqrt{2}. \tag{4.4}$$

Without loss of generality, we may assume that $z_1 = 1$ and $z_2 = -1$. We now relabel Z as Z_0 and extend it to a family of operators $(Z_k)_{k=0,1,\dots,2N-1}$ by fixing an integer $N > 1$, putting $\theta_k = k\pi/2N$, and defining

$$Z_k = [\theta_{k+\pi}] - [\theta_k], \tag{4.5}$$

where, for any angle $\theta \in [0, 2\pi]$, the operator $[\theta] = |\theta\rangle\langle\theta|$ is the orthogonal projection onto the subspace (ray) spanned by the unit vector

$$|\theta\rangle = \sin(\theta/2) \cdot e_1 + \cos(\theta/2) \cdot e_2. \tag{4.6}$$

In the corresponding bipartite setting, we have observables $X_k \equiv Z_k \otimes 1_2$ and $Y_k \equiv 1_2 \otimes Z_k$ on $\mathbb{C}^2 \otimes \mathbb{C}^2$, as well as a maximally correlated (Bell) state $\psi_{AB} \in \mathbb{C}^2 \otimes \mathbb{C}^2$, given by

$$\psi_{AB} = \frac{1}{\sqrt{2}}(e_1 \otimes e_1 + e_2 \otimes e_2). \tag{4.7}$$

Using assumptions **PI** and **SE**, we then have, for $i = 1, 2$, $z_1 = 1$ and $z_2 = -1$,

$$P_\psi(Z = z_i|\lambda) = P_{\psi_{AB}}(X_0 = z_i|\lambda). \tag{4.8}$$

The quantum-mechanical prediction is

$$P_{\psi_{AB}}(X_0 = 1) = P_{\psi_{AB}}(X_0 = -1) = \frac{1}{2}. \tag{4.9}$$

As in Ref. 6, our goal is to show that also

$$P_{\psi_{AB}}(X_0 = 1|\lambda) = P_{\psi_{AB}}(X_0 = -1|\lambda) = \frac{1}{2}. \tag{4.10}$$

To this effect, we introduce the combination of probabilities

$$I^{(N)}(\lambda) = P(X_0 = Y_{2N-1}|\lambda) + \sum_{a \in A_N, b \in B_N, |a-b|=1} P(X_a \neq Y_b|\lambda), \tag{4.11}$$

where $A_N = \{0, 2, \dots, 2N - 2\}$ and $B_N = \{1, 3, \dots, 2N - 1\}$. The inequality⁹

$$\begin{aligned} |P(X_a = x_i|\lambda) - P(Y_b = x_i|\lambda)| &= |P(X_a = x_i, Y_b = x_i|\lambda) + P(X_a = x_i, Y_b \neq x_i|\lambda) \\ &\quad - |P(X_a = x_i, Y_b = x_i|\lambda) - P(X_a \neq x_i, Y_b = x_i|\lambda)| \\ &= |P(X_a = x_i, Y_b \neq x_i|\lambda) - P(X_a \neq x_i, Y_b = x_i|\lambda)| \\ &\leq P(X_a = x_i, Y_b \neq x_i|\lambda) + P(X_a \neq x_i, Y_b = x_i|\lambda) \\ &= P(X_a \neq Y_b|\lambda), \end{aligned} \tag{4.12}$$

where $i = 1, 2$, and we used **PI**, implies a further inequality: since $X_{2N} = -X_0$,

$$\begin{aligned} |P(X_0 = 1|\lambda) - P(X_0 = -1|\lambda)| &= |P(X_0 = 1|\lambda) - P(X_{2N} = 1|\lambda)| \\ &\leq \sum_{a,b, |a-b|=1} |P(X_a = 1|\lambda) - P(Y_b = 1|\lambda)| \\ &\leq \sum_{a,b, |a-b|=1} P(X_a \neq Y_b|\lambda) \leq I^{(N)}(\lambda). \end{aligned}$$

Integrating this with respect to the measure $\mu_{\psi_{AB}}$ and using **CQ** give

$$\int_\Lambda d\mu_{\psi_{AB}}(\lambda) |P(X_0 = 1|\lambda) - P(X_0 = -1|\lambda)| \leq \int_\Lambda d\mu_{\psi_{AB}}(\lambda) I^{(N)}(\lambda) = I_{\psi_{AB}}^{(N)}. \tag{4.13}$$

A routine calculation shows that the quantum-mechanical prediction $I_{\psi_{AB}}^{(N)}$ is given by

$$I_{\psi_{AB}}^{(N)} = 2N \sin^2(\pi/4N), \tag{4.14}$$

so that

$$\lim_{N \rightarrow \infty} I_{\psi_{AB}}^{(N)} = 0. \tag{4.15}$$

Letting $N \rightarrow \infty$ in (4.13) therefore yields (4.10). From (4.8), we then obtain (4.2).

Step 2

Let $H = \mathbb{C}^l$ and let $(e_i)_{i=1}^l$ be an orthonormal basis of eigenvectors of Z , with corresponding eigenvalues z_i , and phase factors for the eigenvectors e_i such that $c_i > 0$ in the expansion

$$\psi = \sum_i c_i e_i. \quad (4.16)$$

Of course, $\sum_i c_i^2 = 1$. The case of interest will be $c_1 = \dots = c_l = 1/l$, but first, we merely assume that $c_1 = c_2$ (the same reasoning applies to any other pair), with $z_1 = 1$ and $z_2 = -1$ (which involves no loss of generality either and just simplifies the notation). The other coefficients c_i ($i > 2$) may or may not be equal to c_1 .

Generalizing (4.10), we will show that

$$P_\psi(Z = 1|\lambda) = P_\psi(Z = -1|\lambda). \quad (4.17)$$

This shows that if two Born probabilities defined by some quantum state ψ are equal, then the underlying hidden variable probabilities (conditioned on ψ) must be equal, too. Eq. (4.3) immediately follows from this result by taking all c_i to be equal.

Given step 1, the derivation of (4.17) is a piece of cake. We again pass to the bipartite setting, introducing two copies $H_A = H_B = \mathbb{C}^l$ of H , and define the correlated state

$$\psi_{AB} = \sum_i c_i \cdot e_i \otimes e_i \quad (4.18)$$

in $H_A \otimes H_B$. Eq. (4.8) again follows from assumptions **PI** and **SE**. Throughout the argument of step 1, we now replace each probability $P(X_a = x, Y_b = y|\lambda)$ by a corresponding probability $P^{(1)}(X_a = x, Y_b = y|\lambda)$, defined as the conditional probability

$$\begin{aligned} P^{(1)}(X_a = x, Y_b = y|\lambda) &= P(X_a = x, Y_b = y | |x| = |y| = 1, \lambda) \\ &= \frac{P(X_a = x, Y_b = y, |x| = |y| = 1|\lambda)}{P(|x| = |y| = 1|\lambda)}, \end{aligned} \quad (4.19)$$

for all λ for which $P(|x| = |y| = 1|\lambda) > 0$, whereas

$$P^{(1)}(X_a = x, Y_b = y|\lambda) = 0 \quad (4.20)$$

whenever $P(|x| = |y| = 1|\lambda) = 0$. The same argument then yields (4.13), with P replaced by $P^{(1)}$ but with the same right-hand side; see (Ref. 9, Section 3.2) for this calculation. As in step 1,

$$P_{\psi_{AB}}^{(1)}(X_0 = 1|\lambda) = P_{\psi_{AB}}^{(1)}(X_0 = -1|\lambda), \quad (4.21)$$

which implies that

$$P_{\psi_{AB}}(X_0 = 1|\lambda) = P_{\psi_{AB}}(X_0 = -1|\lambda), \quad (4.22)$$

either because both sides vanish (if $P(|x| = |y| = 1|\lambda) = 0$), or because (in the opposite case) the denominator $P(|x| = |y| = 1|\lambda)$ cancels from both sides of (4.21).

Combined with (4.8), Eq. (4.22) proves (4.17) and hence establishes step 2.

Step 3

We continue to use the notation established at the beginning of step 2, especially (4.16). As in step 1, we introduce two copies $H_A = H_B = \mathbb{C}^l$ of H , as well as two states

$$\psi_{AB} = \sum_i c_i \cdot e_i \otimes e_i \in H_A \otimes H_B, \quad (4.23)$$

$$\psi_{AB}''' = \kappa_n \otimes e'_1 \otimes e'_1 \otimes \psi_{AB} \in H_A''' \otimes H_B''', \quad (4.24)$$

where κ_n is given by (A4), $H''' = H'' \otimes H' \otimes H$, and we have notationally ignored the obvious permutations of factors in the tensor product.

For any $\varepsilon > 0$ and given coefficients c_i , pick c'_i in \mathbb{R}^+ such that $(c'_i)^2 \in \mathbb{Q}^+$ and

$$|c'_i - c_i| < \varepsilon / \dim(H), \tag{4.25}$$

which implies that, in the sense of (2.4), $\sum_i c'_i e_i \stackrel{\varepsilon/2}{\approx} \sum_i c_i e_i$. Suppose $c'_i = \sqrt{p_i/q_i}$, with $p_i, q_i \in \mathbb{N}$ and $\gcd(p_i, q_i) = 1$, and define

$$m_i = p_i \prod_{i' \neq i} q_{i'}. \tag{4.26}$$

Consequently, writing $q = 1/\sqrt{\sum_i m_i}$, the following quotient is independent of i :

$$\frac{c'_i}{\sqrt{m_i}} = q. \tag{4.27}$$

Given the integers m_i thus obtained, we define a unitary operator $U : H''' \rightarrow H'''$ by

$$U = \sum_{i=1}^l U^{(m_i)} \otimes P_i, \tag{4.28}$$

where $P_i : H \rightarrow H$ projects onto e_i (that is, $P_i = |e_i\rangle\langle e_i|$ in physics notation) and $U^{(m_i)}$ is defined in (A1). From this definition (with additional labels to denote the copies $U_A : H'''_A \rightarrow H'''_A$ and $U_B : H'''_B \rightarrow H'''_B$) and (A6) and (4.25), we then obtain the relations

$$1_{H'''_A} \otimes 1_{H'''_B}(\psi'''_{AB}) = \kappa_n \otimes \sum_{i=1}^l c_i \cdot \xi_{AA'}^{i1} \otimes \xi_{BB''}^{i1}, \tag{4.29}$$

$$U_A \otimes 1_{H'''_B}(\psi'''_{AB}) = \frac{1}{\sqrt{C(n)}} \sum_{i=1}^l \sum_{k=1}^n \frac{c_i}{\sqrt{k}} \cdot e''_{s_k} \otimes e''_k \otimes \xi_{AA'}^{ij_k} \otimes \xi_{BB''}^{i1}, \tag{4.30}$$

$$1_{H'''_A} \otimes U_B(\psi'''_{AB}) = \frac{1}{\sqrt{C(n)}} \sum_{i=1}^l \sum_{k=1}^n \frac{c_i}{\sqrt{k}} \cdot e''_k \otimes e''_{s_k} \otimes \xi_{AA'}^{i1} \otimes \xi_{BB''}^{ij_k}, \tag{4.31}$$

$$U_A \otimes U_B(\psi'''_{AB}) \stackrel{\varepsilon}{\approx} q \cdot \kappa_n \otimes \sum_{i=1}^l \sum_{j_i=1}^{m_i} \xi_{AA'}^{ij_i} \otimes \xi_{BB''}^{ij_i}. \tag{4.32}$$

Here,

$$\xi^{ij} = e_i \otimes e'_j \in H \otimes H', \tag{4.33}$$

with corresponding copies $\xi_{AA'}^{iji} \in H_A \otimes H'_A$ and $\xi_{BB''}^{iji} \in H_B \otimes H'_B$; the right-hand sides of (4.29)–(4.32) have been arranged so as to obtain vectors in the six-fold tensor product

$$H''_A \otimes H''_B \otimes H_A \otimes H'_A \otimes H_B \otimes H'_B.$$

The following (sub)steps are meant to replace (or justify) the core argument of Ref. 6. We repeatedly invoke the following lemma, whose proof just unfolds the notation (which incorporates the identification of X with $X \otimes 1_{H_2}$ and of Y with $1_{H_1} \otimes Y$ as appropriate).

*Lemma 4.2. Assume **PI** and **UI**. For any pair of unitary operators U_1 on H_1 and U_2 on H_2 , and any unit vector $\psi \in H_1 \otimes H_2$, one has*

$$P_{(U_1 \otimes 1_{H_2})\psi}(Y = y|\lambda) = P_\psi(Y = y|\lambda), \tag{4.34}$$

$$P_{(1_{H_1} \otimes U_2)\psi}(X = x|\lambda) = P_\psi(X = x|\lambda). \tag{4.35}$$

We now introduce some convenient notation. Since we assume that Z is nondegenerate, there is a bijective correspondence between its eigenvalues $Z = z_i$ and its eigenvectors e_i . Instead of $P(Z = z_i)$ dressed with whatever parameters ψ or λ , we may then write $P(e_i)$, where Z is understood, and analogously for the more complicated operators on tensor products of Hilbert space appearing below. We are now in a position to go ahead.

- From step 2, using the notation explained in (4.16),

$$P_{q \cdot \sum_{i=1}^l \sum_{j_i=1}^{m_i} \xi_{BB'}^{ij_i}}(\xi_{BB'}^{ij}|\lambda) = q^2. \tag{4.36}$$

- From (3.6) in PE and (4.36),

$$P_{q \cdot \sum_{i,j_i} \xi_{AA'}^{ij_i} \otimes \xi_{BB'}^{ij_i}}(\xi_{BB'}^{ij}|\lambda) = q^2. \tag{4.37}$$

- From (3.5) in SE and (4.37),

$$P_{q \cdot \kappa_n \otimes \sum_{i,j_i} \xi_{AA'}^{ij_i} \otimes \xi_{BB'}^{ij_i}}(\xi_{BB'}^{ij}|\lambda) = q^2. \tag{4.38}$$

- From (4.38), CP (whose notation we use), and (4.32),

$$P_{U_A \otimes U_B(\psi''_{AB})}(\xi_{BB'}^{ij}|\lambda) \stackrel{\varepsilon}{\approx} q^2. \tag{4.39}$$

- From (4.39) and Lemma 4.2, we have

$$P_{1_{H_A''} \otimes U_B(\psi''_{AB})}(\xi_{BB'}^{ij_i}|\lambda) \stackrel{\varepsilon}{\approx} q^2 (j_i = 1, \dots, m_i), \tag{4.40}$$

whereas the definition of the indices in question gives

$$P_{1_{H_A''} \otimes U_B(\psi''_{AB})}(\xi_{BB'}^{ij_i}|\lambda) \stackrel{\varepsilon}{\approx} 0 (j_i = m_i + 1, \dots, m), \tag{4.41}$$

where the number m (satisfying $m \geq m_i$ for all i) is introduced in the Appendix.

We now start a different argument, to be combined with (4.40)–(4.41) in due course.

- From PE, SE, and (4.16), with $e_A^i \in H_A$ denoting $e_i \in H$, we have

$$P_\psi(Z = z_i|\lambda) \equiv P_\psi(e_i|\lambda) = P_{\kappa_n \otimes \sum_i c_i \xi_{AA'}^{i1} \otimes \xi_{BB'}^{i1}}(e_A^i|\lambda). \tag{4.42}$$

- Using Lemma 4.2, (4.29), and (4.30),

$$P_{\kappa_n \otimes \sum_i c_i \xi_{AA'}^{i1} \otimes \xi_{BB'}^{i1}}(e_A^i|\lambda) = P_{1_{H_A''} \otimes U_B(\psi''_{AB})}(e_A^i|\lambda), \tag{4.43}$$

and hence,

$$P_\psi(Z = z_i|\lambda) = P_{1_{H_A''} \otimes U_B(\psi''_{AB})}(e_A^i|\lambda). \tag{4.44}$$

- From quantum mechanics, notably (2.1) and (4.31), for any $i' \neq i$, we have

$$P_{1_{H_A''} \otimes U_B(\psi''_{AB})}(e_A^{i'} \otimes \xi_{BB'}^{ij_i}) = 0. \tag{4.45}$$

- From CQ and (4.45), for any $i' \neq i$,

$$P_{1_{H_A''} \otimes U_B(\psi''_{AB})}(e_A^{i'} \xi_{BB'}^{ij_i}|\lambda) = 0. \tag{4.46}$$

- From PI,

$$P(e_A^{i'}|\lambda) = \sum_{i,j_i} P(e_A^{i'} \xi_{BB'}^{ij_i}|\lambda), \tag{4.47}$$

$$P(\xi_{BB'}^{ij_i}|\lambda) = \sum_{i'} P(e_A^{i'} \xi_{BB'}^{ij_i}|\lambda). \tag{4.48}$$

- From (4.46)–(4.48),

$$P_{1_{H_A''} \otimes U_B(\psi''_{AB})}(e_A^i|\lambda) = \sum_{j_i} P_{1_{H_A''} \otimes U_B(\psi''_{AB})}(\xi_{BB'}^{ij_i}|\lambda). \tag{4.49}$$

Finally, from (4.44), (4.49), (4.40), (4.41), and (4.27), we obtain

$$P_\psi(Z = z_i|\lambda) \stackrel{\varepsilon}{\approx} \sum_{j_i}^{m_i} q^2 = m_i \cdot q^2 = (c_i')^2. \tag{4.50}$$

Since $c_i > 0$, we have $c_i^2 = |c_i|^2$; using (4.25) and letting $\varepsilon \rightarrow 0$ then prove step 3,

$$P_\psi(Z = z_i|\lambda) = |c_i|^2 = P_\psi(Z = z_i). \quad (4.51)$$

Finally, we remove our standing assumption that the spectrum of Z be nondegenerate. In the degenerate case, one has

$$P_\psi(Z = z_i) = \sum_{j_i} P_\psi(e_{j_i}), \quad (4.52)$$

where the sum is over any orthonormal basis $(e_{j_i})_{j_i}$ of the eigenspace of z_i . Since each state e_{j_i} gives the same numerical outcome $Z = z_i$, probability theory gives for all λ ,

$$P(Z = z_i|\lambda) = \sum_{j_i} P(e_{j_i}|\lambda). \quad (4.53)$$

The nondegenerate case of the theorem (which distinguishes the states e_{j_i}) yields

$$P_\psi(e_{j_i}|\lambda) = P_\psi(e_{j_i}), \quad (4.54)$$

from which (4.1) follows once again,

$$P_\psi(Z = z_i|\lambda) = \sum_{j_i} P_\psi(e_{j_i}|\lambda) = \sum_{j_i} P_\psi(e_{j_i}) = P_\psi(Z = z_i).$$

ACKNOWLEDGMENTS

The author is indebted to Eric Cator, Roger Colbeck, Dennis Dieks, Dennis Hendrikx, and especially Gijs Leegwater for discussions, correspondence, and friendly corrections.

APPENDIX: EMBEZZLEMENT

We only treat the amazing technique of embezzlement for maximally entangled states (cf. Ref. 12 for the general case). We will deal with three Hilbert spaces, namely, $H = \mathbb{C}^l$, $H' = \mathbb{C}^m$, and $H'' = \mathbb{C}^n$ (where $n = m^N$ for some large N , see below), each with some fixed orthonormal basis $(e_i)_{i=1}^l$, $(e'_j)_{j=1}^m$, and $(e''_k)_{k=1}^n$, respectively. Given a further number $m_i \leq m$, we now list the nm basis vectors $e''_k \otimes e'_j$ of $H'' \otimes H'$ in two different orders.

1. $e''_1 \otimes e'_1, \dots, e''_n \otimes e'_1, e''_1 \otimes e'_2, \dots, e''_n \otimes e'_2, \dots, e''_1 \otimes e'_m, \dots, e''_n \otimes e'_m$.
2. $e''_1 \otimes e'_1, \dots, e''_1 \otimes e'_{m_i}, e''_2 \otimes e'_1, \dots, e''_2 \otimes e'_{m_i}, \dots, e''_n \otimes e'_1, \dots, e''_n \otimes e'_{m_i}, \dots$,
where the remaining vectors (i.e., those of the form $e''_k \otimes e'_j$ for $1 \leq k \leq n$ and $j > m_i$) are listed in some arbitrary order.

Define $U^{(m_i)} : H'' \otimes H' \rightarrow H'' \otimes H'$ as the unitary operator that maps the first list on the second. We will need the explicit expression

$$U^{(m_i)}(e''_k \otimes e'_1) = e''_{s_k} \otimes e'_{j_k}, \quad (A1)$$

where for given $k = 1, \dots, n$ the numbers $s_k^i = 1, \dots, n_i$ (where n_i is the smallest integer such that $n_i m_i \geq n$) and $j_k^i = 1, \dots, n_i$ are uniquely determined by the decomposition

$$k = (s_k^i - 1)m_i + j_k^i. \quad (A2)$$

We will actually work with two copies of $H'' \otimes H'$, called $H''_A \otimes H'_A$ and $H''_B \otimes H'_B$, with ensuing copies of $U_A^{(m_i)}$ and $U_B^{(m_i)}$ of $U^{(m_i)}$, and hence, leaving the isomorphism

$$H''_A \otimes H'_A \otimes H''_B \otimes H'_B \cong H''_A \otimes H''_B \otimes H'_A \otimes H'_B$$

implicit, we obtain a unitary operator

$$U_A^{(m_i)} \otimes U_B^{(m_i)} : H''_A \otimes H''_B \otimes H'_A \otimes H'_B \rightarrow H''_A \otimes H''_B \otimes H'_A \otimes H'_B. \quad (A3)$$

The point of all these is that the unit vector $\kappa_n \in H''_A \otimes H''_A$ defined by

$$\kappa_n = \frac{1}{\sqrt{C(n)}} \sum_{k=1}^n e''_k \otimes e''_k, \quad (\text{A4})$$

where $C(n) = \sum_{k=1}^n 1/k$, acts as a catalyst in producing the maximally entangled state

$$\varphi = \frac{1}{\sqrt{m_i}} \sum_{j=1}^{m_i} e'_j \otimes e'_j \quad (\text{A5})$$

in $H'_A \otimes H'_B$ from the uncorrelated state $e'_1 \otimes e'_1 \in H'_A \otimes H'_B$, in that for any $m_i \leq m$,

$$U_A^{(m_i)} \otimes U_B^{(m_i)} (\kappa_n \otimes e'_1 \otimes e'_1)^{\varepsilon/2} \approx \kappa_n \otimes \varphi, \quad (\text{A6})$$

where $\varepsilon = 1/N$ if $n = m^{2N}$. This follows straightforwardly from (A.3)–(A.5).

- ¹ Beltrametti, E. G. and Bugajski, S., “A classical extension of quantum mechanics,” *J. Phys. A* **28**, 3329–3343 (1995).
- ² Braunstein, S. L. and Caves, C. M., “Wringing out better Bell inequalities,” *Ann. Phys.* **202**, 22–56 (1990).
- ³ Bub, J., *Interpreting the Quantum World* (Cambridge University Press, 1997).
- ⁴ Colbeck, R. and Renner, R., “No extension of quantum theory can have improved predictive power,” *Nat. Commun.* **2**, 411 (2011).
- ⁵ Colbeck, R. and Renner, R., “Is a system’s wave function in one-to-one correspondence with its elements of reality?,” *Phys. Rev. Lett.* **108**, 150402 (2012).
- ⁶ Colbeck, R. and Renner, R., “The completeness of quantum theory for predicting measurement outcomes,” [arXiv:1208.4123](https://arxiv.org/abs/1208.4123).
- ⁷ Ghirardi, G. C. and Romano, R., “About possible extensions of quantum theory,” *Found. Phys.* **43**, 881–894 (2013).
- ⁸ Laudisa, F., “Against the no-go philosophy of quantum mechanics,” *Eur. J. Philos. Sci.* **4**, 1–17 (2014).
- ⁹ Leegwater, G., “An impossibility theorem for parameter independent hidden variable theories. Manuscript” (submitted).
- ¹⁰ Leifer, M., “Is the quantum state real? An extended review of ψ -ontology theorems,” *Quanta* **3**, 67–155 (2014).
- ¹¹ Seevinck, M. P. and Uffink, J., “Not throwing out the baby with the bathwater: Bell’s condition of local causality mathematically ‘sharp and clean,’” *Explan., Predict., Confirmation* **2**, 425–450 (2011).
- ¹² van Dam, W. and Hayden, P., “Universal entanglement transformations without communication,” *Phys. Rev. A* **67**, 060302(R) (2003).
- ¹³ von Neumann, J., *Mathematische Grundlagen der Quantenmechanik* (Springer, Berlin, 1932).
- ¹⁴ As the notation indicates, μ_ψ depends on ψ only and hence is independent of Z and z . From the point of view of \mathcal{T} , a quantum state ψ is a probability measure on Λ , so one might even write ψ for μ_ψ .
- ¹⁵ Colbeck and Renner look at the setting c as the value of some random variable C , but this is controversial;¹¹ for us, C is simply the set in which c takes values.
- ¹⁶ Colbeck and Renner treat ψ as a random variable and hence interpret $P_\psi(\vec{Z} = \vec{z}|\lambda)$ as a probability conditioned on knowing (that) ψ . We do not do so, yet our mathematical unfolding of (2.2) is similar.
- ¹⁷ In other words, there is a subset $\Lambda' \subset \Lambda$ such that $\mu_\psi(\Lambda') = 0$ and $P_\psi(\vec{Z} = \vec{z}|\lambda) = \alpha(\lambda)$ holds for any $\lambda \in \Lambda \setminus \Lambda'$. If Λ is finite, this simply means that the equality holds for any λ for which $\mu_\psi(\{\lambda\}) > 0$.
- ¹⁸ In words, this assumptions states that the probabilities for Alice’s measurement outcomes, given λ , are not only independent of Bob’s choice of his observable Y , but are even independent of his existence altogether, as they are given by the expression that \mathcal{T} yields for Alice’s experiment alone (and likewise for Bob). This slightly generalizes the usual Parameter Independence in the context of Bell’s Theorem.³ Note that in our form **PI** only makes sense because (2.1) and (3.1) imply that for $P_\psi(\vec{Z} = \vec{z}|\lambda)$ to be nonzero (in the sense of Sec. II) we must have $z_i \in \sigma(Z_i)$ for each i .
- ¹⁹ This generality, which is not a common feature of hidden variable theories (and as such is already a significant assumption), is necessary for the Colbeck–Renner argument to work.
- ²⁰ This assumption may be replaced by its main consequence, i.e., Lemma 4.2.