

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/151382>

Please be advised that this information was generated on 2018-07-22 and may be subject to change.

**Erratum: Measurement of the W Boson Production Charge Asymmetry
in $p\bar{p} \rightarrow W + X \rightarrow e\nu + X$ Events at $\sqrt{s} = 1.96$ TeV
[Phys. Rev. Lett. 112, 151803 (2014)]**

V. M. Abazov *et al.* ^{*}

(D0 Collaboration)

(Received 10 December 2014; published 29 January 2015)

DOI: 10.1103/PhysRevLett.114.049901

PACS numbers: 13.38.Be, 13.85.Qk, 14.60.Cd, 14.70.Fm, 99.10.Cd

The measurement of the W boson production charge asymmetry published in our recent Letter employed a correction K_{eff}^{\pm} to take into account the relative efficiency difference between electrons and positrons. Based on a recent study [1], we realized that the determination of K_{eff}^{\pm} was incorrect. Instead of taking the ratio of the positron efficiency to the electron efficiency, we took the ratio of the number of reconstructed positrons to the number of electrons. In addition, we had not taken into account the solenoid polarity when determining K_{eff}^{\pm} . These two problems have now been corrected.

The corrected W boson charge asymmetry values measured using the updated efficiency correction [1] are given in Table I. These revised measurements, together with those from the CDF Collaboration [2], are shown in Fig. 1. The asymmetry values have changed relative to those in the original publication by $< 2\%$, with smaller asymmetry values for $|y_W| < 0.6$ and larger asymmetry values for $0.8 < |y_W| < 2.4$, compared to the result in our Letter.




FIG. 1 (color online). Measured W boson charge asymmetry, after CP folding, compared to predictions and the CDF 1 fb^{-1} result. The points show the measured asymmetry, with the horizontal bars delineating the statistical uncertainty component and the vertical lines showing the total uncertainty. The central value and uncertainty from MC@NLO [3] using the NNPDF2.3 [4] PDF sets and the predictions from RESBOS [5] using the CTEQ6.6 [6] central PDF set and MC@NLO using the MSTW2008NLO [7] central PDF set are also shown. The inset focuses on the y_W region from 0 to 1.5.

TABLE I. CP -folded W boson charge asymmetry for data and predictions from MC@NLO using the NNPDF2.3 PDFs tabulated in percent (%) for each $|y_W|$ bin. The $\langle |y_W| \rangle$ is calculated as the cross section weighted average of y_W in each bin from RESBOS with PHOTOS [8]. For data, the first uncertainty is statistical and the second is systematic. The uncertainties on the prediction come from both the PDF uncertainties and α_s uncertainties.

Bin index	$ y_W $	$\langle y_W \rangle$	Data	Prediction
1	0.0–0.2	0.10	$1.39 \pm 0.17 \pm 0.12$	1.61 ± 0.19
2	0.2–0.4	0.30	$4.28 \pm 0.18 \pm 0.19$	5.06 ± 0.33
3	0.4–0.6	0.50	$7.28 \pm 0.19 \pm 0.27$	8.50 ± 0.41
4	0.6–0.8	0.70	$10.59 \pm 0.20 \pm 0.30$	12.05 ± 0.53
5	0.8–1.0	0.90	$14.45 \pm 0.21 \pm 0.32$	15.36 ± 0.66
6	1.0–1.2	1.10	$18.63 \pm 0.22 \pm 0.39$	18.86 ± 0.74
7	1.2–1.4	1.30	$22.50 \pm 0.24 \pm 0.44$	22.52 ± 0.80
8	1.4–1.6	1.50	$26.12 \pm 0.27 \pm 0.42$	26.30 ± 0.85
9	1.6–1.8	1.70	$30.06 \pm 0.31 \pm 0.44$	29.89 ± 0.92
10	1.8–2.0	1.90	$34.73 \pm 0.35 \pm 0.49$	34.04 ± 1.08
11	2.0–2.2	2.10	$40.59 \pm 0.40 \pm 0.54$	39.77 ± 1.31
12	2.2–2.4	2.29	$47.65 \pm 0.44 \pm 0.56$	47.73 ± 1.62
13	2.4–2.7	2.52	$59.04 \pm 0.46 \pm 0.60$	61.81 ± 1.74
14	2.7–3.2	2.81	$77.24 \pm 0.93 \pm 0.66$	78.05 ± 4.36

We thank the staffs at Fermilab and collaborating institutions and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); MON, NRC KI, and RFBR (Russia); CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); NRF (Korea); FOM (The Netherlands); STFC and the Royal Society (United Kingdom); MSMT and GACR (Czech Republic); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); and CAS and CNSF (China).

- [1] V. M. Abazov *et al.* (D0 Collaboration), [arXiv:1412.2862](https://arxiv.org/abs/1412.2862).
- [2] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **102**, 181801 (2009).
- [3] S. Frixione and B. R. Webber, *J. High Energy Phys.* 06 (2002) 029.
- [4] R. D. Ball *et al.*, *Nucl. Phys.* **B867**, 244 (2013).
- [5] C. Balazs and C. P. Yuan, *Phys. Rev. D* **56**, 5558 (1997).
- [6] J. Pumplin, D. R. Stump, J. Huston, H.-L. Lai, P. Nadolsky, and W.-K. Tung, *J. High Energy Phys.* 07 (2002) 012; D. Stump, J. Huston, J. Pumplin, W.-K. Tung, H.-L. Lai, S. Kuhlmann, and J. F. Owens, *J. High Energy Phys.* 10 (2003) 046.
- [7] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, *Eur. Phys. J. C* **63**, 189 (2009).
- [8] P. Golonka and Z. Wař, *Eur. Phys. J. C* **45**, 97 (2006).