Nitrate- and nitrite-dependent anaerobic oxidation of methane

Cornelia U. Welte^{*1,2}, Olivia Rasigraf^{1,3}, Annika Vaksmaa¹, Wouter Versantvoort¹, Arslan Arshad¹, Huub J.M. Op den Camp¹, Mike S.M. Jetten^{1,2,3}, Claudia Lüke¹, Joachim Reimann^{*1}

1 Department of Microbiology, Institute for Water and Wetland Research, Radboud University, Heyendaalseweg 135, 6525AJ Nijmegen, The Netherlands

2 Soehngen Institute of Anaerobic Microbiology, Heyendaalseweg 135, 6525AJ Nijmegen, The Netherlands

3 Netherlands Earth Systems Science Center, Heyendaalseweg 135, 6525AJ Nijmegen, The Netherlands

Running title: Nitrate/nitrite AOM

* Address correspondence to Cornelia U. Welte, c.welte@science.ru.nl and Joachim Reimann, j.reimann@science.ru.nl

Summary

Microbial methane oxidation is an important process to reduce the emission of the greenhouse gas methane. Anaerobic microorganisms couple the oxidation of methane to the reduction of sulfate, nitrate and nitrite, and possibly oxidized iron and manganese minerals. In this article, we review the recent finding of the intriguing nitrate- and nitrite-dependent anaerobic oxidation of methane (AOM). Nitrate-dependent AOM is catalyzed by anaerobic archaea belonging to the ANME-2d clade closely related to *Methanosarcina* methanogens. They were named ‘*Candidatus Methanoperedens nitroreducens*’ and use reverse methanogenesis with the key enzyme methyl-coenzyme M (methyl-CoM) reductase for methane activation. Their major end product is nitrite which can be taken up by nitrite-dependent methanotrophs. Nitrite-dependent AOM is performed by the NC10 bacterium ‘*Candidatus Methyloirablis oxyfera*’ that probably utilizes an intra-aerobic pathway through the dismutation of NO to N₂ and O₂ for aerobic methane activation by methane monooxygenase, yet being a strictly anaerobic microbe. Environmental distribution, physiological and biochemical aspects are discussed in this article as well as the cooperation of the microorganisms involved.

Key words: denitrification, methanotrophy, ANME, NC10, Mcr, archaea, anoxic, n-DAMO

Introduction

Methane is an important greenhouse gas contributing substantially to the greenhouse effect and global warming. It is produced in anoxic ecosystems by methanogenic archaea (Thauer *et al.*, 2008) as well as in the oxic ocean by phosphate-starved bacterioplankton through the demethylation of methyl-phosphonates produced by ammonia oxidizing archaea (Karl *et al.*, 2008; Metcalf *et al.*, 2012). The major part of the produced methane (50-80 %) is oxidized by aerobic and anaerobic methanotrophic microorganisms (Thauer *et al.*, 2008; Conrad, 2009).

In oxic environments, methane is consumed by aerobic bacterial methanotrophs with representatives from alpha- and gammaproteobacteria (Semrau *et al.*, 2010) as well as from the Verrucomicrobia (Op den Camp *et al.*, 2009; van Teeseling *et al.*, 2014). In anoxic environments, however, these bacterial groups are probably not involved in methane oxidation. In 2000, ANaerobic METHanotrophic (ANME) archaea closely related to methanogenic archaea were discovered and proven to be involved in methane oxidation in ecosystems devoid of oxygen (Boetius *et al.*, 2000; Raghoebarsing *et al.*, 2006; Orphan *et al.*, 2001; Haroon *et al.*, 2013). ANME archaea belonging to the clades ANME-1, ANME-2a/b, ANME-2c and ANME-3 have been associated with anaerobic oxidation of methane coupled to sulfate reduction in consortia with deltaproteobacterial sulfate reducers (Knittel and Boetius, 2009). Sulfate is abundant in marine ecosystems but generally quite low in freshwater systems (Reeburgh and Heggie, 1977), for which nitrate – and to some extent nitrite – are more relevant electron acceptors. In 2006, an archaeal-bacterial enrichment culture was obtained that coupled the oxidation of methane to denitrification (Raghoebarsing *et al.*, 2006). The archaeal partner couples anaerobic methane oxidation to the reduction of nitrate to nitrite (Raghoebarsing *et al.*, 2006; Haroon *et al.*, 2013;

Arshad *et al.*, 2015). The archaea belong to the ANME-2d clade and the investigated representative was named '*Candidatus Methanoperedens nitroreducens*'. Its metabolism involves a complete reverse methanogenesis pathway with methyl-CoM reductase as the methane activating enzyme. Cytoplasmic oxidation of methane to CO₂ is linked to an elaborate branched membrane-bound respiratory chain involving many unusual protein complexes (Haroon *et al.*, 2013; Arshad *et al.*, 2015) and a high number of *c*-type cytochromes (Haroon *et al.*, 2013; Arshad *et al.*, 2015; Kletzin *et al.*, 2015). The bacterial partner belongs to the NC10 clade and perform nitrite-dependent methane oxidation (Ettwig *et al.*, 2008). In the previously mentioned consortia, nitrite is provided by '*M. nitroreducens*' which benefits from the removal of its toxic end product. Subsequent metagenome sequencing and physiological experiments of the NC10 bacteria provided strong indication for an intra-aerobic methane oxidation metabolism in which nitrite is first reduced to nitric oxide (NO) which is then putatively dismutated to molecular nitrogen and oxygen (Ettwig *et al.*, 2010). The NC10 bacterium responsible for this process, '*Candidatus Methylomirabilis oxyfera*', uses particulate methane monooxygenase (pMMO) for methane oxidation via the aerobic pathway similar to aerobic methanotrophic bacteria.

Here we give an overview of the current knowledge on nitrate- and nitrite-dependent anaerobic methanotrophs that were first identified as denitrifying consortia: their ecology, distribution and the proposed underlying biochemical pathways.

The microbial ecology and diversity of nitrate- and nitrite-AOM

Nitrate and nitrite are common constituents at oxic/anoxic interfaces where ammonium, diffusing from deeper anoxic layers, is oxidized with oxygen diffusing from overlaying oxic zones (Brune

82 *et al.*, 2000). Here, methane can be used as electron donor to sustain populations of
83 nitrate/nitrite-dependent anaerobic methane oxidizing (N-AOM) microorganisms. In view of
84 expanding eutrophication around the globe (Galloway *et al.*, 2008), hypoxic zones with elevated
85 reactive nitrogen and methane concentrations as potential habitats for N-AOM microorganisms
86 will increase.

87
88 Although the potential of methane as electron donor for denitrification was recognized already
89 more than a decade ago (Sollo *et al.*, 1976; Mason, 1977; Panganiban *et al.*, 1979; Smith *et al.*,
90 1991; Thalasso *et al.*, 1997; Costa *et al.*, 2000; Eisentraeger *et al.*, 2001; Waki *et al.*, 2002; Islas-
91 Lima *et al.*, 2004) the first microbiological evidence for N-AOM came from an enrichment
92 culture originating from a highly eutrophic freshwater sediment of Twentekanaal in the
93 Netherlands (Raghoebarsing *et al.*, 2006). The enrichment culture oxidized methane to CO₂
94 while performing full denitrification to N₂. 16S rRNA gene sequencing enabled the design of
95 specific fluorescence *in situ* hybridization (FISH) probes, DARCH-872 and DARCH-641
96 (Raghoebarsing *et al.*, 2006; Schubert *et al.*, 2011; Ettwig *et al.*, 2016) for archaea and DBACT-
97 193 for bacteria (Raghoebarsing *et al.*, 2006), which were used for visualization of the culture
98 and are still commonly used to date (Figure 1). Molecular characterization revealed that this first
99 enrichment culture was mainly composed of anaerobic methane-oxidizing archaea (affiliated to
100 the GOM-ArcI/ANME-2D clade, Figure 2A) related to known methanogens and bacteria without
101 any cultured members and belonging to a new clade.

102
103 Today we know that the two dominant microorganisms, '*M. nitroreducens*' and '*M. oxyfera*', are
104 both capable of methane oxidation independent of each other (Ettwig *et al.*, 2008; Ettwig *et al.*,

2010; Haroon *et al.*, 2013). ‘*M. nitroreducens*’ and other ‘*Methanoperedens*’-like archaea couple AOM to nitrate reduction to nitrite (Equation 1) while ‘*M. oxyfera*’-like bacteria reduce nitrite to dinitrogen gas (Equation 2).

In contrast to sulfate-dependent AOM (Equation 3), it is evident that N-AOM (Equations 1 and 2) are highly exergonic metabolic processes that are not performed at the thermodynamic limit of life (Krüger *et al.*, 2003; Hallam *et al.*, 2004).

The bacterium performing nitrite-dependent AOM, ‘*M. oxyfera*’, belongs to the NC10 clade and was found to contain a phylogenetically distinct particulate methane monooxygenase which offered the possibility to use one of the encoding genes (*pmoA*) as a functional biomarker for their specific detection in the environment next to its 16S rRNA gene (Luesken *et al.*, 2011c; Han and Gu, 2013). The 16S rRNA genes of NC10 bacteria have been divided previously into four groups (A, B, C, D) (Ettwig *et al.*, 2009, Figure 3), where all to date known N-AOM performing ‘*M. oxyfera*’-like bacteria characterized from enrichment cultures fall into group A (Ettwig *et al.*, 2010; Haroon *et al.*, 2013; He *et al.*, 2015). Very recently, a high-quality draft genome of an NC10 bacterium belonging to group D could be re-constructed from an aquifer sediment metagenome (Hug *et al.*, 2016). Notably, this genome does not contain the *pmo* operon (encoding a methane monooxygenase) nor a quinol-dependent NO reductase so it is lacking the essential genes required for methane activation to methanol. However, it does encode enzymes

associated to methylotrophy such as a methanol dehydrogenase and enzymes involved in formaldehyde and formate oxidation (Hug *et al.*, 2016). The genome therefore suggests that this organism is a methylotroph, but does not share the intra-aerobic pathway of methane oxidation with the group A organisms. Group B and C are to date only represented by environmental sequences and no details are known about the physiology of these organisms. In addition to DNA-based biomarkers, it was shown that '*M. oxyfera*' contains high amounts of 10-methylhexadecanoic acid and a unique monounsaturated 10-methylhexadecenoic acid with a double bond at the $\Delta 7$ position, which comprised up to 10 % of the total membrane fatty acid profile (Kool *et al.*, 2012). These lipids have been successfully recovered from samples of a Dutch peatland harboring substantial amounts of '*M. oxyfera*' cells (Kool *et al.*, 2012; Zhu *et al.*, 2012) and therefore provide an alternative mode of detection. The environments where '*M. oxyfera*' biomarkers have been detected are shown in Table 1. The table shows a wide habitat spectrum including ecosystems from eutrophic to oligotrophic, freshwater to marine, and pristine to hydrocarbon contaminated. Although both DNA and lipids can be used to show the presence of '*M. oxyfera*' bacteria in various anoxic habitats, they cannot be used as a proxy for its contribution to methane oxidation activity. Thus, a variety of complementary methods based on RNA (usually used as cDNA) and proteins (direct shotgun proteomic sequencing) (Hanson and Madsen, 2015; Padilla *et al.*, 2016) have been applied and could demonstrate a link between observed methane disappearance and the presence of '*M. oxyfera*'. Furthermore, other features of '*M. oxyfera*' physiology could be used to detect its activity. '*M. oxyfera*' was shown to be an autotroph and assimilated CO₂ instead of methane actively into specific lipids and total biomass (Rasigraf *et al.*, 2014). So far, most stable isotope methods used labeled methane for the detection of active methanotrophs in field samples (Dumont and Murrell, 2005). A modified

method using ^{13}C -labeled CO_2 could potentially aid in the detection of labeled '*M. oxyfera*' DNA or RNA in environmental samples. A further method for environmental detection is based on ^{13}C fractionation of the environmental methane pool. It has been shown previously that methanotrophic bacteria fractionate methane leading to an enrichment of heavier methane in the remaining pool (Feisthauer *et al.*, 2011; Rasigraf *et al.*, 2012). Depending on the source of methanogenesis and the presence of methanotrophic bacteria a specific signature could be determined and linked to the extent of methanotrophy. So far, this tool of '*M. oxyfera*' activity detection has not been applied.

After nearly a decade, the archaeal partner detected in the initial co-culture of Raghoebarsing *et al.* (2006) was described in detail (Haroon *et al.*, 2013). In an enrichment fed with nitrate, methane and ammonium, a stable co-culture of anammox bacteria and above mentioned archaea was established. The archaeal counterpart was identified as the main organism responsible for coupling nitrate reduction to methane oxidation and was named '*Candidatus Methanoperedens nitroreducens*' (Haroon *et al.*, 2013). The successful enrichment of '*M. nitroreducens*' in this co-culture was due to the differential use of nitrate by '*M. nitroreducens*' vs. nitrite by '*M. oxyfera*'. Co-enrichment of both organisms in the culture described by Raghoebarsing *et al.* in 2006 was most likely due to feeding of both nitrate and nitrite. The differentiating effect of the used electron acceptor on enrichment of N-AOM archaea versus bacteria has been observed previously (Hu *et al.*, 2011). Detailed metagenomic analyses revealed that the genome of '*M. nitroreducens*' encoded pathways involved in the utilization of nitrate as electron acceptor (e.g. by the nitrate reductase subunit NarG) as well as reverse methanogenesis, with methyl-CoM reductase (McrA) as the key enzyme. Based on the available genomic data, CO_2 fixation in '*M.*

174 *nitroreducens*' may proceed via the acetyl-CoA pathway possibly leading to very depleted ¹³C
175 biomarkers. Carbon isotope measurements in archaeal lipids from the original culture described
176 by Raghoebarsing *et al.* (2006) indeed revealed strong depletion compared to methane,
177 indicating methane as carbon source for biomass. The possibility to use '*M. nitroreducens*' lipids
178 for its environmental detection has so far not been explored. '*M. nitroreducens*' forms a new
179 cluster within the ANME 16S rRNA gene phylogeny and was classified as the ANME-2d clade
180 (Figure 2A). Few aspects about the physiology of '*M. nitroreducens*' are known, and its
181 environmental detection has been limited to molecular methods based on 16S rRNA and *mcrA*
182 genes (Ding *et al.*, 2015). The overview of environmental distribution based on those biomarkers
183 is summarized in Table 1. The table shows that *Methanoperedens*-like archaea have been found
184 in a variety of environments including mostly freshwater and some marine habitats. Based on
185 16S rRNA gene classification, the ANME-2d clade is referred to as GOM Arc I in the Silva 16S
186 rRNA gene database (Figure 2A, Quast *et al.*, 2013), as the first sequences were found in
187 environmental samples from the Gulf of Mexico (GOM). To date the GOM Arc I/ANME-2d
188 group consists of 96 high quality sequences, which split into three defined clusters A, B and C
189 (Figure 2B). The 16S rRNA sequences of the two known genomes from enrichment cultures
190 (Haroon *et al.*, 2013; Arshad *et al.*, 2015) cluster into group A, which is the largest and most
191 uniform group. With few exceptions of sequences found in marine and brackish environments,
192 this group consists of sequences detected in freshwater environments such as aquifers, lakes and
193 rivers (Li *et al.*, 2012; Flynn *et al.*, 2013). Group B and C have no cultured representatives so far
194 and consist exclusively of environmental sequences. The sequences of group B and C have been
195 found in extreme environments such as marine and terrestrial mud volcanoes, marine sediment
196 and hydrothermal vents (Inagaki *et al.*, 2006; Pachiadaki *et al.*, 2011; Yang *et al.*, 2012).

Metabolic cooperation and competition of N-AOM microorganisms

Physiological studies showed that nitrite was the main product of nitrate reduction by N-AOM archaea (Haroon *et al.*, 2013; Zhu, 2014). In high concentrations, nitrite becomes toxic and must be removed. The N-AOM archaea encode a membrane-bound nitrite reductase which could convert some of the nitrite into ammonia (Arshad *et al.*, 2015). The presence of nitrate, nitrite and ammonium creates a basis for metabolic co-operation with nitrite and ammonium scavenging organisms (Figure 4). The first described co-culture of '*M. nitroreducens*' contained anaerobic ammonium oxidizing (anammox) bacteria which use nitrite for respiration (Haroon *et al.*, 2013). The original N-AOM culture described in 2006 also contained archaea closely related to '*M. nitroreducens*'. The 16S rRNA gene sequence of the 2006 enrichment and the '*Methanoperedens*' sp. BLZ1 are 99.2 % identical and cannot be resolved from each other in the phylogenetic tree in Figure 2B. They were enriched together with '*M. oxyfera*' bacteria, the latter being known by now to use nitrite as electron acceptor (Raghoebarsing *et al.*, 2006). Thus, anammox and '*M. oxyfera*'-like bacteria are most likely common metabolic partners of N-AOM archaea as both methane and ammonium derived from anaerobic food chains are often present at oxic/anoxic interfaces. The co-occurrence of anammox and '*M. oxyfera*'-like bacteria with '*M. nitroreducens*' would lead to a competition of the two for available nitrite. Previous studies have shown that anammox bacteria can be co-enriched and form a stable co-culture with '*M. oxyfera*' in a bioreactor system upon gradual increase of ammonium concentrations in the influent medium (Luesken *et al.*, 2011b; Ding *et al.*, 2014; Zhu *et al.*, 2011). In contrast, Hu *et al.* (2015) found that anammox bacteria successfully outcompeted '*M. oxyfera*' in bioreactor systems fed with ammonium and methane and amended either with nitrate or nitrite. Environmental

220 molecular studies have shown that both co-occur in anoxic environments (Wang *et al.*, 2012;
221 Shen *et al.*, 2014c; Shen *et al.*, 2015). It is likely that different substrate affinities of different '*M.*
222 *oxyfera*' and anammox species/strains would determine the success of competition as well as
223 tolerance to harmful nitrogen oxide species (e.g. NO and NH₂OH). In natural settings, the
224 interactions will become more complex due to the activity of nitrifying bacteria and archaea.
225 Ammonium concentrations shape the community composition of nitrifying organisms with
226 ammonia oxidizing bacteria (AOB) typically dominating at higher concentrations and archaea
227 (AOA) mostly occurring at lower concentrations (Yan *et al.*, 2012). Moreover, higher
228 concentrations of nitrite would lead to the presence of nitrite oxidizing bacteria (NOB), with
229 *Nitrobacter/Nitrococcus* dominating at higher and *Nitrospira/Nitrospina* at lower nitrite
230 concentrations (Nowka *et al.*, 2015). Recently, a complete nitrification process has been
231 described in a *Nitrospira*-like organism (comammox), which seems to predominate at very low
232 substrate concentrations and thus become competitive with the "classical" two stage process
233 (Daims *et al.*, 2015; van Kessel *et al.*, 2015). The comammox process produces nitrate and
234 bypasses the release of nitrite and could directly provide substrate to '*M. nitroreducens*'. This
235 scenario seems likely for oligotrophic environments with overall low concentration of
236 nitrogenous compounds and high methane (e.g. drinking water wells, Gülay *et al.*, 2016; Palomo
237 *et al.*, 2016; Pinto *et al.*, 2016). Thus, the co-occurrence of anammox and comammox bacteria
238 with N-AOM organisms might be a common scenario. The presence of other electron donors in
239 the environment (e.g. organic carbon, reduced iron and sulfur species) would potentially
240 intensify the competition for nitrate (and nitrite) in the form of denitrification and dissimilatory
241 nitrate/nitrite reduction to ammonium. Thus, various primary and secondary factors can
242 determine the outcome of each particular competition.

Biochemistry and metabolism of N-AOM microorganisms

Methane is quite inert due to the absence of functional groups and breaking the first C-H bond poses an energetic barrier of $\Delta H_{298} = 439$ kJ/mol (Blanksby and Ellison, 2003). Therefore, oxidation of methane requires it to be activated first. Until now there are only two biological processes known to activate methane, incorporation of oxygen by methane monooxygenases utilized by aerobic methanotrophic bacteria and formation of methyl-CoM employing methyl-CoM reductase in a reverse manner utilized by anaerobic methanotrophic archaea (Figure 5).

M. oxyfera is so far unique in its ability to couple anaerobic methane oxidation to nitrite reduction. The biochemistry and general metabolism of *M. oxyfera* is not yet well explored. The current metabolic model of nitrite-dependent methane conversion is therefore largely inferred from the genome and based on homology. As there are no organisms sharing this metabolism, a global comparative analysis is not available. Most of the metabolic modules, however, are shared with canonical methanotrophs and denitrifiers, which allowed a metabolic prediction for *M. oxyfera* (Ettwig *et al.*, 2010). Although *M. oxyfera* was cultivated under strictly anaerobic conditions and displayed severe oxygen intolerance (Luesken *et al.*, 2012), its genome encodes the complete aerobic methane oxidation pathway and is postulated to employ an intra-aerobic pathway for the degradation of methane. Candidate enzymes for oxygen generation are two nitric oxide reductase-like proteins that were hypothesized to disproportionate two molecules of NO into N₂ and O₂ (Ettwig *et al.*, 2012). This dismutation reaction is highly exergonic ($\Delta G^0 = -173$ kJ/mol O₂) but due to complex bond reorganizations is expected to present the rate limiting step in *M. oxyfera*'s energy metabolism. Activation of methane by

either NO or N₂O directly is thermodynamically feasible, but incompatible with the measured substrate stoichiometries (Ettwig *et al.*, 2010; Reimann *et al.*, 2015). As in most aerobic methanotrophs '*M. oxyfera*' employs a particulate methane monooxygenase for the activation of methane into methanol (Figure 5). Methane has a high octanol-water partition coefficient and accumulates in the hydrophobic membrane core in a ~12:1 ratio (Hansch *et al.*, 1995), making it available in high effective concentrations to the particulate methane monooxygenase (pMMO). Amino acid sequence comparison of the PmoA, PmoB and PmoC subunits from '*M. oxyfera*' to canonical pMMOs suggested a similar overall architecture and conserved function for this enzyme. Alternative reaction mechanisms involving NO, N₂O or NO₂ in methane activation are difficult to justify in this context (Ettwig *et al.*, 2010; Reimann *et al.*, 2015).

'*M. oxyfera*' has three PQQ-dependent methanol dehydrogenases (MDH) at its disposal for the conversion of methanol to formaldehyde. One gene cluster encodes for a calcium-dependent MDH, which harbors all accessory genes, next to the canonical alpha (MxaF) and beta (MxaI) subunits. The two additional MDHs belong to the recently described class of lanthanide-dependent XoxF MDHs (Keltjens *et al.*, 2014). The XoxF methanol dehydrogenase from the methanotroph *Methylophilum fumariolicum* SolV was isolated as a homodimer and shown to incorporate the rare earth element cerium believed to confer superior catalytic activity (Pol *et al.*, 2014). Purification of the dominant MDH from '*M. oxyfera*' resolved a unique combination of the XoxF1 large subunit and the MxaI small subunit forming a heterodimeric complex ($\alpha_2\beta_2$) (Wu *et al.*, 2015). It remains to be shown whether a rare earth element is indeed bound in the enzyme. Although PQQ-biosynthesis genes were mostly absent in the '*M. oxyfera*' genome (Wu *et al.*, 2011) spectroscopy on the purified MDH clearly confirmed the presence of the PQQ

cofactor (Wu *et al.*, 2015). It thus appears that formation of the holoprotein requires PQQ acquisition from the environment as has been previously observed for glucose dehydrogenase in PQQ-deficient enteric bacteria (Hommes *et al.*, 1991). Dependence on other microorganisms for the production of this crucial cofactor could possibly explain why '*M. oxyfera*' has thus far not been obtained as a pure culture. Formaldehyde is further oxidized to formate via two possible pathways, a highly expressed methanopterin (H₄MPT) route likely used for energy conservation and a lowly expressed folate (H₄F) route, where folate or methanopterin function as C1 carriers for biosynthetic purposes (Reimann *et al.*, 2015). Three enzymes are available to '*M. oxyfera*' for the oxidation of formate to CO₂, a highly expressed formyl-MFR dehydrogenase and two minor expressed NAD(P)⁺-dependent formate dehydrogenases (FDH) in which the extended N-terminal parts show homology with bacterial complex I subunits NuoG for both FdhA subunits and NuoE for the FdhB2 subunit. These alternative FDHs might provide extra reducing equivalents in the form of NADH (Reimann *et al.*, 2015).

Although two nitrate reductases, NarGHI and NapAB, are present in the genome of '*M. oxyfera*' low transcription and translation levels suggest that neither of the two systems is highly active. Nitrite reduction to NO is catalyzed by cytochrome *cd*₁-type nitrite reductase (NirS), the only nitrite reductase present in the genome. Produced NO is dismutated into N₂ and O₂ by two putative NO dismutases (NOD). The genome does not code for a recognizable N₂O reductase and N₂O was only measured in trace amounts in methane-driven nitrite reduction experiments of '*M. oxyfera*' enrichments. The proposed NODs are homologous to the quinol-dependent NO reductases, but display amino acid alterations in the catalytic site, the quinol-binding site and the proposed proton channel. These changes hamper electron and proton entry into the active site

and could facilitate the disproportionation of NO to N₂ and O₂ rather than its reduction to N₂O (Ettwig *et al.*, 2012). In addition to these two putative NO dismutases three nitric oxide reductases (NOR) are encoded in the '*M. oxyfera*' genome, one canonical qNOR, one gNOR and one sNOR (Hemp and Gennis, 2008). The product of these enzymes, N₂O, was only detected in trace amounts under standard conditions. It therefore remains an open question what the redundancy in NO reductases offers to '*M. oxyfera*'. NORs might be present to quickly respond to external nitrosative stress and to ensure that concentrations of the metabolic intermediate NO are kept below toxic levels. The NORs may also play a role in oxygen respiration. Since only three of the four O₂ molecules produced from NO disproportionation are consumed during methane activation the remaining O₂ molecule might be reduced to water, a side reactivity that has been demonstrated for both c- and qNORs, with rates that could match the overall metabolic rates of methane conversion (Reimann *et al.*, 2015).

The only known microorganisms capable of oxidizing methane with nitrate as electron acceptor are '*M. nitroreducens*' and *Methanoperedens*-like archaea (Raghoebarsing *et al.*, 2006; Haroon *et al.*, 2013). In contrast to '*M. oxyfera*' they do not use oxygen for methane activation but instead utilizes the reverse reaction of methyl-CoM reductase (Figure 5, Krüger *et al.*, 2003; Hallam *et al.*, 2004; Scheller *et al.*, 2010; Haroon *et al.*, 2013). Metabolic reconstructions from environmental genomes (Haroon *et al.*, 2013; Arshad *et al.*, 2015) suggested that '*M. nitroreducens*' oxidizes methane via reverse methanogenesis to CO₂. One of the key questions is how electrons from methane oxidation are transferred to the final electron acceptor nitrate. During reverse methanogenesis, electrons are transferred to yield cofactor F₄₂₀H₂, reduced ferredoxin and the thiol cofactors coenzyme M (CoM-SH) and coenzyme B (CoB-SH). F₄₂₀H₂

335 and reduced ferredoxin can be re-oxidized by a canonical $F_{420}H_2$ dehydrogenase (Fqo) and an
336 Ech hydrogenase, respectively (Welte and Deppenmeier, 2014). CoM-SH and CoB-SH are either
337 oxidized via the reverse reaction of the membrane-bound heterodisulfide reductase (HdrDE) or
338 via the cytoplasmic heterodisulfide reductase (HdrABC). The latter enzyme has been
339 exemplified to perform electron bifurcation in methanogens (Costa *et al.*, 2010; Kaster *et al.*,
340 2011) and due to thermodynamic limitations provided by the reversal of methanogenesis in '*M.*
341 *nitroreducens*' would have to act as an electron confurcation enzyme here (Arshad *et al.*, 2015).
342 In methanogens, both HdrDE and Fqo interact with methanophenazine, a membrane-integral
343 electron carrier; in '*M. nitroreducens*', however, methanophenzine could not be detected but
344 instead a menaquinone biosynthesis pathway was encoded and expressed (Arshad *et al.*, 2015). It
345 is therefore likely that HdrDE and Fqo interact with menaquinones in this organism. The genome
346 also encodes a nitrate reductase subunit harboring the active site (NarG) for nitrate reduction to
347 nitrite. Electron transport to nitrate reductase seems to happen via a Rieske-cytochrome *b*
348 complex. The gene cluster encoding the Rieske-cytochrome *b* complex contains additional genes
349 for cytochrome *c* proteins whose function is unclear but may be connected to the electron
350 transport to nitrate reductase. The nitrate reductase gene cluster shows a highly unusual
351 composition that to our knowledge has not been observed in other prokaryotes (Arshad *et al.*,
352 2015). As in other archaea, NarG contains a signal peptide for the translocation of NarGH into
353 the pseudoperiplasm and nitrate may therefore be reduced non-cytoplasmically (Yoshimatsu *et*
354 *al.*, 2000; Martinez-Espinosa *et al.*, 2007; de Vries *et al.*, 2010). All other nitrate-reducing
355 archaea studied to date harbor NarM as a membrane anchor for the soluble NarGH complex (de
356 Vries *et al.*, 2010) that is absent in '*M. nitroreducens*'. Instead, a NapH like membrane anchor
357 together with membrane-integral heme-copper oxidase subunits was encoded in the same gene

cluster, along with a cytochrome *b* protein that is also found in the nitrate reductase complex of halophilic archaea (Martinez-Espinosa *et al.*, 2007). A small part of the formed nitrite can be further reduced to ammonium by a NrfAH type cytochrome *c* nitrite reductase, possibly to prevent toxic accumulation of nitrite. The '*M. nitroreducens*' genomes do not encode other denitrification enzymes illustrating that neither NO, N₂O nor N₂ are final products; instead, nitrite is the main product of nitrate reduction with about 10 % of the nitrite reduced to ammonium (Haroon *et al.*, 2013; Ettwig *et al.*, 2016). Another unusual feature of the ANME-2d genomes are that they encode for a high number of cytochrome *c* proteins (Haroon *et al.*, 2013; Arshad *et al.*, 2015; Kletzin *et al.*, 2015). The role of cytochrome *c* proteins in '*M. nitroreducens*' remains enigmatic, as the metabolism of nitrate-dependent AOM does not require electron transfer to a syntrophic partner microorganism as found for ANME-2a (McGlynn *et al.*, 2015; Wegener *et al.*, 2015). A recent publication detected ANME-2d archaea in a culture that coupled Cr(VI) reduction to anaerobic methane oxidation (Lu *et al.*, 2016) which may require *c*-type cytochromes. As many other microorganisms manage nitrate reduction without the excessive use of cytochrome *c*, and furthermore closely related *Methanosarcina* strains harbor only a few – if any – *c*-type cytochromes, the role of these proteins in N-AOM has to be further investigated.

Concluding remarks

Methane oxidizing microorganisms play an essential role in counteracting biological methane production and its release to the atmosphere. The widespread occurrence and substantial size of potential habitats suggests an important role for nitrate- and nitrite-dependent methane oxidizers that link the biogeochemical carbon and nitrogen cycles (Figure 5). Application of more specific

detection methods are needed and will hopefully broaden our insight into the environmental significance of N-AOM microorganisms. Physiological experiments with co-cultures of various nitrogen cycle organisms need to be further explored. Competition for nitrate and nitrite as well as composition of microbial communities in natural habitats is likely determined by the availability and relative concentrations of electron donors and acceptors. Further laboratory studies and environmental data sets are needed to understand substrate fluxes and microbial community development in relevant ecosystems to ultimately understand and possibly predict the fate of involved substrates.

Models describing the metabolic pathways for methane and nitrate/nitrite conversion and involved enzyme systems in *Methanoperedens* and *Methyloirabilis*-like microorganisms have been proposed (Figure 5). It is interesting to note that the degree of genetic innovation required to catalyze the two processes appears to be limited. Nitrite-dependent methane oxidation by '*M. oxyfera*' mostly employs enzymatic modules commonly found in denitrifiers and aerobic methanotrophs. The key novelty that seems to enable these organisms to respire methane with nitrite is the alteration of a canonical nitric oxide reductase into a nitric oxide disproportionating enzyme. Nitrate-dependent methane oxidation by '*M. nitroreducens*' is based on the reversal of the methyl-CoM reductase reaction and subsequent steps from the Wood-Ljungdahl/methanogenesis pathway. The key innovation is the acquisition of a nitrate reductase and accessory proteins. The exceptionally large number of Cyt_c present in ANME organisms suggests the additional need to rewire the electron transfer routes to accommodate this metabolism or additional metabolic capacities that have not yet been discovered. Furthermore,

403 physiological and detailed biochemical studies are needed to test the current models for these
404 fascinating processes.

Acknowledgements

The authors are thankful for financial support by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek through the SIAM Gravitation Grant 024.002.002 and the NESSC Gravitation Grant 024.002.001. MSMJ, AV, AA, JR, WV and CL are further supported by ERC AG EcoMoM 339880 and HOdC by ERC AG Volcano 669371. The funding agencies had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

References

- Arshad, A., Speth, D.R., De Graaf, R.M., Op den Camp, H.J.M., Jetten, M.S.M., and Welte, C.U. (2015) A metagenomics-based metabolic model of nitrate-dependent anaerobic oxidation of methane by *Methanoperedens*-like archaea. *Front Microbiol* 6: 1423.
- Blanksby, S.J., and Ellison, G.B. (2003) Bond dissociation energies of organic molecules. *Acc Chem Res* 36: 255-263.
- Boetius, A., Ravensschlag, K., Schubert, C.J., Rickert, D., Widdel, F., Gieseke, A. et al. (2000) A marine microbial consortium apparently mediating anaerobic oxidation of methane. *Nature* 407: 623-626.
- Boyd, E.S., Skidmore, M., Mitchell, A.C., Bakermans, C., and Peters, J.W. (2010) Methanogenesis in subglacial sediments. *Environ Microbiol Rep* 2: 685-692.
- Brune, A., Frenzel, P., and Cypionka, H. (2000). Life at the oxic–anoxic interface: microbial activities and adaptations. *FEMS Microbiol Rev* 24: 691-710.
- Cadillo-Quiroz, H., Yashiro, E., Yavitt, J.B., and Zinder, S.H. (2008) Characterization of the archaeal community in a minerotrophic fen and terminal restriction fragment length polymorphism-directed isolation of a novel hydrogenotrophic methanogen. *Appl Environ Microbiol* 74: 2059-2068.

430 Cadillo-Quiroz, H., Brauer, S., Yashiro, E., Sun, C., Yavitt, J., and Zinder, S. (2006) Vertical
 431 profiles of methanogenesis and methanogens in two contrasting acidic peatlands in central New
 432 York State, USA. *Environ Microbiol* 8: 1428-1440.

433 Chang, Y.H., Cheng, T.W., Lai, W.J., Tsai, W.Y., Sun, C.H., Lin, L.H., and Wang, P.L. (2012)
 434 Microbial methane cycling in a terrestrial mud volcano in eastern Taiwan. *Environ Microbiol* 14:
 435 895-908.

436 Chen, J., Zhou, Z., and Gu, J.-D. (2014a) Complex community of nitrite-dependent anaerobic
 437 methane oxidation bacteria in coastal sediments of the Mai Po wetland by PCR amplification of
 438 both 16S rRNA and *pmoA* genes. *Appl Microbiol Biotechnol* 99: 1463-1473.

439 Chen, J., Zhou, Z.-C., and Gu, J.-D. (2014b) Occurrence and diversity of nitrite-dependent
 440 anaerobic methane oxidation bacteria in the sediments of the South China Sea revealed by
 441 amplification of both 16S rRNA and *pmoA* genes. *Appl Microbiol Biotechnol* 98: 5685-5696.

442 Chin, K.J., Lueders, T., Friedrich, M.W., Klose, M., and Conrad, R. (2004) Archaeal community
 443 structure and pathway of methane formation on rice roots. *Microb Ecol* 47: 59-67.

444 Conrad, R. (2009) The global methane cycle: recent advances in understanding the microbial
 445 processes involved. *Environ Microbiol Rep* 1: 285-292.

446 Conrad, R., Klose, M., Noll, M., Kemnitz, D., and Bodelier, P.L.E. (2008) Soil type links
 447 microbial colonization of rice roots to methane emission. *Glob Chang Biol* 14: 657-669.

448 Costa, C., Dijkema, C., Friedrich, M., García-Encina, P., Fernández-Polanco, F., and Stams,
 449 M.A.J. (2000) Denitrification with methane as electron donor in oxygen-limited bioreactors.
 450 Appl Microbiol Biotechnol 53: 754-762.

451 Costa, K.C., Wong, P.M., Wang, T., Lie, T.J., Dodsworth, J.A., Swanson, I. et al. (2010) Protein
 452 complexing in a methanogen suggests electron bifurcation and electron delivery from formate to
 453 heterodisulfide reductase. Proc Natl Acad Sci U S A 107: 11050-11055.

454 Daims, H., Lebedeva, E.V., Pjevac, P., Han, P., Herbold, C., Albertsen, M. et al. (2015)
 455 Complete nitrification by *Nitrospira* bacteria. Nature 528: 504-509.

456 de Vries, S., Momcilovic, M., Strampraad, M.J., Whitelegge, J.P., Baghai, A., and Schroder, I.
 457 (2010) Adaptation to a high-tungsten environment: *Pyrobaculum aerophilum* contains an active
 458 tungsten nitrate reductase. Biochemistry 49: 9911-9921.

459 Deutzmann, J.S., and Schink, B. (2011) Anaerobic oxidation of methane in sediments of Lake
 460 Constance, an oligotrophic freshwater lake. Appl Environ Microbiol 77: 4429-4436.

461 Deutzmann, J.S., Stief, P., Brandes, J., and Schink, B. (2014) Anaerobic methane oxidation
 462 coupled to denitrification is the dominant methane sink in a deep lake. Proc Natl Acad Sci U S A
 463 111: 18273-18278.

464 Ding, J., Ding, Z.-W., Fu, L., Lu, Y.-Z., Cheng, S.H., and Zeng, R.J. (2015) New primers for
 465 detecting and quantifying denitrifying anaerobic methane oxidation archaea in different
 466 ecological niches. *Appl Microbiol Biotechnol* 99: 9805-9812.

467 Ding, Z.-W., Ding, J., Fu, L., Zhang, F., and Zeng, R.J. (2014) Simultaneous enrichment of
 468 denitrifying methanotrophs and anammox bacteria. *Appl Microbiol Biotechnol* 98: 10211-10221.

469 Dumont, M.G., and Murrell, J.C. (2005) Stable isotope probing - linking microbial identity to
 470 function. *Nat Rev Micro* 3: 499-504.

471 Eisentraeger, A., Klag, P., Vansbotter, B., Heymann, E., and Dott, W. (2001) Denitrification of
 472 groundwater with methane as sole hydrogen donor. *Water Res* 35: 2261-2267.

473 Ettwig, K.F., van Alen, T., van de Pas-Schoonen, K.T., Jetten, M.S., and Strous, M. (2009)
 474 Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10
 475 phylum. *Appl Environ Microbiol* 75: 3656-3662.

476 Ettwig, K.F., Speth, D.R., Reimann, J., Wu, M.L., Jetten, M.S., and Keltjens, J.T. (2012)
 477 Bacterial oxygen production in the dark. *Front Microbiol* 3: 273.

478 Ettwig, K.F., Shima, S., van de Pas-Schoonen, K.T., Kahnt, J., Medema, M.H., Op den Camp,
 479 H.J. et al. (2008) Denitrifying bacteria anaerobically oxidize methane in the absence of *Archaea*.
 480 *Environ Microbiol* 10: 3164-3173.

481 Ettwig, K.F., Butler, M.K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M.M. et al.
 482 (2010) Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* 464: 543-548.

483 Ettwig, K.F., Zhu, B., Speth, D.R., Keltjens, J.T., Jetten, M.S.M., Kartal, B. (2016) Archaea
 484 catalyze iron-dependent anaerobic oxidation of methane. *Proc Natl Acad Sci U S A*, in press.

485 Feisthauer, S., Vogt, C., Modrzynski, J., Szlenkier, M., Krüger, M., Siebert, M., and Richnow,
 486 H.-H. (2011) Different types of methane monooxygenases produce similar carbon and hydrogen
 487 isotope fractionation patterns during methane oxidation. *Geochim Cosmochim Acta* 75: 1173-
 488 1184.

489 Flynn, T.M., Sanford, R.A., Ryu, H., Bethke, C.M., Levine, A.D., Ashbolt, N.J., and Santo
 490 Domingo, J.W. (2013) Functional microbial diversity explains groundwater chemistry in a
 491 pristine aquifer. *BMC Microbiology* 13: 1-15.

492 Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R. et al. (2008)
 493 Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science*
 494 320: 889-892.

495 Gülay, A., Musovic, S., Albrechtsen, H.J., Al-Soud, W.A., Sorensen, S.J., and Smets, B.F.
 496 (2016) Ecological patterns, diversity and core taxa of microbial communities in groundwater-fed
 497 rapid gravity filters. *ISME J* doi: 10.1038/ismej.2016.16. [Epub ahead of print]

498 Hallam, S.J., Putnam, N., Preston, C.M., Detter, J.C., Rokhsar, D., Richardson, P.M., and
 499 DeLong, E.F. (2004) Reverse methanogenesis: testing the hypothesis with environmental
 500 genomics. *Science* 305: 1457-1462.

501 Han, P., and Gu, J.-D. (2013) A newly designed degenerate PCR primer based on *pmoA* gene for
 502 detection of nitrite-dependent anaerobic methane-oxidizing bacteria from different ecological
 503 niches. *Appl Microbiol Biotechnol* 97: 10155-10162.

504 Hansch, C., Leo, A., and Hoekman, D. (1995) - Hydrophobic, electronic, and steric constants. In
 505 *Exploring QSAR*, p. 3.

506 Hanson, B.T., and Madsen, E.L. (2015) *In situ* expression of nitrite-dependent anaerobic
 507 methane oxidation proteins by *Candidatus Methylomirabilis oxyfera* co-occurring with
 508 expressed anammox proteins in a contaminated aquifer. *Environ Microbiol Rep* 7: 252-264.

509 Haroon, M.F., Hu, S., Shi, Y., Imelfort, M., Keller, J., Hugenholtz, P. et al. (2013) Anaerobic
 510 oxidation of methane coupled to nitrate reduction in a novel archaeal lineage. *Nature* 500: 567-
 511 570.

512 Hatamoto, M., Kimura, M., Sato, T., Koizumi, M., Takahashi, M., Kawakami, S. et al. (2014)
 513 Enrichment of denitrifying methane-oxidizing microorganisms using up-flow continuous
 514 reactors and batch cultures. *PLoS ONE* 9: e115823.

515 He, Z., Geng, S., Cai, C., Liu, S., Liu, Y., Pan, Y. et al. (2015) Halophilic anaerobic oxidation of
 516 methane coupled to nitrite reduction by marine NC10 bacteria. *Appl Environ Microbiol* 81:
 517 5538-5545.

518 Hemp, J., and Gennis, R.B. (2008) Diversity of the heme–copper superfamily in archaea:
 519 insights from genomics and structural modeling. In *Bioenergetics: Energy Conservation and*
 520 *Conversion*. Schäfer, G., and Penefsky, H.S. (eds). Berlin, Heidelberg: Springer Berlin
 521 Heidelberg, pp. 1-31.

522 Hommes, R.W.J, Simons, J.A., Snoep, J.L., Postma, P.W., Tempest, D.W., and Neijssse, O.M.
 523 (1991) Quantitative aspects of glucose metabolism by *Escherichia coli* B/r, grown in the
 524 presence of pyrroloquinoline quinone. *Antonie van Leeuwenhoek* 60: 373-382.

525 Hu, S., Zeng, R.J., Keller, J., Lant, P.A., and Yuan, Z. (2011) Effect of nitrate and nitrite on the
 526 selection of microorganisms in the denitrifying anaerobic methane oxidation process. *Environ*
 527 *Microbiol Rep* 3: 315-319.

528 Hu, S., Zeng, R.J., Haroon, M.F., Keller, J., Lant, P.A., Tyson, G.W., and Yuan, Z. (2015) A
 529 laboratory investigation of interactions between denitrifying anaerobic methane oxidation
 530 (DAMO) and anammox processes in anoxic environments. *Sci Rep* 5: 8706.

531 Hug, L.A., Thomas, B.C., Sharon, I., Brown, C.T., Sharma, R., Hettich, R.L. et al. (2016)
 532 Critical biogeochemical functions in the subsurface are associated with bacteria from new phyla
 533 and little studied lineages. *Environ Microbiol* 18: 159-173.

534 Inagaki, F., Nunoura, T., Nakagawa, S., Teske, A., Lever, M., Lauer, A. et al. (2006)
 535 Biogeographical distribution and diversity of microbes in methane hydrate-bearing deep marine
 536 sediments on the Pacific Ocean Margin. *Proc Natl Acad Sci U S A* 103: 2815-2820.

537 Islas-Lima, S., Thalasso, F., and Gómez-Hernandez, J. (2004) Evidence of anoxic methane
 538 oxidation coupled to denitrification. *Water Res* 38: 13-16.

539 Kampman, C., Temmink, H., Hendrickx, T.L.G., Zeeman, G., and Buisman, C.J.N. (2014)
 540 Enrichment of denitrifying methanotrophic bacteria from municipal wastewater sludge in a
 541 membrane bioreactor at 20°C. *J Hazard Mater* 274: 428-435.

542 Karl, D.M., Beversdorf, L., Bjorkman, K.M., Church, M.J., Martinez, A., and Delong, E.F.
 543 (2008) Aerobic production of methane in the sea. *Nature Geosci* 1: 473-478.

544 Kasai, Y., Takahata, Y., Hoaki, T., and Watanabe, K. (2005) Physiological and molecular
 545 characterization of a microbial community established in unsaturated, petroleum-contaminated
 546 soil. *Environ Microbiol* 7: 806-818.

547 Kaster, A.K., Moll, J., Parey, K., and Thauer, R.K. (2011) Coupling of ferredoxin and
 548 heterodisulfide reduction via electron bifurcation in hydrogenotrophic methanogenic archaea.
 549 *Proc Natl Acad Sci U S A* 108: 2981-2986.

550 Kato, S., Chan, C., Itoh, T., and Ohkuma, M. (2013) Functional gene analysis of freshwater iron-
 551 rich flocs at circumneutral pH and isolation of a stalk-forming microaerophilic iron-oxidizing
 552 bacterium. *Appl Environ Microbiol* 79: 5283-5290.

553 Keltjens, J.T., Pol, A., Reimann, J., and Op den Camp, H.J. (2014) PQQ-dependent methanol
 554 dehydrogenases: rare-earth elements make a difference. *Appl Microbiol Biotechnol* 98: 6163-
 555 6183.

556 Kletzin, A., Heimerl, T., Flechsler, J., van Niftrik, L., Rachel, R., and Klingl, A. (2015)
 557 Cytochromes *c* in *Archaea*: distribution, maturation, cell architecture, and the special case of
 558 *Ignicoccus hospitalis*. *Front Microbiol* 6: 439.

559 Knittel, K., and Boetius, A. (2009) Anaerobic oxidation of methane: progress with an unknown
 560 process. *Annu Rev Microbiol* 63: 311-334.

561 Kojima, H., Tsutsumi, M., Ishikawa, K., Iwata, T., Mußmann, M., and Fukui, M. (2012)
 562 Distribution of putative denitrifying methane oxidizing bacteria in sediment of a freshwater lake,
 563 Lake Biwa. *Syst Appl Microbiol* 35: 233-238.

564 Kool, D.M., Zhu, B., Rijpstra, W.I.C., Jetten, M.S.M., Ettwig, K.F., and Sinninghe Damsté, J.S.
 565 (2012) Rare branched fatty acids characterize the lipid composition of the intra-aerobic methane
 566 oxidizer “*Candidatus Methylomirabilis oxyfera*”. *Appl Environ Microbiol* 78: 8650-8656.

567 Krüger, M., Meyerdierks, A., Glockner, F.O., Amann, R., Widdel, F., Kube, M. et al. (2003) A
 568 conspicuous nickel protein in microbial mats that oxidize methane anaerobically. *Nature* 426:
 569 878-881.

570 Lever, M.A., Rouxel, O., Alt, J.C., Shimizu, N., Ono, S., Coggon, R.M. et al. (2013) Evidence
 571 for microbial carbon and sulfur cycling in deeply buried ridge flank basalt. *Science* 339: 1305-
 572 1308.

573 Li, Q., Wang, F., Chen, Z., Yin, X., and Xiao, X. (2012) Stratified active archaeal communities
 574 in the sediments of Jiulong River Estuary, China. *Front Microbiol* 3: 311.

575 Liu, Y., Zhang, J., Zhao, L., Li, Y., Yang, Y., and Xie, S. (2015) Aerobic and nitrite-dependent
 576 methane-oxidizing microorganisms in sediments of freshwater lakes on the Yunnan Plateau.
 577 *Appl Microbiol Biotechnol* 99: 2371-2381.

578 Lomakina, A.V., Pogodaeva, T.V., Morozov, I.V., and Zemskaya, T.I. (2014) Microbial
 579 communities of the discharge zone of oil- and gas-bearing fluids in low-mineral Lake Baikal.
 580 *Mikrobiologiya* 83: 355-365.

581 Lu, Y.Z., Fu, L., Ding, J., Ding, Z.W., Li, N., and Zeng, R.J. (2016) Cr(VI) reduction coupled
 582 with anaerobic oxidation of methane in a laboratory reactor. *Water Res* 102: 445-452.

583 Luesken, F., van Alen, T., van der Biezen, E., Frijters, C., Toonen, G., Kampman, C. et al.
 584 (2011a) Diversity and enrichment of nitrite-dependent anaerobic methane oxidizing bacteria
 585 from wastewater sludge. *Appl Microbiol Biotechnol* 92: 845-854.

586 Luesken, F.A., Sánchez, J., van Alen, T.A., Sanabria, J., Op den Camp, H.J.M., Jetten, M.S.M.,
 587 and Kartal, B. (2011b) Simultaneous nitrite-dependent anaerobic methane and ammonium
 588 oxidation processes. *Appl Environl Microbiol* 77: 6802-6807.

589 Luesken, F.A., Wu, M.L., Op den Camp, H.J., Keltjens, J.T., Stunnenberg, H., Francoijs, K.J. et
 590 al. (2012) Effect of oxygen on the anaerobic methanotroph '*Candidatus* Methyloirabilis
 591 oxyfera': kinetic and transcriptional analysis. *Environ Microbiol* 14: 1024-1034.

592 Luesken, F.A., Zhu, B., van Alen, T.A., Butler, M.K., Diaz, M.R., Song, B. et al. (2011c) *pmoA*
 593 primers for detection of anaerobic methanotrophs. *Appl Environ Microbiol* 77: 3877-3880.

594 Martinez-Espinosa, R.M., Dridge, E.J., Bonete, M.J., Butt, J.N., Butler, C.S., Sargent, F., and
 595 Richardson, D.J. (2007) Look on the positive side! The orientation, identification and
 596 bioenergetics of 'Archaeal' membrane-bound nitrate reductases. *FEMS Microbiol Lett* 276: 129-
 597 139.

598 Mason, I. (1977) Methane as a carbon source in biological denitrification. *J Water Pollut Control*
 599 *Fed* 49: 855-857.

600 McGlynn, S.E., Chadwick, G.L., Kempes, C.P., and Orphan, V.J. (2015) Single cell activity
601 reveals direct electron transfer in methanotrophic consortia. *Nature* 526: 531-535.

602 Metcalf, W.W., Griffin, B.M., Cicchillo, R.M., Gao, J., Janga, S.C., Cooke, H.A. et al. (2012)
603 Synthesis of methylphosphonic acid by marine microbes: a source for methane in the aerobic
604 ocean. *Science* 337: 1104-1107.

605 Nowka, B., Daims, H., and Spieck, E. (2015) Comparison of oxidation kinetics of nitrite-
606 oxidizing bacteria: nitrite availability as a key factor in niche differentiation. *Appl Environ*
607 *Microbiol* 81: 745-753.

608 Op den Camp, H.J.M., Islam, T., Stott, M.B., Harhangi, H.R., Hynes, A., Schouten, S. et al.
609 (2009) Environmental, genomic and taxonomic perspectives on methanotrophic
610 Verrucomicrobia. *Environ Microbiol Rep* 1: 293-306.

611 Orphan, V.J., House, C.H., Hinrichs, K.U., McKeegan, K.D., DeLong, E.F. (2001) Methane-
612 consuming archaea revealed by directly coupled isotopic and phylogenetic analysis. *Science* 293:
613 484-487.

614 Pachiadaki, M.G., Kallionaki, A., Dählmann, A., De Lange, G.J., and Kormas, K.A. (2011)
615 Diversity and spatial distribution of prokaryotic communities along a sediment vertical profile of
616 a deep-sea mud volcano. *Microb Ecol* 62: 655-668.

617 Padilla, C.C., Bristow, L.A., Sarode, N., Garcia-Robledo, E., Gomez Ramirez, E., Benson, C.R.
618 et al. (2016) NC10 bacteria in marine oxygen minimum zones. ISME J doi:
619 10.1038/ismej.2015.262. [Epub ahead of print].

620 Palomo, A., Jane Fowler, S., Gulay, A., Rasmussen, S., Sicheritz-Ponten, T., and Smets, B.F.
621 (2016) Metagenomic analysis of rapid gravity sand filter microbial communities suggests novel
622 physiology of *Nitrospira* spp. ISME J doi: 10.1038/ismej.2016.63. [Epub ahead of print].

623 Panganiban, A.T., Patt, T.E., Hart, W., and Hanson, R.S. (1979) Oxidation of methane in the
624 absence of oxygen in lake water samples. Appl Environ Microbiol 37: 303-309.

625 Pinto, A.J., Marcus, D.N., Ijaz, U.Z., Bautista-de Iose Santos, Q.M., Dick, G.J., and Raskin, L.
626 (2016) Metagenomic evidence for the presence of comammox *Nitrospira*-like bacteria in a
627 drinking water system. mSphere 1: 1-8.

628 Pol, A., Barends, T.R., Dietl, A., Khadem, A.F., Eygensteyn, J., Jetten, M.S., and Op den Camp,
629 H.J. (2014) Rare earth metals are essential for methanotrophic life in volcanic mudpots. Environ
630 Microbiol 16: 255-264.

631 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P. et al. (2013) The SILVA
632 ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic
633 Acids Res 41: D590-596.

634 Raghoebarsing, A.A., Pol, A., van de Pas-Schoonen, K.T., Smolders, A.J.P., Ettwig, K.F.,
635 Rijpstra, W.I.C. et al. (2006) A microbial consortium couples anaerobic methane oxidation to
636 denitrification. *Nature* 440: 918-921.

637 Rasigraf, O., Vogt, C., Richnow, H.-H., Jetten, M.S.M., and Ettwig, K.F. (2012) Carbon and
638 hydrogen isotope fractionation during nitrite-dependent anaerobic methane oxidation by
639 *Methyloirabilis oxyfera*. *Geochim Cosmochim Acta* 89: 256-264.

640 Rasigraf, O., Kool, D.M., Jetten, M.S.M., Sinninghe Damsté, J.S., and Ettwig, K.F. (2014)
641 Autotrophic carbon dioxide fixation via the Calvin-Benson-Bassham cycle by the denitrifying
642 methanotroph "*Candidatus Methyloirabilis oxyfera*". *Appl Environ Microbiol* 80: 2451-2460.

643 Rastogi, G., Sani, R.K., Peyton, B.M., Moberly, J.G., and Ginn, T.R. (2009) Molecular studies
644 on the microbial diversity associated with mining-impacted Coeur d'Alene River sediments.
645 *Microb Ecol* 58: 129-139.

646 Reeburgh, W. S., and Heggie, D. T. (1977). Microbial methane consumption reactions and their
647 effect on methane distributions in freshwater and marine environments. *Limnol Oceanogr* 22: 1-
648 9.

649 Reimann, J., Jetten, M.S., and Keltjens, J.T. (2015) Metal enzymes in "impossible"
650 microorganisms catalyzing the anaerobic oxidation of ammonium and methane. In *Met Ions Life*
651 *Sci*, pp. 257-313.

652 Scheller, S., Goenrich, M., Boecher, R., Thauer, R.K., Jaun, B. (2010) The key nickel enzyme of
653 methanogenesis catalyses the anaerobic oxidation of methane. *Nature* 465: 606-608.

654 Schubert, C.J., Vazquez, F., Lösekann-Behrens, T., Knittel, K., Tonolla, M., and Boetius, A.
655 (2011) Evidence for anaerobic oxidation of methane in sediments of a freshwater system (Lago
656 di Cadagno). *FEMS Microbiol Ecol* 76: 26-38.

657 Semrau, J.D., DiSpirito, A.A., and Yoon, S. (2010) Methanotrophs and copper. *FEMS Microbiol*
658 *Rev* 34: 496-531.

659 Shen, L.-D., Zhu, Q., Liu, S., Du, P., Zeng, J.-N., Cheng, D.-Q. et al. (2014a) Molecular
660 evidence for nitrite-dependent anaerobic methane-oxidising bacteria in the Jiaojiang Estuary of
661 the East Sea (China). *Appl Microbiol Biotechnol* 98: 5029-5038.

662 Shen, L.-D., Liu, S., Zhu, Q., Li, X.-Y., Cai, C., Cheng, D.-Q. et al. (2014b) Distribution and
663 diversity of nitrite-dependent anaerobic methane-oxidising bacteria in the sediments of the
664 Qiantang River. *Microb Ecol* 67: 341-349.

665 Shen, L.-D., Liu, S., He, Z.-F., Lian, X., Huang, Q., He, Y.-F. et al. (2015) Depth-specific
666 distribution and importance of nitrite-dependent anaerobic ammonium and methane-oxidising
667 bacteria in an urban wetland. *Soil Biol Biochem* 83: 43-51.

668 Shen, L.-D., Liu, S., Huang, Q., Lian, X., He, Z.-F., Geng, S. et al. (2014c) Evidence for the
 669 cooccurrence of nitrite-dependent anaerobic ammonium and methane oxidation processes in a
 670 flooded paddy field. *Appl Environ Microbiol* 80: 7611-7619.

671 Shen, L.-D., Wu, H.S., Gao, Z.q., Li, J., and Liu, X. (2016) Presence of diverse *Candidatus*
 672 *Methylomirabilis oxyfera*-like bacteria of NC10 phylum in agricultural soils. *J Appl Microbiol*
 673 120: 1552-1560.

674 Smith, R.L., Howes, B.L., and Garabedian, S.P. (1991) *In situ* measurement of methane
 675 oxidation in groundwater by using natural-gradient tracer tests. *Appl Environ Microbiol* 57:
 676 1997-2004.

677 Sollo, F.W., Mueller, H.F., and Larson, T.E. (1976) Denitrification of wastewater effluents with
 678 methane. *J Water Pollut Control Fed* 48: 1840-1842.

679 Stein, L.Y., La Duc, M.T., Grundl, T.J., and Nealson, K.H. (2001) Bacterial and archaeal
 680 populations associated with freshwater ferromanganous micronodules and sediments. *Environ*
 681 *Microbiol* 3: 10-18.

682 Thalasso, F., Vallecillo, A., García-Encina, P., and Fdz-Polanco, F. (1997) The use of methane
 683 as a sole carbon source for wastewater denitrification. *Water Res* 31: 55-60.

684 Thauer, R.K., Kaster, A.K., Seedorf, H., Buckel, W., and Hedderich, R. (2008) Methanogenic
 685 archaea: ecologically relevant differences in energy conservation. *Nat Rev Microbiol* 6: 579-591.

686 van Kessel, M.A.H.J., Speth, D.R., Albertsen, M., Nielsen, P.H., Op den Camp, H.J.M., Kartal,
687 B. et al. (2015) Complete nitrification by a single microorganism. *Nature* 528: 555-559.

688 van Teeseling, M.C.F., Pol, A., Harhangi, H.R., van der Zwart, S., Jetten, M.S.M., Op den Camp,
689 H.J.M., and van Niftrik, L. (2014) Expanding the verrucomicrobial methanotrophic world:
690 description of three novel species of *Methyloacidimicrobium* gen. nov. *Appl Environ Microbiol*
691 80: 6782-6791.

692 Waki, M., Tanaka, Y., Osada, T., and Suzuki, K. (2002) Effects of nitrite and ammonium on
693 methane-dependent denitrification. *Appl Microbiol Biotechnol* 59: 338-343.

694 Wang, Y., Zhu, G., Harhangi, H.R., Zhu, B., Jetten, M.S.M., Yin, C., and Op den Camp, H.J.M.
695 (2012) Co-occurrence and distribution of nitrite-dependent anaerobic ammonium and methane-
696 oxidizing bacteria in a paddy soil. *FEMS Microbiol Lett* 336: 79-88.

697 Wegener, G., Krukenberg, V., Riedel, D., Tegetmeyer, H.E., and Boetius, A. (2015) Intercellular
698 wiring enables electron transfer between methanotrophic archaea and bacteria. *Nature* 526: 587-
699 590.

700 Welte, C., and Deppenmeier, U. (2014) Bioenergetics and anaerobic respiratory chains of
701 acetoclastic methanogens. *Biochim Biophys Acta* 1837: 1130-1147.

702 Wrede, C., Brady, S., Rockstroh, S., Dreier, A., Kokoschka, S., Heinzelmann, S.M. et al. (2012)
703 Aerobic and anaerobic methane oxidation in terrestrial mud volcanoes in the Northern
704 Apennines. *Sediment Geol* 263-264: 210-219.

705 Wu, M.L., Ettwig, K.F., Jetten, M.S.M., Strous, M., Keltjens, J.T., and van Niftrik, L. (2011) A
706 new intra-aerobic metabolism in the nitrite-dependent anaerobic methane-oxidizing bacterium
707 *Candidatus* 'Methylomirabilis oxyfera'. *Biochem Soc Trans* 39: 243-248.

708 Wu, M.L., Wessels, J.C., Pol, A., Op den Camp, H.J., Jetten, M.S., and van Niftrik, L. (2015)
709 XoxF-type methanol dehydrogenase from the anaerobic methanotroph "*Candidatus*
710 *Methylomirabilis oxyfera*". *Appl Environ Microbiol* 81: 1442-1451.

711 Xu, Y., Ma, K., Huang, S., Liu, L., and Lu, Y. (2012) Diel cycle of methanogen *mcrA* transcripts
712 in rice rhizosphere. *Environ Microbiol Rep* 4: 655-663.

713 Yan, J., Haaijer, S.C.M., Op den Camp, H.J.M., van Niftrik, L., Stahl, D.A., Könneke, M. et al.
714 (2012) Mimicking the oxygen minimum zones: stimulating interaction of aerobic archaeal and
715 anaerobic bacterial ammonia oxidizers in a laboratory-scale model system. *Environ Microbiol*
716 14: 3146-3158.

717 Yang, H.M., Lou, K., Sun, J., Zhang, T., and Ma, X.L. (2012) Prokaryotic diversity of an active
718 mud volcano in the Usu City of Xinjiang, China. *J Basic Microbiol* 52: 79-85.

719 Yoshimatsu, K., Sakurai, T., and Fujiwara, T. (2000) Purification and characterization of
 720 dissimilatory nitrate reductase from a denitrifying halophilic archaeon, *Haloarcula marismortui*.
 721 FEBS Lett 470: 216-220.

722 Zhang, G., Tian, J., Jiang, N., Guo, X., Wang, Y., and Dong, X. (2008) Methanogen community
 723 in Zoige wetland of Tibetan plateau and phenotypic characterization of a dominant uncultured
 724 methanogen cluster ZC-I. Environ Microbiol 10: 1850-1860.

725 Zhu, B., Sanchez, J., van Alen, T.A., Sanabria, J., Jetten, M.S., Ettwig, K.F., and Kartal, B.
 726 (2011) Combined anaerobic ammonium and methane oxidation for nitrogen and methane
 727 removal. Biochem Soc Trans 39: 1822-1825.

728 Zhu, B., van Dijk, G., Fritz, C., Smolders, A.J.P., Pol, A., Jetten, M.S.M., and Ettwig, K.F.
 729 (2012) Anaerobic oxidation of methane in a minerotrophic peatland: enrichment of nitrite-
 730 dependent methane-oxidizing bacteria. Appl Environ Microbiol 78: 8657-8665.

731 Zhu, G., Zhou, L., Wang, Y., Wang, S., Guo, J., Long, X.-E. et al. (2015) Biogeographical
 732 distribution of denitrifying anaerobic methane oxidizing bacteria in Chinese wetland ecosystems.
 733 Environ Microbiol Rep 7: 128-138.

734 **Table legends**

735 **Table 1:** Overview of the environmental distribution and relevant detection methods for N-AOM

736 microorganisms.

Figure legends

Figure 1: Microscopic image of a co-culture catalysing nitrite- and nitrate-dependent anaerobic oxidation of methane that was subjected to fluorescence *in situ* hybridization. The epifluorescence micrograph was obtained after hybridization with the ARCH-641 probe targeting *Methanoperedens*-like archaea (green) and the *M. oxyfera* specific DBACT-193 probe (red). The scale bar represents 20 μm .

Figure 2: Phylogenetic overview of *Methanoperedens*-like archaea based on 16S rRNA gene sequences. (A) Phylogenetic positioning of GOM Arc 1/ANME-2D within other ANME groups and methanogens. (B) Clustering of ANME-2D into groups A-C. Cultured representatives are marked in bold. The classification of the groups was performed with all available 16S rRNA gene sequences of ANME-2D and confirmed by Neighbour-joining and maximum likelihood algorithms. The representative Neighbour-joining phylogenetic tree was calculated using the Jukes Cantor correction, filter over 290 bp and ANME 1 as an outgroup. The full 16S rRNA sequences of the two cultured representatives '*M. nitroreducens* ANME2D' and '*Methanoperedens* sp. BLZ1' are 95.2 % identical.

Figure 3: Phylogenetic overview of NC10 bacteria based on 16S rRNA gene sequences. Depicted is the clustering of the NC10 clade into groups A-D. *Candidatus* 'Methylomirabilis oxyfera' of the group A is marked in bold. The calculation of the tree was carried out by Neighbour-joining algorithm using the Jukes Cantor correction and filter over 1158bp and Acidobacteria as an outgroup.

Figure 4: Simplified overview of how different bacterial and archaeal physiological groups depend on or compete with each other including the anaerobic methanotrophs described in this article. The nitrate-dependent methanotrophs ANME-2d compete with ‘*M. oxyfera*’ for methane yet ‘*M. oxyfera*’ requires the provisioning of nitrite which is the final product of nitrate reduction by ANME-2d. A key competitor for ‘*M. oxyfera*’ seems to be anammox bacteria that take up nitrite very efficiently. Dotted arrow, diffusion. Solid line, metabolic conversion. For the description of the individual groups, please see main text.

Figure 5: Schematic overview of central metabolism of the archaeon ‘*Methanoperedens*’ (A) and the bacterium ‘*Methyloirabilis oxyfera*’ (B). Key enzymes in methane activation and nitrogen conversion reactions are indicated with their encoding genes. Biochemical (solid arrows) and electron transfer reactions (dashed arrows) are depicted schematically and do not indicate stoichiometries. fdh, formate dehydrogenase; fmd, formylmethanofuran dehydrogenase; ftr, formyl transferase; mch, methenyltetrahydromethanopterin cyclohydrolase; mcr, methyl-CoM reductase; mdh, methanol dehydrogenase (XoxF and MxaFI type); mer, methylene tetrahydromethanopterin reductase; mtd, methylene tetrahydromethanopterin dehydrogenase; mtr, Na⁺ translocating methyl transferase; nar, nitrate reductase; nir, *cd*₁ nitrite reductase; nod, NO dismutase; nrf, ammonium-producing nitrite reductase; pmo, particulate methane monooxygenase.

Table 1:

N-AOM organism (Environment type)	Environment	Location	Nutrient conditions	Detection method	Reference
<i>Methylomirabilis / Methanoperedens</i>					
Freshwater ditch	Freshwater ditch	Ooijpolder, Netherlands Twentekanaal, Netherlands	Eutrophic, high nitrite, high methane	Enrichment culture, 16S rRNA and <i>pmoA</i> gene clone libraries, FISH	Raghoebarsing <i>et al.</i> , 2006 Ettwig <i>et al.</i> , 2009
<i>Methylomirabilis</i>					
Wetlands	Various wetlands	China	Oligotrophic-eutrophic, temperatures -25°-80°, pH 5-9,	<i>pmoA</i> and 16S rRNA gene clone libraries, qPCR	Zhu <i>et al.</i> , 2015
	Minerotrophic peatland	Brunsummer- heide, Netherlands	Oligotrophic, influence by agricultural groundwater	<i>pmoA</i> and 16S rRNA gene clone libraries, qPCR, lipids, FISH	Kool <i>et al.</i> , 2012 Zhu <i>et al.</i> , 2012
Freshwater environments	Freshwater lakes	Yunnan Plateau, China	Various levels of reactive N and C	<i>pmoA</i> gene clone libraries, qPCR	Liu <i>et al.</i> , 2015
	Lake sediment	Lake Biwa, Japan	Mesotrophic	16S rRNA and <i>pmoA</i> gene clone libraries, DGGE, qPCR, CARD-FISH	Kojima <i>et al.</i> , 2012
	Lake sediment	Lake Constance, Germany	Oligotrophic	qPCR of 16S rRNA and <i>pmoA</i> genes, FISH	Deutzmann and Schink, 2011 Deutzmann <i>et al.</i> , 2014
	River sediment	Qiantang River, China	Eutrophic, polluted by domestic and industrial effluents, high reactive N and P contents	16S rRNA and <i>pmoA</i> gene clone libraries	Shen <i>et al.</i> , 2014b
	Agricultural soils	Nanjing City,	Slightly acidic pH,	16S rRNA gene amplicon	Shen <i>et al.</i> , 2016

	Paddy field soils	China	moderate N loading	sequencing, qPCR	
	Paddy field soils	Japan	Fertilized soils	enrichment cultures, <i>pmoA</i> gene clone library, FISH	Hatamoto <i>et al.</i> , 2014
	Paddy field soils	China	Fertilized soils, eutrophic, high N and C load	<i>pmoA</i> and 16S rRNA gene clone libraries, qPCR	Wang <i>et al.</i> , 2012 Shen <i>et al.</i> , 2014c
Wastewater & contaminated sites	Waste water treatment plants	Netherlands China	Long sludge retention times, low biological oxygen demand/N rations	Enrichment culture, <i>pmoA</i> and 16S rRNA gene clone libraries, FISH	Luesken <i>et al.</i> , 2011a Luesken <i>et al.</i> , 2011b Ding <i>et al.</i> , 2014 Kampman <i>et al.</i> , 2014
	Coal tar contaminated aquifer	South Glens Falls, USA	High CH ₄ content, moderate reactive N load, contamination with naphthalene	Metaproteomic libraries	Hanson and Madsen, 2015
	Contaminated aquifers	Cape Cod, USA Banisveld, Netherlands	Elevated N load, contaminated with leachate and waste water	<i>pmoA</i> gene clone library	Luesken <i>et al.</i> , 2011b
Brackish environments	Coastal wetland	Mai Po, China	Brackish, eutrophic	16S rRNA and <i>pmoA</i> gene clone libraries	Chen <i>et al.</i> , 2014b
	Coastal sediments	Xiaogan Island, China	Marine	Enrichment culture, <i>pmoA</i> and 16S rRNA gene clone libraries, qPCR, FISH	He <i>et al.</i> , 2015
	Estuary	Jiaojiang, East Sea, China	Eutrophic, moderate salinity, high pollution with polycyclic aromatic hydrocarbons	<i>pmoA</i> and 16S rRNA gene clone libraries, qPCR	Shen <i>et al.</i> , 2014a
Marine environments	Sea sediments	South China Sea, China	n.a.	<i>pmoA</i> and 16S rRNA gene clone libraries	Chen <i>et al.</i> , 2014a
	Oceanic oxygen minimum zone	Mexico and Costa Rica	Anoxic core zone with detectable CH ₄ and NO ₂ ⁻	<i>pmoA</i> gene clone library, 16S rRNA gene qPCR, metatranscriptome	Padilla <i>et al.</i> , 2016

Methanoperedens

Peatlands	Peatland	Michigan Hollow, USA	Minerotrophic fen	16S rRNA gene clone library	Cadillo-Quiroz <i>et al.</i> , 2008
	Peatland	Ithaca, USA	Acidic ombrotrophic bog	16S rRNA gene clone library	Cadillo-Quiroz <i>et al.</i> , 2006
	Peatland	Zoige, Tibet	Neutral peatland	16S rRNA and <i>mcrA</i> gene clone libraries	Zhang <i>et al.</i> , 2008
Freshwater environments	Aquifer	Illinois, USA	High CH ₄ , low SO ₄ ²⁻	16S rRNA gene clone library	Flynn <i>et al.</i> , 2013
	Aquifer	Tokyo, Japan	NO ₃ ⁻ ~7 µM	16S rRNA gene clone library	Kato <i>et al.</i> , 2013
	Subglacial sediment	Alberta, Canada	High C and N	16S rRNA and <i>mcrA</i> gene clone libraries	Boyd <i>et al.</i> , 2010
	Freshwater sediment	Green Bay, USA	Mn-/Fe-rich	16S rRNA gene clone library	Stein <i>et al.</i> , 2001
	River sediment from mining district	Idaho, USA	Heavy metal contaminated	16S rRNA gene clone library	Rastogi <i>et al.</i> , 2009
	Lake sediment	Lago di Cadagno, Switzerland	High SO ₄ ²⁻ and S ²⁻ , no NO ₃ ⁻	16S rRNA gene clone library	Schubert <i>et al.</i> , 2011
	Paddy field	Vercelli, Italy	Nitrogen loaded	<i>mcrA</i> gene clone libraries	Conrad <i>et al.</i> , 2008 Chin <i>et al.</i> , 2004 Xu <i>et al.</i> , 2012
Wastewater & contaminated sites	Lake sediment of oil seep	Baikal lake, Russia	Hydrocarbon rich, low NO ₃ ⁻ and SO ₄ ²⁻	16S rRNA gene clone library	Lomakina <i>et al.</i> , 2014
	Contaminated soil	Shizuoka, Japan	Petroleum contaminated, Mn- and Fe-rich	16S rRNA gene clone library	Kasai <i>et al.</i> , 2005
	Freshwater lake sediment,	Australia	n.a.	Enrichment, metagenome, FISH	Haroon <i>et al.</i> , 2013

digester sludge
and activated
sludge

Mud volcanoes	Mud volcano	Usu City of Xinjiang, China	Alkaline, brackish	16S rRNA gene clone library	Yang <i>et al.</i> , 2012
	Mud volcano	Salse di Nirano, Italy	Hydrocarbon rich, brackish	16S rRNA gene clone library	Wrede <i>et al.</i> , 2012
	Mud volcano	Lei-Gong-Huo, Taiwan	Hydrocarbon rich, Mn- and Fe-rich, moderately saline	16S rRNA gene clone library	Chang <i>et al.</i> , 2012
Brackish environments	Estuary	Fujian, China	High CH ₄ and SO ₄ ²⁻	16S rRNA and <i>mcrA</i> gene clone library	Li <i>et al.</i> , 2012
Marine environments	Seafloor sediment	Juan de Fuca Ridge, USA	High organic carbon	<i>mcrA</i> gene clone library	Lever <i>et al.</i> , 2013

Figure 1

Figure 2

Figure 3

Figure 4

Figure 4