

Basal Ganglia Structure in Tourette's Disorder and/or Attention-Deficit/Hyperactivity Disorder

Natalie J. Forde, MSc,^{1,2*} Marcel P. Zwiers, PhD,² Jilly Naaijen, MSc,² Sophie E. A. Akkermans, MSc,² Thaira J. C. Openneer, MSc,¹ Frank Visscher, MD,³ Andrea Dietrich, PhD,¹ Jan K. Buitelaar, MD, PhD,^{2,4} and Pieter J. Hoekstra, MD, PhD¹

¹University of Groningen, University Medical Center Groningen, Department of Psychiatry, Groningen, The Netherlands ²Radboud University Medical Center, Donders Institute for Brain, Cognition and Behaviour, Department of Cognitive Neuroscience, Nijmegen, The Netherlands ³Admiraal De Ruyter Ziekenhuis, Department of Neurology, Goes, The Netherlands ⁴Karakter Child and Adolescent Psychiatry University Centre, Nijmegen, The Netherlands

ABSTRACT

Background: Tourette's disorder and attention-deficit/hyperactivity disorder often co-occur and have both been associated with structural variation of the basal ganglia. However, findings are inconsistent and comorbidity is often neglected.

Methods: T1-weighted magnetic resonance images from children (n = 141, 8 to 12 years) with Tourette's disorder and/or attention-deficit/hyperactivity disorder and controls were processed with the Oxford Centre for Functional MRI [Magnetic resonance imaging] of the Brain (FMRIB) integrated registration and segmentation tool to determine basal ganglia nuclei volume and shape. Across all participants, basal ganglia nuclei volume and shape were estimated in relation to Tourette's disorder (categorical), attention-deficit/hyperactivity disorder severity (continuous across all participants), and their interaction.

Results: The analysis revealed no differences in basal ganglia nuclei volumes or shape between children with and without Tourette's disorder, no association with attention-deficit/hyperactivity disorder severity, and no interaction between the two.

*Corresponding author: Natalie J. Forde, Department of Psychiatry, University Medical Center Groningen, Postbus 660, 9700 AR Groningen, The Netherlands; j.n.forde@umcg.nl

Funding agency: This study received funding from the European Union under the Seventh Framework People Programme (TS-EUROTRAIN, grant agreement number 316978, and TACTICS, grant agreement number: 278948).

Relevant conflicts of interests/financial disclosures: Nothing to report.

Received: 5 July 2016; **Revised:** 14 September 2016; **Accepted:** 15 September 2016

Published online 10 November 2016 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.26849

Conclusion: We found no evidence that Tourette's disorder, attention-deficit/hyperactivity disorder severity, or a combination thereof are associated with structural variation of the basal ganglia in 8- to 12-year-old patients. © 2016 International Parkinson and Movement Disorder Society.

Key Words: Tourette's Disorder; attention-deficit/hyperactivity disorder; basal ganglia; children; comorbidity

Although Tourette's disorder (TD) is characterized by the presence of both motor and vocal tics,¹ there are also frequently concurrent psychiatric comorbidities. These occur in up to 86% of those with TD during their lifetime.² Attention-deficit/hyperactivity disorder (ADHD) is the most common of these, occurring in approximately 40% of TD cases.³ Even more patients with TD have ADHD symptoms without meeting full diagnostic criteria.⁴

Structural neuroimaging studies of both disorders have highlighted alterations in the nuclei of the basal ganglia (BG), caudate nucleus (CN), putamen (Pu), and globus pallidus (GP).⁵⁻⁷ However, the literature of TD and ADHD neuroimaging research is inconsistent, and determining whether BG structural abnormalities are unique or common to the respective disorders is difficult because few studies have examined groups with TD and ADHD together in 1 study. Furthermore, those studies have been underpowered and inconsistent in their findings.⁸⁻¹²

In the present study, we aimed to elucidate whether BG structural abnormalities are associated with TD in children and whether they occur in relation to comorbid ADHD symptoms. This was done by investigating BG volumes and shape in participants with TD and/or ADHD and healthy controls side by side in a well-sized pediatric sample, thereby focusing on the age range when tics are most frequently present.¹³

Participants and Methods

All participants (Table 1) satisfied the following inclusion criteria: 8 to 12 years of age, IQ >70, White, no previous head injuries or neurological disorders, no contraindications for MRI assessment, no major physical illness, and available good-quality MR scan. Participants of the TD group met *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.) criteria for TD (n = 46) or persistent (chronic) motor or vocal tic disorder (with motor tics only; n = 1); psychiatric comorbidities (eg, ADHD and obsessive compulsive disorder [OCD]) were not excluded. Participants of the ADHD group had a diagnosis of ADHD or subthreshold ADHD, those with tics and/or OCD were excluded. Healthy controls had no mental disorders as screened for by the Child Behaviour

TABLE 1. Demographic description of participants

	Control	ADHD	TD	Test statistic	P value
n	55	39	47		
ADHD, diagnosis/subthreshold	0	35/4	22/10		
Age years, mean (SD)	11.0 (1.0)	10.7 (1.3)	10.5 (1.4)	2.48	.29
Sex, male/female	39/16	21/18	41/6	11.72	.003**
IQ, mean (SD) ^a	109 (12)	102 (13)	105 (11)	3.18	.04
Handed, right/left	50/5	36/3	42/5	0.22	.89
ADHD severity, mean (SD) ^b	45.5 (4.9)	70.9 (10.7)	62.9 (11.3)	86.98	<.0001***
Tic severity, mean (SD) ^c	-	-	T = 20.6 (8.6) M = 13.4 (5.0) V = 7.2 (5.5)		
Age tic onset years, mean (SD) ^c	-	-	5.6 (1.7)		
Duration since tic onset years, mean (SD) ^c	-	-	5.0 (1.8)		
OCD, n ^d	-	-	9		
Medication ^e		6			
Stimulant		26	11		
Strattera		1	0		
Antipsychotic		1	8		
Clonidine		0	2		

ADHD, attention-deficit/hyperactivity disorder; K-W, Kruskal-Wallis; OCD, obsessive compulsive disorder; SD, standard deviation; TD, Tourette's disorder.

^aEstimated from a subtest of the Wechsler Intelligence Scale for Children-III¹⁹ rating.

^bT-scores from the Conners' Parent Rating Scale-Revised long version.¹⁸

^cDetermined with the Yale Global Tic Severity Scale.¹⁵ Total (T), motor (M), and vocal (V) severity ratings exclude impairment score.

^dTotal score ≥ 16 on the Children's Yale-Brown Obsessive Compulsive Scale.¹⁷

^eCurrent medication status, determined from parental report.

P < .01, *P < 0.001.

Checklist.¹⁴ Parents or guardians of all participants gave written informed consent; in addition, participants who were 12 years of age provided written assent. This study was approved by the regional ethics board (Commissie Mensgebonden Onderzoek (CMO) Regio Arnhem-Nijmegen).

The following various instruments were used to determine the presence and severity of disorders: for TD, the Yale Global Tic Severity Scale (using 0-50 ratings, not considering overall impairment)¹⁵; for ADHD and/or other psychiatric disorders, the Schedule for Affective Disorders and Schizophrenia for School-Age Children¹⁶ screening interview plus appropriate modules if required; for OCD, Children's Yale-Brown Obsessive Compulsive Scale¹⁷; for ADHD severity, Conners' Parent Rating Scale-Revised long version.¹⁸ Full-scale IQ was estimated by 4 subtests of the Wechsler Intelligence Scale for Children-III.¹⁹ Medication status was determined from parental reports.

T1-weighted anatomical images were acquired on a 3T Siemens Prisma scanner (Siemens, Erlangen, Germany) with a transversal, three-dimensional magnetisation-prepared rapid gradient-echo (MPRAGE) parallel imaging sequence (parameters: echo time (TE) = 2.98 ms, inversion time (TI) = 900 ms, repetition time (TR) = 2300 ms, flip angle = 9°, voxel size = 1 × 1 × 1.2 mm, acquisition time = 5.30 minutes). Datasets were processed with the automated Oxford Centre for Functional MRI of the Brain (FMRIB) integrated registration and segmentation tool²⁰⁻²² standard procedure to generate volumetric data and surface meshes for subcortical structures. BG nuclei surfaces were reconstructed in native space (useReconNative,

First_utils), aligned to the average structure shape for the cohort (useRigidAlign) and scaled to account for size differences (useScale). Probability maps for gray and white matter tissue types were estimated using the VBM8 toolbox²³ of Statistical Parametric Mapping (Wellcome Department of Imaging Neuroscience, University College London). Total brain volume (TBV) was calculated as the voxel-wise sum of both probability maps.

A repeated-measures analysis of covariance was used to determine the effect of TD and ADHD severity (continuous measure across all groups) and their interaction on BG nuclei volumes. A categorical factor for TD was deemed more robust than using symptom severity because of the fluctuating nature of tics in TD, whereas a continuous measure for ADHD severity was used because multiple participants, particularly within the TD group, displayed ADHD symptoms without meeting criteria for diagnosis. The hemisphere was included as the repeated measure with hemisphere-by-TD and hemisphere-by-ADHD-severity interaction terms as measures of asymmetry difference associated with the respective disorders. TBV, sex, age, and IQ were used as covariates. BG nuclei shape were analyzed using FMRIB Software Library (FSL) randomize²⁴ with 5,000 random permutations and threshold-free cluster enhancement.²⁵ Sex, age, and IQ were entered as covariates. Contrasts tested shape differences between those with and without TD and positive or negative associations between shape and ADHD severity. A separate model was used to assess the interaction effect of TD and ADHD severity on shape using the same covariates. The effect of current stimulant medication use on volumes or shape within patients (TD and/or ADHD) was

investigated in similar models including TD, ADHD severity, age, sex, and IQ as covariates. In addition, for the volume analysis, TBV and hemisphere were included. With similar models, current antipsychotic use within the TD group was also analyzed.

Results

There were no main effects of TD, ADHD severity, or significant interactions between TD and ADHD severity on BG volume (Table 2) or shape.

In the volume analysis, hemisphere was seen to have a significant effect in the CN (left smaller than right) and a trend (uncorrected) toward an interaction with TD. This was not mirrored in the Pu and GP; no significant effects of hemisphere or hemisphere-by-TD interaction were seen (Table 2). No hemisphere-by-ADHD-severity interactions were seen. TBV was significant in each region. It did not differ with TD ($t = -0.90, P = .37$), ADHD severity ($t = -0.90, P = .37$), or their interaction ($t = 0.78, P = .44$). No effect of stimulant or antipsychotic medication was seen on either BG nuclei volume or shape.

Discussion

This is the largest pediatric study of BG structures to investigate TD and ADHD together. Complementary volume and shape analyses of BG nuclei revealed no structural alterations associated with either the presence of TD, ADHD severity, or their interaction.

Although both TD and ADHD have been previously associated with volume alterations in the BG,⁵⁻⁷ the literature to date is heterogeneous. The current null findings replicate in a larger sample a small number of studies that found no associations between BG nuclei volume and TD ($n = 13-38$)^{26,27} and no association between either TD or ADHD and BG nuclei volumes ($n = 14-37$)⁸⁻¹¹ in children. Furthermore, consistent with previous findings in children, the current study also found no asymmetry abnormalities associated with TD.^{27,28} The few studies that have reported associations between BG volume and

TD in children have been inconsistent regarding the regions implicated and direction of change: increased Pu volume bilaterally ($n = 49$ and 14 TD cases, respectively),^{29,30} reduced CN volume bilaterally ($n = 154$, child and adult sample),²⁸ or left only ($n = 23$).³¹ Although all of these studies relate to children, there may still be consequential demographics differences. The few studies that have shown associations between BG structure and TD had a slightly higher mean age³⁰ or a wider age range^{28,29,31} than the current study. It is possible, therefore, that the former studies in question identified differences that occur later in development. BG abnormalities have been more consistently reported in adult samples (eg, refs. ^{28,32}). Along with the results here, this implies that BG abnormalities in TD reflect compensatory mechanisms or an effect of the illness, but do not relate to TD etiology.

Two (overlapping) meta-analyses of BG structure in ADHD^{6,7} showed right BG nuclei volume reductions in children with ADHD. However, similar to the current findings, the large NeuroIMAGE study of ADHD ($n = 307$) in adolescence found no main effect of ADHD on BG volumes either by voxel-based morphometry (VBM)³³ or automated segmentation analyses.³⁴ The second of these studies did, however, show an age-by-diagnosis interaction. This suggests that differences become apparent with increasing age and may account for the current null findings as the cohort was young (8-12 years). Another possible source of the discrepancy between the current and former studies is the use of a continuous measure for ADHD instead of a categorical group for analysis. The dimensional approach is favored³⁵ opposed to categorizing participants with arbitrary thresholds, especially considering the large number of participants with TD that display ADHD or subthreshold ADHD.

In addition to volume analysis, we also applied shape analysis, which is more sensitive to subtle morphological differences. In TD, only 1 small study to date examined BG morphology differences in children, and they found no relation between TD and shape,²⁷ which is in accordance with the findings presented here. In ADHD studies, inward deformation of the BG has been seen in children and adolescents with ADHD when compared with controls.³⁶⁻³⁸ The current study did not corroborate these findings. This discrepancy may relate to demographic differences as discussed earlier.

In the current study, we found no stimulant-dependent associations with BG structure in line with the large ($n = 540$) longitudinal study of BG development in ADHD by Shaw and colleagues,³⁸ who reported no association between stimulant treatment history and developmental trajectories. However, other studies in ADHD reported normalizing effects of stimulants on BG structure.^{6,7,36} Our study was underpowered to determine the effect of antipsychotic treatment on brain structure during development, and this warrants further study.

Our findings should be considered in light of certain limitations. Females were underrepresented in the TD sample, as is expected because males are more frequently

TABLE 2. Results from basal ganglia volume analysis

	Caudate nucleus		Putamen		Globus pallidus	
	<i>F</i>	<i>P</i> value	<i>F</i>	<i>P</i> value	<i>F</i>	<i>P</i> value
TD	0.29	.59	0.27	.60	3.60	.06
ADHD severity	0.001	.98	0.04	.85	0.06	.81
TD × ADHD Severity	2.55	.11	0.14	.71	0.80	.37
Hemisphere	27.02	<.001	2.47	.12	3.50	.06
Hemisphere × TD	3.13	.08	0.80	.37	0.92	.34
Hemisphere × ADHD Severity	0.45	.50	0.19	.67	0.25	.62
TBV	55.09	<.001	44.81	<.001	100.19	<.001

F statistics and uncorrected *P* values are presented from the analysis of basal ganglia volumes in relation to Tourette's disorder (TD), attention-deficit/hyperactivity disorder (ADHD) severity, and hemisphere, including their interaction terms (×). TBV, total brain volume. Statistically significant *p* values are highlighted in bold.

affected than females.² There were, however, no interactions with sex, suggesting the sex imbalance did not confound our findings. A total of 9 participants from the TD group also had OCD. ADHD and OCD were recently shown to have opposing associations with BG structure, although comorbidity was not considered.³⁹ Here we saw that OCD severity was not predictive of BG nuclei volumes within the TD group, and removal of the participants with OCD did not alter our findings.

In conclusion, we found no evidence that TD, ADHD, or their combination are associated with BG structure in patients 8 to 12 years of age. ■

Acknowledgments: We gratefully acknowledge and thank all the participants and their families for their enthusiastic involvement in the study. The authors also thank Nicole Driessen, Leonie Hennissen, and Saskia de Ruiter for their assistance with data collection and Paul Gaalman for his technical support. Furthermore, we wish to acknowledge the numerous clinicians who played a vital role in recruiting participants to the study, most notably we thank Dr Deborah Sival and Dr Els van den Ban.

References

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th ed. Washington, DC: American Psychiatric Association; 2013:81-82.
- Hirschtritt ME, Lee PC, Pauls DL, et al. Lifetime prevalence, age of risk, and genetic relationships of comorbid psychiatric disorders in Tourette syndrome. *JAMA Psychiatry* 2015;72(4):325.
- Rickards H. Republished review: Tourette's syndrome and other tic disorders. *Postgrad Med J* 2011;87(1024):142-149.
- Robertson MM. Tourette syndrome, associated conditions and the complexities of treatment. *Brain* 2000;123(Pt 3):425-462.
- Felling RJ, Singer HS. Neurobiology of tourette syndrome: current status and need for further investigation. *J Neurosci* 2011;31(35):12387-12395.
- Frodl T, Skokauskas N. Meta-analysis of structural MRI studies in children and adults with attention deficit hyperactivity disorder indicates treatment effects. *Acta Psychiatr Scand* 2012;125(2):114-126.
- Nakao T, Radua J, Rubia K, Mataix-Cols D. Gray matter volume abnormalities in ADHD: voxel-based meta-analysis exploring the effects of age and stimulant medication. *Am J Psychiatry* 2011;168(11):1154-1163.
- Zimmerman AM, Abrams MT, Giuliano JD, Denckla MB, Singer HS. Subcortical volumes in girls with tourette syndrome: support for a gender effect. *Neurology* 2000;54(12):2224-2229.
- Jeppesen SS, Debes NM, Simonsen HJ, Rostrup E, Larsson HBW, Skov L. Study of medication-free children with Tourette syndrome do not show imaging abnormalities. *Mov Disord* 2014;29(9):1212-1216.
- Singer HS, Reiss AL, Brown JE, et al. Volumetric MRI changes in basal ganglia of children with Tourette's syndrome. *Neurology* 1993;43(5):950.
- Castellanos FX, Giedd JN, Hamburger SD, Marsh WL, Rapoport JL. Brain morphometry in Tourette's syndrome: the influence of comorbid attention-deficit/hyperactivity disorder. *Neurology* 1996;47(6):1581-1583.
- Peterson B, Riddle MA, Cohen DJ, et al. Reduced basal ganglia volumes in Tourette's syndrome using three-dimensional reconstruction techniques from magnetic resonance images. *Neurology* 1993;43(5):941-949.
- Bloch MH, Leckman JF. Clinical course of Tourette syndrome. *J Psychosom Res* 2009;67(6):497-501.
- Bordin IA, Rocha MM, Paula CS, et al. Child Behavior Checklist (CBCL), Youth Self-Report (YSR) and Teacher's Report Form (TRF): an overview of the development of the original and Brazilian versions. *Cad Saude Publica* 2013;29(1):13-28.
- Leckman JF, Riddle MA, Hardin MT, et al. The Yale Global Tic Severity Scale: initial testing of a clinician-rated scale of tic severity. *J Am Acad Child Adolesc Psychiatry* 1989;28(4):566-573.
- Kaufman J, Birmaher B, Brent D, et al. Schedule for Affective Disorders and Schizophrenia for School-Age Children—Present and Lifetime Version (K-SADS-PL): initial reliability and validity data. *J Am Acad Child Adolesc Psychiatry* 1997;36(7):980-988.
- Scahill L, Riddle MA, McSwiggin-Hardin M, et al. Children's Yale-Brown Obsessive Compulsive Scale: reliability and validity. *J Am Acad Child Adolesc Psychiatry* 1997;36(6):844-852.
- Conners CK, Wells KC, Parker JD, Sitarenios G, Diamond JM, Powell JW. A new self-report scale for assessment of adolescent psychopathology: factor structure, reliability, validity, and diagnostic sensitivity. *J Abnorm Child Psychol* 1997;25(6):487-497.
- Wechsler D. WISC-III Handleiding. London: The Psychological Corporation; 2002.
- Woolrich MW, Jbabdi S, Patenaude B, et al. Bayesian analysis of neuroimaging data in FSL. *Neuroimage* 2009;45(1 suppl):S173-S186.
- Smith SM, Jenkinson M, Woolrich MW, et al. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 2004;23(suppl 1):S208-S219.
- Patenaude B, Smith SM, Kennedy DN, Jenkinson M. A Bayesian model of shape and appearance for subcortical brain segmentation. *Neuroimage* 2011;56(3):907-922.
- Ashburner J, Friston KJ. Voxel-based morphometry—the methods. *Neuroimage* 2000;11(6):805-821.
- Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation inference for the general linear model. *Neuroimage* 2014;92:381-397.
- Smith SM, Nichols TE. Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage* 2009;44(1):83-98.
- Roessler V, Overlack S, Baudewig J, Dechent P, Rothenberger A, Helms G. No brain structure abnormalities in boys with Tourette's syndrome: a voxel-based morphometry study. *Mov Disord* 2009;24(16):2398-2403.
- Williams AC, McNeely ME, Greene DJ, et al. A pilot study of basal ganglia and thalamus structure by high dimensional mapping in children with Tourette syndrome. *F1000Research* 2013;2:207.
- Peterson BS, Thomas P, Kane MJ, et al. Basal ganglia volumes in patients with Gilles de la Tourette syndrome. *Arch Gen Psychiatry* 2003;60(4):415-424.
- Roessler V, Overlack S, Schmidt-Samoa C, et al. Increased putamen and callosal motor subregion in treatment-naive boys with Tourette syndrome indicates changes in the bihemispheric motor network. *J Child Psychol Psychiatry Allied Discip* 2011;52(3):306-314.
- Ludolph AG. Grey-matter abnormalities in boys with Tourette syndrome: magnetic resonance imaging study using optimised voxel-based morphometry. *Br J Psychiatry* 2006;188(5):484-485.
- Makki MI, Behen M, Bhatt A, Wilson B, Chugani HT. Microstructural abnormalities of striatum and thalamus in children with Tourette syndrome. *Mov Disord* 2008;23(16):2349-2356.
- Müller-Vahl KR, Kaufmann J, Grosskreutz J, Dengler R, Emrich HM, Peschel T. Prefrontal and anterior cingulate cortex abnormalities in Tourette Syndrome: evidence from voxel-based morphometry and magnetization transfer imaging. *BMC Neurosci* 2009;10:47.
- Bralten J, Greven CU, Franke B, et al. Voxel-based morphometry analysis reveals frontal brain differences in participants with ADHD and their unaffected siblings. *J Psychiatry Neurosci* 2015;41(2):140377.
- Greven CU, Bralten J, Mennes M, et al. Developmentally stable whole-brain volume reductions and developmentally sensitive caudate and putamen volume alterations in those with attention-deficit/hyperactivity disorder and their unaffected siblings. *JAMA Psychiatry* 2015;72(5):490-499.
- Robbins TW, Gillan CM, Smith DG, de Wit S, Ersche KD. Neurocognitive endophenotypes of impulsivity and compulsivity: Towards dimensional psychiatry. *Trends Cogn Sci* 2012;16(1):81-91.
- Sobel LJ, Bansal R, Maia T V, et al. Basal ganglia surface morphology and the effects of stimulant medications in youth with attention deficit hyperactivity disorder. *Am J Psychiatry* 2010;167(8):977-986.
- Qiu A, Crocetti D, Adler M, et al. Basal ganglia volume and shape in children with attention deficit hyperactivity disorder. *Am J Psychiatry* 2009;166(1):74-82.
- Shaw P, De Rossi P, Watson B, et al. Mapping the development of the basal ganglia in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2014;53(7):780-789.e11.
- Norman LJ, Carlisi C, Lukito S, et al. Structural and functional brain abnormalities in attention-deficit/hyperactivity disorder and obsessive-compulsive disorder. *JAMA Psychiatry* 2016;73(8):815.