

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/178446>

Please be advised that this information was generated on 2019-11-20 and may be subject to change.

Erratum: Upper Limits on the Stochastic Gravitational-Wave Background from Advanced LIGO's First Observing Run [Phys. Rev. Lett. 118, 121101 (2017)]

B. P. Abbott *et al.* and *et al.*

(Received 5 June 2017; published 11 July 2017)

DOI: [10.1103/PhysRevLett.119.029901](https://doi.org/10.1103/PhysRevLett.119.029901)

In Fig. 1 of the Letter, we have plotted an estimator for $\Omega_0 h_0^2$, rather than Ω_0 , where $h_0 = 0.68$ is the Hubble constant divided by 100 km/s/Mpc. This is inconsistent with the legend of Fig. 1, which states that an estimator for Ω_0 is plotted, as well as with the conventions used throughout the rest of the Letter. A new figure showing the estimator for Ω_0 with $h_0 = 0.68$ is shown below. This does not affect any other result in the Letter.




FIG. 1. We show the estimator for Ω_0 in each frequency bin, along with $\pm 2\sigma$ error bars, in the frequency band that contains 99% of the sensitivity for $\alpha = 0$. The loss of sensitivity at around 65 Hz is due to a zero in the overlap reduction function. There are several lines associated with known instrumental artifacts which do not lead to excess cross-correlation. The data are consistent with Gaussian noise, as described in the Results section.