

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/182620>

Please be advised that this information was generated on 2019-04-21 and may be subject to change.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF NIJMEGEN The Netherlands

**TWO SIMPLE DERIVATIONS OF UNIVERSAL
BOUNDS FOR THE C.B.S.
INEQUALITY CONSTANT**

Owe Axelsson, Radim Blaheta

Report No. 0133 (December 2001)

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF NIJMEGEN
Toernooiveld
6525 ED Nijmegen
The Netherlands

Two simple derivations of universal bounds for the C.B.S. inequality constant

Owe Axelsson¹, Radim Blaheta²

Abstract

Universal bounds for the constant in the strengthened Cauchy-Bunyakowski-Schwarz inequality for piecewise linear-linear and piecewise quadratic-linear finite element spaces in 2 space dimensions are derived. The bounds hold for arbitrary shaped triangles, or equivalently, arbitrary matrix coefficients for both scalar diffusion problems and the elasticity theory equations.

Keywords: finite element method, h- and p- refinement, strengthened Cauchy-Bunyakowski-Schwarz inequality

MSC 1991: 65N30, 65N22, 65F10

1 Introduction

This paper deals with the estimates of the constant γ , which appears in the strengthened Cauchy - Bunyakowski - Schwarz (C.B.S.) inequality

$$|a(u, v)| \leq \gamma \sqrt{a(u, u)} \sqrt{a(v, v)} \quad \forall u \in U, v \in V,$$

where U, V are two (finite dimensional) linear spaces, $U \cap V = \{0\}$ and $a(., .)$ is a symmetric positive definite or semidefinite bilinear form. It follows that γ equals the cosine of the angle between the subspaces U, V of $U \cup V$ in a metric defined by $\sqrt{a(u, u)}$.

More precisely, we are interested in the cases, where a comes from the variational formulation of an elliptic boundary value problem in a domain $\Omega \subset R^2$, U is a finite element space of piecewise linear functions and V is the complement of U in another finite element space $U \oplus V$, which arises by h - or p - refinement of U .

¹Faculty of Mathematics and Computer Science, University of Nijmegen, Toornveld 1, 6525 ED Nijmegen, The Netherlands, axelsson@sci.kun.nl

²Department of Applied Mathematics, Institute of Geonics AS CR, Studentská 1768, Ostrava, The Czech Republic, blaheta@ugn.cas.cz. The work was supported by the grant INCO KIT 977006 of the European Commission and the grant OK 383 of the Ministry of Education of the Czech Republic.

In such cases the value of γ can be estimated locally. Namely, if \mathcal{T} is a triangulation of Ω , which is used for the definition of U , then

$$\gamma = \max_{E \in \mathcal{T}} \gamma_E,$$

$$|a_E(u, v)| \leq \gamma_E \sqrt{a_E(u, u)} \sqrt{a_E(v, v)} \quad \forall u \in U_E, v \in V_E$$

where U_E and V_E are linear spaces of functions, which are restrictions of functions from U and V to $E \in \mathcal{T}$. By a_E we denote the restriction of the bilinear form a to E .

The estimates of γ can be used for the convergence analysis of many numerical procedures connected with the application of the finite element method, let us mention two-level and multi-level iterative methods and preconditioners, local refinement composite grid methods, a posteriori error estimates etc.. Therefore, much effort has been devoted to the estimation of γ via local estimates, we mention here the papers by Axelsson [3], Axelsson and Gustafsson [6], Maitre and Musy [11], Margenov [13], Achchab and Maitre [2], Jung and Maitre [10], Axelsson [4], Achchab, et al. [1] and the references therein.

From the estimates presented in the literature, it is seen that γ generally depends on the bilinear form a , i.e. also on the problem coefficients and the type and shape of the finite element used. We can also see that in some cases it is possible to have *universal bounds*, which do not depend on problem coefficients and shape of the finite elements.

This paper will be devoted to the derivation of such universal bounds of the C.B.S constant. We shall present results concerning the bilinear forms corresponding to anisotropic Laplacian and anisotropic elasticity operators. Moreover, we show two ways of simple derivations of the bounds. The resulting bounds generalize and extend the results obtained previously in the literature. The new estimate concerns the case of general anisotropic elasticity.

The paper is organized as follows. In Section 2, we describe more precisely the bilinear forms corresponding to the anisotropic Laplacian and the elasticity operator and we formulate the $P_1 - P_1$ and $P_1 - P_2$ strengthened C.B.S. constant estimation problems. Then in Section 3, we prove a universal bound for the $P_1 - P_1$ problem in 2D. Another way of obtaining the universal bounds, using a relation between $P_1 - P_1$ and $P_1 - P_2$ problems will be shown in Section 4. The paper ends with some concluding remarks.

2 Formulation of $P_1 - P_1$ and $P_1 - P_2$ problems for anisotropic Lagrangian and anisotropic elasticity operators

We shall consider two general types of bilinear forms. The first one can be written in the form

$$a(u, v) = \int_{\Omega} \sum_{i,j=1}^2 a_{ij} \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_j} dx = \int_{\Omega} \langle Dd(u), d(v) \rangle dx. \quad (2.1)$$

Here $u, v \in H^1(\Omega)$, $D = [a_{ij}]$ is a matrix of problem coefficients, which is assumed to be symmetric positive definite, $d(u) = \left(\frac{\partial u}{\partial x_1}, \frac{\partial u}{\partial x_2} \right)^T$ is the gradient of u and $\langle x, y \rangle = x^T y$ for $x, y \in R^2$. This bilinear form corresponds to an anisotropic Laplacian in R^2 .

The second bilinear form corresponds to a general anisotropic elasticity operator. It has the form

$$a(u, v) = \int_{\Omega} \sum_{i,j,k,l=1}^2 c_{ijkl} \varepsilon_{ij}(u) \varepsilon_{kl}(v) dx \quad (2.2)$$

where $u, v \in [H^1(\Omega)]^d$, c_{ijkl} are elasticity moduli, which are nonnegative and possess the following symmetries

$$c_{ijkl} = c_{jikl} = c_{ijlk} = c_{klij}, \quad (2.3)$$

see e.g. [12] for further details. The quantities

$$\varepsilon_{ij}(u) = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right) \quad (2.4)$$

are the small strain tensor components. From (2.3) and (2.4), it follows that the bilinear form can be rewritten in the following form,

$$a(u, v) = \int_{\Omega} \langle Cd(u), d(v) \rangle dx \quad (2.5)$$

where for $u \in [H^1(\Omega)]^2$, $u = (u_1, u_2)$,

$$d(u) = \left(\frac{\partial u_1}{\partial x_1}, \frac{\partial u_1}{\partial x_2}, \frac{\partial u_2}{\partial x_1}, \frac{\partial u_2}{\partial x_2} \right)^T \quad (2.6)$$

and C is the matrix consisting of the elasticity moduli,

$$C = \begin{bmatrix} c_{1111} & c_{1112} & c_{1121} & c_{1122} \\ c_{1211} & c_{1212} & c_{1221} & c_{1222} \\ c_{2111} & c_{2112} & c_{2121} & c_{2122} \\ c_{2211} & c_{2212} & c_{2221} & c_{2222} \end{bmatrix}. \quad (2.7)$$

As follows from (2.3), the matrix C is symmetric and also positive semidefinite, see e.g. [12]. In particular, it is positive definite for vectors $w = (w_{11}, w_{12}, w_{21}, w_{22})^T$ with a symmetry relation $w_{12} = w_{21}$. As an example, the matrix C corresponding to the case of plane strain with isotropic material has the form

$$C = \begin{bmatrix} \lambda + 2\mu & 0 & 0 & \lambda \\ 0 & \mu & \mu & 0 \\ 0 & \mu & \mu & 0 \\ \lambda & 0 & 0 & \lambda + 2\mu \end{bmatrix}, \quad (2.8)$$

where λ, μ are the Lamè moduli, which are positive numbers.

Further, we shall assume that \mathcal{T}_H is a finite element triangulation of Ω and that the coefficients a_{ij} and c_{ijkl} are constant on each element in τ_H . Now we can formulate two strengthened C.B.S. constant estimation problems.

$P_1 - P_1$ problem: The h -refinement gives a new division \mathcal{T}_h of Ω , which arises by dividing each $E \in \mathcal{T}_H$ into smaller triangles. How this is done will be outlined in the next section. For each element $E \in \mathcal{T}_H$, we consider the spaces

$$U_E = \{v \in C(E) : v \in P_1\} \quad (2.9)$$

$$U_E^h = \{v \in C(E) : v|_e \in P_1 \quad \forall e \in \mathcal{T}_h, e \subset E\} \quad (2.10)$$

$$V_E = \{v \in U_E^h : v(x) = 0 \quad \text{for all vertices } x \text{ of } E\}, \quad (2.11)$$

$$U_E^h = U_E \oplus V_E. \quad (2.12)$$

Here $C(E)$ denotes the set of continuous functions on E .

Then the $P_1 - P_1$ problem for anisotropic Laplacian is to find a nontrivial bound for the C.B.S. constant $\gamma_{E,1}$ such that for all $u \in U_E, v \in V_E$

$$|a_E(u, v)| \leq \gamma_{E,1} \sqrt{a_E(u, u)} \sqrt{a_E(v, v)}. \quad (2.13)$$

Here,

$$a_E(u, v) = \int_E \langle Dd(u), d(v) \rangle. \quad (2.14)$$

For the corresponding $P_1 - P_1$ problem for the anisotropic elasticity problem, (2.13) holds with

$$a_E(u, v) = \int_E \langle Cd(u), d(v) \rangle dx \quad (2.15)$$

and $u \in \hat{U}_E, v \in \hat{V}_E$, where

$$\hat{U}_E = \{v = (v_1, v_2) : v_i \in U_E \quad \text{for } i = 1, 2\}$$

$$\hat{V}_E = \{v = (v_1, v_2) : v_i \in V_E \quad \text{for } i = 1, 2\}$$

$P_1 - P_2$ problem: Consider now p -refinement, i.e. piecewise linear and quadratic functions over the elements of \mathcal{T}_H . For each element $E \in \mathcal{T}_H$, we consider the spaces




Figure 1: Division of the triangle into m^2 congruent ones, $h = H/m$, $m = 3$.

$$\begin{aligned}
 U_E &= \{v \in C(E) : v \in P_1\} \\
 U_E^p &= \{v \in C(E) : v \in P_2\} \\
 V_E &= \{v \in U_E^p : v(x) = 0 \text{ for all vertices } x \text{ of } E\} \\
 U_E^p &= U_E \oplus V_E.
 \end{aligned}$$

The $P_1 - P_2$ problem for anisotropic Laplacian is again to find a nontrivial bound for the C.B.S. constant $\gamma_{E,2}$ such that

$$|a_E(u, v)| \leq \gamma_{E,2} \sqrt{a_E(u, u)} \sqrt{a_E(v, v)}$$

for all $u \in U_E$, $v \in V_E$ and a_E defined by (2.14) The corresponding $P_1 - P_2$ problem for the elasticity can be defined in the same way as before.

3 Universal estimates for the $P_1 - P_1$ problem

For 2D problems, we shall consider triangular elements and the h -refinement of the type, which is illustrated in Figure 1. It means that now each coarse triangle is divided into m^2 smaller congruent triangles with edges which are m times shorter than the edges of the original coarse triangle. Moreover, each edge of the small triangle is parallel to some side of the original triangle. By the described division, we get from the original coarse triangulation with the mesh parameter H a new triangulation with the mesh parameter $h = H/m$.

The aim of this section is to prove the following theorem

Theorem 3.1 *Consider the bilinear forms (2.14) and (2.15) corresponding respectively to a general 2D anisotropic Laplacian or a general 2D anisotropic elasticity operator on Ω . Further let \mathcal{T}_H be a triangulation of Ω and assume that the problem*




Figure 2: The reference triangle and values of $\frac{\partial v}{\partial x_i}$.

coefficients are constant on the coarse elements $E \in \mathcal{T}_H$. Assume also that each element $E \in \mathcal{T}_H$ is divided into m^2 smaller congruent triangles in the described way. Then

$$\gamma_{E,1} \leq \sqrt{\frac{m^2 - 1}{m^2}}. \quad (3.1)$$

We shall prove Theorem 1 in several sub-steps in the following subsections. First we restrict our attention to a right angle isosceles reference triangle. For it, we first prove a universal estimate in the case of anisotropic Laplacian and a division of the coarse triangles into four smaller ones, i.e. for $m = 2$. This simple case is for illustration only, and is then extended to $m \geq 2$ and both anisotropic Laplacian and anisotropic elasticity. Finally, we show how to extend the estimates to the case of general triangles by an affine mapping of these triangles to the reference one.

3.1 Universal estimate of γ for a reference triangle and $m = 2$

As an illustration of our approach, we shall start with the simplest case of a reference triangle E with two axiparallel sides, see Figure 2. The triangles are ordered as indicated. We denote

$$\delta_i = \frac{\partial u}{\partial x_i}, \quad i = 1, 2, \quad d(u) = \delta = (\delta_1, \delta_2)^T \quad \text{for } u \in U_E, \quad (3.2)$$

$$\delta_i^{(k)} = \frac{\partial v}{\partial x_i} \Big|_{T_k}, \quad i = 1, 2, \quad d(v)|_{T_k} = d^{(k)} = (d_1^{(k)}, d_2^{(k)})^T \quad \text{for } v \in V_E. \quad (3.3)$$

Note that all quantities $\delta_i, d_i^{(k)}$ are constants. We shall exploit certain relations between $d_i^{(k)}$. These relations are induced by the fact that v is zero in the vertices (it gives $d_1^{(2)} = -d_1^{(1)}$ and $d_2^{(4)} = -d_2^{(1)}$) and the fact that some triangles share an axiparallel side (it gives $d_1^{(3)} = d_1^{(4)}$ and $d_2^{(2)} = d_2^{(3)}$). These relations are illustrated in Figure 2.

Now we can write

$$\begin{aligned}
a_E(u, v) &= \sum_{k=1}^4 \int_{T_k} \langle D\delta, d^{(k)} \rangle dx = \sum_{k=1}^4 \langle \hat{\delta}, d^{(k)} \rangle \Delta \\
&= [\hat{\delta}_1 (d_1^{(1)} - d_1^{(1)} + 2d_1^{(3)}) + \hat{\delta}_2 (d_2^{(1)} - d_2^{(1)} + 2d_2^{(2)})] \Delta \\
&= 2\Delta \langle \hat{\delta}, \hat{d} \rangle = 2\Delta \langle D\delta, \hat{d} \rangle \leq 2\Delta \|\delta\|_D \|\hat{d}\|_D
\end{aligned} \tag{3.4}$$

where $\hat{\delta} = D\delta = (\hat{\delta}_1, \hat{\delta}_2)^T$. Here we have used the fact that $\hat{\delta}_i$ are constant on E due to δ and D being constant on E . Above, we introduced $\|w\|_D = \sqrt{\langle Dw, w \rangle}$ for $w \in R^2$ and Δ denotes the area of the smaller triangles, i.e. $|E| = 4\Delta$.

In (3.4), we have also introduced an auxiliary vector $\hat{d} = (d_1^{(3)}, d_2^{(2)})^T$. Thus $\hat{d} = d^{(3)}$ and moreover

$$\hat{d} = d^{(1)} + d^{(2)} + d^{(4)}, \quad \|\hat{d}\|_D^2 \leq 3 \sum_{k \neq 3} \|d^{(k)}\|_D^2. \tag{3.5}$$

Thus

$$a_E(v, v) = \sum_{k=1}^4 \|d^{(k)}\|_D^2 \Delta \geq (1 + \frac{1}{3}) \|\hat{d}\|_D^2 \Delta, \tag{3.6}$$

$$a_E(u, u) = \|\delta\|_D^2 4\Delta. \tag{3.7}$$

From (3.4), (3.6) and (3.7), we therefore get

$$a_E(u, v) \leq \sqrt{\frac{3}{4}} \sqrt{a_E(u, u)} \sqrt{a_E(v, v)} \quad \forall u \in U_E, v \in V_E. \tag{3.8}$$

This means that (2.13) holds with $\gamma_{E,1} = \sqrt{\frac{3}{4}}$.

This estimate is in accordance with the estimates in [11] and [4].

Note 3.1 *The estimate (3.8) is valid for all coefficient matrices D . For a specific D , we can of course get a better estimate. For example, for $D = I$, it follows that*

$$a_E(v, v) = 2 \left[(d_1^{(1)})^2 + (d_1^{(3)})^2 + (d_2^{(1)})^2 + (d_2^{(2)})^2 \right] \Delta \geq 2\Delta \|\hat{d}\|_D^2. \tag{3.9}$$

From (3.4), (3.7) and (3.9), we now get

$$|a_E(u, v)| \leq \sqrt{\frac{1}{2}} \sqrt{a_E(u, u)} \sqrt{a_E(v, v)} \quad \forall u \in U_E, v \in V_E. \tag{3.10}$$

This estimate (3.10) is in accordance with the results in [3] and [11].




Figure 3: The reference triangle and $m = 3$. $I_1^0 = \{1, 2, 3\}$, $I_2^0 = \{1, 6, 9\}$, $I^* = \{4, 5, 8\}$

3.2 Universal estimates of γ for a reference triangle and $m \geq 2$.

The estimate obtained in Subsection 3.1 can be simply generalized to denser grid refinements. Let us consider the reference triangle and its division into m^2 triangles in the way, which is illustrated in Figure 3.

We shall again consider the derivatives of the functions $u \in U_E$ and $v \in V_E$. By considering the x_1 -parallel sides of the small triangles, we can see that

- there are m triangles T_k , $k \in I_1^0$ which do not share this side with other triangles. Because $v \in V_H$ is zero in the vertices, we get $\sum_{k \in I_1^0} d_1^{(k)} = 0$.
- the remaining triangles can be divided into pairs which share one x_1 -parallel side. Let I^* be the set of indices of the lower triangles from each pair.

For the x_2 -parallel sides of the small triangles, we observe a similar structure

- there are m triangles T_k , $k \in I_2^0$ which do not share this side with other triangles. Because $v \in V_H$ is zero in the vertices, we get $\sum_{k \in I_2^0} d_2^{(k)} = 0$.
- the remaining triangles can be divided into pairs which share one x_2 -parallel side. The set I^* introduced above now gives the indices of the left triangles from each pair.

For $u \in U_E$ and $v \in V_E$, we can now write

$$\begin{aligned}
 a_E(u, v) &= \sum_{k=1}^{m^2} \int_{T_k} \langle D\delta, d^{(k)} \rangle dx = \sum_{k=1}^{m^2} \langle \hat{\delta}, d^{(k)} \rangle \Delta \\
 &= 2\Delta \langle \hat{\delta}, \hat{d} \rangle = 2\Delta \langle D\delta, \hat{d} \rangle \leq 2\Delta \|\delta\|_D \|\hat{d}\|_D \quad (3.11)
 \end{aligned}$$

where the meaning of $\hat{\delta}$, Δ is the same as in Subsection 3.1, $\hat{\delta}$ is constant on E and

$$\hat{d} = \left(\sum_{k \in I^*} d_1^{(k)}, \sum_{k \in I^*} d_2^{(k)} \right)^T. \quad (3.12)$$

Thus,

$$\hat{d} = \sum_{k \in I^*} d^{(k)} \quad \text{and also} \quad \hat{d} = \sum_{k \in (I^*)^c} d^{(k)}, \quad (3.13)$$

where $(I^*)^c = \{1, \dots, m^2\} \setminus I^*$ is the complementary set to I^* .

Noting that the sets I^* and $(I^*)^c$ contain $(m^2 - m)/2$ and $(m^2 + m)/2$ indices, respectively, this gives the estimates

$$\|\hat{d}\|_D^2 \leq \frac{m(m-1)}{2} \sum_{k \in I^*} \|d^{(k)}\|_D^2, \quad (3.14)$$

$$\|\hat{d}\|_D^2 \leq \frac{m(m+1)}{2} \sum_{k \in (I^*)^c} \|d^{(k)}\|_D^2. \quad (3.15)$$

Thus,

$$a_E(v, v) = \sum_{k=1}^{m^2} \|d^{(k)}\|_D^2 \Delta \geq \left[\frac{2}{m(m-1)} + \frac{2}{m(m+1)} \right] \|\hat{d}\|_D^2 \Delta \quad (3.16)$$

$$a_E(u, u) = \|\delta\|_D^2 m^2 \Delta. \quad (3.17)$$

From (3.11), (3.16) and (3.17), we get then

$$a_E(u, v) \leq \sqrt{\frac{m^2-1}{m^2}} \sqrt{a_E(u, u)} \sqrt{a_E(v, v)}. \quad (3.18)$$

which implies the general result $\gamma_{E,1} \leq \sqrt{\frac{m^2-1}{m^2}}$.

3.3 Universal estimate of γ for the elasticity operator

In this subsection the reference triangle and its division will be the same as in the previous subsection but we shall consider the bilinear form and spaces corresponding to the elasticity operator. The expression of the elasticity bilinear form in formula (2.5) allows us to readily extend the previous results.

Let us denote

$$\delta = d(u) = (\delta_{11}, \delta_{12}, \delta_{21}, \delta_{22})^T, \quad \delta_{ij} = \frac{\partial u_i}{\partial x_j} \quad \text{for } u \in \hat{U}_E, \quad (3.19)$$

$$d^{(k)} = d(v)|_{T_k} = (d_{11}^{(k)}, d_{12}^{(k)}, d_{21}^{(k)}, d_{22}^{(k)})^T, \quad d_{ij}^{(k)} = \frac{\partial v_i}{\partial x_j} \Big|_{T_k} \quad \text{for } v \in \hat{V}_E. \quad (3.20)$$

Then all quantities δ_{ij} , $d_{ij}^{(k)}$ are constants and for $i, j = 1, 2$ there hold the same relations between the derivatives $d_{ij}^{(k)}$ as in Subsection 3.2. Thus

$$\begin{aligned} a_E(u, v) &= \sum_{k=1}^{m^2} \int_{T_k} \langle C\delta, d^{(k)} \rangle dx = \sum_{k=1}^{m^2} \langle \hat{\delta}, d^{(k)} \rangle \Delta \\ &= \sum_{i,j=1}^2 \hat{\delta}_{ij} 2 \sum_{k \in I^*} d_{ij}^{(k)} \cdot \Delta = \\ &= 2\Delta \langle \hat{\delta}, \hat{d} \rangle = 2\Delta \langle C\delta, \hat{d} \rangle \leq 2\Delta \|\delta\|_C \|\hat{d}\|_C \end{aligned} \quad (3.21)$$

where $\hat{\delta} = C\delta$ is again constant on E due to δ and C being constant, $\|z\|_C = \sqrt{\langle Cz, z \rangle}$ is the seminorm induced by C and

$$\hat{d} = \sum_{k \in I^*} d^{(k)} \quad \text{and also} \quad \hat{d} = \sum_{k \in (I^*)^c} d^{(k)}. \quad (3.22)$$

The above expressions for \hat{d} lead to the estimates

$$\|\hat{d}\|_C^2 \leq \frac{m(m-1)}{2} \sum_{k \in I^*} \|d^{(k)}\|_C^2 \quad (3.23)$$

$$\|\hat{d}\|_C^2 \leq \frac{m(m+1)}{2} \sum_{k \in (I^*)^c} \|d^{(k)}\|_C^2. \quad (3.24)$$

Thus,

$$a_E(v, v) = \sum_{k=1}^{m^2} \|d^{(k)}\|_C^2 \Delta \geq \left[\frac{2}{m(m+1)} + \frac{2}{m(m-1)} \right] \|\hat{d}\|_C^2 \Delta \quad (3.25)$$

$$a_E(u, u) = \|\delta\|_C^2 m^2 \Delta. \quad (3.26)$$

From (3.21), (3.25) and (3.26), we get again that $\gamma_{E,1} \leq \sqrt{\frac{m^2-1}{m^2}}$.




Figure 4: Mapping the reference triangle \tilde{E} to a general triangle E .

3.4 Universal estimate of γ for a general triangle

Now, we shall consider a general triangle E and its division to m^2 congruent smaller triangles with each side parallel to some side of the original triangle as well as the reference triangle \tilde{E} with vertices $(0,0)$, $(1,0)$, $(0,1)$ and its division in the same way. It is important to note that it is possible to find an affine mapping $F : \tilde{E} \rightarrow E$. This mapping will also map each smaller triangle from the division of \tilde{E} into a corresponding smaller triangle from the division of E , see Figure 4.

Let $p_i = (p_{i1}, p_{i2})$ be the vertices of E then the mapping $F : \tilde{x} \rightarrow x$ can be described analytically by the following relations

$$\begin{aligned} x_1 &= p_{11} + (p_{21} - p_{11})\tilde{x}_1 + (p_{31} - p_{11})\tilde{x}_2, \\ x_2 &= p_{12} + (p_{22} - p_{12})\tilde{x}_1 + (p_{32} - p_{12})\tilde{x}_2. \end{aligned}$$

Now, let us consider $u \in U_E$, $v \in V_E$. Then $\tilde{u} = u \circ F \in U_{\tilde{E}}$ and $\tilde{v} = v \circ F \in V_{\tilde{E}}$. Moreover, $a_E(u, u)$, $a_E(v, v)$ and $a_E(u, v)$ can be transformed to $\tilde{a}_{\tilde{E}}(\tilde{u}, \tilde{u})$, $\tilde{a}_{\tilde{E}}(\tilde{v}, \tilde{v})$ and $\tilde{a}_{\tilde{E}}(\tilde{u}, \tilde{v})$, see also [4]. The transformation $a \rightarrow \tilde{a}$ starts with the transformation of the derivatives. If $\tilde{u} = u \circ F$ then

$$\frac{\partial \tilde{u}}{\partial \tilde{x}_j}(\tilde{x}) = \frac{\partial u}{\partial x_1}(x) (p_{j+1,1} - p_{1,1}) + \frac{\partial u}{\partial x_2}(x) (p_{j+1,2} - p_{1,2}).$$

It gives $d(\tilde{u}) = G^T d(u)$ where G is the Jacobian matrix of F ,

$$G = \begin{bmatrix} p_{21} - p_{11} & p_{31} - p_{11} \\ p_{22} - p_{12} & p_{32} - p_{12} \end{bmatrix}.$$

Now, we can write

$$\begin{aligned} a_E(u, u) &= \int_E \langle Dd(u), d(u) \rangle dx = \int_{\tilde{E}} \langle DG^{-T}d(\tilde{u}), G^{-T}d(\tilde{u}) \rangle |\det(G)| d\tilde{x} \\ &= \int_{\tilde{E}} \langle \tilde{D}d(\tilde{u}), d(\tilde{u}) \rangle d\tilde{x} = \tilde{a}_{\tilde{E}}(\tilde{u}, \tilde{u}), \quad \tilde{u} = u \circ F, \end{aligned}$$

where $\tilde{D} = |\det(G)| G^{-1}DG^{-T}$ is a transformed coefficient matrix, which is symmetric and positive definite, and

$$\tilde{a}_{\tilde{E}}(\tilde{u}, \tilde{v}) = \int_{\tilde{E}} \langle \tilde{D}d(\tilde{u}), d(\tilde{v}) \rangle d\tilde{x} \quad \forall \tilde{u} \in U_{\tilde{E}}, \tilde{v} \in V_{\tilde{E}}. \quad (3.27)$$

Similarly, we can show that

$$a_E(v, v) = \tilde{a}_{\tilde{E}}(\tilde{v}, \tilde{v}) \quad \text{and} \quad a_E(u, v) = \tilde{a}_{\tilde{E}}(\tilde{u}, \tilde{v}).$$

For $u \in U_E, v \in V_E$, we get $\tilde{u} \in U_{\tilde{E}}$ and $\tilde{v} \in V_{\tilde{E}}$. Using now the universal estimate of γ , which has been proved for the case of the reference triangle and arbitrary symmetric positive definite coefficient matrix, we get

$$\begin{aligned} |a_E(u, v)| &= |\tilde{a}_{\tilde{E}}(\tilde{u}, \tilde{v})| \\ &\leq \sqrt{\frac{m^2-1}{m^2}} \sqrt{\tilde{a}_{\tilde{E}}(\tilde{u}, \tilde{u})} \sqrt{\tilde{a}_{\tilde{E}}(\tilde{v}, \tilde{v})} \\ &= \sqrt{\frac{m^2-1}{m^2}} \sqrt{a_E(u, u)} \sqrt{a_E(v, v)}. \end{aligned}$$

The extension of this transformation technique to the case of general elasticity with the bilinear form (2.5) is straightforward. The transformed matrix \tilde{C} will be in the form $\tilde{C} = |\det(G)| G_2^{-1}CG_2^{-T}$, where G_2 is a block diagonal 4×4 matrix with the diagonal blocks equal to G .

Remark 3.1 The universal estimates of the C.B.S. constant have been proven for a special grid refinement. For other types of division of triangles, it may be impossible to derive a nontrivial universal estimate of the C.B.S. inequality. For example, for the division of triangles illustrated in Figure 5 and 2D elasticity (plane strain), we do not get a nontrivial estimate, as the C.B.S. constant γ depends on the Poisson ratio ν and γ tends to unity as ν goes to $1/2$, see [8], [10].




Figure 5: A different type of division of the triangle.

The technique of derivation of the universal estimates of the C.B.S. constant exploited in Section 3, can be used also for 3D anisotropic Laplacian and 3D elasticity assuming the use of linear tetrahedral finite elements, see [9].

4 An algebraic approach to derive the $P1-P2$ C.B.S. constant

Given a triangular element E with angles α, β, γ which has been regularly refined in four subtriangles, the local assembled $P1$ and $P2$ matrices have the form (see e.g. [5])

$$A_{H/2}^{(1)} = \frac{1}{2} \begin{bmatrix} 2d & -2c & -2b & 0 & -a & -a \\ -2c & 2d & -2a & -b & 0 & -b \\ -2b & -2a & 2d & -c & -c & 0 \\ 0 & -b & -c & b+c & 0 & 0 \\ -a & 0 & -c & 0 & a+c & 0 \\ -a & -b & 0 & 0 & 0 & a+b \end{bmatrix}$$

which is the matrix assembled from the element matrix for the four subtriangles corresponding to piecewise linear basis functions, and

$$A_H^{(2)} = \frac{1}{6} \begin{bmatrix} 8d & -8c & -8b & 0 & -4a & -4a \\ -8c & 8d & -8a & -4b & 0 & -4b \\ -8b & -8a & 8d & -4c & -4c & 0 \\ 0 & -4b & -4c & 3(b+c) & c & b \\ -4a & 0 & -4c & a & 3(a+c) & a \\ -4a & -4b & 0 & b & a & 3(a+b) \end{bmatrix}$$

which is the local finite element matrix corresponding to quadratic basis functions on E . Here $a = \cot \alpha$, $b = \cot \beta$, $c = \cot \gamma$ and $d = a + b + c$.

From these matrices follows the relation

$$A_H^{(2)} = \frac{4}{3} A_{H/2}^{(1)} - N, \quad (4.28)$$

where

$$N = \frac{1}{3} \begin{bmatrix} 0 & 0 \\ 0 & A_H^{(1)} \end{bmatrix} \quad (4.29)$$

and

$$A_H^{(1)} = \frac{1}{2} \begin{bmatrix} (b+c) & -c & -b \\ -c & (a+c) & -a \\ -b & -a & (a+b) \end{bmatrix}$$

which latter is the local finite element matrix for the vertex nodes of E , corresponding to linear basis functions.

Remark 4.1 The relation

$$A_H^{(2)} = \frac{4}{3}A_{H/2}^{(1)} = \frac{1}{3} \begin{bmatrix} 0 & 0 \\ 0 & A_H^{(1)} \end{bmatrix}$$

could have been derived in an alternate way, using one step of the classical Richardson H -extrapolation to derive a locally third order accurate scheme ($A_H^{(2)}$) from the two locally second order accurate schemes ($A_{H/2}^{(1)}$ and $A_H^{(1)}$), by elimination of the $O(H^2)$ error components.

Now relation (4.28) can be used to derive a relation between the C.B.S. constants for the $P1 - P2$ and $P1 - P1$ hierarchical finite element spaces combinations.

Theorem 4.1 For any regularly refined finite element mesh into congruent elements, for which (4.28) holds one has

$$\gamma_2^2 = \frac{4}{3}\gamma_1^2 \quad (4.30)$$

where γ_1, γ_2 are the C.B.S. constants for piecewise linear and piecewise quadratic finite elements, respectively.

Proof. Since all block parts of the first two matrices in (4.28), except the lower right block are equal we can take Schur complements, and (4.29) shows that

$$S^{(2)} = \frac{4}{3}S^{(1)} - \frac{1}{3}A_H^{(1)}.$$

Hence

$$\frac{x_2^T S^{(2)} x_2}{x_2^T A_H^{(1)} x_2} = \frac{4}{3} \frac{x_2^T S^{(1)} x_2}{x_2^T A_H^{(1)} x_2} - \frac{1}{3}$$

i.e.

$$\begin{aligned} 1 - \gamma_2^2 &= \min_{x_2} \frac{x_2^T S^{(2)} x_2}{x_2^T A_H^{(1)} x_2} = \frac{4}{3} \min_{x_2} \frac{x_2^T S^{(1)} x_2}{x_2^T A_H^{(1)} x_2} - \frac{1}{3} \\ &= \frac{4}{3}(1 - \gamma_1^2) - \frac{1}{3} = 1 - \frac{4}{3}\gamma_1^2. \end{aligned}$$

□

Remark 4.2 The relation (4.30) has been shown previously in [11], [4] using a more involved derivation. Using the already derived expressions for γ_1 in Section 3, we have then also a general expression for $\gamma_2 = \frac{2}{\sqrt{3}}\gamma_1$. We note that γ_2 can take values arbitrarily close to the unit number for degenerate triangles, or equivalently, for certain anisotropies of the coefficients in the differential operator.

5 Concluding remarks

The matrices of triangular finite elements derived from the two-level methods can be partitioned in a two-by-two block matrix from $[A_{ij}]_{i,j=1}^2$, where A_{11} corresponds to the node-points added in the refinement process, A_{22} corresponds to the original vertex nodes and A_{12} (A_{21}) corresponds to the coupling between the two finite element subspaces used.

The following matrix relation holds

$$u^T A_{12} v \leq \gamma \{u^T A_{11} u v^T A_{22} v\}^{\frac{1}{2}}$$

where $u = (\alpha_1, \alpha_2, \alpha_3, 0, 0, 0)^T$, $v = (0, 0, 0, \beta_1, \beta_2, \beta_3)^T$ are the corresponding orthogonal vectors.

Using the hierarchical form of the matrices one can precondition the block matrix by its block-diagonal part, $\begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix}$, in which case the condition number becomes $(1 + \gamma)/(1 - \gamma)$. Alternatively, one can precondition the reduced (Schur complement) system, $A_{22} - A_{21}A_{11}^{-1}A_{12}$ with A_{22} , in which case the condition number becomes $1/(1 - \gamma^2)$. Here it should be noted that the Schur complements will be the same as for the hierarchical basis even if the reduction takes place from the standard basis function matrix (i.e. from $A_H^{(2)}$ or $A_{H/m}^{(1)}$). For further details, see e.g. [5]. Systems with the matrix A_{22} which occur in this preconditioning can be solved using the same method recursively, unless one finds that the matrix $A_{11}^{(1)}$ is already sufficiently coarse to use a direct solution method or a simpler iterative method

Matrix A_{11} is frequently well-conditioned and systems with it can be solved efficiently by some simple iteration method. However, for nearly degenerate triangles or, equivalently, strongly anisotropic coefficients it becomes very ill-conditioned and requires some special element by element preconditioner, see e.g. [7] for details.

References

- [1] B. Achchab, O. Axelsson L. Laayouni and A. Souissi, Strengthened Cauchy-Bunyakowski-Schwarz inequality for a three dimensional elasticity system, Numer. Linear Algebra Appl. 8 (2001), 191-205.
- [2] B. Achchab and J. Maitre, Estimate of the constant in two strengthened C.B.S. inequalities for F.E.M. systems of 2D elasticity. Applications to multilevel methods and a posteriori error estimators, Numer. Linear Algebra Appl., 3(1996), 147-159.

- [3] O. Axelsson, On multigrid methods of the two-level type, In: W. Hackbusch, U Trottenberg eds., *Multigrid Methods, Lecture Notes in Math. 960*, Springer-Verlag, Berlin 1982, pp. 352-367.
- [4] O. Axelsson, Stabilization of algebraic multilevel iteration methods; additive methods, *Numerical Algorithms* 21(1999) 23-47.
- [5] O. Axelsson and V.A. Barker, *Finite Element Solution of Boundary Value Problems. Theory and Computation*, Academic Press, Orlando, 1984. Reprinted as *SIAM Classics in Applied Mathematics* 35, SIAM, Philadelphia, 2001.
- [6] O. Axelsson and I. Gustafsson, Preconditioning and two-level factorization methods of arbitrary degree of approximation, [Report 8120 (July 1981), Department of Mathematics, University of Nijmegen, The Netherlands], *Math. Comp.* 40 (1983), 219-242.
- [7] O. Axelsson and A. Padiy, On the additive version of the algebraic multilevel iteration method for anisotropic elliptic problems, *SIAM J. Sci. Comp.*, 20 (1999), 1807-1830.
- [8] R. Blaheta: Adaptive composite grid methods for problems of plasticity. *Mathematics and Computers in Simulation* 50 (1999) 123 - 134.
- [9] R. Blaheta: Estimates of the C.B.S. inequality constant for 3D anisotropic Laplace and elasticity operators, in progress, partly presented at the Numerical Analysis Seminar, KUN Nijmegen, November 2000.
- [10] M. Jung, J.F. Maitre: Some remarks on the constant in the strengthened C.B.S. inequality: Applications to h- and p- hierarchical FE discretizations of elasticity problems, Preprint SFB 383/97-15, TU Chemnitz - Zwickau 1997.
- [11] J.F. Maitre and F. Musy, The contraction number of a class of two-level methods, an exact evaluation for some finite element subspaces and model problems. In: W. Hackbusch, U Trottenberg eds., *Multigrid Methods, Lecture Notes in Math. 960*, Springer-Verlag, Berlin 1982, pp. 535-544.
- [12] J. Nečas, I. Hlaváček: *Mathematical Theory of Elastic and Elasto-plastic Bodies: An Introduction*, Elsevier, Amsterdam 1981.
- [13] S.D. Margenov: Upper bound on the constant in the strengthened C.B.S. inequality for FEM 2D elasticity equations, *Numer. Linear Algebra Appl.* 1(1994), 65-74.