

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/197720>

Please be advised that this information was generated on 2021-09-27 and may be subject to change.

Synchronizing non-deterministic finite automata

Henk Don¹ and Hans Zantema^{2,3}

¹Department of Mathematics, Vrije Universiteit Amsterdam, De Boelelaan 1081a,
1081 HV Amsterdam, The Netherlands, email: h.don@vu.nl

²Department of Computer Science, TU Eindhoven, P.O. Box 513,, 5600 MB
Eindhoven, The Netherlands, email: h.zantema@tue.nl

³Radboud University Nijmegen, P.O. Box 9010, , 6500 GL Nijmegen, The
Netherlands

October 23, 2018

Abstract

In this paper, we show that every D3-directing CNFA can be mapped uniquely to a DFA with the same synchronizing word length. This implies that Černý's conjecture generalizes to CNFAs and that the general upper bound for the length of a shortest D3-directing word is equal to the Pin-Frankl bound for DFAs. As a second consequence, for several classes of CNFAs sharper bounds are established. Finally, our results allow us to detect all critical CNFAs on at most 6 states. It turns out that only very few critical CNFAs exist.

1 Introduction and preliminaries

In this paper we study synchronization of non-deterministic finite automata (NFAs). As is the case for deterministic finite automata (DFAs), symbols define functions on the state set Q . However, in an NFA symbols are allowed to send a state to a subset of Q , rather than to a single state. An NFA is called complete if these subsets are non-empty. This basically says that in every state, every symbol has at least one out-going edge. Formally, a *complete non-deterministic finite automaton (CNFA)* \mathcal{A} over a finite alphabet Σ consists of a finite set Q of states and a map $\delta : Q \times \Sigma \rightarrow 2^Q \setminus \{\emptyset\}$. We denote the number of states by $|\mathcal{A}|$ or by $|Q|$.

A DFA is called synchronizing if there exists a word that sends every state to the same fixed state. In 1964 Černý [4] conjectured that a synchronizing DFA on n states always admits a synchronizing (or directing, reset) word of length at most $(n - 1)^2$. He gave a sequence C_n of DFAs in which the shortest synchronizing word attains this bound. In this paper, we denote

the maximal length of a shortest synchronizing word in an n -state DFA by $d(n)$. The best known bounds for $d(n)$ are


$$(n - 1)^2 \leq d(n) \leq \frac{n^3 - n}{6}. \quad (1)$$

For a proof of the upper bound, we refer to [14]. A DFA on n states is *critical* if its shortest synchronizing word has length $(n - 1)^2$; it is *super-critical* if its shortest synchronizing word has length $> (n - 1)^2$. So Černý's conjecture states that no super-critical DFAs exist. It turns out that there are not too many critical DFAs. Investigation of all critical DFAs with less than 7 states but unrestricted alphabet was recently completed [8, 6]. DFAs without copies of the same symbol and without the identity are called *basic*. For $n = 3, 4, 5, 6$, only 31 basic critical DFAs exist up to isomorphism. So critical DFAs are very infrequent, as the total number of basic DFAs on n states is $2^{n^n - 1}$, including isomorphisms. For $n \geq 7$, the only known examples are from Černý's sequence.

For $S \subseteq Q$ and $w \in \Sigma^*$, let Sw be the set of all states where one can end when starting in some state $q \in S$ and reading the symbols in w consecutively. Write qw for $\{q\}w$. Formal definitions will be given in Section 1.1. A DFA is synchronizing if there exists $w \in \Sigma^*$ and $q_s \in Q$ such that $qw = q_s$ for all $q \in Q$. There are several ways to generalize this concept of synchronization to CNFAs, see [12]. In this paper, we study CNFAs known in the literature as D3-directing. This notion is defined as follows:

Definition 1. A CNFA (Q, Σ, δ) is called *D3-directing* if there exists a word $w \in \Sigma^*$ and a state q_s such that $q_s \in qw$ for all $q \in Q$. The word w is called a *D3-directing word*.

An example of a D3-directing CNFA is depicted below. There exist several D3-directing words of length four, but no shorter ones. An example is $w = baba$, which gives $1w = \{1, 3\}$, $2w = \{1, 2\}$ and $3w = \{1, 2, 3\}$. The synchronizing state for this word is 1. Another D3-directing word is $v = aabb$, for which $1v = 3v = \{1, 2, 3\}$ and $2v = 2$. Here the synchronizing state is 2.



A word is D3-directing if starting in any state q , there exists a path labelled by w that ends in q_s . For DFAs this notion coincides with a synchronizing word. If a CNFA \mathcal{A} is D3-directing, a natural question is to find the length of a shortest D3-directing word. We denote this length by $d_3(\mathcal{A})$. Furthermore, we denote by $cd_3(n)$ the worst case, i.e. we let $\text{CDir}(3)$ be the collection of all D3-directing CNFAs and define

$$cd_3(n) = \max \{d_3(\mathcal{A}) : \mathcal{A} \in \text{CDir}(3), |\mathcal{A}| = n\}. \quad (2)$$

In [12] it is shown that for all $n \geq 1$,

$$(n-1)^2 \leq cd_3(n) \leq \frac{1}{2}n(n-1)(n-2) + 1. \quad (3)$$

The lower bound follows from the fact that every DFA is also a CNFA and that for DFAs the notions of synchronization and D3-directability coincide. As far as we are aware, these bounds are still the sharpest known for CNFAs, although sharper results were recently obtained for the essentially equivalent problem of bounding lengths of column-primitive products of matrices [5]. Analogous to DFAs, a D3-directing CNFA is called critical if its shortest D3-directing word has length $(n-1)^2$, and super-critical if it has length $> (n-1)^2$. In the current paper, we will prove that in fact $cd_3(n) = d(n)$, which immediately sharpens the upper bound for $cd_3(n)$ to $(n^3 - n)/6$. Our result also implies that Černý's conjecture is equivalent to the following:

Conjecture 1. *Every D3-directing CNFA with n states admits a D3-directing word of length at most $(n-1)^2$.*

The main ingredient to prove that $cd_3(n) = d(n)$ is a splitting transformation **Split** that maps a CNFA to a DFA. Every D3-directing CNFA \mathcal{A} is transformed into a synchronizing DFA $\text{Split}(\mathcal{A})$, preserving the shortest D3-directing word length. For several classes of DFAs, the Černý conjecture has been established, or sharper bounds than the general bounds have been proven, see for example [1, 2, 7, 9, 10, 13, 16]. If **Split**(\mathcal{A}) satisfies the properties for one of these classes, then the sharper results for **Split**(\mathcal{A}) also apply to the CNFA \mathcal{A} . This observation gives rise to generalize several properties of DFAs into notions for CNFAs and to check if these generalized properties are preserved under **Split**. In this way, we derive sharper upper bounds on the maximal D3-directing word length for several classes of CNFAs.

Finally, in this paper we search for examples of critical D3-directing CNFAs. Note that the number of CNFAs without identical symbols on n states is huge, namely $2^{(2^n - 1)^n}$ when we include isomorphisms. Therefore an exhaustive search is problematic, even for small n . However, since we know that every critical CNFA can be transformed into a critical DFA, we can try to find critical examples by reversing the transformation **Split**. Since all critical DFAs on ≤ 6 states are known, this approach allows us to identify all critical CNFAs on ≤ 6 states. Applying this strategy to the other known critical

DFA, the only critical CNFAs we find are small modifications of Černý's sequence.

1.1 Preliminaries

In this section we present our formal definitions and notation which will be slightly different from the traditional notation, as we avoid the use of the transition function. A symbol (or letter, label) a in a CNFA will be a function $a : Q \rightarrow 2^Q \setminus \{\emptyset\}$, and we denote $a(q)$ by qa . A symbol extends (denoting the extension by a as well) to a function $a : 2^Q \rightarrow 2^Q \setminus \{\emptyset\}$ by $Sa = \bigcup_{q \in S} qa$. The set of all letters on Q that can be obtained in this way is denoted $T(Q)$, which is a strict subset of the set of all functions from 2^Q to $2^Q \setminus \{\emptyset\}$. The set of all possible symbols in a DFA on its turn is a subset of $T(Q)$:

$$T^d(Q) = \{a \in T(Q) : \forall q \in Q |qa| = 1\}.$$

A CNFA \mathcal{A} is defined to be a pair (Q, Σ) , where $\Sigma \subseteq T(Q)$. Similarly a DFA is a pair (Q, Σ) with $\Sigma \subseteq T^d(Q)$. Note that these definitions do not allow for two symbols that act exactly in the same way: if a is a possible symbol, then either $a \in \Sigma$ or $a \notin \Sigma$.

A symbol $a \in T(Q)$ induces a directed graph G_a with vertex set Q and can therefore be viewed as a subset of $Q \times Q$:

$$a = \{(q, p) : q \in Q, p \in qa\},$$

so a is identified with the set of all edges in G_a . This point of view is used to define set relations and operations like inclusion and union on $T(Q)$. For example, if $a, b \in T(Q)$, then

$$a \cup b = \{(q, p) : q \in Q, p \in qa \cup qb\}.$$

Suppose $\mathcal{A} = (Q, \Sigma)$ is a CNFA and $a, b \in \Sigma$ are such that $a \subseteq b$. If \mathcal{A} is D3-directing, then the automaton $(Q, \Sigma \setminus \{a\})$ is D3-directing as well with the same shortest synchronizing word length. Also the identity symbol has no influence on synchronization. Therefore, a CNFA is called *basic* if it has no identity symbol and no symbol is contained in another one. For DFAs this coincides with the existing notion of basic.

If $\mathcal{A} = (Q, \Sigma)$ and $\mathcal{B} = (Q, \Gamma)$ are CNFAs, we say that \mathcal{B} is contained in \mathcal{A} and write $\mathcal{B} \subseteq \mathcal{A}$ if for all $b \in \Gamma$ there exists $a \in \Sigma$ such that $b \subseteq a$. Alternatively, we say that \mathcal{A} is an extension of \mathcal{B} . If $\mathcal{B} \subseteq \mathcal{A}$ and $\mathcal{B} \neq \mathcal{A}$, we say that \mathcal{A} strictly contains (or is a strict extension of) \mathcal{B} . A critical CNFA is *minimal* if it is not the strict extension of another critical CNFA; it is *maximal* if it does not admit a basic critical strict extension.

Finally, for $w = w_1 \dots w_k \in \Sigma^*$ and $S \subseteq Q$, define Sw inductively by $S\epsilon = S$ and $Sw = (Sw_1 \dots w_{k-1})w_k$. So a word also is a function on 2^Q , being the composition of the transformations by each of its letters. Therefore also the transition monoid Σ^* is contained in $T(Q)$.

2 Transforming a CNFA into a DFA, preserving D3-directing word length

In this section we present the transformation `Split` and explore some of its properties. We note that similar but less explicit ideas were recently used in [3, 11] to give bounds on the length of a positive product in a primitive set of matrices. We start by introducing a parametrized version of our transformation:

Definition 2. Let $\mathcal{A} = (Q, \Sigma)$ be a CNFA. Fix $q_{split} \in Q, a \in \Sigma$ and denote the set $q_{split}a$ by $\{q_1, \dots, q_m\}$. Define new symbols a_1, \dots, a_m on Q as follows:

$$q_{split}a_i := q_i, \quad \text{and} \quad qa_i := qa \quad \text{for } q \neq q_{split},$$

and a new alphabet $\Gamma = \{a_1, \dots, a_m\} \cup (\Sigma \setminus \{a\})$. The CNFA (Q, Γ) will be denoted $\text{Split}(\mathcal{A}, q_{split}, a)$.

The idea of this transformation is that we want to make a CNFA \mathcal{A} ‘more deterministic’. If $|q_{split}a| \geq 2$, then multiple outgoing edges in the state q_{split} are labelled by the symbol a . So we could say that a offers a choice in q_{split} . For each possible choice, we introduce a new symbol that is deterministic in q_{split} and behaves as a in all other states. If $|q_{split}a| = 1$, then \mathcal{A} is not changed by the transformation. This definition immediately implies the following properties:

Lemma 1. Let $\mathcal{A} = (Q, \Sigma)$ be a CNFA. Fix $q_{split} \in Q, a_s \in \Sigma$ and let (Q, Γ) be $\text{Split}(\mathcal{A}, q_{split}, a_s)$. Let $c \in T^d(Q)$ be an arbitrary deterministic symbol. Then

1. for all $b \in \Gamma$, there exists $a \in \Sigma$ such that $b \subseteq a$,
2. (there exists $a \in \Sigma$ such that $c \subseteq a$) \iff (there exists $b \in \Gamma$ such that $c \subseteq b$).

Proof. Let $b \in \Gamma$, we will find a with the property claimed in the lemma. If $b \in \Gamma \cap \Sigma$, take $a = b$. If $b \in \Gamma \setminus \Sigma$, then b is one of the new symbols a_1, \dots, a_m . In this case take $a = a_s$. Then $qb = qa$ for $q \neq q_{split}$ and $qb \in qa$ for $q = q_{split}$. This proves the first statement.

Suppose $a \in \Sigma$ such that $c \subseteq a$, so $qc \in qa$ for all q . Then $q_{split}c \in q_{split}a$. By Definition 2, there exists $b \in \Gamma$ such that $q_{split}b = q_{split}c$ and $qb = qa$ for $q \neq q_{split}$. This means $c \subseteq b$. Now suppose $b \in \Gamma$ such that $c \subseteq b$. By statement 1 of the lemma, there exists $a \in \Sigma$ such that $b \subseteq a$, which implies $c \subseteq a$. \square

The parametrized `Split` preserves synchronization properties, as is shown in the next lemma.

Lemma 2. Let $\mathcal{A} = (Q, \Sigma)$ be a CNFA and let $\mathcal{B} = \text{Split}(\mathcal{A}, q_{\text{split}}, a)$ for some $q_{\text{split}} \in Q, a \in \Sigma$. Then

1. \mathcal{A} is D3-directing if and only if \mathcal{B} is D3-directing,
2. If \mathcal{A} and \mathcal{B} are D3-directing, then $d_3(\mathcal{A}) = d_3(\mathcal{B})$.

Proof. Let Γ be the alphabet of \mathcal{B} and denote the new labels by a_1, \dots, a_m . First assume that \mathcal{A} is D3-directing. There exist $w \in \Sigma^*$ and $q_s \in Q$ such that $q_s \in qw$ for all $q \in Q = \{q_1, \dots, q_n\}$. From each state q there exists a path labelled by $w = w_1 \dots w_{|w|}$ that ends in q_s :

$$\begin{array}{ccccccc} q_1 = q_1^0 & \xrightarrow{w_1} & q_1^1 & \xrightarrow{w_2} & \dots & \xrightarrow{w_{|w|}} & q_1^{|w|} = q_s \\ q_2 = q_2^0 & \xrightarrow{w_1} & q_2^1 & \xrightarrow{w_2} & \dots & \xrightarrow{w_{|w|}} & q_2^{|w|} = q_s \\ & & \vdots & & & & \\ q_n = q_n^0 & \xrightarrow{w_1} & q_n^1 & \xrightarrow{w_2} & \dots & \xrightarrow{w_{|w|}} & q_n^{|w|} = q_s \end{array}$$

We will construct a word $\tilde{w} = \tilde{w}_1 \dots \tilde{w}_{|w|} \in \Gamma^*$ which follows the same paths. We may assume that paths do not diverge again once they have met, i.e. if $q_i^t = q_j^t$, then $q_i^{t+1} = q_j^{t+1}$.

Let $Q_t = \{q_1^t, \dots, q_n^t\}$ and suppose $w_t = a$ for some $1 \leq t \leq |w|$. If $q_{\text{split}} \notin Q_{t-1}$, define \tilde{w}_t to be a_1 . Then $qw_t = q\tilde{w}_t$ for all $q \in Q_{t-1}$. If $q_{\text{split}} \in Q_{t-1}$, then $q_i^{t-1} = q_{\text{split}}$ for some $1 \leq i \leq n$. This means $q_i^t \in q_{\text{split}}a$, so there exists $\tilde{a} \in \Gamma$ such that $q_i^t = q_{\text{split}}\tilde{a}$. Define \tilde{w}_t to be \tilde{a} . Finally, for all $w_t \neq a$, let $\tilde{w}_t = w_t$. Then $q_s \in q\tilde{w}$ for all $q \in Q$, so \tilde{w} is a D3-directing word for \mathcal{B} .

Now assume that \mathcal{B} is D3-directing with D3-directing word $w \in \Gamma^*$ and synchronizing state q_s . By repeated application of Lemma 1 (replacing every symbol of w that is not in Σ by a), it follows that there exists $\tilde{w} \in \Sigma^*$ such that $qw \subseteq q\tilde{w}$ for all q . Therefore $q_s \in q\tilde{w}$ and the word \tilde{w} is D3-directing for \mathcal{A} .

The above arguments prove the first statement of the lemma. Clearly, rewriting a D3-directing word from Σ^* to Γ^* and vice versa preserves the length. This implies the second statement. \square

Next we investigate the result of applying consecutive parametrized Split transformations to all non-deterministic symbols in a CNFA \mathcal{A} . We will show that this terminates and that the result is a uniquely defined DFA.

Lemma 3. Let \mathcal{A}_0 be a CNFA, and repeat the following. If $\mathcal{A}_k = (Q, \Sigma_k, \delta)$ is not a DFA, choose $a_k \in \Sigma_k$ and $q_k \in Q$ for which $|q_k a_k| \geq 2$. Let $\mathcal{A}_{k+1} = \text{Split}(\mathcal{A}_k, q_k, a_k)$. Then

1. There exists $k \geq 0$ where this process ends such that \mathcal{A}_k is a DFA.

2. The resulting DFA does not depend on the choices of a_k and q_k .

Proof. If \mathcal{A}_k is a DFA, then $|qa| = 1$ for all $q \in Q$ and $a \in \Sigma_k$. If \mathcal{A}_k is not a DFA, then

$$\prod_{q \in Q} \prod_{a \in \Sigma_k} |qa| > \prod_{q \in Q} \prod_{a \in \Sigma_{k+1}} |qa| \geq 1.$$

As this integer sequence is strictly decreasing, it ends in 1, i.e. there exists $k \geq 0$ such that the k th term is equal to 1. This is equivalent to the first claim of the lemma.

For the second claim, choose k such that \mathcal{A}_k is a DFA. Let $c \in T^d(Q)$ be an arbitrary deterministic symbol. By repeated application of the second statement of Lemma 1, it follows that $c \in \Sigma_k$ if and only if there exists $a \in \Sigma_0$ for which $c \subseteq a$. Therefore the DFA \mathcal{A}_k does not depend on the splitting choices, proving the second statement. \square

Definition 3. Let $\mathcal{A} = (Q, \Sigma)$ be a CNFA. The unique DFA that is produced by repeated application of the parametrized `Split` will be called `Split`(\mathcal{A}).

Lemma 3 guarantees that `Split`(\mathcal{A}) is well-defined. Extension of Lemma 1 leads to the following characterization:

Lemma 4. Let $\mathcal{A} = (Q, \Sigma)$ be a CNFA. Denote the DFA `Split`(\mathcal{A}) by (Q, Γ) . Let $b \in T^d(Q)$ be an arbitrary deterministic symbol. Then $b \in \Gamma$ if and only if there exists $a \in \Sigma$ such that $b \subseteq a$.

Proof. If $b \in \Gamma$, repeatedly apply the first statement of Lemma 1. If there exists $a \in \Sigma$ such that $b \subseteq a$, then repeated application of the second statement of Lemma 1 proves existence of $b' \in \Gamma$ such that $b \subseteq b'$. Since both b and b' are deterministic symbols, $b = b'$, so $b \in \Gamma$. \square

Moreover, we have the following:

Corollary 1. If $\mathcal{A} = (Q, \Sigma)$ is a D3-directing CNFA, then $d_3(\mathcal{A}) = d(\text{Split}(\mathcal{A}))$.

Proof. This is an immediate consequence of Lemma 2. \square

Now also the main result of this section is straightforward:

Theorem 1. The maximal shortest D3-directing word length for DFAs is the same as for CNFAs, i.e.

$$d(n) = cd_3(n).$$

Proof. Since every DFA is also a CNFA and the notions of synchronization and D3-directedness coincide for DFAs, it follows that $d(n) \leq cd_3(n)$. By Corollary 1 every CNFA \mathcal{A} has a corresponding DFA `Split`(\mathcal{A}) with the same shortest D3-directing word length. Therefore $cd_3(n) \leq d(n)$. \square

This theorem establishes equivalence of Černý's conjecture to Conjecture 1. It also implies the following sharpening of the upper bound for $cd_3(n)$:

Corollary 2. $cd_3(n) \leq \frac{n^3 - n}{6}$.

3 Sharper bounds for several classes of CNFAs

For several classes of DFAs the Černý conjecture has been settled, or at least better upper bounds than the cubic one for the general case have been obtained. If the **Split** transform reduces a CNFA to a DFA that belongs to one of these classes, then as a direct consequence we obtain improved bounds for the D3-directing length in the CNFA. In this section we present a couple of results of this type.

The general pattern of the arguments in this section is as follows. First we give the definition of a property \mathcal{P} for DFAs, together with references to the best known upper bound $u_{\mathcal{P}}$ for synchronization lengths in DFAs satisfying \mathcal{P} . Then we give a natural extension of \mathcal{P} to the class of CNFAs. Finally, we show that every CNFA \mathcal{A} satisfying \mathcal{P} is reduced to a DFA $\text{Split}(\mathcal{A})$ satisfying \mathcal{P} . Corollary 1 then guarantees that the length of the shortest D3-directing word in \mathcal{A} is at most $u_{\mathcal{P}}$.

3.1 Cyclic automata

A DFA $\mathcal{A} = (Q, \Sigma)$ is *cyclic* if one of the letters in Σ acts as a cyclic permutation on Q .

Definition 4. A DFA $\mathcal{A} = (Q, \Sigma)$ with $|Q| = n$ is called *cyclic* if there exists $a \in \Sigma$ such that for all $q \in Q$

$$qa^n = q \quad \text{and} \quad qa^k \neq q \quad \text{for} \quad 1 \leq k \leq n-1.$$

Equivalently, the states can be indexed in such a way that $q_i a = q_{i+1}$ for $1 \leq i \leq n-1$, and $q_n a = q_1$. Examples of cyclic automata include the well-known sequence \mathcal{C}_n discovered by Černý.

Dubuc [9] proved that a synchronizing cyclic DFA has a synchronizing word of length at most $(n-1)^2$, as predicted by Černý's conjecture. We define non-deterministic cyclic automata as follows.

Definition 5. A CNFA $\mathcal{A} = (Q, \Sigma)$ is called *cyclic* if there exists $a \in \Sigma$ and an indexing of the states such that

$$q_{i+1} \in q_i a \quad \text{for} \quad 1 \leq i \leq n-1, \quad \text{and} \quad q_1 \in q_n a.$$

Note that with this definition, a CNFA is cyclic if and only if it is the extension of a cyclic DFA.

Proposition 1. *If \mathcal{A} is a D3-directing cyclic CNFA, then \mathcal{A} has a shortest D3-directing word of length at most $(n - 1)^2$.*

Proof. Denote $\text{Split}(\mathcal{A})$ by (Q, Γ) . Choose $a \in \Sigma$ and an indexing of the states as in Definition 5. Define b such that $q_i b = q_{i+1}$ for $1 \leq i \leq n - 1$, and $q_n b = q_1$. Then $b \subseteq a$ so Lemma 4 gives $b \in \Gamma$. Therefore $\text{Split}(\mathcal{A})$ is a cyclic DFA and the result follows. \square

3.2 One-cluster automata

A DFA $\mathcal{A} = (Q, \Sigma)$ is called *one-cluster* if for some letter $a \in \Sigma$, there is only one cycle (possibly a self-loop) labelled a . For every $q \in Q$, the path $qaaa \dots$ eventually ends in this cycle. One way to formally define this is:

Definition 6. *A DFA $\mathcal{A} = (Q, \Sigma)$ is called one-cluster if there exists $a \in \Sigma$ and $p \in Q$ such that for all $q \in Q$*

$$qa^k = p \quad \text{for some } k \in \mathbb{N}.$$

Note that cyclic DFAs are contained in the class of one-cluster automata. Béal, Berlinkov and Perrin [2] proved that in a synchronizing one-cluster DFA, the length of the shortest synchronizing word is at most $2n^2 - 7n + 7$. We define one-cluster CNFAs in the following way.

Definition 7. *A CNFA $\mathcal{A} = (Q, \Sigma)$ is called one-cluster if there exists $a \in \Sigma$ and $p \in Q$ such that for all $q \in Q$*

$$p \in qa^k \quad \text{for some } k \in \mathbb{N}.$$

With this definition, cyclic CNFAs are a special case of one-cluster CNFAs. Like for the cyclic case, the CNFA is one-cluster if and only if it is an extension of a one-cluster DFA.

Proposition 2. *If \mathcal{A} is a D3-directing one-cluster CNFA, then \mathcal{A} has a shortest D3-directing word of length at most $2n^2 - 7n + 7$.*

Proof. Let $\text{Split}(\mathcal{A}) = (Q, \Gamma)$. Choose a and p as in Definition 7 and denote the states by q_1, \dots, q_n . For $i = 1, \dots, n$, let $k_i \geq 1$ be the smallest integer such that $p \in q_i a^{k_i}$ (note that this can also be done if $q_i = p$). Choose $q'_i \in q_i a$ such that $p \in q'_i a^{k_i - 1}$. Define a symbol b by $q_i b = q'_i$ for all i . Then $b \subseteq a$ and $q_i b^{k_i} = p$. By Lemma 4, $b \in \Gamma$ and therefore $\text{Split}(\mathcal{A})$ is a cyclic DFA. \square

Remark 1. To see if a CNFA is one-cluster, it is sufficient to check pairs of states. A CNFA $\mathcal{A} = (Q, \Sigma)$ is one-cluster if and only if there exists $a \in \Sigma$ such that for any $q, q' \in Q$ there exist $r, s \in \mathbb{N}$ for which $qa^r \cap q'a^s \neq \emptyset$.

3.3 Monotonic automata

Definition 8. A DFA $\mathcal{A} = (Q, \Sigma)$ is called *monotonic* if Q admits a linear order $<$ such that for each $a \in \Sigma$ the map $a : Q \rightarrow Q$ preserves the order $<$, i.e.

$$qa \leq q'a \quad \text{whenever} \quad q \leq q'.$$

Ananichev and Volkov [1] proved that the length of the shortest synchronizing word in a synchronizing monotonic DFA is at most $n - 1$. The following definition extends the notion of monotonicity to non-deterministic automata:

Definition 9. A CNFA $\mathcal{A} = (Q, \Sigma)$ is called *monotonic* if Q admits a linear order \leq such that for each $a \in \Sigma$

$$\max \{qa\} \leq \min \{q'a\} \quad \text{whenever} \quad q \leq q'.$$

Every DFA contained in a monotonic CNFA is monotonic. But for a CNFA to be monotonic, it is not sufficient that it contains a monotonic DFA. One can easily extend a monotonic DFA to a CNFA \mathcal{A} for which $\text{Split}(\mathcal{A})$ is not monotonic. Just add to the DFA a symbol that sends every state to the full state set Q .

Proposition 3. If \mathcal{A} is a $D3$ -directing monotonic CNFA, then \mathcal{A} has a shortest $D3$ -directing word of length at most $n - 1$.

Proof. Let $\mathcal{A} = (Q, \Sigma)$ and $\text{Split}(\mathcal{A}) = (Q, \Gamma)$. Choose $b \in \Gamma$. By Lemma 4 there exists $a \in \Sigma$ such that $b \subseteq a$, i.e. $qb \in qa$ for all $q \in Q$. The monotonicity of \mathcal{A} implies

$$qb \leq \max \{qa\} \leq \min \{q'a\} \leq q'b \quad \text{whenever} \quad q \leq q',$$

demonstrating that $\text{Split}(\mathcal{A})$ is a monotonic DFA. This implies the result. \square

3.4 Orientable automata

A DFA $\mathcal{A} = (Q, \Sigma)$ is called *orientable* if Q admits a cyclic order

$$q_1 \prec q_2 \prec \dots \prec q_n \prec q_1$$

on Q , such that for each letter $a \in \Sigma$ the sequence q_1a, q_2a, \dots, q_na is (after removal of duplicates) a subsequence of a cyclic shift of q_1, q_2, \dots, q_n . One can think of this as the order of the states on the circle being preserved under a . This can equivalently be formalized in terms of a linear order:

Definition 10. A DFA $\mathcal{A} = (Q, \Sigma)$ is called *orientable* if Q admits a strict linear order $q_1 < q_2 < \dots < q_n$ such that for each $a \in \Sigma$ at most one of the following inequalities is violated:

$$q_1a \leq q_2a \leq \dots \leq q_na \leq q_1a.$$

Eppstein [10] proved that orientable automata satisfy the conjecture of Černý: if an orientable DFA is synchronizing, then the shortest synchronizing word has length at most $(n-1)^2$. Černý's own examples $\mathcal{C}_n, n = 2, 3, \dots$ are orientable. We extend the notion of orientability as follows to the non-deterministic case:

Definition 11. A CNFA $\mathcal{A} = (Q, \Sigma)$ is called orientable if Q admits a strict linear order $q_1 < q_2 < \dots < q_n$ on Q such that for each $a \in \Sigma$ at most one of the following inequalities is violated:

$$\max\{q_1a\} \leq \min\{q_2a\} \leq \dots \leq \min\{q_na\} \leq \max\{q_na\} \leq \min\{q_1a\}.$$

As is the case for DFAs, the class of orientable CNFAs contains the monotonic CNFAs. Like in the monotonic case, every DFA contained in an orientable CNFA is orientable, but not every extension of an orientable DFA is an orientable CNFA.

Proposition 4. If \mathcal{A} is a D3-directing orientable CNFA, then \mathcal{A} has a shortest D3-directing word of length at most $(n-1)^2$.

Proof. Suppose $\mathcal{A} = (Q, \Sigma)$ is an orientable CNFA. Let $\text{Split}(\mathcal{A}) = (Q, \Gamma)$ and choose $b \in \Gamma$. Then by Lemma 4 there exists $a \in \Sigma$ such that $b \subseteq a$, i.e. $q_k b \in q_k a$ for all $1 \leq k \leq n$. It follows that at most one of the inequalities

$$q_1 b \leq q_2 b \leq \dots \leq q_n b \leq q_1 b$$

is violated so that $\text{Split}(\mathcal{A})$ is an orientable DFA. □

3.5 Automata with underlying Eulerian digraph

Every DFA $\mathcal{A} = (Q, \Sigma)$ has an underlying directed graph (digraph) with vertex set Q and with edges corresponding to actions of elements of Σ . Formally, $G = (V, E)$ with $V = Q$ and $E = \bigcup_{a \in \Sigma} \{(p, q) \in Q^2 : pa = q\}$, where we consider E to be a multiset, which means that each edge has a multiplicity.

A digraph is called Eulerian if it is strongly connected and all indegrees and outdegrees are the same. Kari [13] proved that a synchronizing DFA for which the underlying graph is Eulerian admits a synchronizing word of length at most $(n-2)(n-1) + 1$.

For a CNFA \mathcal{A} with underlying Eulerian digraph G it is not necessarily the case that the underlying digraph of the DFA $\text{Split}(\mathcal{A})$ is Eulerian as well. However, we can decompose G into directed graphs for each of the symbols in Σ : define $G_a = (V, E_a)$ with $V = Q$ and $E_a = \{(p, q) \in Q^2 : pa = q\}$. The following stronger property is preserved under Split :

Definition 12. A CNFA $\mathcal{A} = (Q, \Sigma)$ is called strongly Eulerian if for all $a \in \Sigma$ the corresponding digraph G_a is Eulerian.

Proposition 5. *If $\mathcal{A} = (Q, \Sigma)$ is a D3-directing strongly Eulerian CNFA, then \mathcal{A} has a shortest D3-directing word of length at most $(n-2)(n-1)+1$.*

Proof. Let $\text{Split}(\mathcal{A}) = (Q, \Gamma)$ and denote its underlying digraph by H . Let's first assume that Σ is a singleton $\{a\}$ and that all in- and outdegrees of G_a are equal to k . Suppose $q, q' \in Q$ are such that $q' \in qa$. Since all $p \neq q$ have a -outdegree k , there will be exactly k^{n-1} letters $\tilde{a}_i \in \Gamma, i = 1, \dots, k^{n-1}$ for which $q' = q\tilde{a}_i$. Doing this for all $q' \in qa$ and using that $|qa| = k$ we obtain that q has outdegree k^n in H . An analogous argument gives the same for the indegrees.

Now suppose Σ is not a singleton, i.e. $\Sigma = \{a_1, \dots, a_m\}$. Assume that all degrees in G_{a_i} are equal to k_i . Then by repeatedly applying the above argument, we obtain that all degrees in H are equal to $\sum_{i=1}^m k_i^n$. Clearly H is also strongly connected. Therefore H is Eulerian which implies the result. \square

3.6 Aperiodic automata

One way to define aperiodic DFAs is the following:

Definition 13. *A DFA $\mathcal{A} = (Q, \Sigma)$ is called aperiodic if for all $w \in \Sigma^*$ and $q \in Q$ there exists $k \geq 0$ such that $qw^k = qw^{k+1}$.*

An aperiodic DFA with strongly connected underlying digraph is synchronizing and has a synchronizing word of length at most $\lfloor \frac{n(n+1)}{6} \rfloor$, see [16].

The definition could also be written down for CNFAs, but this property is not preserved under the Split transformation. For example, let $\mathcal{A} = (Q, \Sigma)$ with $|Q| \geq 2$ and $\Sigma = \{a\}$, where a is defined by $qa = Q$ for all $q \in Q$. Then clearly $qw^k = qw^{k+1}$ for all $w \in \Sigma^*$ and $q \in Q$. We will show that $\mathcal{B} = \text{Split}(\mathcal{A}) = (Q, \Gamma)$ admits periodic words. Let $q_1, q_2 \in Q, q_1 \neq q_2$ and let $b \in T^d(Q)$ be such that $q_1b = q_2$ and $q_2b = q_1$. Then $b \subseteq a$ (as a contains every possible symbol), so by Lemma 4 it follows that $b \in \Gamma$. However, $q_1b^k = q_1$ if k even and $q_1b^k = q_2$ if k odd. Therefore \mathcal{B} fails to be aperiodic.

Proposition 6. *Suppose $\mathcal{A} = (Q, \Sigma)$ is a CNFA with the following property: for all $w \in \Sigma^*$ and $q \in Q$ there exists $k \geq 0$ such that $qw^k = qw^{k+1}$ and $|qw^k| = 1$. If its underlying digraph is strongly connected, then \mathcal{A} is D3-directing and has a D3-directing word of length at most $\lfloor \frac{n(n+1)}{6} \rfloor$.*

Proof. Let $\mathcal{B} = \text{Split}(\mathcal{A}) = (Q, \Gamma)$. Let $w = w_1w_2 \dots w_l \in \Gamma^*$. By Lemma 4 there exist $v = v_1v_2 \dots v_l \in \Sigma^*$ such that $w_i \subseteq v_i$ for all i . Let $q \in Q$ and choose k such that $qv^k = qv^{k+1}$ and $|qv^k| = 1$. Since q_w^k is not empty and $qw^k \subseteq qv^k$ it follows that $qw^k = qv^k$ and similarly $qw^{k+1} = qv^{k+1}$. Consequently, \mathcal{B} is an aperiodic DFA. Furthermore, if \mathcal{A} has a strongly connected underlying digraph, then so has \mathcal{B} . \square

CNFAs with the property of Proposition 6 transform by **Split** into an aperiodic DFA. However, having this property is not necessary for being transformed into an aperiodic DFA, as the next example shows. Let $\mathcal{A} = (\{1, 2\}, \{a\})$ where a is defined by $1a = \{1, 2\}$ and $2a = 2$. Then $|1a^k| = 2$ for all $k \geq 1$. Nevertheless $\text{Split}(\mathcal{A})$ is aperiodic, as can be easily verified.

4 Investigating critical CNFAs

In [8] all critical DFAs on 3 or 4 states were identified, and for $n \geq 5$ states all critical extensions of known critical DFAs. Recently it was confirmed [6] that for $n = 5$ and 6 no more critical DFAs exist. From Corollary 1 we know that $\text{Split}(\mathcal{A})$ is a critical DFA for every critical CNFA \mathcal{A} . So if for any DFA \mathcal{D} we can investigate which CNFAs map to \mathcal{D} by **Split**, we can combine these observations to identify all critical CNFAs with < 7 states. For ≥ 7 states the investigation restricts to resulting known critical DFAs. In doing so, first we concentrate on investigating which CNFAs map by **Split** to a given DFA, independent of size or being critical.

For a DFA $\mathcal{D} = (Q, \Sigma)$ we define a graph structure on Σ . More precisely, we define $G(\mathcal{D}) = (\Sigma, E)$ to be the undirected graph of which Σ is the set of nodes and the set E of edges is defined by


$$\{a, b\} \in E \iff \exists q \in Q : qa \neq qb \wedge \forall r \neq q : ra = rb.$$

Next we show that if $G(\mathcal{D}) = (\Sigma, E)$ then any $E' \subseteq E$ gives rise to a CNFA $N(\mathcal{D}, E')$ such that $\text{Split}(N(\mathcal{D}, E')) = \mathcal{D}$. The CNFA $N(\mathcal{D}, E') = (Q, \Sigma')$ is defined by


$$\Sigma' = \{a \cup b \mid \{a, b\} \in E'\} \cup \{a \mid \nexists b : \{a, b\} \in E'\}.$$

So symbols not connected by an edge in E' remain unchanged, and any two symbols a, b that are connected by an edge in E' are joined into the new symbol $a \cup b$. It is defined by $q(a \cup b) = \{qa, qb\}$ for the single state q with $qa \neq qb$, and $r(a \cup b) = ra = rb$ for $r \neq q$. In particular, every symbol in $N(\mathcal{D}, E')$ is non-deterministic in at most one state, and in that state only two choices are possible.

As an example consider the following DFA A_3^+ on three states, and six symbols a, b, c, d, e, f . In fact it is the DFA A_3 from [8], extended by an extra symbol f that acts as the identity. On the right its graph $G(A_3^+)$ is shown: the nodes are a, b, c, d, e, f , and there are three edges $\{b, c\}$, $\{b, f\}$ and $\{d, e\}$. These are exactly the pairs of symbols acting in the same way on two of the three states.



The edge set E of the graph $G(A_3^+)$ has size 3, so there are 8 possible choices for E' . As an example we show the resulting CNFA $N(\mathcal{D}, E')$ for $E' = \{\{b, c\}, \{d, e\}\}$, in which there are 4 symbols $a, b \cup c, d \cup e, f$:



The next theorem states that under some conditions for a given DFA \mathcal{D} these CNFAs $N = N(\mathcal{D}, E')$ are exactly all *pre-basic* CNFAs N for which $\text{Split}(N) = \mathcal{D}$. Here a CNFA is called *pre-basic* if no symbol is contained in another; the difference with basic is that now the identity symbol is allowed. Note that by definition every CNFA of the shape $N(\mathcal{D}, E')$ is pre-basic, and also every DFA is pre-basic.

Theorem 2. *Let $\mathcal{D} = (Q, \Sigma)$ be a DFA for which $G(\mathcal{D}) = (\Sigma, E)$ does not admit cycles of length 3 or 4. Then a pre-basic CNFA $\mathcal{N} = (Q, \Sigma')$ satisfies $\text{Split}(\mathcal{N}) = \mathcal{D}$ if and only if $\mathcal{N} = N(\mathcal{D}, E')$ for some $E' \subseteq E$.*

Proof. For the ‘if’-part, let $\mathcal{N} = N(\mathcal{D}, E')$ for $E' \subseteq E$. If $E' = \emptyset$ then $\mathcal{N} = \mathcal{D}$, so $\text{Split}(\mathcal{N}) = \mathcal{D}$. Otherwise, let $\{a, b\} \in E'$ and $E'' = E' \setminus \{\{a, b\}\}$. Since $\{a, b\} \in E$ there exists $q \in Q$ such that $qa \neq qb$ and $ra = rb$ for all $r \neq q$. Now it is straightforward from the definitions that $\text{Split}(N(\mathcal{D}, E'), q, a) = N(\mathcal{D}, E'')$. By repeating this process by removing all elements from E' one by one, after applying a number of Split operations on $\mathcal{N} = N(\mathcal{D}, E')$ we obtain $N(\mathcal{D}, \emptyset) = \mathcal{D}$, hence $\text{Split}(\mathcal{N}) = \mathcal{D}$.

For the ‘only if’-part we have to show that no other pre-basic CNFA \mathcal{N} satisfies $\text{Split}(\mathcal{N}) = \mathcal{D}$. So let $\mathcal{N} = (Q, \Sigma')$ be an arbitrary CNFA satisfying $\text{Split}(\mathcal{N}) = \mathcal{D} = (Q, \Sigma)$. We will prove by induction on the length of the shortest $\text{Split}(\mathcal{N}, -, -)$ -path from $\mathcal{N} = (Q, \Sigma')$ to \mathcal{D} that $\mathcal{N} = N(\mathcal{D}, E')$ for some $E' \subseteq E$. Let the first step be $\text{Split}(\mathcal{N}, q, a) = \mathcal{N}'$ in which qa consists of at least two states for $q \in Q, a \in \Sigma'$; by the induction hypothesis we assume $\mathcal{N}' = N(\mathcal{D}, E'')$ for some $E'' \subseteq E$.

First assume that qa consists of at least three states. Then according to Lemma 4 every $b \in T^d(Q)$ satisfying $b \subseteq a$ is in Σ . Among these there are three symbols $b_1, b_2, b_3 \in \Sigma$ such that the states qb_i are all distinct for $i = 1, 2, 3$ and for every $r \neq q$ the states rb_i are all equal for $i = 1, 2, 3$. But then b_1, b_2, b_3 form a 3-cycle in $G(\mathcal{D})$, contradicting the assumption of the theorem.

Hence qa consists of exactly two states qa_1, qa_2 , where a_1, a_2 are symbols in $\text{Split}(\mathcal{N}, q, a) = \mathcal{N}'$ for which qa_1, qa_2 are single states and $ra_1 = ra_2 = ra$ for all other states r . Now we claim that ra consists of exactly one state for every $r \neq q$. If not, then choose $r \in Q$ for which $r_1, r_2 \in ra, r_1 \neq r_2$. For all other states s choose $s' \in sa$. For $i, j = 1, 2$ define $qa_{i,j} = a_i, ra_{i,j} = r_j$ and $sa_{i,j} = s'$ for all other states s . Then for $i, j = 1, 2$ we obtain $a_{i,j} \subset a_i$, so $a_{i,j} \in \Sigma$ by Lemma 4. But this yields a 4-cycle $a_{1,1}, a_{1,2}, a_{2,2}, a_{2,1}, a_{1,1}$ in $G(\mathcal{D})$, contradicting the assumption of the theorem. Hence indeed ra consists of exactly one state for every $r \neq q$. Hence $a_1, a_2 \in T^d(Q)$, and $a = a_1 \cup a_2$, and $\{a_1, a_2\} \in E$. Since $\text{Split}(\mathcal{N}, q, a) = \mathcal{N}' = N(\mathcal{D}, E'')$, the non-deterministic symbols of both \mathcal{N} and $N(\mathcal{D}, E'' \cup \{\{a_1, a_2\}\})$ are exactly $a = a_1 \cup a_2$ and the non-deterministic symbols of \mathcal{N}' . Since $\text{Split}(\mathcal{N}) = \mathcal{D} = (Q, \Sigma)$ and \mathcal{N} is pre-basic, the deterministic symbols of \mathcal{N} are exactly the symbols from Σ that are not covered by $E'' \cup \{\{a_1, a_2\}\}$. As the same holds for $N(\mathcal{D}, E'' \cup \{\{a_1, a_2\}\})$, we conclude that all symbols of \mathcal{N} and $N(\mathcal{D}, E'' \cup \{\{a_1, a_2\}\})$ coincide. Hence $\mathcal{N} = N(\mathcal{D}, E'' \cup \{\{a_1, a_2\}\})$, concluding the proof. \square

The proof demonstrates that the requirement concerning cycles is only needed for one of the implications in Theorem 2. If $G(\mathcal{D})$ does contain a 3- or 4-cycle, it is still possible to detect CNFAs that are mapped to \mathcal{D} by Split . As before, every set of edges in $G(\mathcal{D})$ corresponds to such a CNFA. However, there might exist other CNFAs which are mapped to \mathcal{D} as well.

The following examples show that for the other implication in Theorem 2 it is essential to disallow cycles of length both 3 and 4 in $G(\mathcal{D})$. Let \mathcal{N} be defined by $Q = \{1, 2, 3\}, \Sigma = \{a\}, 1a = \{1, 2, 3\}, 2a = \{2\}, 3a = \{3\}$. Then in $\mathcal{D} = \text{Split}(\mathcal{N}) = \text{Split}(\mathcal{N}, 1, a)$ we have three symbols that form a 3-cycle in $G(\mathcal{D})$, and \mathcal{N} is not of the shape $N(\mathcal{D}, E')$ for some set E' of edges of $G(\mathcal{D})$.

As a next example let \mathcal{N} be defined by $Q = \{1, 2\}, \Sigma = \{a\}, 1a = 2a = \{1, 2\}$. Then in $\mathcal{D} = \text{Split}(\mathcal{N})$ we have four symbols that form a 4-cycle in $G(\mathcal{D})$, and \mathcal{N} is not of the shape $N(\mathcal{D}, E')$ for some set E' of edges of $G(\mathcal{D})$.

The role of being pre-basic is illustrated in the following example. Let \mathcal{N} be defined by $Q = \{1, 2\}$, $\Sigma = \{a, b\}$, $1a = \{1, 2\}$, $2a = 1b = 2b = \{2\}$. Then \mathcal{N} is not pre-basic since $b \subseteq a$. In $\mathcal{D} = \text{Split}(\mathcal{N}, 1, a)$ the symbol a is split into two symbols a_1, a_2 , in which $a_2 = b$. So by Split the symbol b disappears, and in $N(\mathcal{D}, \{\{a_1, a_2\}\})$ there is only one symbol a .

Note that if \mathcal{D} is basic, then $N(\mathcal{D}, E')$ is basic too. However, the converse does not hold. For instance, if $\Sigma = \{a\}$, $Q = \{1, 2\}$, $1a = \{1, 2\}$, $2a = \{2\}$, then the CNFA $\mathcal{N} = (Q, \Sigma)$ is basic, but $\text{Split}(\mathcal{N})$ is not since one of the symbols acts as the identity.

As in [8] basic critical DFAs were investigated, we want to combine this by Theorem 2 to investigate basic critical CNFAs. For doing so, we need the following lemma. For any CNFA $\mathcal{N} = (Q, \Sigma)$ we write \mathcal{N}^+ for $(Q, \Sigma \cup \{id\})$ for id being the identity function on Q .

Lemma 5. *A basic CNFA \mathcal{N} is critical if and only if $\text{Split}(\mathcal{N}^+) = \mathcal{D}^+$ for some basic critical DFA \mathcal{D} .*

Proof. By Corollary 1 and the fact that adding id does not influence synchronization, the CNFA \mathcal{N} is critical if and only if $\text{Split}(\mathcal{N}^+)$ is critical. The 'if'-part follows since \mathcal{D}^+ is critical if \mathcal{D} is critical. For the 'only if'-part let \mathcal{N} be a basic critical CNFA. Then $\text{Split}(\mathcal{N}^+)$ is a critical DFA. Since id is contained in \mathcal{N}^+ , by Lemma 4 we obtain that id is also contained in $\text{Split}(\mathcal{N}^+)$. Hence $\text{Split}(\mathcal{N}^+) = \mathcal{D}^+$ for some basic DFA \mathcal{D} , which is critical since $\text{Split}(\mathcal{N}^+) = \mathcal{D}^+$ is critical. \square

So basic critical CNFAs can be obtained by taking a basic critical DFA \mathcal{D} and compute $G(\mathcal{D}^+)$. If it does not contain 3- or 4-cycles, all pre-basic CNFAs \mathcal{N} with $\text{Split}(\mathcal{N}) = \mathcal{D}^+$ can be obtained by Theorem 2. Finally, by Lemma 5 the resulting basic critical CNFAs are obtained by removing id wherever it occurs. As long as no 3- or 4-cycles occur in $G(\mathcal{D}^+)$, all basic critical CNFAs can be obtained in this way. The number of resulting basic critical CNFAs is equal to the number of pre-basic critical CNFAs obtained by Theorem 2 since by removing the possible occurrence of id any pre-basic CNFA is transformed to a basic CNFA, and no two map to the same basic CNFA since for a basic CNFA \mathcal{N} with $\text{Split}(\mathcal{N}) = \mathcal{D}^+$, the CNFA \mathcal{N}^+ is not pre-basic since it has a symbol in which id is contained.

For instance, in the cyclic graph $G(A_3^+)$ we have $f = id$, and there are exactly 3 edges, yielding exactly 8 sets of edges. Hence there are exactly 8 basic CNFAs \mathcal{N} with $\text{Split}(\mathcal{N}^+) = A_3^+$, all obtained from $N(A_3^+, E')$ for a set E' of edges. If $\{b, f\} \in E'$ then this CNFA is already basic since the identity $f = id$ is joined with b , and otherwise the symbol $f = id$ is removed.

For $n \leq 6$ all critical DFAs are known; next for all the $n \leq 6$ we apply the above approach to find and count all basic critical CNFAs.

4.1 Analyzing 2 states

In [8] no analysis of automata with two states was made, since that is a quite degenerate case in which the maximal shortest synchronizing word is only one single symbol. On two states 1, 2 there are three possible deterministic non-identity symbols: a mapping both states to 1, b mapping both states to 2, and s swapping 1 and 2. A DFA is critical if and only if at least one of the symbols a and b occurs. These yield exactly 6 basic critical DFAs, having alphabets $\{a\}$, $\{b\}$, $\{a, b\}$, $\{a, s\}$, $\{b, s\}$, $\{a, b, s\}$, among which $\{a\}$ is isomorphic to $\{b\}$ and $\{a, s\}$ is isomorphic to $\{b, s\}$, so up to isomorphism there are exactly 4. To each of these DFAs, we add $f = id$ and apply the above approach. Computing the graph and counting the edges gives the following numbers of basic critical CNFAs corresponding to sets of edges

symbols DFA	nr of CNFAs	symbols DFA	nr of CNFAs
a, b, s	$2^4 = 16$	a, b	$2^2 = 4$
a, s	$2^2 = 4$	a	$2^1 = 2$
b, s	$2^2 = 4$	b	$2^1 = 2$

so yielding 32 CNFAs corresponding to sets of edges. The DFA \mathcal{D} with all three symbols a, b, s together with id yields a graph that is a 4-cycle: the edges are (a, s) , (b, s) , (a, id) , (b, id) . Theorem 2 does not apply to \mathcal{D} due to this 4-cycle, which means that there might be more CNFAs that are mapped to \mathcal{D} by Split . It turns out that apart from the 16 in the table there is one more CNFA \mathcal{N} for which $\text{Split}(\mathcal{N}) = \mathcal{D}$: the CNFA on $\{1, 2\}$ with one symbol c satisfying $1c = 2c = \{1, 2\}$. This makes 33 basic critical CNFAs on two states in total; up to isomorphism the number is 20.

4.2 Analyzing 3 states

From [8] we recall that there are exactly 15 basic critical DFAs on 3 states, namely the restrictions of A_3 to the 15 sets of symbols indicated in the table below. For each of them we add $f = id$ and proceed as before. Since $G(A_3^+)$ does not contain 3- or 4-cycles, also its subgraphs don't. Therefore Theorem 2 applies and by the above approach we obtain the following numbers 2^k of basic critical CNFAs in which k is the number of edges in the corresponding


graph.

symbols DFA	nr of CNFAs	symbols DFA	nr of CNFAs
a, b, c, d, e	$2^3 = 8$	a, b, e	$2^1 = 2$
a, b, c, d	$2^2 = 4$	a, c, d	$2^0 = 1$
a, b, c, e	$2^2 = 4$	a, d, e	$2^1 = 2$
a, b, d, e	$2^2 = 4$	b, c, e	$2^2 = 4$
a, c, d, e	$2^1 = 2$	c, d, e	$2^1 = 2$
b, c, d, e	$2^3 = 8$	a, b	$2^1 = 2$
a, b, c	$2^2 = 4$	a, d	$2^0 = 1$
a, b, d	$2^1 = 2$		

So we conclude that there are exactly 50 basic critical CNFAs on three states, including the 15 basic critical DFAs, corresponding to the empty set of edges. No two of them are isomorphic.

4.3 Analyzing 4 states

From [8] we recall that there are exactly 12 basic critical DFAs on 4 states, namely C_4 and T4-2 depicted below and the restrictions of A_4 to the 10 sets of symbols $\{a, b, c, d, e\}$, $\{a, b, c, d\}$, $\{a, b, c, e\}$, $\{a, b, d, e\}$, $\{b, c, d, e\}$, $\{a, b, c\}$, $\{a, b, d\}$, $\{a, b, e\}$, $\{b, d, e\}$ and $\{a, b\}$. Here A_4 is the restriction of A_4^+ to $abcde$ depicted below. In fact, A_4^+ was obtained from A_4 by adding an extra symbol $f = id$.



It is straightforward that $G(C_4^+)$, is a graph with one edge: b connected to id , hence yielding exactly two basic CNFAs. Doing the same for T4-2⁺ yields the empty graph, so there are no more CNFAs for which the Split is T4-2 then only T4-2 itself.

For A_4^+ the corresponding graph $G(A_4^+)$ is indicated, so on all its subgraphs Theorem 2 applies. For the 10 relevant subgraphs we obtain the following numbers of basic critical CNFAs.

symbols DFA	nr of CNFAs	symbols DFA	nr of CNFAs
a, b, c, d, e	$2^2 = 4$	a, b, c	$2^1 = 2$
a, b, c, d	$2^2 = 4$	a, b, d	$2^0 = 1$
a, b, c, e	$2^1 = 2$	a, b, e	$2^0 = 1$
a, b, d, e	$2^0 = 1$	b, d, e	$2^0 = 1$
b, c, d, e	$2^2 = 4$	a, b	$2^0 = 1$

So we conclude that there are exactly 21 basic critical CNFAs \mathcal{N} for which the $\text{Split}(\mathcal{N}^+)$ is A_4^+ . Together with the two related to C_4 and the single one T4-2 this yields exactly 24 basic critical CNFAs on four states (none of them isomorphic), including the 12 basic critical DFAs.

4.4 Analyzing ≥ 5 states

On ≥ 5 states the only known basic critical DFAs are C_n for every $n \geq 5$, and two more: one from Roman on 5 states and 3 symbols, and one from Kari on 6 states and 2 symbols, for their definitions and references we refer to [8]. It was shown by Trahtman [15] that this investigation is complete for $n \leq 10$ when restricting to at most two symbols; recently in [6] it was shown that for $n \leq 6$ indeed no more critical DFAs exists, with no restrictions on the numbers of symbols.

For both the Kari and Roman DFA the corresponding graph is empty, yielding no other basic critical CNFAs. The DFA C_n is defined to be $(\{1, 2, \dots, n\}, \{a, b\})$ for a, b defined by $qa = q + 1$ for $q < n$, $na = 1$, $1b = 2$, $qb = q$ for $q > 1$. After adding the identity id its graph consists of a single edge $\{b, id\}$, yielding one more basic critical CNFA on n states: C_n to which a b -self-loop is added to 1, yielding $1b = \{1, 2\}$. Summarizing, for $n \leq 6$ we have up to isomorphism the following numbers of basic critical DFAs and CNFAs:

nr of states	nr of DFAs	nr of CNFAs
2	4	20
3	15	50
4	12	24
5	2	3
6	2	3

while for $n > 6$ the only known basic critical DFA is C_n , to be extended to exactly one more basic critical CNFA.

5 Conclusions

The central result of this paper is that every D3-directing CNFA can be transformed to a synchronizing DFA with the same synchronizing word length. In this paper we present this **Split** transformation and explore its properties. An immediate consequence is that the maximal shortest D3-directing length for CNFAs is equal to the maximal shortest synchronizing length for DFAs, which means that the famous Černý conjecture for DFAs extends to CNFAs.

For several classes of DFAs with some additional properties, tighter bounds for synchronization lengths have been established. If a CNFA is transformed into a DFA belonging to such a class, then the tighter bound also applies to the CNFA. This observation is used to define properties for CNFAs that guarantee improvements over the general cubic bound.

In the last part of the paper, we investigate critical CNFAs. The tight connection between critical DFAs and critical CNFAs, combined with the fact that critical DFAs are extremely rare, implies that also a very small fraction of CNFAs is critical. All critical DFAs on at most 6 states are known. By essentially inverting **Split**, we identify all critical CNFAs on at most 6 states. Furthermore, for all $n \geq 3$ there is exactly one critical CNFA which is a strict extension of Černý's DFA C_n .

References

- [1] D.S. Ananichev and M.V. Volkov. Synchronizing monotonic automata. *Theoretical Computer Science*, 327:225–239, 2004.
- [2] M.-P. Béal, M.V. Berlinkov, and D. Perrin. A quadratic upper bound on the size of a synchronizing word in one-cluster automata. *International Journal of Foundations of Computer Science*, 22:277–288, 2011.
- [3] V. D. Blondel, R. M. Jungers, and A. Olshevsky. On primitivity of sets of matrices. *Automatica*, 61(C):80–88, 2015.
- [4] J. Černý. Poznámka k homogénnym experimentom s konečnými automatmi. *Matematicko-fyzikálny časopis, Slovensk. Akad. Vied*, 14(3):208–216, 1964.
- [5] P. Chevalier, J. M. Hendrickx, and R. M. Jungers. Reachability of consensus and synchronizing automata. *54th IEEE Conference on Decision and Control*, pages 4139–4144, 2015.
- [6] M. de Bondt, H. Don, and H. Zantema. DFAs and PFAs with long shortest synchronizing word length. Available at <http://arxiv.org/abs/1703.07618>, 2017.

- [7] H. Don. The Černý conjecture and 1-contracting automata. *Electronic Journal of Combinatorics*, 23(3):P3.12, 2016.
- [8] H. Don and H. Zantema. Finding DFAs with maximal shortest synchronizing word length. In Drewes F., Martín-Vide C., and Truthe B., editors, *Language and Automata Theory and Applications*, volume 10168 of *Springer Lecture Notes in Computer Science*. Springer, Cham, 2017.
- [9] L. Dubuc. Sur les automates circulaires et la conjecture de Černý. *RAIRO Inform. Theor. Appl.*, 32:21–34, 1998.
- [10] D. Eppstein. Reset sequences for monotonic automata. *SIAM Journal on Computing*, 19:500–510, 1990.
- [11] B. Gerencsér, V. V. Gusev, and R. M. Jungers. Primitive sets of nonnegative matrices and synchronizing automata. Available at <https://arxiv.org/abs/1602.07556>, 2016.
- [12] B. Imreh and M. Steinby. Directable non-deterministic automata. *Acta Cybernetica*, 14:105–115, 1999.
- [13] J. Kari. Synchronizing finite automata on eulerian digraphs. *Theoretical Computer Science*, 295(1–3):223–232, 2003.
- [14] J.-E. Pin. On two combinatorial problems arising from automata theory. *Annals of Discrete Mathematics*, 17:535–548, 1983.
- [15] A. N. Trahtman. An efficient algorithm finds noticeable trends and examples concerning the Černý conjecture. In Rastislav Kráľovič and Paweł Urzyczyn, editors, *Mathematical Foundations of Computer Science 2006: 31st International Symposium, MFCS 2006*, pages 789–800. Springer Berlin Heidelberg, 2006.
- [16] M.V. Volkov. Synchronizing automata preserving a chain of partial orders. *Theoretical Computer Science*, 410(37):3513–3519, 2009.