

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/204444>

Please be advised that this information was generated on 2021-09-29 and may be subject to change.

$SU_q(3)$ corepresentations and bivariate q -Krawtchouk polynomials

Cite as: J. Math. Phys. 60, 051701 (2019); doi: 10.1063/1.5088985

Submitted: 15 January 2019 • Accepted: 28 March 2019 •

Published Online: 8 May 2019



View Online



Export Citation



CrossMark

G. Bergeron,^{1,a)}  E. Koelink,^{2,b)} and L. Vinet^{1,c)}

AFFILIATIONS

¹Centre de Recherches Mathématiques, Université de Montréal, P.O. Box 6128, Centre-ville Station, Montréal, Québec H3C 3J7, Canada

²IMAPP, Radboud Universiteit, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands

^{a)}Electronic mail: bergerog@crm.umontreal.ca

^{b)}Electronic mail: e.koelink@math.ru.nl

^{c)}Electronic mail: vinet@crm.umontreal.ca

ABSTRACT

The matrix elements of unitary $SU_q(3)$ corepresentations, which are analogs of the symmetric powers of the natural representation, are shown to be the bivariate q -Krawtchouk orthogonal polynomials, thus, providing an algebraic interpretation of these polynomials in terms of quantum groups.

Published under license by AIP Publishing. <https://doi.org/10.1063/1.5088985>

I. INTRODUCTION

A fruitful connection exists between Lie groups and algebras and the theory of orthogonal polynomials. Algebraic interpretations for these orthogonal polynomials enable simple derivations of their properties and often lead to new identities. Similar connections between the theory of quantum groups and (mostly univariate) q -orthogonal polynomials have been established.¹ The results of this paper pursue such a connection in multivariate situations by giving an algebraic interpretation of the quantum bivariate q -Krawtchouk polynomials in terms of the quantum group $SU_q(3)$.

In the classical case, the Krawtchouk polynomials of the Tratnik type form a family of multivariate Krawtchouk polynomials constructed from univariate Krawtchouk polynomials using a construction developed in Ref. 2 that applies to all polynomials of the ($q = 1$) Askey scheme. These are orthogonal polynomials with respect to the multinomial distribution. Many Lie-theoretic interpretations have been given for the Krawtchouk polynomials. The multivariate Krawtchouk polynomials in d variables were shown³ to be matrix elements of the $SO(d + 1)$ Lie group and identified as well^{4,5} as overlaps of anti-automorphisms of $\mathfrak{sl}(d + 1)$ -modules.

In the context of quantum groups and algebras, interpretations analogous to the classical ones have been given for the q -Krawtchouk polynomials, which are orthogonal with respect to the q -binomial distribution. Koornwinder obtained⁶ the univariate q -Krawtchouk polynomials as the matrix elements of unitary corepresentations of the $SU_q(2)$ quantum group. In a complementary way based on the quantum algebras, the q -Krawtchouk polynomials were seen^{7,8} to arise as matrix elements of a class of $U_q(\mathfrak{sl}(2))$ automorphisms. These two approaches are essentially dual one to another.⁹

A family of multivariate q -Krawtchouk polynomials was first derived by Gasper and Rahman in Ref. 10 where they constructed q -deformations of Tratnik's polynomials. We will thus refer to these multivariate extensions of the q -Krawtchouk as being of the Tratnik type. An interpretation of the bivariate and multivariate q -Krawtchouk polynomials based on the quantum algebra viewpoint was obtained in Ref. 11. With an eye to generalizations and in view of the fact that for $q = 1$, Lie groups rather than algebras provide a most natural framework, it seems appropriate to examine how the multivariate q -Krawtchouk polynomials can be obtained and analyzed in a quantum group framework. In this paper, we build upon the quantum group approach of Koornwinder to obtain the bivariate quantum q -Krawtchouk polynomials of the Tratnik type as the matrix element of unitary $SU_q(3)$ corepresentations. Within this quantum group approach, the structure of the unitary elements of $U_q(\mathfrak{sl}(3))$ constructed in Ref. 11 is explained from the representation theory of $SU_q(3)$.

This paper is organized as follows: In Sec. I, a presentation of the $SU_q(3)$ algebra is first given and the construction of its unitary representations is reviewed. Symmetric $SU_q(3)$ corepresentations are then constructed at the beginning of Sec. II, followed by the derivation of their matrix elements and a proof of the unitarity of the corepresentations. A generating function for the matrix elements is then obtained. In Sec. III, the matrix elements are evaluated in irreducible $SU_q(3)$ representations and identified as bivariate q -Krawtchouk polynomials, which follows from Soibelman’s tensor product theorem. Finally, Sec. IV illustrates how evaluating the matrix elements in reducible $SU_q(3)$ representations leads to identities for orthogonal polynomials. This is followed by a brief conclusion.

II. THE $SU_q(3)$ HOPF ALGEBRA AND ITS REPRESENTATIONS

We first give in this section a presentation of the $SU_q(3)$ quantum group, a Hopf $*$ -algebra, and discuss how its representations are constructed.

A. The coordinate ring $A(SU_q(3))$

The coordinate ring $A(SL_q(3; \mathbb{C}))$ is a \mathbb{C} -algebra $A = \mathbb{C}[x_{ij}; 1 \leq i, j \leq 3]$ with the relations

$$\begin{aligned} x_{ik}x_{jk} &= qx_{jk}x_{ik}, & x_{ki}x_{kj} &= qx_{kj}x_{ki}, & \forall i < j, & (1) \\ x_{il}x_{jk} &= x_{jk}x_{il}, & x_{ik}x_{jl} - qx_{il}x_{jk} &= x_{jl}x_{ik} - \frac{1}{q}x_{jk}x_{il}, & \forall i < j, \quad k < l, \\ \sum_{\sigma \in S_3} (-q)^{l(\sigma)} x_{1\sigma(1)}x_{2\sigma(2)}x_{3\sigma(3)} &= 1. \end{aligned}$$

A Hopf algebra structure is given with the following coproduct Δ , counit ε , and antipode S :

$$\Delta(x_{ij}) = \sum_{k=1}^3 x_{ik} \otimes x_{kj}, \quad \varepsilon(x_{ij}) = \delta_{ij}, \quad S(x_{ij}) = (-q)^{i-j} \xi_{ji}, \quad 1 \leq i, j \leq 3, \quad (2)$$

where ξ_{ij} denotes the (i, j) quantum minor, that is, the quantum determinant of x with the i th row and the j th column removed

$$\xi_{ij} = \sum_{\tau \in S_2} (-q)^{l(\tau)} x_{i_1\tau(1)} x_{i_2\tau(2)}, \quad i_1 < i_2 \in \{1, 2, 3\} \setminus \{i\}, \quad j_1 < j_2 \in \{1, 2, 3\} \setminus \{j\}.$$

Moreover, a unique conjugate linear antihomomorphism $*$: $A(SL_q(3; \mathbb{C})) \rightarrow A(SL_q(3; \mathbb{C}))$: $x \mapsto x^*$ exists such that

$$x_{ij}^* = S(x_{ji}) = (-q)^{j-i} \xi_{ij}, \quad \forall i, j. \quad (3)$$

This $*$ -operation makes $A(SL_q(3))$ into the $*$ -Hopf algebra $A(SU_q(3))$ which we will refer to as simply $SU_q(3)$.

B. $SU_q(3)$ representations

The $SU_q(3)$ $*$ -representations used in this paper were constructed in Ref. 12. However, to obtain our main results, we will make use of a theorem of Soibelman¹³ on the construction of modules over quantum groups. Thus, we review how $*$ -representations of unitary quantum groups are constructed¹⁴ using this result. Following Ref. 15, one first defines¹⁶ the infinite dimensional representations τ_α with $\alpha \in U(1)$, of $SU_q(2)$ ¹⁷ where its generators $\{t_{ij}; i, j = 1, 2\}$ act on a Hilbert space H with orthonormal basis $\{|n\rangle, n \in \mathbb{N}\}$, as follows:

$$\begin{aligned} \tau_\alpha(t_{12})|n\rangle &= -q^{n+1} \alpha^{-1}|n\rangle, & \tau_\alpha(t_{21})|n\rangle &= q^n \alpha|n\rangle, & \tau_\alpha(t_{11})|0\rangle &= 0, \\ \tau_\alpha(t_{11})|n\rangle &= \sqrt{1 - q^{2n}}|n - 1\rangle, & \tau_\alpha(t_{22})|n\rangle &= \sqrt{1 - q^{2(n+1)}}|n + 1\rangle. \end{aligned}$$

From these representations, one can build $SU_q(3)$ representations. Indeed, consider the two canonical embeddings $\varphi_i : U_q(\mathfrak{sl}(2)) \hookrightarrow U_q(\mathfrak{sl}(3))$, $i = 1, 2$. These embeddings define by duality the projections $\varphi_i^* : SU_q(3) \rightarrow SU_q(2)$ such that irreducible $*$ -representations of $SU_q(3)$ are given by the maps $\pi_i \equiv \tau_\alpha \circ \varphi_i^*$ with $\alpha = -1$ acting on $V_{s_i} \cong H$. Explicitly, these elementary representations are specified by

$$\pi_1 \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix} \cdot |k\rangle = \begin{pmatrix} \sqrt{1 - q^{2k}}|k - 1\rangle & q^{k+1}|k\rangle & 0 \\ -q^k|k\rangle & \sqrt{1 - q^{2k+2}}|k + 1\rangle & 0 \\ 0 & 0 & |k\rangle \end{pmatrix} \quad (4)$$

and

$$\pi_2 \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix} \cdot |k\rangle = \begin{pmatrix} |k\rangle & 0 & 0 \\ 0 & \sqrt{1 - q^{2k}}|k - 1\rangle & q^{k+1}|k\rangle \\ 0 & -q^k|k\rangle & \sqrt{1 - q^{2k+2}}|k + 1\rangle \end{pmatrix}. \quad (5)$$

All $SU_q(3)$ representations can be constructed from the elementary representations π_1 and π_2 using the following tensor product theorem (Ref. 15, Theorem 6.2.7). Denoting by $\mathbb{C}[G]_q$ the quantized algebra of functions¹⁸ on G , a connected and simply connected simple compact Lie group with associated Weyl group W , one has the following theorem.

Theorem 1 (Tensor product theorem). *For any unitarizable irreducible $\mathbb{C}[G]_q$ \ast -representation V , there exists a unique element $w \in W$ of the Weyl group and a unique element $\tau \in T$ of the distinguished maximal torus such that*

$$V \cong V_w \otimes V_\tau, \quad V_w = V_{s_{i_1}} \otimes V_{s_{i_2}} \otimes \cdots \otimes V_{s_{i_k}},$$

where $w = s_{i_1} s_{i_2} \dots s_{i_k}$ is a reduced decomposition of w . The tensor product does not depend (up to a unitary equivalence) on the choice of reduced decomposition for w .

The map $\pi_w: SU_q(3) \rightarrow \text{End}(V_w)$ associated with the representation in the above theorem is specified by

$$\pi_w = (\pi_{i_1} \otimes \pi_{i_2} \otimes \cdots \otimes \pi_{i_k}) \circ \Delta^{(k-1)}, \quad w = s_{i_1} s_{i_2} \dots s_{i_k},$$

where the repeated coproduct $\Delta^{(k-1)}$ is defined through

$$\Delta^{(1)} = \Delta, \quad \Delta^{(k)} = \underbrace{1 \otimes \cdots \otimes 1}_{k-1 \text{ times}} \otimes \Delta \circ \Delta^{(k-1)}.$$

The representation on V_τ is a one-dimensional representation of the form $\rho_\tau: SU_q(3) \rightarrow \mathbb{C}: x_{ij} \mapsto \alpha_i(\tau) \delta_{ij}$ with $\alpha_i(\tau) \in U(1)$ such that $\alpha_1(\tau) \alpha_2(\tau) \alpha_3(\tau) = 1$. Thus, it only contributes a global phase to our result and will not be further considered. Take now the case where $w = s_2 s_1$. One has

$$\pi_{21} \equiv \pi_{s_2 s_1} = (\pi_2 \otimes \pi_1) \circ \Delta. \tag{6}$$

The explicit action on the generators is easily obtained from (6) using (2), (4), and (5) such that acting on the $SU_q(3)$ generators yields

$$\pi_{12} \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix} |n, m\rangle = \begin{pmatrix} \sqrt{1-q^2} |m-1, n\rangle & q^{m+1} \sqrt{1-q^{2n}} |m, n-1\rangle & q^{m+n+2} |m, n\rangle \\ -q^m |m, n\rangle & \sqrt{1-q^{2m+2}} \sqrt{1-q^{2n}} |m+1, n-1\rangle & q^{n+1} \sqrt{1-q^{2m+2}} |m+1, n\rangle \\ 0 & -q^n |m, n\rangle & \sqrt{1-q^{2n+2}} |m, n+1\rangle \end{pmatrix}.$$

The representation π_{121} is obtained similarly and, upon tensoring with one-dimensional representations, is the most general representation in the sense that the intersection of the kernels of these representations is trivial, cf., e.g., Ref. 14 (Sec. 5). It follows from Theorem 1 that π_{121} and π_{212} are equivalent, so it suffices to consider one of them.

III. UNITARY $SU_q(3)$ COREPRESENTATIONS

We now turn to the construction of unitary $SU_q(3)$ corepresentations in analogy with the $GL(3)$ coaction on functions on \mathbb{C}^3 . Consider the space $\mathcal{F}^{(1)}(\mathbb{C}^3)$ of linear functions on \mathbb{C}^3 with orthonormal basis $\{z_i\}_{i=1,2,3}$. By identifying these basis elements with a fixed column of the $SU_q(3)$ quantum group as follows

$$z_i = x_{ij}, \quad \text{for } j \text{ fixed}, \tag{7}$$

a natural¹⁹ coaction is defined using the coproduct (2) as

$$\Delta: \mathcal{F}^{(1)}(\mathbb{C}^3) \rightarrow SU_q(3) \otimes \mathcal{F}^{(1)}(\mathbb{C}^3): z_i \mapsto \Delta(x_{ij}) = \sum_{k=1}^3 x_{ik} \otimes x_{kj} \equiv \sum_{j=1}^3 x_{ik} \otimes z_k. \tag{8}$$

The algebra $\mathcal{F}(\mathbb{C}^3)$ of polynomial functions on \mathbb{C}^3 is identified with the tensor algebra of $\mathcal{F}^{(1)}(\mathbb{C}^3)$ as follows

$$\mathcal{F}(\mathbb{C}^3) = T(\mathcal{F}^{(1)}(\mathbb{C}^3)), \quad T(V) \equiv \bigoplus_{n=0}^{\infty} V^{\otimes n}, \quad V^{\otimes n} \equiv \underbrace{V \otimes \cdots \otimes V}_n, \quad V^{\otimes 0} \cong \mathbb{C}. \tag{9}$$

The identification (7) together with (1) establishes the following relations:

$$\mathcal{R} = \{z_i z_j \sim q z_j z_i \mid i < j, i, j = 1, 2, 3\}. \tag{10}$$

Denote the quotient (Ref. 18, Chap. 7) of the tensor algebra (9) by the relations (10) as $\text{Sym}_q(\mathbb{C}^3)$. A natural grading on $\text{Sym}_q(\mathbb{C}^3)$ is inherited from the one of $\mathcal{F}(\mathbb{C}^3)$ as the relations (10) preserve this grading. Explicitly,

$$\text{Sym}_q(\mathbb{C}^3) = \bigoplus_{n=0}^{\infty} \mathcal{F}_q^{(n)}(\mathbb{C}^3), \quad \mathcal{F}_q^{(n)}(\mathbb{C}^3) \equiv \left(\mathcal{F}^{(1)}(\mathbb{C}^3)\right)^{\otimes n} / \mathcal{R}. \tag{11}$$

The coproduct being an homomorphism, the coaction (8) is extended to $\text{Sym}_q(\mathbb{C}^3)$ as follows:

$$\Delta(z_1^{m_1} z_2^{m_2} z_3^{m_3}) = \Delta(z_1)^{m_1} \Delta(z_2)^{m_2} \Delta(z_3)^{m_3}, \quad \forall z_1^{m_1} z_2^{m_2} z_3^{m_3} \in \text{Sym}_q(\mathbb{C}^3). \quad (12)$$

Being constructed from the coproduct, (12) defines an $SU_q(3)$ corepresentation. From (8), it is easily seen that the coaction (12) preserves the natural grading (11) on $\text{Sym}_q(\mathbb{C}^3)$. Thus, $\text{Sym}_q(\mathbb{C}^3)$ as an $SU_q(3)$ corepresentation decomposes as a direct sum of corepresentations as follows:

$$\Delta : \text{Sym}_q(\mathbb{C}^3) \longrightarrow \bigoplus_{n=0}^{\infty} \Delta(\mathcal{F}_q^{(n)}(\mathbb{C}^3)).$$

In view of (10), a basis for $\mathcal{F}_q^{(N)}(\mathbb{C}^3)$ is given by

$$\mathcal{B}_N = \left\{ z^{\vec{m}} \mid |\vec{m}| = N \right\}, \quad \text{for } z^{\vec{m}} \equiv z_1^{m_1} z_2^{m_2} z_3^{m_3} \quad \text{and} \quad |\vec{m}| \equiv m_1 + m_2 + m_3. \quad (13)$$

A. Matrix elements

On the basis (13), the matrix elements of the corepresentation $\mathcal{F}_q^{(N)}(\mathbb{C}^3)$ are given in the following proposition.

Proposition 1 (Matrix elements). Let $0 < N \in \mathbb{N}$. The matrix elements $h_{\vec{m}, \vec{n}}^{(N)}$ of the $SU_q(3)$ corepresentation $\mathcal{F}_q^{(N)}(\mathbb{C}^3)$ with coaction Δ are given in the basis \mathcal{B}_N by

$$\Delta(z^{\vec{m}}) = \sum_{|\vec{n}|=N} h_{\vec{m}, \vec{n}}^{(N)} \otimes z^{\vec{n}}, \quad h_{\vec{m}, \vec{n}}^{(N)} = \sum_{\substack{|\vec{a}_p|=m_p \\ |\vec{a}_p|=n_p}} Q(a) \begin{bmatrix} \vec{m} \\ a \end{bmatrix}_{q^{-2}} \prod_{k=1}^3 \left(\prod_{i=1}^3 x_{ik}^{a_{ik}} \right), \quad (14)$$

with $|\vec{a}_p| \equiv \sum_{j=1}^3 a_{pj}$, $|a_p| \equiv \sum_{i=1}^3 a_{ip}$, and

$$\begin{bmatrix} \vec{m} \\ a \end{bmatrix}_q \equiv \begin{bmatrix} m_1 \\ \bar{a}_1 \end{bmatrix}_q \begin{bmatrix} m_2 \\ \bar{a}_2 \end{bmatrix}_q \begin{bmatrix} m_3 \\ \bar{a}_3 \end{bmatrix}_q, \quad \begin{bmatrix} m \\ \bar{a}_i \end{bmatrix}_q \equiv \frac{(q; q)_m}{(q; q)_{a_i} (q; q)_{a_{i2}} (q; q)_{a_{i3}}},$$

and where

$$Q(a) = q^{-f(a)}, \quad f(a) = a_{13}(a_{21} + a_{22} + a_{32}) + a_{31}(a_{12} + a_{22} + a_{23}) + a_{12}a_{21} + a_{13}a_{31} + a_{23}a_{32}, \quad (15)$$

is a manifestly symmetric function of the matrix of indices.

Proof. First, compute the matrix elements of the coaction on powers of a single generator of $\mathcal{F}_q^{(1)}(\mathbb{C}^3)$. From (8) and (12), one has

$$\Delta(z_i^m) = \Delta(x_{ij})^m = \left(\sum_{k=1}^3 x_{ik} \otimes x_{kj} \right)^m,$$

and, knowing that $(x_{ik} \otimes x_{kj})(x_{il} \otimes x_{lj}) = q^2(x_{il} \otimes x_{ij})(x_{ik} \otimes x_{kj})$, $\forall k < l$, one has that⁶

$$\Delta(z_i^m) = \sum_{|\vec{a}_i|=m} \begin{bmatrix} m \\ \vec{a}_i \end{bmatrix}_{q^{-2}} x_{i1}^{a_{i1}} x_{i2}^{a_{i2}} x_{i3}^{a_{i3}} \otimes z_1^{a_{i1}} z_2^{a_{i2}} z_3^{a_{i3}} = \sum_{|\vec{a}_i|=m} \begin{bmatrix} m \\ \vec{a}_i \end{bmatrix}_{q^{-2}} \prod_{k=1}^3 (x_{ik} \otimes x_{kj})^{a_{ik}},$$

with the understanding that $\vec{a}_i \equiv (a_{i1}, a_{i2}, a_{i3})$ and $|\vec{a}_i| \equiv \sum_{j=1}^3 a_{ij}$. The coaction (12) of a generic basis elements in (13) is then

$$\Delta(z^{\vec{m}}) = \prod_{i=1}^3 \Delta(z_i^{m_i}) = \sum_{\substack{|\vec{a}_1|=m_1 \\ |\vec{a}_2|=m_2 \\ |\vec{a}_3|=m_3}} \begin{bmatrix} \vec{m} \\ a \end{bmatrix}_{q^{-2}} \prod_{i=1}^3 \left(\prod_{k=1}^3 (x_{ik} \otimes x_{kj})^{a_{ik}} \right), \quad (16)$$

where all the products are to be expanded from left to right and from innermost to outermost. Then, observe that

$$\prod_{i=1}^3 \left(\prod_{k=1}^3 x_{ik}^{a_{ik}} \right) = x_{11}^{a_{11}} x_{12}^{a_{12}} x_{13}^{a_{13}} x_{21}^{a_{21}} x_{22}^{a_{22}} x_{23}^{a_{23}} x_{31}^{a_{31}} x_{32}^{a_{32}} x_{33}^{a_{33}} = x_{11}^{a_{11}} x_{21}^{a_{21}} x_{31}^{a_{31}} x_{12}^{a_{12}} x_{22}^{a_{22}} x_{32}^{a_{32}} x_{13}^{a_{13}} x_{23}^{a_{23}} x_{33}^{a_{33}} = \prod_{k=1}^3 \left(\prod_{i=1}^3 x_{ik}^{a_{ik}} \right). \quad (17)$$

Writing $n_k = \sum_{i=1}^3 a_{ik} \equiv |a_k|$ for $k = 1, 2, 3$ with $\vec{n} \equiv (n_1, n_2, n_3)$, one finds with (15) that

$$\prod_{i=1}^3 \left(\prod_{k=1}^3 x_{kj}^{a_{ik}} \right) = Q(a) \prod_{k=1}^3 \left(x_{kj}^{a_{1k}+a_{2k}+a_{3k}} \right) = Q(a) z_1^{n_1} z_2^{n_2} z_3^{n_3} \equiv Q(a) z^{\vec{n}}. \quad (18)$$

Using (17) and (18) in (16) yields

$$\Delta(z^{\vec{m}}) = \sum_{|\vec{n}|=N} \left(\sum_{\substack{|\vec{a}_p|=m_p \\ |\vec{a}_j|=n_j}} Q(a) \left[\begin{matrix} \vec{m} \\ a \end{matrix} \right]_{q^{-2}} \prod_{k=1}^3 \left(\prod_{i=1}^3 x_{ik}^{a_{ik}} \right) \otimes \prod_{k=1}^3 x_{kj}^{n_k} \right),$$

where we introduced $\underline{a}_j \equiv (a_{1j}, a_{2j}, a_{3j})$ with the sums over $|\vec{a}_p| = m_p$ and $|\vec{a}_j| = n_j$ being sums over all the $\{a_{ij}\}_{i,j=1,2,3}$ satisfying $\sum_{j=1}^3 a_{ij} = m_i$ and $\sum_{i=1}^3 a_{ij} = n_j$. From this expression, one directly identifies the matrix elements of $\mathcal{F}_q^{(N)}(\mathbb{C}^3)$. \square

B. Unitarity

Unitary corepresentations can be constructed from the above corepresentations through normalization. We have the following theorem:

Theorem 2 (Unitarity). *The following $SU_q(3)$ corepresentation is unitary:*

$$\Delta : z^{\vec{m}} \left[\begin{matrix} N \\ \vec{m} \end{matrix} \right]_{q^{-2}}^{\frac{1}{2}} \mapsto \sum_{|\vec{n}|=N} t_{\vec{m},\vec{n}}^{(N)} \otimes z^{\vec{n}} \left[\begin{matrix} N \\ \vec{n} \end{matrix} \right]_{q^{-2}}^{\frac{1}{2}},$$

with the matrix elements given by

$$t_{\vec{m},\vec{n}}^{(N)} = \sqrt{\left[\begin{matrix} N \\ \vec{m} \end{matrix} \right]_{q^{-2}} \left[\begin{matrix} N \\ \vec{n} \end{matrix} \right]_{q^{-2}}^{-1}} \sum_{\substack{|\vec{a}_j|=m_j \\ |\vec{a}_j|=n_j}} Q(a) \left[\begin{matrix} \vec{m} \\ a \end{matrix} \right]_{q^{-2}} \prod_{k=1}^3 \left(\prod_{i=1}^3 x_{ik}^{a_{ik}} \right), \tag{19}$$

and $Q(a)$ as in (15).

Proof. A right $SU_q(3)$ comodule is constructed similarly as the left comodule constructed in (12). Identifying the right comodule generators as $w_j = x_{ij}$ for i fixed, one has now

$$\Delta(w^{\vec{m}}) = \prod_{j=1}^3 \Delta(x_{ij}^{m_j}) = \sum_{\substack{|\vec{b}_p|=m_p \\ |\vec{b}_p|=m_p}} \left[\begin{matrix} \vec{m} \\ b \end{matrix} \right]_{q^{-2}} \prod_{j=1}^3 \left(\prod_{k=1}^3 (x_{ik} \otimes x_{kj})^{b_{kj}} \right),$$

where the notation for the indices b is the same as in Proposition 1. Writing $n_k = b_{k1} + b_{k2} + b_{k3}$ such that

$$\prod_{j=1}^3 \left(\prod_{k=1}^3 x_{ik}^{b_{kj}} \right) = Q(b^T) \prod_{k=1}^3 \left(\prod_{j=1}^3 x_{ik}^{b_{kj}} \right) = Q(b^T) \prod_{k=1}^3 x_{ik}^{n_k} = Q(b^T) w^{\vec{n}},$$

to obtain, given that (15) is symmetric,

$$\Delta(w^{\vec{m}}) = \sum_{|\vec{n}|=N} w^{\vec{n}} \otimes \tilde{h}_{\vec{n},\vec{m}}^{(N)}, \quad \tilde{h}_{\vec{n},\vec{m}}^{(N)} = \sum_{\substack{|\vec{b}_p|=n_p \\ |\vec{b}_p|=m_p}} Q(b) \left[\begin{matrix} \vec{m} \\ b \end{matrix} \right]_{q^{-2}} \prod_{j=1}^3 \left(\prod_{k=1}^3 x_{kj}^{b_{kj}} \right). \tag{20}$$

The generators of the left and right comodules are related by the $*$ operation in the following way:

$$(w_j)^* = (x_{ij})^* = S(x_{ji}) = S(z_j) \implies (w^{\vec{m}})^* = S(z^{\vec{m}}).$$

Thus, knowing that $\Delta \circ S(x) = (S \otimes S) \circ \tau \circ \Delta$ for $\tau(x \otimes y) \equiv y \otimes x$, one has on the one hand

$$\Delta(w^{\vec{m}})^* = \Delta \circ S(z^{\vec{m}}) \mapsto \sum_{|\vec{n}|=N} S(z^{\vec{n}}) \otimes S(h_{\vec{m},\vec{n}}) = \sum_{|\vec{n}|=N} (w^{\vec{n}})^* \otimes S(h_{\vec{m},\vec{n}}). \tag{21}$$

On the other hand, Δ being a $*$ -homomorphism, one has

$$\Delta(w^{\vec{m}})^* \mapsto \sum_{|\vec{n}|=N} (w^{\vec{n}})^* \otimes (\tilde{h}_{\vec{n},\vec{m}})^*. \tag{22}$$

Knowing that the $w^{\vec{n}}$ are linearly independent, it follows from (21) and (22) that

$$S(h_{\vec{m},\vec{n}}^{(N)}) = (\tilde{h}_{\vec{n},\vec{m}}^{(N)})^*. \tag{23}$$

Comparing (20) with the matrix elements $h_{\vec{n},\vec{m}}^{(N)}$ of the left coaction (14), one can see that they only differ by the q -trinomial coefficient. However, it is easy to show that

$$\begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}} \begin{bmatrix} \vec{m} \\ a \end{bmatrix}_{q^{-2}} = \begin{bmatrix} N \\ \vec{n} \end{bmatrix}_{q^{-2}} \begin{bmatrix} \vec{n} \\ a \end{bmatrix}_{q^{-2}}$$

so that with the proper normalization of the matrix elements, one has

$$t_{\vec{n},\vec{m}}^{(N)} \equiv \sqrt{\begin{bmatrix} N \\ \vec{n} \end{bmatrix}_{q^{-2}} \begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}}^{-1}} h_{\vec{n},\vec{m}}^{(N)} = \sum_{\substack{|\vec{a}_p|=n_p \\ |\vec{a}_p|=m_p}} Q(a) \sqrt{\begin{bmatrix} \vec{n} \\ a \end{bmatrix}_{q^{-2}} \begin{bmatrix} \vec{m} \\ a \end{bmatrix}_{q^{-2}}} \prod_{k=1}^3 \left(\prod_{i=1}^3 x_{ik}^{a_{ik}} \right) = \sqrt{\begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}} \begin{bmatrix} N \\ \vec{n} \end{bmatrix}_{q^{-2}}^{-1}} \tilde{h}_{\vec{n},\vec{m}}^{(N)} \equiv \tilde{t}_{\vec{n},\vec{m}}^{(N)}.$$

This normalization is equivalent to normalizing the basis elements of the left and right corepresentations as follows:

$$\underline{w}^{\vec{m}} \equiv \sqrt{\begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}}} w^{\vec{m}}, \quad \underline{z}^{\vec{m}} \equiv z^{\vec{m}} \sqrt{\begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}}}. \tag{24}$$

With this normalization and (23), one has

$$S(t_{\vec{m},\vec{n}}^{(N)}) = (t_{\vec{n},\vec{m}}^{(N)})^*, \tag{25}$$

which establishes (Ref. 18, Chap. 4.1) the unitarity of the corepresentations. \square

A direct corollary of Theorem 2 is the orthonormality of the matrix elements. Indeed, writing the $SU_q(3)$ product as $\nabla: SU_q(3) \otimes SU_q(3) \rightarrow SU_q(3)$, the hexagonal relation from the Hopf algebra structure gives

$$\sum_{|\vec{n}|=N} t_{\vec{m},\vec{n}} S(t_{\vec{n},\vec{p}}) = \nabla \circ (1 \otimes S) \circ \Delta(t_{\vec{m},\vec{p}}) = \eta \circ \varepsilon(t_{\vec{m},\vec{p}}) = \delta_{\vec{m},\vec{p}}, \tag{26}$$

where the last equality relies on the fact that the counit ε vanishes on off-diagonal generators of $SU_q(3)$ and also that the single term in (19) containing only diagonal generators has coefficient one. Upon using (25) in (26), one obtains

$$\sum_{|\vec{n}|=N} t_{\vec{m},\vec{n}}^* t_{\vec{p},\vec{n}}^* = \delta_{\vec{m},\vec{p}}, \quad \sum_{|\vec{m}|=N} t_{\vec{m},\vec{n}}^* t_{\vec{m},\vec{p}} = \delta_{\vec{n},\vec{p}}, \tag{27}$$

where a similar argument is used to obtain the second identity.

C. Generating function

A two-sided generating function for the matrix elements is easily obtained. To do so, first observe that

$$(w_i \otimes \Delta(z_i))(w_j \otimes \Delta(z_j)) = q^2 (w_j \otimes \Delta(z_j))(w_i \otimes \Delta(z_i)), \quad \forall i < j,$$

so that

$$\left(\sum_{i=1}^3 w_i \otimes \Delta(z_i) \right)^N = \sum_{|\vec{m}|=N} \begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}} w^{\vec{m}} \otimes \Delta(z^{\vec{m}}) = \sum_{\substack{|\vec{m}|=N \\ |\vec{n}|=N}} \underline{w}^{\vec{m}} \otimes t_{\vec{m},\vec{n}}^{(N)} \otimes \underline{z}^{\vec{n}}.$$

Expanding the left-hand side, one finds

$$\left(\sum_{i,j=1}^3 w_i \otimes x_{ij} \otimes z_j \right)^N = \sum_{\substack{|\vec{m}|=N \\ |\vec{n}|=N}} \underline{w}^{\vec{m}} \otimes t_{\vec{m},\vec{n}}^{(N)} \otimes \underline{z}^{\vec{n}}, \tag{28}$$

with the normalized basis elements defined in (24). From this generating function, one can obtain a structure relation for the matrix elements as follows: With (24), one has that

$$w_i \underline{w}^{\vec{m}} = q^{-2 \sum_{k=1}^{i-1} m_k} \left[\frac{1 - q^{2m_i - 2}}{1 - q^{2N - 2}} \right]^{1/2} \underline{w}^{\vec{m} + \vec{e}_i},$$

and likewise for $z_j z^{\vec{n}}$. Using (28) and the above, with the convention that the matrix elements of negative indices vanish, one has

$$\begin{aligned} \sum_{\substack{|\vec{m}|=N+1 \\ |\vec{n}|=N+1}} \underline{w}^{\vec{m}} \otimes t_{\vec{m},\vec{n}}^{(N+1)} \otimes \underline{z}^{\vec{n}} &= \left(\sum_{i,j=1}^3 w_i \otimes x_{ij} \otimes z_j \right) \sum_{\substack{|\vec{m}|=N \\ |\vec{n}|=N}} \underline{w}^{\vec{m}} \otimes t_{\vec{m},\vec{n}}^{(N)} \otimes \underline{z}^{\vec{n}} \\ &= (1 - q^{2N-2})^{-1} \sum_{\substack{|\vec{m}|=N+1 \\ |\vec{n}|=N+1}} \underline{w}^{\vec{m}} \otimes \left[\sum_{i,j=1}^3 q^{-2 \sum_{k=1}^{i-1} m_k - 2 \sum_{l=1}^{j-1} n_l} \times \sqrt{(1 - q^{2m_i-2})(1 - q^{2n_j-2})} x_{ij} t_{\vec{m}-\vec{e}_i, \vec{n}-\vec{e}_j}^{(N)} \right] \otimes \underline{z}^{\vec{n}}. \end{aligned}$$

Finally, as the basis element is linearly independent, one arrives at

$$(1 - q^{2N-2}) t_{\vec{m},\vec{n}}^{(N+1)} = \sum_{i,j=1}^3 q^{-2 \sum_{k=1}^{i-1} m_k - 2 \sum_{l=1}^{j-1} n_l} \sqrt{(1 - q^{2m_i-2})(1 - q^{2n_j-2})} x_{ij} t_{\vec{m}-\vec{e}_i, \vec{n}-\vec{e}_j}^{(N)},$$

which is a structure relation for the matrix elements (19). Note that a similar structure relation exists where the multiplication by x_{ij} is to the right.

IV. MATRIX ELEMENTS IN $SU_q(3)$ REPRESENTATIONS

In this section, the matrix elements (19) are shown to be q -Krawtchouk polynomials. Following Ref. 6, we first identify the matrix elements in the elementary representations π_1 and π_2 as univariate quantum q -Krawtchouk polynomials. Then, the matrix elements in the π_{21} representation are identified as the Tratnik type bivariate quantum q -Krawtchouk polynomials. Finally, it is shown the matrix elements in the π_{121} can be expressed in terms of the same polynomials.

A. Elementary representations

Consider the matrix elements (19) in the π_1 and π_2 representations. That these matrix elements are given in terms of univariate quantum q -Krawtchouk polynomials is a corollary of a previous result of Koornwinder,⁶ as these $SU_q(3)$ representations were constructed from an $SU_q(2)$ representation. However, as we now work in an $SU_q(3)$ representation instead of the algebra itself, a derivation in the current context is given. The univariate quantum q -Krawtchouk polynomials²⁰ are defined²¹ by

$$k_n(x; p, N, q) = (-1)^n (q^{-N}; q)_n q^{n(n-1)/2} {}_2\phi_1 \left(q^{-n}, q^{-x}; q^{-N}; q, pq^{n+1} \right) \quad (29)$$

and are orthogonal with respect to the weight $w_x(p)^2$ for $x \in \{0, \dots, N\}$, where

$$w_x(p) = \left[(-1)^{N-x} q^{x(x-1)/2} \begin{bmatrix} N \\ x \end{bmatrix}_q \frac{(pq; q)_{N-x}}{(q; q)_N} p^{-N} q^{-N(N+1)/2} \right]^{1/2}, \quad (30)$$

with normalization

$$\Theta_n(p) = \frac{q^{-n(n-1)/2}}{(q^{-N}; q)_n} \left[(-1)^n q^{n(n+1)/2 - Nn} \begin{bmatrix} N \\ n \end{bmatrix}_q \frac{(q; q)_N}{(qp; q)_n} \right]^{1/2}. \quad (31)$$

One has the following proposition.

Proposition 2. The matrix elements (19) evaluated in the elementary representations π_i with $i = 1, 2$ as defined in (4) and (5), are shift operators given by

$$\pi_i \left(t_{\vec{m},\vec{n}}^{(N)} \right) |k\rangle = (\delta_{i,2} \delta_{m_1, n_1} + \delta_{i,1} \delta_{N-m_1-m_2, N-n_1-n_2}) t_{m_i, n_i, T_i}(k) |k + T_i - m_i - n_i\rangle, \quad (32)$$

$$t_{m,n,T}(k - T + n) \equiv (-1)^{n-m} w_n(q^{-2(k+1)}) \Theta_m(q^{-2(k+1)}) k_m(q^{-2n}; q^{-2(k+1)}, T, q^2). \quad (33)$$

Proof. Looking at (4), the monomials in the $SU_q(3)$ generators in (19) evaluate to

$$\pi_1 \left(\prod_{i,j=1}^3 x_{ij}^{a_{ij}} \right) |k\rangle = (-1)^{a_{21}} \delta_{a_{33}, m_3} \prod_{i=1}^2 \delta_{a_{3i}, 0} \delta_{a_{i3}, 0} q^{a_{12}(k+a_{22}+1) + a_{21}(k+a_{22})} \sqrt{(q^{2k+2}, q^2)_{a_{22}} (q^{2(k+a_{22})}, q^{-2})_{a_{11}}} |k + a_{22} - a_{11}\rangle.$$

Using the above and writing $T = N - m_3 = N - n_3$, the matrix elements simplifies to

$$\pi_1 \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |k\rangle = \sum_{a_{21}=0}^{m_2} \begin{bmatrix} m_1 \\ a_{11} \end{bmatrix}_{q^{-2}} \begin{bmatrix} m_2 \\ a_{21} \end{bmatrix}_{q^{-2}} q^{-a_{21}(a_{12})} (-1)^{a_{21}} q^{a_{12}(k+a_{22}+1)+a_{21}(k+a_{22})} \\ \times \sqrt{\begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}} \begin{bmatrix} N \\ \vec{n} \end{bmatrix}_{q^{-2}}^{-1}} \sqrt{(q^{2k+2}; q^2)_{a_{22}} (q^{2(k+a_{22})}; q^{-2})_{a_{11}}} |k + a_{22} - a_{11}\rangle.$$

Taking the following parametrization of the summation indices

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} T - n_2 - i & n_2 - m_2 + i \\ i & m_2 - i \end{pmatrix},$$

one has

$$\pi_1 \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |k\rangle = (-1)^{T-n_2} q^{(n_2-m_2)(k+m_2+1)+n_1(n_1+1)} \\ \times \sqrt{(q^{2(k+1)}; q^2)_{m_2} (q^{2(k+m_2-n_1+1)}; q^2)_{n_1}} \frac{(q^{-2m_1}; q^2)_{n_1}}{(q^2; q^2)_{n_1}} |k - n_1 + m_2\rangle \\ \times \sqrt{\begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}} \begin{bmatrix} N \\ \vec{n} \end{bmatrix}_{q^{-2}}^{-1}} \sum_{i=0}^{m_2} \frac{(q^{-2n_1}; q^2)_i (q^{-2m_2}; q^2)_i}{(q^{2(n_2-m_2+1)}; q^2)_i (q^{-2(k+m_2)}; q^2)_i (q^2; q^2)_i} q^{2i}.$$

Using Jackson's identity (Ref. 22, Sec. III.5),

$$\left(q^{-n} \frac{z}{b}; q \right)_n {}_3\phi_2 \left(\begin{matrix} q^{-n}, b, 0 \\ qb/z, c \end{matrix}; q, q \right) = {}_2\phi_1 \left(\begin{matrix} q^{-n}, c/b \\ c \end{matrix}; q, z \right),$$

one can write the ${}_3\phi_2$ as a ${}_2\phi_1$,

$$\pi_1 \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |k\rangle = (-1)^{T-n_2} q^{(n_2-m_2)(k+m_2+1)+n_1(n_1+1)} \sqrt{\frac{(q^{2(k+m_2-n_1+1)}; q^2)_{n_1}}{(q^{2(k+1)}; q^2)_{m_2}} \begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}} \begin{bmatrix} N \\ \vec{n} \end{bmatrix}_{q^{-2}}^{-1}} \\ \times \frac{(q^{-2m_1}; q^2)_{n_1}}{(q^2; q^2)_{n_1}} {}_2\phi_1 \left(\begin{matrix} q^{-2m_2}, q^{2(n_1+n_2-m_2+1)} \\ q^{2(n_2-m_2+1)} \end{matrix}; q^2, q^{2(k+1-n_1+m_2)} \right) |k - n_1 + m_2\rangle.$$

Finally, one reverses the order of summation using²² [Exercise 1.4 (ii)]

$${}_2\phi_1 \left(\begin{matrix} q^{-n}, b \\ c \end{matrix}; q, z \right) = (-1)^n q^{-n(n+1)/2} \frac{(b; q)_n}{(c; q)_n} z^n {}_2\phi_1 \left(\begin{matrix} q^{-n}, q^{-n+1} c^{-1} \\ q^{-n+1} b^{-1} \end{matrix}; q, \frac{q^{n+1} c}{bz} \right),$$

which, after shifting the parameter k by $-n_2$ and using q -Pochhammer symbol identities, leads to

$$\pi_1 \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |k\rangle = (-1)^{m_2} q^{k(n_2+m_2)+n_2(m_2+1)-2n_1n_2} \sqrt{\begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^{-2}} \begin{bmatrix} N \\ \vec{n} \end{bmatrix}_{q^{-2}}^{-1}} \frac{(q^{2(k-n_1+1)}; q^2)_{n_1}}{(q^{2(k-n_1+1)}; q^2)_{m_2}} \\ \times \begin{bmatrix} T \\ n_1 \end{bmatrix}_{q^2} {}_2\phi_1 \left(\begin{matrix} q^{-2m_2}, q^{-2n_2} \\ q^{-2(n_1+n_2)} \end{matrix}; q^2, q^{-2k} \right) |k - n_1 + m_2\rangle.$$

We now use the following of Heine's transformation formulas (Ref. 22, Sec. III.3)

$${}_2\phi_1 \left(\begin{matrix} q^{-2m}, q^{-2p} \\ q^{-2N} \end{matrix}; q^2, q^{-2k+2p} \right) = \frac{1}{(q^{-2k+2p}; q^2)_{N-m-p}} {}_2\phi_1 \left(\begin{matrix} q^{-2(N-m)}, q^{-2(N-p)} \\ q^{-2N} \end{matrix}; q^2, q^{-2k+2(N-m)} \right),$$

to obtain

$$\begin{aligned} \pi_1 \left(t_{\bar{m}, \bar{n}}^{(N)} \right) |k - T + n_1\rangle &= \delta_{N-m_1-m_2, N-n_1-n_2} (-1)^{n_1} q^{n_1^2 - n_1 T} \\ &\times \left[(-1)^{T-n_1+m_1} q^{2T(k+1)-T(T+1)-m_1(m_1-1)+n_1(n_1-1)} \frac{(q^{-2k}, q^2)_{T-n_1}}{(q^{-2k}, q^2)_{m_1}} \left[\begin{matrix} N \\ \bar{m} \end{matrix} \right]_{q^{-2}} \left[\begin{matrix} N \\ \bar{n} \end{matrix} \right]_{q^{-2}}^{-1} \right]^{1/2} \\ &\times \left[\begin{matrix} T \\ n_1 \end{matrix} \right]_{q^2} {}_2\phi_1 \left(q^{-2m_1}, q^{-2n_1}; q^2, q^{-2k+2m_1} \right) |k - m_1\rangle. \end{aligned} \quad (34)$$

Proceeding similarly, one obtains for π_2

$$\begin{aligned} \pi_2 \left(t_{\bar{m}, \bar{n}}^{(N)} \right) |k - T + n_2\rangle &= \delta_{m_1, n_1} (-1)^{n_2} q^{n_2^2 - n_2 T} \\ &\times \left[(-1)^{T-n_2+m_2} q^{2T(k+1)-T(T+1)-m_2(m_2-1)+n_2(n_2-1)} \frac{(q^{-2k}, q^2)_{T-n_2}}{(q^{-2k}, q^2)_{m_2}} \left[\begin{matrix} N \\ \bar{m} \end{matrix} \right]_{q^{-2}} \left[\begin{matrix} N \\ \bar{n} \end{matrix} \right]_{q^{-2}}^{-1} \right]^{1/2} \\ &\times \left[\begin{matrix} T \\ n_2 \end{matrix} \right]_{q^2} {}_2\phi_1 \left(q^{-2m_2}, q^{-2n_2}; q^2, q^{-2k+2m_2} \right) |k - m_2\rangle. \end{aligned} \quad (35)$$

From (34) and (35), one directly finds (32) and (33), which concludes the proof. \square

1. Orthogonality

The orthogonality relation of the univariate q -Krawtchouk polynomials of the quantum type follows from the unitarity of the corepresentation, upon using (27). The adjoint of (32) is calculated easily as

$$\pi_1 \left(t_{\bar{m}, \bar{n}}^{(N)} \right)^* |k\rangle = (-1)^{n_1 - m_1} t_{m_1, n_1, T}(k - T + m_1 + n_1) |k - T + m_1 + n_1\rangle.$$

Then, from (27), one has

$$\begin{aligned} \delta_{\bar{n}, \bar{p}} \langle k' | k \rangle &= \sum_{\bar{m}} \langle k' | \pi_1 \left(t_{\bar{m}, \bar{p}}^{(N)} \right)^* \pi_1 \left(t_{\bar{m}, \bar{n}}^{(N)} \right) |k\rangle \\ &= \sum_{\bar{m}} \langle k' | \pi_1 \left(t_{\bar{m}, \bar{p}}^{(N)} \right)^* |k + T - m_1 - n_1\rangle (-1)^{n_1 - m_1} t_{m_1, n_1, T}(k) \\ &= (-1)^{n_1 - p_1} \langle k' | k + p_1 - n_1 \rangle \sum_{\bar{m}} t_{m_1, p_1, T}(k - n_1 + p_1) t_{m_1, n_1, T}(k) \end{aligned} \quad (36)$$

so that

$$\delta_{p_1, n_1} = (-1)^{n_1 - p_1} \sum_{\bar{m}} t_{m_1, p_1, T}(k + p_1 - T) t_{m_1, n_1, T}(k + n_1 - T), \quad (37)$$

where in the last line, the parameter k was shifted by $n_1 - T$. Writing $m_1 = m$, $n_1 = n$ and $p_1 = p$ and setting $N = T$, one uses (33) in (37) while rearranging to obtain

$$\begin{aligned} (-1)^{T-n} \left[\begin{matrix} T \\ n \end{matrix} \right]_{q^2}^{-1} \frac{(q^2; q^2)_T}{(q^{-2k}, q^2)_{T-n}} q^{-n(n-1)} \delta_{p, n} &= \sum_{m=0}^T (-1)^m q^{2T(k+1)} \left[\begin{matrix} T \\ m \end{matrix} \right]_{q^2} \frac{(q^2; q^2)_T}{(q^{-2k}, q^2)_m} q^{-T(T+1)+m(m+1)-2mT} \\ &\times {}_2\phi_1 \left(q^{-2m}, q^{-2p}; q^2, q^{-2k+2m} \right) {}_2\phi_1 \left(q^{-2m}, q^{-2n}; q^2, q^{-2k+2m} \right), \end{aligned}$$

which one can identify as the dual orthogonality relation of the univariate quantum q -Krawtchouk polynomials as given in Ref. 20.

B. Product representations

We now demonstrate that the matrix elements in the representation π_{21} are expressed in terms of bivariate quantum q -Krawtchouk polynomials of the Tratnik type. These polynomials are defined in Ref. 10 as

$$K_{n, m}(x, y; u, v, N, q) = k_n(x; v^{-2}, x + y, q) k_m(x + y - n; u^{-2}, N - n, q), \quad (38)$$

with $k_n(x; p, N, q)$ as in (29). They are orthogonal with respect to the weight $W_{n_1, n_2}^{(N)}(u, v)^2$ where

$$W_{n_1, n_2}^{(N)}(u, v) = \left[(-1)^{N-n_1} q^{2v(n_1+n_2)} q^{n_1(n_1-1)} \left[\begin{matrix} N \\ \bar{n} \end{matrix} \right]_{q^2} (q^{-2v}; q^2)_{n_2} (q^{-2u}; q^2)_{N-n_1-n_2} q^{2N(u+1)} q^{-N(N+1)} \right]^{1/2}, \quad (39)$$

with the normalization given by

$$N_{m_1, m_2}^{(N)}(u, v) = q^{-m_1(m_1-1) - m_2(m_2-1)} q^{-m_1(u+1)} \frac{(q^2; q^2)_{N-m_1-m_2}}{(q^2; q^2)_N} \times \left[(-1)^{m_1+m_2} \begin{bmatrix} N \\ \vec{m} \end{bmatrix}_{q^2} \frac{q^{2N(m_1+m_2)+4m_1+2m_2-2m_1m_2-m_1(m_1-1)-m_2(m_2-1)}}{(q^{-2u}; q^2)_{m_2} (q^{-2v}; q^2)_{m_1}} \right]^{1/2}. \quad (40)$$

With these notations, one has the following proposition.

Proposition 3. The matrix elements (19) evaluated in π_{21} as defined in (6) are shift operators specified by

$$\pi_{21} \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |u, v\rangle = t_{\vec{m}, \vec{n}}^{(21)}(u, v) |u - m_2 + N - n_1 - n_2, v + n_2 - m_1\rangle, \quad (41)$$

where

$$t_{\vec{m}, \vec{n}}^{(21)}(u - N + n_1 + n_2, v - n_2) = (-1)^{m_1 - n_2} W_{n_1, n_2}^{(N)}(u, v) N_{m_1, m_2}^{(N)}(u, v) K_{m_1, m_2}(n_1, n_2; q^{u+1}, q^{v+1}, N, q^2) \quad (42)$$

are the normalized and weighted bivariate quantum q -Krawtchouk polynomials of the Tratnik type.

Proof. From (6), one has

$$\pi_{21} \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |u, v\rangle = (\pi_2 \otimes \pi_1) \circ \Delta \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |u, v\rangle = \sum_{|\vec{k}|=N} \pi_2 \left(t_{\vec{m}, \vec{k}}^{(N)} \right) |u\rangle \otimes \pi_1 \left(t_{\vec{k}, \vec{n}}^{(N)} \right) |v\rangle. \quad (43)$$

Using (32) of Proposition 2 in (43), one sees that the Kronecker deltas δ_{m_1, k_1} and $\delta_{k_1+k_2, n_1+n_2}$ remove the sums, and one has

$$k_1 = m_1, \quad k_2 = (n_1 + n_2 - m_1),$$

which implies $T_1 = n_1 + n_2$ and $T_2 = N - m_1$. One can, thus, identify (43) as the shift operator in (41). Then, shifting u by $n_1 + n_2 - N$ and v by n_2 in (43), one has that

$$t_{\vec{m}, \vec{n}}^{(21)}(u - N + n_1 + n_2, v - n_2) = t_{m_2, n_1+n_2-m_1, N-m_1}(u - N + n_1 + n_2) t_{m_1, n_1, n_1+n_2}(v - n_2). \quad (44)$$

Using (33) from Proposition 2 in (44), one obtains by involved but direct computations

$$t_{\vec{m}, \vec{n}}^{(21)}(u - N + n_1 + n_2, v - n_2) = (-1)^{m_1 - n_2} W_{n_1, n_2}^{(N)}(u, v) N_{m_1, m_2}^{(N)}(u, v) K_{m_1, m_2}(n_1, n_2; q^{u+1}, q^{v+1}, N, q^2),$$

which concludes the proof. \square

The expression in (42) for the scalar factor of the matrix elements (19) evaluated in π_{21} has been obtained in a related but different approach in Ref. 3; see also Ref. 23. There, (42) arises as an expression for the matrix elements in symmetric representations of unitary elements of $U_q(\mathfrak{su}(3))$ constructed from q -exponentials. This correspondence is expected in view of the duality (Refs. 24 and 18, Chap. 7) between quantum algebras and groups and has been discussed⁹ in the context of q -special functions. The parameters of the polynomials in (42) are discrete. However, one recovers the usual q -Krawtchouk polynomials with continuous parameters by extension with analytic continuation.

C. Representation corresponding to the longest Weyl group element

We now study the matrix elements in π_{121} . In this case, one has

$$\pi_{121} \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |t, u, v\rangle = (\pi_1 \otimes \pi_{21}) \circ \Delta \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |t, u, v\rangle = \sum_{|\vec{k}|=N} \pi_1 \left(t_{\vec{m}, \vec{k}}^{(N)} \right) |t\rangle \otimes \pi_{21} \left(t_{\vec{k}, \vec{n}}^{(N)} \right) |u, v\rangle.$$

Using (32), (33), (41), and (42) in the above, one obtains

$$\pi_{121} \left(t_{\vec{m}, \vec{n}}^{(N)} \right) |t, u, v\rangle = \sum_{k_1=0}^{m_1+m_2} (-1)^{k_1-m_1} t_{m_1, k_1, T_1}(t) t_{\vec{k}, \vec{n}}^{(21)}(u, v) |t + m_2 - k_1, u + N - m_1 - m_2 - n_1 - n_2 + k_1, v + n_2 - k_1\rangle. \quad (45)$$

As $t_{m_1, k_1, T_1}(t)$ does not depend on the variables, one can identify the scalar coefficients of (45) as bivariate q -Krawtchouk polynomials normalized by a factor expressed as a univariate q -Krawtchouk polynomials. Thus, studying the matrix elements in π_{121} leads to the same polynomials. Indeed, the unitarity of the corepresentations is equivalent to the orthogonality relation of the Tratnik bivariate q -Krawtchouk polynomials (38)

$$\delta_{\vec{n}, \vec{p}} \langle a', b', c' | a, b, c \rangle = \sum_{|\vec{m}|=N} \langle a', b', c' | \pi_{121} \left(t_{\vec{m}, \vec{n}}^{(N)} \right)^* \pi_{121} \left(t_{\vec{m}, \vec{p}}^{(N)} \right) | a, b, c \rangle,$$

which, after shifting b by $p_1 + p_2$ and c by $-p_2$, is given by

$$\delta_{\vec{n}, \vec{p}}(a', b', c' | a, b + p_1 + p_2, c - p_2) = \sum_{|\vec{m}|=N} \sum_{k_1=0}^{m_1+m_2} \sum_{l_1=0}^{m_1+m_2} (-1)^{k_1+l_1} t_{m_1, k_1, T_1}(a) t_{m_1, l_1, T_1}(a - k_1 + l_1) t_{\vec{k}, \vec{p}}^{(21)}(b + p_1 + p_2, c - p_2) \times t_{\vec{l}, \vec{n}}^{(21)}(b + n_1 + n_2 + k_1 - l_1, c - n_2 - k_1 + l_1) \langle a', b', c' | a - k_1 + l_1, b + n_1 + n_2 + k_1 - l_1, c - n_2 - k_1 + l_1 \rangle.$$

Noticing that the overlap $\langle a' | a - k_1 + l_1 \rangle$ fixes the difference $l_1 - k_1 \equiv s$, one can rearrange the sums to obtain

$$\delta_{\vec{n}, \vec{p}}(a - s, b', c' | a, b + p_1 + p_2, c - p_2) = \sum_{|\vec{k}|=N} \sum_{m_1=0}^{k_1+k_2} t_{\vec{k}, \vec{p}}^{(21)}(b + p_1 + p_2, c - p_2) t_{\vec{k}+s(\vec{e}_1-\vec{e}_2), \vec{n}}^{(21)}(b + n_1 + n_2 - s, c - n_2 + s) \times (-1)^s t_{m_1, k_1, T_1}(a) t_{m_1, k_1-s, T_1}(a - s) \langle b', c' | b + n_1 + n_2 - s, c - n_2 + s \rangle.$$

Using the univariate q -Krawtchouk dual orthogonality relation²⁰ in the sum over m_1 forces s to vanish so that one has

$$\delta_{\vec{n}, \vec{p}}(b', c' | b + p_1 + p_2, c - p_2) = \sum_{|\vec{k}|=N} t_{\vec{k}, \vec{p}}^{(21)}(b + p_1 + p_2, c - p_2) t_{\vec{k}, \vec{n}}^{(21)}(b + n_1 + n_2, c - n_2) \langle b', c' | b + n_1 + n_2, c - n_2 \rangle,$$

which one can recognize as the dual orthogonality relation of the bivariate q -Krawtchouk polynomials of the Tratnik type.

V. REDUCIBLE TENSOR PRODUCTS

The algebraic interpretation of the multivariate q -Krawtchouk polynomials presented in this paper can be used to derive identities for these polynomials. Consider, for example, the reducible tensor product of the $SU_q(2)$ representations $\tau_\alpha \otimes \tau_\beta$ with $\alpha, \beta \in S^1$. It is known²⁵ to decompose into the direct integral

$$\rho = \int_{S^1}^{\oplus} \tau_\gamma d\gamma,$$

with the intertwiner $\Lambda: \tau_\alpha \otimes \tau_\beta \rightarrow \rho$ acting as follows on the basis vectors²⁶

$$\Lambda |v, t\rangle = \sum_{w \in \mathbb{N}} \alpha^{t-w} \beta^{w-v} \bar{p}_v(q^{2w}; q^{2|t-v|}; q^2) \frac{\gamma^{-v}}{\sqrt{2\pi}} \otimes |w\rangle,$$

where the Clebsch-Gordan coefficients, given by

$$\bar{p}_v(q^{2w}; q^{2|t-v|}; q^2) = (-1)^{v+w} \sqrt{\frac{q^{2(w-v)(|t-v|+1)} (q^{2|t-v|+2}; q^2)_\infty (q^{2|t-v|+2}; q^2)_v}{(q^2; q^2)_v (q^2; q^2)_w}} {}_2\phi_1\left(\begin{matrix} q^{-2v}, 0 \\ q^{2|t-v|+2} \end{matrix}; q^2, q^{2w+2}\right),$$

are the weighted and normalized Wall polynomials.²⁰ This result can be used in two ways to calculate the matrix elements in the π_{211} representation. One has, on one hand,

$$\pi_{211}\left(t_{\vec{m}, \vec{n}}^{(N)}\right) |u, v, t\rangle = \int_{S^1} d\kappa \sum_{k_1=0}^N \sum_{w' \in \mathbb{N}} (\sqrt{2\pi})^{-1/2} |u - m_2 + n_3, w'\rangle \alpha^{t+n_2-k_1-w'} \beta^{w'-v-n_1-n_2+k_1+m_1} \kappa^{m_1+k_1-n_1-n_2-v} \times t_{\vec{m}, (k_1, n_1+n_2-k_1)}^{(21)}(u, v) t_{k_1, n_1, T_1}(t) \bar{p}_{v+n_1+n_2-k_1-m_1}(q^{2w'}; q^{2|t+m_1-v-n_1|}; q^2), \quad (46)$$

and, on the other hand,

$$\begin{aligned} \pi_{211}\left(t_{\vec{m}, \vec{n}}^{(N)}\right) |u, v, t\rangle &= \int_{S^1} d\gamma \sum_{w \in \mathbb{N}} \frac{\alpha^{t-w} \beta^{w-v} \gamma^{-v}}{\sqrt{2\pi}} \bar{p}_v(q^{2w}; q^{2|t-v|}; q^2) \pi_{21}\left(t_{\vec{m}, \vec{n}}^{(N)}\right) |u, w\rangle \\ &= \int_{S^1} d\gamma \sum_{w \in \mathbb{N}} \frac{\alpha^{t-w} \beta^{w-v} \gamma^{-v}}{\sqrt{2\pi}} \bar{p}_v(q^{2w}; q^{2|t-v|}; q^2) t_{\vec{m}, \vec{n}}^{(21)}(u, w) |u - m_2 + n_3, w + n_2 - m_1\rangle. \end{aligned} \quad (47)$$

Taking the inner product on $S^1 \otimes V_\alpha \otimes V_\gamma$ of the left-hand sides of (46) and (47) with

$$\frac{\gamma^{-v}}{\sqrt{2\pi}} \otimes |u - m_2 + n_3, w + n_2 - m_1\rangle,$$

for a fixed but arbitrary w and equating the two results gives

$$\begin{aligned} \tilde{p}_v(q^{2w}; q^{2|t-v|}; q^2) t_{\tilde{m}, \tilde{n}}^{(21)}(u, w) &= \sum_{k_1=0}^N \alpha^{m_1-k_1} \beta^{k_1-n_1} t_{k_1, n_1, m_1+n_2}(t) \\ &\times \tilde{p}_{v+n_1+n_2-k_1-m_1}(q^{2(w+n_2-m_1)}; q^{2|t+m_1-v-n_1|}; q^2) t_{\tilde{m}, (k_1, n_1+n_2-k_1)}^{(21)}(u, v). \end{aligned}$$

Upon taking $\alpha = \beta = \gamma = -1$ while shifting u by $n_1 + n_2 - N$, v by $-n_1 - n_2$, w by $-n_2$, and t by $-n_2$, one can use (33) and (42) to express the above in terms of the polynomials, obtaining

$$\begin{aligned} \tilde{p}_{v-n_1-n_2}(q^{2w-2n_2}; q^{2|t+n_1-v|}; q^2) K_{m_1, m_2}(n_1, n_2; q^{u+1}, q^{w+1}, N, q^2) &= \\ \sum_{j=0}^N C_{\tilde{m}, \tilde{n}, j}^{(N)}(u, v, t) \tilde{p}_{v-j-m_1}(q^{2(w-n_2-m_1)}; q^{2|t+m_1-v|}; q^2) & \\ \times k_j(q^{-2n_1}; q^{-2(t+1)}, n_1 + n_2, q^2) K_{m_1, m_2}(j, n_1 + n_2 - j; q^{u+1}, q^{v-j+1}, N, q^2), & \quad (48) \end{aligned}$$

where

$$C_{\tilde{m}, \tilde{n}, j}^{(N)}(u, v, t) = (-1)^{m_1-n_1} w_{n_1}(q^{-2(t+1)}) \Theta_j(q^{-2(t+1)}) \frac{W_{j, n_1+n_2-j}^{(N)}(u, v-j)}{W_{n_1, n_2}^{(N)}(u, w)} \sqrt{\frac{(q^{-2w}; q^2)_{m_1}}{(q^{-2v+2j}; q^2)_{m_1}}},$$

with w , Θ , and W as in (30), (31), and (39), respectively. Consequently, (48) provides an identity for the product of a bivariate quantum q -Krawtchouk polynomial and a Wall polynomial.

VI. CONCLUSION

This paper identified the matrix elements of the $SU_q(3)$ symmetric corepresentations as the bivariate quantum q -Krawtchouk polynomials of the Tratnik type. This was done by first constructing the symmetric unitary corepresentations and obtaining abstract expressions for the matrix elements and then evaluating these expressions in $SU_q(3)$ representations. These results, thus, provide an algebraic interpretation for the bivariate q -Krawtchouk polynomials within the quantum group setting. Moreover, this identification of the matrix elements evaluated in $SU_q(3)$ representations is complete in the sense that it held for the most general irreducible $*$ -representation. Finally, this paper illustrated how the quantum group interpretation could be exploited to obtain properties of the q -Krawtchouk polynomials.

The results presented here should admit a generalization to the generic case of $SU_q(N)$ that would yield a similar algebraic interpretation of the multivariate quantum q -Krawtchouk polynomials of the Tratnik type with valuable outcomes. In view of the relation between reducibility of $SU_q(3)$ representations and an identity for orthogonal polynomials, this algebraic interpretation might prove useful to obtain identities for the polynomials.

ACKNOWLEDGMENTS

G.B. would like to thank the Institute for Mathematics, Astrophysics, and Particle Physics at Radboud Universiteit for its hospitality and Paul Terwilliger for stimulating discussions. E.K. thanks the Centre de Recherches Mathématiques at the Université de Montréal for its hospitality. G.B. holds a scholarship of the Natural Science and Engineering Research Council of Canada (NSERC). The research of L.V. is supported by a Discovery Grant from NSERC.A.

REFERENCES

- ¹T. H. Koornwinder, "Orthogonal polynomials in connection with quantum groups," *Orthogonal Polynomials* **294**, 257–292 (1990).
- ²M. V. Tratnik, "Some multivariable orthogonal polynomials of the Askey tableau-discrete families," *J. Math. Phys.* **32**, 2337–2342 (1991).
- ³V. X. Genest, L. Vinet, and A. Zhedanov, "The multivariate Krawtchouk polynomials as matrix elements of the rotation group representations on oscillator states," *J. Phys. A: Math. Theor.* **46**, 505203 (2013).
- ⁴P. Iliev, "A Lie-theoretic interpretation of multivariate hypergeometric polynomials," *Compositio Math.* **148**, 991–1002 (2012); e-print [arXiv:1101.1683](https://arxiv.org/abs/1101.1683).
- ⁵P. Iliev and P. Terwilliger, "The Rahman polynomials and the Lie algebra $\mathfrak{sl}_3(\mathbb{C})$," *Trans. Am. Math. Soc.* **364**, 4225–4238 (2012).
- ⁶T. Koornwinder, "Representations of the twisted $SU(2)$ quantum group and some q -hypergeometric orthogonal polynomials," *Indagationes Math.* **92**, 97–117 (1989).
- ⁷A. S. Zhedanov, "Q rotations and other Q transformations as unitary nonlinear automorphisms of quantum algebras," *J. Math. Phys.* **34**, 2631–2647 (1993).
- ⁸V. X. Genest, S. Post, L. Vinet, G.-F. Yu, and A. Zhedanov, " q -Rotations and Krawtchouk polynomials," *Ramanujan J.* **40**, 335–357 (2016).
- ⁹R. Floreanini and L. Vinet, "On the quantum group and quantum algebra approach to q -special functions," *Lett. Math. Phys.* **27**, 179–190 (1993).
- ¹⁰G. Gasper and M. Rahman, "Some systems of multivariable orthogonal q -Racah polynomials," *Ramanujan J.* **13**, 389–405 (2007).
- ¹¹V. X. Genest, S. Post, and L. Vinet, "An algebraic interpretation of the multivariate q -Krawtchouk polynomials," *Ramanujan J.* **43**, 415–445 (2017); e-print [arXiv:1508.07770](https://arxiv.org/abs/1508.07770).
- ¹²K. Bragić, "The twisted $SU(3)$ group. Irreducible $*$ -representations of the C^* -Algebra $C(S_\mu U(3))$," *Lett. Math. Phys.* **17**, 37–44 (1989).

- ¹³Y. S. Soibelman, “Algebra of functions on a compact quantum group and its representations,” *Algebra Analiz* **2**(1), 190–212 (1990) [*Leningrad. Math. J.* **2**(1), 161–178 (1991)].
- ¹⁴H. T. Koelink, “On $*$ -representations of the Hopf $*$ -algebra associated with the quantum group $U_q(n)$,” *Compositio Math.* **77**(2), 199–231 (1991).
- ¹⁵L. I. Korogodski and Y. S. Soibelman, “Algebras of functions on quantum groups: Part I,” in *Mathematical Surveys and Monographs* (American Mathematical Society, 1998), Vol. 56, p. 150.
- ¹⁶L. L. Vaksman and Y. S. Soibelman, “Algebra of functions on the quantum group $SU(2)$,” *Funct. Anal. Appl.* **22**, 170–181 (1988).
- ¹⁷S. L. Woronowicz, “Compact matrix pseudogroups,” *Commun. Math. Phys.* **111**, 613–665 (1987).
- ¹⁸V. Chari and A. Pressley, *A Guide to Quantum Groups* (Cambridge University Press, 1994), p. 651.
- ¹⁹M. Noumi, H. Yamada, and K. Mimachi, “Finite dimensional representations of the quantum group $GL_q(n; \mathbb{C})$ and the zonal spherical functions on $U_q(n-1) \backslash U_q(n)$,” *Jpn. J. Math.* **19**, 31–80 (1993).
- ²⁰R. Koekoek, P. A. Lesky, and R. F. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their q -Analogues*, Springer Monographs in Mathematics Series (Springer, 2010).
- ²¹Part of the normalization of Ref. 20 is here included in the polynomial $k_n(x; p, N, q)$ themselves.
- ²²G. Gasper and M. Rahman, *Encyclopedia of Mathematics and its Application*, 2nd ed. (Cambridge University Press, 2004), Vol. 96, p. 428.
- ²³W. Groenevelt, “A quantum algebra approach to multivariate Askey-Wilson polynomials,” e-print [arXiv:1809.04327](https://arxiv.org/abs/1809.04327) (2018).
- ²⁴N. Burroughs, “Relating the approaches to quantised algebras and quantum groups,” *Commun. Math. Phys.* **133**, 91–117 (1990).
- ²⁵T. H. Koornwinder, “The addition formula for little q -Legendre polynomials and the $SU(2)$ quantum group,” *SIAM J. Math. Anal.* **22**, 295–301 (1991).
- ²⁶W. Groenevelt, “Coupling coefficients for tensor product representations of quantum $SU(2)$,” *J. Math. Phys.* **55**, 101702 (2014).