

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/215390>

Please be advised that this information was generated on 2020-10-21 and may be subject to change.

## Research Article

Blood-derived dendritic cell vaccinations induce immune responses that correlate with clinical outcome in patients with chemo-naive castration-resistant prostate cancer

1. Harm Westdorp<sup>1,2</sup>,
2. Jeroen H. A. Creemers<sup>1</sup>,
3. Inge M. van Oort<sup>3</sup>,
4. Gerty Schreibelt<sup>1</sup>,
5. Mark A. J. Gorris<sup>1</sup>,
6. Niven Mehra<sup>1,2</sup>,
7. Michiel Simons<sup>4</sup>,
8. Anna L. de Goede<sup>5</sup>,
9. Michelle M. van Rossum<sup>6</sup>,
10. Alexandra J. Croockewit<sup>7</sup>,
11. Carl G. Figdor<sup>1</sup>,
12. J. Alfred Witjes<sup>3</sup>,
13. Erik H. J. G. Aarntzen<sup>8</sup>,
14. Roel D. M. Mus<sup>8</sup>,
15. Mareke Brüning<sup>9</sup>,
16. Katja Petry<sup>9</sup>,
17. Martin Gotthardt<sup>8</sup>,
18. Jelle O. Barentsz<sup>8</sup>,
19. I. Jolanda M. de Vries<sup>1,2,s</sup> and
20. Winald R. Gerritsen<sup>1,2</sup>

## Author affiliations

### Abstract

**Background** Clinical benefit of cellular immunotherapy has been shown in patients with castration-resistant prostate cancer (CRPC). We investigated the immunological response and clinical outcome of vaccination with blood-derived CD1c<sup>+</sup> myeloid dendritic cells (mDCs; cDC2) and plasmacytoid DCs (pDCs).

**Methods** In this randomized phase IIa trial, 21 chemo-naive CRPC patients received maximally 9 vaccinations with mature mDCs, pDCs or a combination of mDCs plus pDCs. DCs were stimulated with protamine/mRNA and loaded with tumor-associated antigens NY-ESO-1, MAGE-C2 and MUC1. Primary endpoint was the immunological response after DC vaccination, which was monitored in peripheral blood and in T cell cultures of biopsies of post-treatment delayed-type hypersensitivity-skin tests. Main secondary endpoints were safety, feasibility, radiological PFS (rPFS) and overall survival. Radiological responses were assessed by MRIs and contrast-enhanced <sup>68</sup>Ga-prostate-specific membrane antigen PET/CT, according to RECIST 1.1, PCWG2 criteria and immune-related response criteria.

**Results** Both tetramer/dextramer-positive ( $dm^+$ ) and IFN- $\gamma$ -producing (IFN- $\gamma^+$ ) antigen specific T cells were detected more frequently in skin biopsies of patients with radiological non-progressive disease (5/13 patients; 38%) compared to patients with progressive disease (0/8 patients; 0%). In these patients with vaccination enhanced  $dm^+$  and IFN- $\gamma^+$  antigen-specific T cells median rPFS was 18.8 months ( $n = 5$ ) vs. 5.1 months ( $n = 16$ ) in patients without IFN- $\gamma$ -producing antigen-specific T cells ( $p = 0.02$ ). The overall median rPFS was 9.5 months. All DC vaccines were well tolerated with grade 1–2 toxicity.

**Conclusions** Immunotherapy with blood-derived DC subsets was feasible and safe and induced functional antigen-specific T cells. The presence of functional antigen-specific T cells correlated with an improved clinical outcome.

**Trial registration** ClinicalTrials.gov identifier [NCT02692976](https://clinicaltrials.gov/ct2/show/study/NCT02692976), registered 26 February 2016, retrospectively registered.

- Abbreviations:

combiDCs

combined CD1c<sup>+</sup> myeloid and plasmacytoid dendritic cells

CRPC

Castration-resistant prostate cancer

CTCAE

Common Terminology Criteria for Adverse Events

DCs

Dendritic cells

$dm^+$

tetramer or dextramer-positive

DTH

Delayed-type hypersensitivity

ECOG

Eastern Cooperative Oncology Group

GMP

Good Manufacturing Practices

IFN- $\gamma^+$

IFN- $\gamma$ -producing

KLH

Keyhole limpet hemocyanin

mDCs

CD1c<sup>+</sup> myeloid dendritic cells/cDC2

OS

Overall survival

PBMCs

Peripheral blood mononuclear cells

PCa

Prostate cancer


PCWG2

Prostate Cancer Clinical Trials Working Group 2  
pDCs  
plasmacytoid dendritic cells  
PSA  
Prostate-specific antigen  
PSAdt  
Prostate-specific antigen doubling time  
PSMA  
Prostate-specific membrane antigen  
RECIST  
Response Evaluation Criteria In Solid Tumours  
rPFS  
radiological progression-free survival  
SKILs  
Skin-test infiltrating lymphocytes  
TAA  
Tumor-associated antigen  
Th1  
T helper 1; Th2: T helper 2

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

[View Full Text](#)  
<http://dx.doi.org/10.1186/s40425-019-0787-6>

Statistics from Altmetric.com



[See more details](#)

Tweeted by 7

4 readers on Mendeley

- [Request Permissions](#)

## Background

Prostate cancer (PCa) remains the most common non-cutaneous malignancy and the second leading cause of cancer-related death in men [1]. For years, docetaxel-based chemotherapy was the only effective treatment for castration-resistant prostate cancer (CRPC) [2–4]. This changed with the approval of multiple agents, including androgen-signaling-targeted inhibitors abiraterone and enzalutamide [5–8], the cell-based vaccine sipuleucel-T [9], the radionuclide radium-223 [10] and second-line taxane cabazitaxel [11, 12]. These new agents extend overall survival (OS) with approximately 3–4 months [5–10, 13].

Recent advances in the field of cancer immunotherapy led to growing interest in prostate cancer immunotherapy. Immune-checkpoint inhibitor ipilimumab failed to show survival benefit in advanced PCa in phase III trials [14, 15]. Sipuleucel-T is still the only FDA-approved cellular immunotherapy for men with minimally symptomatic metastatic CRPC [9]. In Europe, sipuleucel-T is not available since its marketing authorization was withdrawn in 2015 at the request of the manufacturer [16]. Sipuleucel-T is an autologous antigen-presenting cell-based vaccination strategy, targeting prostatic acid phosphatase on prostate adenocarcinomas. The proposed mechanism of sipuleucel-T is induction of antigen-specific immune responses against PCa cells [17]. However, a complete understanding of the mechanism of action of sipuleucel-T is lacking. It remains unclear whether sipuleucel-T acts via priming of naive T cells through antigen presentation since the sipuleucel-T products contained more than 60% CD3<sup>+</sup> T cells and < 20% cells expressing the co-stimulatory molecule CD54, indicated as dendritic cells (DCs) [18]. It remains unclear whether sipuleucel-T harbors mature DC properties necessary for priming of naive T cells. Therefore, vaccination with antigen-specific blood-derived DCs may be a more potent alternative.

DCs are the most potent antigen-presenting cells of the immune system. They are crucial for inducing adaptive immune responses [19] and are widely studied in clinical trials, predominantly in advanced melanoma patients [20–24]. Antigen-loaded autologous DCs are given to patients with the intention of inducing functional tumor-associated antigen (TAA)-specific T cell responses. There are two major types of naturally occurring DCs that circulate in the blood [25], myeloid DCs (mDCs) and plasmacytoid DCs (pDCs). These subsets can be distinguished by the presence of different surface markers. mDCs can be further subdivided into two populations, based on their differential surface

expression of CD1c (BDCA-1; cDC2) and CD141 (BDCA-3; cDC1) [25]. mDCs act in particular against bacteria [26] and have the capacity to prime cytotoxic T cell responses [27]. pDCs produce high amounts of type I interferons, mainly in response to viral stimuli [28, 29].

mDCs and pDCs express different pattern recognition receptors, respond differently to stimuli and have different migration patterns [30]. This suggests that mDCs and pDCs have unique functional characteristics and may act synergistically by bi-directional crosstalk between the subsets and T cells [28, 30, 31]. Previously, we studied the safety, immunogenicity and clinical efficacy of pDC and CD1c<sup>+</sup> mDC vaccinations in stage IV melanoma patients [23, 24]. In these studies, promising tumor-specific T cell responses, cytokine production profiles and clinical responses were observed. This supports the use of both pDCs and cDC2 for evaluation in a phase IIa clinical trial in patients with CRPC .

## Materials and methods

### Patients

In this open-label, randomized, phase IIa study we screened 44 chemotherapy-naïve patients with CRPC. Patients with rising prostate-specific antigen (PSA) were closely monitored to detect early biochemical progression. Patients were screened for study eligibility as soon as patients met the criteria for CRPC [32]. Since there is no clear consensus on the correct timing for CRPC treatment, this window was used to screen asymptomatic or minimally symptomatic CRPC patients. Twenty-two of the screened patients were HLA-A\*0201. One of these patients was excluded because a second primary malignancy was detected (Additional file 1: Figure S1). All 21 included patients had histologically confirmed adenocarcinoma of the prostate. Eligible patients had not received any immunotherapy, docetaxel, cabazitaxel or treatment with the RANKL-inhibitor denosumab. Concurrent use of glucocorticoids up to 10 mg per day or a prednisone equivalent was permitted. Patients requiring opioids for cancer-related pain at screening were excluded. Patients had no visceral metastases. Other eligibility criteria were: an Eastern Cooperative Oncology Group (ECOG) performance status grade of 0 or 1; ongoing luteinizing hormone-releasing analogue therapy or status after bilateral orchidectomy; serum testosterone level of < 1.73/l (< 50 ng/dl); absence of active autoimmune diseases, absence of an active viral infection or allergy to shell fish; and no significant laboratory abnormalities (hemoglobin > 5.6 mmol/l (9.0 g/dl); white blood cell count of > 3.0 × 10<sup>9</sup>/l; platelets > 100 × 10<sup>9</sup>/l; serum creatinine < 150 µmol/l; AST/ALT < 3 x ULN, and serum bilirubin < 25 µmol/l, exception Gilbert's syndrome). Baseline disease sites were assessed using <sup>68</sup>Ga- prostate-specific membrane antigen (PSMA) PET/CT scans [33], including thin-section diagnostic CT (3 mm) and ferumoxtran-10-enhanced MRIs [34, 35] and regular MRI of bones and lymph nodes. Response evaluation was assessed according to Response Evaluation Criteria In Solid Tumours

(RECIST) version 1.1 [36] and the Prostate Cancer Clinical Trials Working Group 2 (PCWG2) criteria [37]. The immune-related response criteria and the iRECIST criteria were used to assess immune unconfirmed progressive disease [38–41]. Response evaluation was assessed by using contrast-enhanced <sup>68</sup>Ga-PSMA PET/CT scans and ferumoxtran-10-enhanced MRIs at 3 months, and for patients with long-term clinical benefit after 12 and 24 months. Regular follow-up MRI of lymph nodes and bones was performed at 6, 9, 15, 18 and 21 months. Measurable lesions were measured in at least one dimension with longest diameter ≥ 10 mm. Small lesions (longest diameter < 10 mm or pathological lymph nodes with < 15 mm short axis) are considered non-measurable disease. Bone metastases were documented and assessed according to PCWG2 criteria. Patients with absence of disease progression, defined as patients with a radiological complete or partial response or stable disease for > 6 months, were eligible for a maintenance cycle of three biweekly vaccinations. Patients without progressive disease after 12 months were eligible for a final vaccination cycle (Additional file 2: Figure S2A). Baseline characteristics and prior therapies are presented in Table 1.

- [VIEW INLINE](#)

- [VIEW POPUP](#)

## **Baseline characteristics of patients treated with blood-derived DC vaccinations**

### **Study design and objectives**

Patients with CRPC were randomly assigned in a 1:1:1 ratio to receive CD1c<sup>+</sup> mDC vaccinations ( $2\text{--}5 \times 10^6$  cells per injection; arm A), pDC vaccinations ( $1\text{--}3 \times 10^6$  cells; arm B), or combined CD1c<sup>+</sup> mDC and pDC vaccinations (combiDC;  $3\text{--}8 \times 10^6$  cells; arm C). One cycle of vaccinations consisted of three biweekly vaccinations administered intranodally in a clinically tumor-free lymph node by our expert radiologist or nuclear medicine physician. One to two weeks after the third vaccination a delayed-type hypersensitivity (DTH)-skin test was performed after intradermal administration of  $1\text{--}10 \times 10^5$  cells [42]. Adverse events were defined in accordance with the Common Terminology Criteria for Adverse Events (CTCAE) version 4.0. Primary endpoint of the study was the immunological response after DC vaccinations. Secondary objectives were safety, feasibility, quality of life and clinical efficacy (radiological progression-free survival (rPFS), OS, prostate-specific antigen doubling time (PSAdt), time to opiate use for cancer-related pain, time to SRE, time to decline in WHO/ECOG performance score by ≥1 point and time to the initiation of docetaxel chemotherapy). rPFS was defined as the time from apheresis to radiological progression of soft-tissue lesions or two or more new bone lesions or death from any cause. The event date of the

unconfirmed progression was used for calculation of rPFS. OS was defined as the time from apheresis to death from any cause. The PSA<sub>dt</sub> was calculated according to the Memorial Sloan-Kettering Cancer Center guidelines (<http://nomograms.mskcc.org/Prostate/PsaDoublingTime.aspx>). An SRE was defined as a pathologic fracture, palliative radiotherapy to a bone lesion, spinal-cord compression or surgery involving bone.

### Statistical analysis

Paired t-tests were performed to evaluate immunological responses before and after vaccination and independent-samples t-tests (Mann-Whitney U tests) were used to evaluate differences between groups. Statistical significance was defined as  $p < 0.05$  (two-tailed significance level). Time-to-event data were evaluated using the Kaplan-Meier method. Statistical significance was evaluated using the two-sided log-rank test and was defined as  $p < 0.05$ . Differences between treatment arms were evaluated using one-way ANOVA. Statistical analysis was performed using SPSS® Statistics version 22 software (SPSS Inc., Chicago, IL, USA) and GraphPad Prism 5.03 (GraphPad Software, Inc., San Diego, CA, USA).

### Vaccine preparation and features

CD1c<sup>+</sup> mDCs and pDCs were manufactured according to Good Manufacturing Practices (GMP). DCs were directly isolated from apheresis products using the fully automated and enclosed immunomagnetic CliniMACS Prodigy isolation system (Miltenyi Biotec, Bergisch-Gladbach, Germany). GMP-grade magnetic bead-coupled antibodies were used, following the manufacturer's guidelines. For mDC isolation, first, CD19<sup>+</sup> and CD14<sup>+</sup> cells were depleted, followed by positive selection of BDCA1<sup>+</sup> cells with biotin-coated CD1c (BDCA-1) antibodies and anti-biotin coated magnetic beads (arm A). pDC were selected with anti-CD304 (BDCA-4) coupled beads (arm B). When patients were randomized for vaccination with both mDCs and pDCs (arm C), first pDC were selected with anti-CD304 coupled beads, followed by depletion of CD19<sup>+</sup> and CD14<sup>+</sup> cells and positive selection of CD1c<sup>+</sup> cells. mDCs were cultured overnight at a concentration of  $1.5 \times 10^6$  cells/ml with 800 IU/ml recombinant human GM-CSF in TexMACS GMP medium (both Miltenyi Biotec) supplemented with 2% human serum (Sanquin) and 10 µg/ml keyhole limpet hemocyan (KLH; Immucothel, Biosyn Arzneimittel GmbH) for immunomonitoring purposes. pDCs were cultured overnight at a concentration of  $1.5 \times 10^6$  cells/ml with 10 ng/ml recombinant human IL-3 in TexMACS GMP medium (both Miltenyi Biotec) supplemented with 2% pooled human serum. mDCs and pDCs were loaded with HLA-A\*0201 binding peptides of NY-ESO-1:157–165 (SLLMWITQC) and MAGE-C2:336–344 (ALKDVEERV) [43] as well as NY-ESO-1 and MUC1 PepTivators (Miltenyi Biotec, Bergisch-Gladbach, Germany)) at a concentration of 1 µM. PepTivators consist of overlapping long peptides that cover the complete protein and bind multiple HLA-types, both MHC class I and II (Additional file 2: Figure S2B).

NY-ESO-1 and MUC1 PepTivators were added during overnight culturing. Thereafter, mDCs and pDCs were activated with premixed protamine HCl



(Meda Pharma) and mRNA (gp100, Universitätsklinik Erlangen) for 6 h. Premix ratio 10 µg protamine + 5 µg mRNA, 10 µl premix per ml cell suspension [44]. During the last 3 h of maturation NY-ESO-1 and MAGE-C2 peptides were added at final concentration of 1 µM. This isolation and culture procedure gave rise to mature mDC and pDC meeting the release criteria: sterile, endotoxin level < 7 EU/ml, more than 50% viability, more than 50% purity, expression of CD80 > 50% on pDCs and expression of CD83 > 50% on mDCs. Expression of MHC class I, MHC class II, CD86 and CCR7 was reported, but no release criterium (Additional file 3: Figure S3A-D). Protamine/mRNA complex activated both mDCs and pDCs into mature functional DCs that secrete IFN-α, TNF-α (pDCs only), IL-12p70, and IL-6 (both mDCs and pDCs, but mainly pDCs) (Additional file 3: Figure S3E). Cells were frozen in TexMACS medium containing 10% dimethyl sulfoxide (DMSO; WAK Chemie Medical GmbH) and 40% Albuman (Sanquin), stored at < - 80 °C for max. 2 years and thawed on the day of vaccination. For combined pDC and mDC vaccines, both subsets were pooled in one syringe after thawing.

After apheresis, sufficient amounts of blood-derived DCs could be obtained for at least one vaccination cycle. In two patients randomized for treatment with combiDCs the final CD1c<sup>+</sup> mDC product did not fulfill the release criteria. Therefore, these patients were vaccinated with pDCs only. Because the primary endpoint of the study was immunological, two extra patients were randomized within the combination arm. In patient pDC-06 the pDC purity was initially only 43%, which increased to 54% after overnight culture and maturation. In patient combiDC-06 CD1c<sup>+</sup> mDC purity was 41% and thus out-of-specification. Nevertheless, the product was released and administered after accounting for the lower purity by administering at least  $2.4 \times 10^6$  cells. Consequently, patient received at least the minimum required dose of  $2 \times 10^6$  CD1c<sup>+</sup> mDCs per vaccination.

### Flow cytometry

Purity and phenotype of mDCs and pDCs after CliniMACS isolation were determined by flow cytometry with a FACSVerse (BD Biosciences, San Jose, CA, USA) or MACS Quant (Miltenyi Biotec). The following primary monoclonal antibodies and the appropriate isotype or fluorescence minus one controls were used: anti-CD1c-Viobright FITC, anti-BDCA-2-PE, anti-CD20-PE-Vio770, anti-CD123-APC, anti-CD45-APC-Vio770, anti-CD14-VioGreen, anti-FcεRI-VioBlue, anti-CD14-FITC, anti-CD15-PE, anti-CD56-APC, anti-CD3-BioBlue, anti-HLA-ABC-APC, anti-HLA-DR,DP,DQ-APC, anti-CCR7-APC, anti-CD80-APC, anti-CD83-APC and anti-CD86-APC (all Miltenyi Biotec).

### Skin-test infiltrating lymphocyte culture and PBMC analyses

DTH challenges were performed 2 weeks after each vaccination cycle to assess TAA-specific immune response in DC vaccinated patients [42, 45]. DCs used for the DTH-skin test were produced accordingly to the vaccinated cells, except that no KLH was added to the culture medium. At four different sites at the patient's back maximally  $5.0 \times 10^5$  peptide-loaded blood-derived DCs were injected intradermally. After 48 h, 6 mm punch biopsies were taken. The

biopsies were manually cut and half of the tissue was stored at  $-150\text{ }^{\circ}\text{C}$ ; the other half was cultured as described previously [42]. After 2 to 4 weeks of culturing, skin-test infiltrating lymphocytes (SKILs) were tested for the presence of tumor antigen-specific T cells. SKILs and peripheral blood mononuclear cells (PBMCs) were stained with 1) anti-CD8-FITC and tetrameric PE- and APC-coupled MHC complexes containing NY-ESO-1 (SLLMWITQC), MAGE-C2 (ALKDVEERV) and MUC1 (LLLLTVLTV) HLA-A\*0201 epitopes (all Sanquin, Amsterdam, The Netherlands); or 2) anti-CD8-BV421, anti-CD19-FITC and dextrameric PE- and APC-coupled MHC complexes containing the indicated epitopes (all Immudex, Copenhagen, Denmark). Dextrameric HLA-B\*0801 (AAKGRGAAL) and tetrameric and dextrameric HIV (SLYNTVATL) were used as a negative control. Cells were analyzed by flow cytometry. To test peptide recognition, SKILs were challenged with autologous PBMCs loaded with the indicated peptides and PepTivators, **phorbol myristate acetate** (positive control), carcinoembryonic antigen peptide or no peptide (both negative control). Production of interferon- $\gamma$  (IFN- $\gamma$ ), IL-2, IL-5 and IL-10 was measured in the supernatants after overnight co-culture by cytometric bead array according to the manufacturer's instructions (BD Biosciences).

#### Proliferative and humoral response to KLH

Cellular responses against KLH were measured in a proliferation assay. PBMCs were isolated from blood samples after each vaccination.  $1 \times 10^5$  PBMCs were plated per well of a 96-well tissue culture microplate either in the presence or absence of KLH. After 4 days of culture,  $1\text{ }\mu\text{Ci}$ /well of tritiated thymidine was added, incorporation of tritiated thymidine was measured in a beta-counter. A proliferation index (proliferation with KLH/proliferation without KLH) of  $> 2$  was considered positive. Antibodies against KLH were measured in the serum of DC vaccinated patients by ELISA. KLH antibodies were detected with mouse anti-human IgG, IgA, or IgM antibodies labeled with horseradish peroxidase. 3,3',5,5'-Tetramethylbenzidine was used as a substrate. Plates were measured with a microtiter plate reader at 450 nm. An isotype-specific calibration curve for the KLH response was included in each microtiter plate.

#### Immunohistochemistry

Formalin-fixed, paraffin-embedded tissue blocks of prostate biopsies or radical prostatectomy at time of diagnosis were collected from primary treatment centers located in the Netherlands and sections of  $4\text{-}\mu\text{m}$  thickness were cut. Slides were deparaffinized using xylene and rehydrated with ethanol. Antigen retrieval was performed by boiling in EnVision™ FLEX target retrieval solution (pH 9, K8004, Dako) for 10 min for MUC1 staining or in Citrate buffer (pH 6, CBB999, ScyTek Laboratories) for 15 min for NY-ESO-1 and MAGE-C2 staining. After cooling down, endogenous peroxidase was blocked using 3% hydrogen peroxidase (76,051,800.1000, EMD Millipore) in PBS (4391.9010, Klinipath). Primary antibodies MUC1 (M0613, clone E26, Dako, dilution: 1/250), NY-ESO-1 (MABC1151, clone D8.38, Merck, dilution: 1/200) and

MAGE-C2 (HPA062230, rabbit polyclonal, Merck, dilution: 1/200) were diluted in Normal Antibody diluent (VWRKBD09–999, Immunologic) and were incubated at room temperature for 1 h. Slides were washed between steps with EnVision™ FLEX Wash Buffer (DM831, Dako). Next, incubation with BrightVision poly-HRP-anti-Ms/Rb/Rt IgG (DPVO999HRP, ImmunoLogic) was performed at room temperature for 30 min. Visualization was performed with EnVision™ FLEX DAB Buffered Substrate and EnVision™ FLEX Substrate Buffer (K5207 and SM803; DAKO) for 7 min at room temperature. After dehydration, slides were counterstained with hematoxylin and were enclosed with Quick-D mounting medium (7281, Klinipath). Observed staining was cytoplasmic. Immunoreactivity was evaluated by a pathologist using a semi-quantitative, stepwise scoring system: negative (0% of cells stained), weak (1–10% of cells stained), moderate (11–50% of cells stained) and strong (51 to 100% of cells stained). Representative slides were scanned using the PerkinElmer Vectra (Vectra 3.0.4, PerkinElmer). Testicular or tonsil tissue (positive control) was used for antibody validation (Additional file [4](#): Figure S4).

## Results

### Patient characteristics

In this prospective study twenty-one eligible patients with CRPC were enrolled. Participants were treated with blood-derived DC vaccines from November 2015 until May 2018. Baseline demographic, disease characteristics and prior therapies for hormone-sensitive PCa and CRPC are listed in Table [1](#). The described results are based on the cut-off date of 6th of March 2019. The median follow-up is 27.2 months (range 10.7–41.2\* months). All twenty-one patients, seven per arm, received at least one cycle of three biweekly DC vaccinations and a DTH-skin test. Thirteen patients also received a second cycle and seven patients a third vaccination cycle.

### **Safety and adverse events**

DC vaccinations were well tolerated. In all vaccinated patients only low-grade toxicity (CTCAE grade 1–2) was noticed. Most frequent grade 1–2 toxicity included flu-like symptoms, fatigue, upper respiratory infections, dizziness, vaccination-induced hematomas and injection site reactions. Also, some low-grade laboratory adverse events were seen (Table [2](#)).

- [VIEW INLINE](#)

- [VIEW POPUP](#)

### Adverse events

#### **Cellular and humoral responses to KLH**

mDCs (arm A and C) were loaded with KLH as a control antigen. Since pDCs cannot take up KLH-protein, pDCs were not cultured in the presence of KLH [46]. None of the patients had a KLH-specific proliferation index > 2 at baseline. KLH-specific proliferation increased significantly after one vaccination cycle. In 5 of 7 mDC-treated patients ( $p = 0.01$ ) and 3 of 7 patients in the combiDC group ( $p = 0.04$ ), a T cell response against KLH was observed (Additional file 5: Figure S5A). This indicates that KLH-exposed DCs were indeed able to induce de novo T cell responses to KLH. Humoral responses to KLH were determined in serum before treatment and after each cycle of vaccinations. A significant increase in total IgG titer was seen in mDC vaccinated patients (arm A and C) (Additional file 5: Figure S5B). There was no significant induction of IgA and IgM.

#### ***Tumor antigen-specific responses in DTH skin-test and blood***

DTH skin-tests were performed after each cycle of DC vaccinations to study NY-ESO-1-, MAGE-C2- and MUC1-specific T cell responses (Fig. 1a). NY-ESO-1-specific CD8<sup>+</sup> T cells were detected in skin biopsies in 15 patients (71%). MAGE-C2- and MUC1-specific CD8<sup>+</sup> T cells were found in 12 patients (57%) and 5 patients (24%), respectively. There were no significant differences in TAA-specific responses between patients vaccinated with mDCs, pDCs or combiDCs (Fig. 1b). In 15 of 21 patients (71%), tetramer- or dextramer-positive skin-derived T cells were observed for at least one TAA (Fig. 1c). In 7 of 20 patients (35%), these antigen-specific T cells were already detected after the first cycle of vaccinations. In 5 patients antigen-specific T cells were found against all 3 TAAs.

- [Download figure](#)
- [Open in new tab](#)
- [Download powerpoint](#)

## Fig. 1

Immunological responses in the DTH skin-test and in blood. a Example of flow cytometric analysis of SKILs of patients combiDC-04. SKILs were stained with dextramers encompassing HLA-A0201-specific peptides of NY-ESO-1, MAGE-C2 and Mucin-1 (MUC1) or with a negative control (HLA-B\*0801) and with anti-CD8. Tumor antigen-specific T cells were detected against all 3 tumor-associated antigens. b Tumor-associated antigen-specific responses in DTH skin-tests. NY-ESO-1-, MAGE-C2 and MUC1-specific T cell responses are presented per study arm and in total. c Number of antigen-specific responses in DTH skin-tests and in blood. Results are presented per vaccination cycle and in total. d Radiological non-progressive patients ( $n = 13$ ) are defined as patients with the absence of disease progression within 6 months. Radiological progressive patients ( $n = 8$ ) are defined as patients with progressive disease within 6 months. Presented are percentages of non-progressive and progressive patients with a positive DTH skin-test (tetramer/dextramer positive, dm<sup>+</sup>) for at least one epitope, IFN- $\gamma$  producing SKILs (IFN- $\gamma$ <sup>+</sup>), presence of both dm<sup>+</sup> and IFN- $\gamma$ <sup>+</sup> SKILs, and dominant IL-5<sup>+</sup>- or IL-10<sup>+</sup>-skewed immune responses, demonstrated by higher IL-5 or IL-10 production compared to IFN- $\gamma$  production in supernatant of antigen-challenged SKILs. e The presence of dm<sup>+</sup> antigen-specific T cells and IFN- $\gamma$ -producing (IFN- $\gamma$ <sup>+</sup>) SKILs are shown for patients with non-progressive disease ( $n = 13$ ) and progressive patients ( $n = 8$ ). +: 1 epitope; ++: 2 epitopes; +++: 3 epitopes recognized. DTH: delayed-type hypersensitivity; dm: dextramer; PBMCs: peripheral blood mononuclear cells; PE: phyco erytrin; SKILs: skin-test infiltrating lymphocytes

In peripheral blood prior to the start of DC vaccinations, in 7 of 21 patients (33%) NY-ESO-1-specific CD8<sup>+</sup> T cells were detected. No MAGE-C2- or MUC1-specific T cells were found before DC vaccinations. Post-vaccination antigen-specific T cells could be detected in peripheral blood in 12 of 21 patients (57%). NY-ESO-1-, MAGE-C2- and MUC-1 specific T cells were detected in blood of 10 of 21 (48%), 4 of 21 (19%) and 2 of 21 (10%) patients, respectively. In 4 patients antigen-specific T cells in blood were found against more than one TAA (Fig. [1c](#)).

SKILs were tested for their capacity to produce T helper 1 (Th1) cell cytokines (IFN- $\gamma$  and IL-2) or T helper 2 (Th2) cell cytokines (IL-5 and IL-10) upon co-culture with the tumor antigen peptides. Th1-type cytokines are proinflammatory, whereas Th2-type cytokines have a suppressive action and dampen the immune responses. IFN- $\gamma$  production (IFN- $\gamma^+$ ) was detected in 8 of 21 patients (31%). In radiological non-progressive patients, both induced tumor antigen-specific T cells (tetramer/dextramer<sup>+</sup> (dm<sup>+</sup>)) and functionality (IFN- $\gamma^+$ ) was observed in 5 of 13 patients (38%) compared to 0 of 8 in radiological progressive patients (0%) (Fig. 1d). Recognition of multiple epitopes by induced antigen-specific T cells and IFN- $\gamma^+$  was seen more frequently in patients with non-progressive disease (Fig. 1e). In 5 of 8 patients (63%) with radiological progression, we found a dominant IL-5- or IL-10-skewed immune response, compared to 3 of 13 patients (23%) with non-progressive disease (Fig. 1d).

### **Clinical outcome**

Of the 21 included patients, in 1 patient (5%) a partial radiological response was observed. Stable disease that persisted > 6 months was seen in 12 patients (57%). In 8 patients (38%) disease progression was observed within 6 months. Median rPFS for all patients was 9.5 months (range: 3.2–24.8\* months). The 6- and 12-months rPFS was 62% and 29%, respectively (Fig. 2a). There was no significant difference between the three treatment arms; in the mDC group the rPFS was 12.0 months (range 3.4–24.8\* months), in the pDC group 10.7 months (range 3.4–23.9\* months) and 4.2 months (range 3.2–12.0 months) in the combiDC group. The presence of functional antigen-specific T cells correlated with longer rPFS. In dm<sup>+</sup> and IFN- $\gamma^+$  patients ( $n = 5$ ) median rPFS was found to be 18.8 months compared to 5.1 months in dm<sup>-</sup> patients or patients without IFN- $\gamma$ -producing antigen-specific T cells ( $n = 16$ ;  $p = 0.02$ , Fig. 2b). Dm<sup>+</sup> and IFN- $\gamma^+$  patients showed longer PSA<sub>dt</sub> at 6 months compared to dm<sup>-</sup> patients or patients without IFN- $\gamma$ -producing antigen-specific T cells (mean PSA<sub>dt</sub> 12.9 months vs. 8.6 months, Fig. 2c). A decrease in PSA level was detected only in 2 of 21 patients. One of these patients (combiDC-07) showed a > 99% PSA-decrease which co-occurred with a partial radiological response (Fig. 3). Median OS was not reached. The median follow-up of all patients is 27.2 months (range 10.7–41.2\* months). Reversed Kaplan Meier estimate of the median follow-up was not reached, to take censored cases into account. To date 8 patients deceased during the study period, 7 PCa-related deaths occurred and there was one non-PCa-related death due to a ruptured type A aortic dissection (Table 3). OS appeared longer in dm<sup>+</sup> and IFN- $\gamma^+$  patients ( $n = 5$ ) versus dm<sup>-</sup> patients/patients without IFN- $\gamma$ -producing antigen-specific T cells ( $n = 16$ ) (Additional file 7: Figure S7). 5 patients (3 mDC, 1 pDC and 1 combiDC treated) had a skeletal-related event (SRE). The median time to SRE was not reached (range 3.6–21.8 months after apheresis). These 5 patients had bone pain secondary to bone metastases and were treated with palliative radiotherapy. Four of them were post radiotherapy treated with docetaxel-based chemotherapy. In 7 patients, docetaxel was initiated (range 3.7–

29.2 months after apheresis) (Additional file [6](#): Figure S6). Median time to opiate use for cancer-related pain and median time to ECOG performance score deterioration was also not reached. Seven patients started with opioids (range 1.4–20.1 months after apheresis). Ten patients had a decline in ECOG performance score (3 mDC, 3 pDC and 4 combiDC treated; range 1.5–20.1 months after apheresis). Details of clinical, immunological, immunohistochemical and sequencing outcomes are presented in Table [3](#), Additional file [6](#): Figure S6, Additional file [7](#): Figure S7 and Additional file [8](#): Table S1.

- [Download figure](#)
- [Open in new tab](#)
- [Download powerpoint](#)

## Fig. 2

**Radiological progression-free survival and biochemical responses. a** Kaplan-Meier analysis of rPFS of all patient determined by a log-rank test. **b** Kaplan-Meier analysis of rPFS of patients with ( $dm^+$  and  $IFN-\gamma^+$ ) or without ( $dm^-$  or  $IFN-\gamma^-$ ) the presence of functional antigen-specific T cells in skin biopsies was determined by a log-rank test. **c** PSA doubling tome during DC vaccination therapy in  $dm^+$  and  $IFN-\gamma^+$  patients ( $n = 5$ ) and  $dm^-$  or  $IFN-\gamma^-$  patients ( $n = 16$ )

- [Download figure](#)
- [Open in new tab](#)
- [Download powerpoint](#)

### **Fig. 3**

**Biochemical and radiological response upon first DC vaccination cycle of patient combiDC-07. a Biochemical analysis shows a PSA normalization upon the first cycle of DC vaccinations. b Fused  $^{68}\text{Ga}$ -prostate-specific membrane antigen PET/CT images showed a significant reduction of bilateral para-iliac and para-aortic lymph node metastases, right inguinal node metastases and a left supraclavicular lymph node metastasis after the 1st cycle of DC vaccinations. Lymph nodes are indicated with white arrows. c Maximal intensity projection images. Lymph nodes are indicated with red arrows**

- [VIEW INLINE](#)

- [VIEW POPUP](#)



## Clinical and immunological outcome

### ***Clinical outcome related to immunohistochemical results***

To study the effect of DC vaccination on TAA expression by the primary tumor, TAA expression was assessed on available prostate biopsies or radical prostatectomy tissue (Fig. 4a-d and Additional file 7: Table S1). Patients with TAA-specific T cells whose tumor expressed the same TAA (dm<sup>+</sup> and tumor<sup>+</sup>; *n* = 5) had a median rPFS of 10.7 months (range: 9.5–24.8\*). Patients that did not have matching TAA-specific T cells and TAA-expression of the tumor (dm<sup>+/-</sup> and tumor<sup>-</sup>; *n* = 16), had a median rPFS of 5.2 months (range: 3.2–24.3\*) (Fig. 4b). This difference was not statistically significant. In two patients who progressed after DC vaccination, loss of MUC1 expression by the tumor was observed. In one of these patients, MUC-1-specific T cells were detected. Tumor PD-L1 expression was studied in 10 patients. In two of these patients tumor PD-L1 expression post-vaccination was ≥1%. One of them was a dm<sup>+</sup> and IFN-γ<sup>+</sup> patient showing tumor PD-L1 expression of 60%. Tumors of all biopsied patients were microsatellite stable (Additional file 7: Table S1).

- [Download figure](#)
- [Open in new tab](#)
- [Download powerpoint](#)

## **Fig. 4**

**Expression of NY-ESO-1, MAGE-C2 and MUC1 and its relation to antigen-specific T cells in skin biopsies. a-d Representative immunohistochemical images showing (a) haematoxylin and eosin stain (H&E stain) and the expression of (b) NY-ESO-1, (c) MAGE-C2 and (d) MUC1. e Kaplan-Meier curve of rPFS in patients with or without the presence of antigen-specific T cells (dm<sup>+</sup>) in skin biopsies and expression of the same tumor-associated antigen in the tumor (dm<sup>+</sup> and tumor<sup>+</sup>)**

## Discussion

Patients with CRPC were vaccinated with DCs isolated directly from blood with a fully closed semi-automated system. Patients received mature mDCs (cDC2) and/or pDCs in order to induce tumor antigen-specific immune responses. We showed that vaccination with blood-derived DCs is safe and leads to the induction of antigen-specific T cells in the majority of patients. The induction of both antigen specific and functional T cells correlates with beneficial clinical outcome. In these small cohorts, no significant differences between DC subsets were observed, although responses to mDCs might be most promising (Additional File 6: Figure S6). Clinical efficacy of single DC subset vaccination or the combination of mDCs and pDCs will be assessed further in follow-up phase II/III studies.

DTH-skin test-derived and IFN- $\gamma$ -producing antigen-specific T cells were detected more frequently in SKIL cultures of patients with non-progressive disease compared to those with progressive disease. Thus, the presence of functional antigen-specific T cells might be indicative for a clinical beneficial response to DC vaccination. This is in line with our previous study in stage IV melanoma patients vaccinated with CD1c<sup>+</sup> DCs (cDC2), in whom the presence of functional tumor antigen-specific T cells in SKIL cultures coincided with improved clinical outcome [24]. Despite the fact that our study is not designed for clinical outcome assessment, we found a difference in median rPFS between patients with functional antigen-specific T cells (18.8 months;  $n = 5$ ) and patients with no functional antigen-specific T cells (5.1 months;  $n = 16$ ). In addition, observed survival times of the patients might potentially indicate a difference in OS in favor of dm<sup>+</sup> and IFN- $\gamma$ <sup>+</sup> patients. However, this finding has to be interpreted with caution. Firstly, the limited size renders our trial underpowered concerning reliable statements on the OS. Secondly, the number and type of subsequent therapies likely influenced the OS of these patients, clouding the direct effects of DC vaccination hereon (Additional File 6: Figure S6).

Following DC vaccination the rPFS of patients with functional antigen-specific T cells appeared comparable to median rPFS reported for abiraterone- (16.5 months) and enzalutamide-treated (20.0 months) men with metastatic CRPC who were chemotherapy naive [47, 48]. It is important to note that there is a considerable risk for guarantee-time bias [49] when correlating immunological responses of multiple vaccination cycles to clinical responses since patients who completed more than one vaccination cycle had a higher chance of developing IFN- $\gamma$ -producing antigen-specific SKILs [50]. The clinical impact of DC vaccination and validation of an immunological response readout as a surrogate endpoint will have to be studied in a larger phase II or III clinical trial.

In contrast to conventional response assessment using contrast enhanced-CT scans and radioisotope bone scans, we used <sup>68</sup>Ga-PSMA PET/CT scans [33], including thin-section diagnostic CT (3 mm) and ferumoxtran-10-enhanced MRIs [34, 35] for disease evaluation according to RECIST version 1.1 [36] and PCWG2 criteria [37, 51]. To assess immune unconfirmed progressive disease

immune-related response criteria and the iRECIST criteria were used [38 41]. We introduced both contrast-enhanced <sup>68</sup>Ga-PSMA PET/CT scans and ferumoxtran-10-enhanced MRIs to be able to study disease distribution, disease biology and host reaction within the tumor microenvironment of both measurable as non-measurable lesions [52]. Therefore, we have decided to use the best imaging modalities since it is very likely that these imaging modalities will become standard of care in the next five years. The functional imaging data will be reported elsewhere (manuscript in preparation).

Vaccination with blood-derived DCs resulted in only low-grade toxicity, that was similar to our previous studies [23, 24]. There were four patients who experienced possible vaccine-related symptoms of a grade 2 upper respiratory tract infection. These patients clinically recovered after treatment with oral antibiotics. Therefore, in our opinion, these were not related in retrospect, but this adverse event has to be monitored during ongoing and future trials with DCs.

The relation between the presence of antigen-specific T cells and tumor-antigen expression on PCa tissue was not obvious. PCa biopsies were obtained years before patients developed CRPC. From literature is known that in localized PCa the expression of MAGE-C2 is significantly lower (3%), compared to the CRPC setting (23%) [53]. The same accounts for NY-ESO-1 expression, which is positive in 3% of patients with localized PCa and 15% of patients with CRPC [54]. Also, an association of MUC1 upregulation with the development of CRPC is previously reported [55]. Retrospectively, compulsory tumor biopsies taken in the CRPC setting, prior to start of DC vaccination, would have been most informative for assessing associations between antigen expression and induction of antigen-specific T cells. In follow-up trials, tumor antigen-expression in fresh biopsies will be included, which may serve as either a selection criterion, or as an exploratory endpoint.

Our study is the second trial worldwide investigating immune responses upon vaccination with blood-derived DCs in advanced PCa. Previously, Prue and colleagues performed a phase I trial with HLA-A\*0201 peptide-loaded CD1c<sup>+</sup> DCs in 12 prostate cancer patients [56]. This vaccine was also very well tolerated, showing only grade 1–2 adverse events. In contrast to our study, in none of the patients in the study of Prue et al. tumor antigen-specific immune responses were observed and only 25% of patients developed a DTH skin-test response to the control antigens after vaccination. This might be due to the difference in administration route. We vaccinated patients intranodally, Prue and colleagues vaccinated their patients intradermally and intravenously. Due to the low numbers of DCs available, intradermally and intravenously administered blood-derived DCs might not have reached the lymph nodes in sufficient numbers. Although only low numbers of DCs are necessary to induce an immune response [57], direct intranodal injection of these scarce DCs might be more effective for T cell priming.

The therapeutic landscape for patients with CRPC is changing drastically with the vast number of potential single agent therapies and combination therapies that have been approved and are under investigation for CRPC. Until now, the clinical outcome of immune checkpoint inhibitors is disappointing in advanced PCa [14, 15]. However, the PD-1 inhibitor pembrolizumab shows antitumor activity in patients with evidence of progression on enzalutamide [58] and in the docetaxel-refractory setting (Keynote-199 trial; [NCT02787005](#)). Several trials with immune checkpoint inhibitors as a single agent treatment or as combination therapy are currently ongoing in both unselected as in immunogenic subtypes, such as those harboring microsatellite instability, high tumor mutational load or biallelic inactivation of CDK12 or BRCA2 [59].

Cellular immunotherapy with sipuleucel-T showed OS benefit in the phase III IMPACT trial [9]. On the contrary, the GM-CSF secreting GVAX cell line did not improve clinical outcome, but had similar survival data when compared to docetaxel in a phase III clinical trial [60, 61]. Since the study was designed as a superiority trial no statement could be made that these treatments were equally effective. The phase III trial of pox-virus-based co-stimulatory molecule-assisted vaccine PROSTVAC +/- granulocyte-macrophage colony-stimulating factor was stopped early. It had no effect on OS [62]. The VIABLE trial (docetaxel +/- DC vaccination, [NCT02111577](#)) is currently undergoing phase III evaluation.

Our DC vaccination strategy aims at inducing cytotoxic T lymphocytes. However, immune cell recruitment to the tumor and efficient tumor cell killing by cytotoxic T cells is probably less effective in patients with CRPC compared to patients with localized cancer. Indeed, recent studies show that in advanced cancer patients, the immunosuppressive state of the tumor, caused by regulatory T cells, myeloid-derived suppressor cells, expression of PD-L1 and production of immunosuppressive cytokines hampers the immune response towards the tumor [63–66]. Induced T cell responses frequently fail to fully eliminate cancer, because of an exhausted or dysfunctional state of the T cells [67]. This can be caused by an imbalance between T cell invigoration and reinvigoration and tumor burden [64]. We showed promising preliminary clinical outcome for patients with functional antigen-specific T cells. However, most single agent immunotherapies will fail to completely eliminate cancer cells in the majority of advanced cancer patients. Future trials could focus on combination therapies, such as a cellular-based immunotherapy and immune checkpoint inhibition. Another strategy is treating early-stage PCa in order to precede cancer-induced immunosuppressive mechanisms.

This trial was not designed to study the effects of sequential follow-up therapies for patients with CRPC. Most frequent subsequent treatments were abiraterone acetate, enzalutamide and docetaxel (Additional file 6: Figure S6). Current data showed no hampering of the therapeutic effect of these agents. In addition, it is hypothesized that immunotherapy has a delayed effect on the tumor growth curve evoking durable and adaptable anti-cancer immune responses over an extended time period [68]. However, it remains a major

challenge how to position the different therapies in the current treatment strategy of patients with CRPC. There are still unanswered questions regarding the preferred therapy approach (sequence or combination therapy), the timing of therapies and the relative efficacy of every single treatment. However, harboring vaccination-induced functional antigen-specific T cells might be beneficial even after disease progression upon DC vaccination. This relative efficacy might have clinical benefit during subsequent therapies .

In conclusion, we demonstrated that vaccination with blood-derived mDCs (cDC2) and/or pDCs induced functional tumor antigen-specific immune responses in patients with CRPC. Patients harboring functional antigen-specific T cells showed a significantly increased median rPFS and might have an OS benefit compared to patients without these cells. This immune correlate might be indicative for a beneficial response to DC vaccination and opens up new opportunities for future immunotherapy trials with the intention of long-term cancer control.

## Acknowledgments

The authors thank the patients and their families for their participation in the study. The authors also acknowledge the technicians of the department of Tumor Immunology for their excellent work: Nicole M. Scharenborg, Annemiek J. de Boer, Mandy W.M.M. van de Rakt, Tjitske Duiveman-de Boer, Jeanette M. Pots, Tom G.M. van Oorschot, Kevin J.H. Bos, Valeska de Ruiter, Ruud P.J. Hendrikx, Carlijn J. Popelier and Kiek N. Verrijp.

## Supplementary information

[\[40425\\_2019\\_787\\_MOESM1\\_ESM.tif\]](#)

[\[40425\\_2019\\_787\\_MOESM2\\_ESM.zip\]](#)

[\[40425\\_2019\\_787\\_MOESM3\\_ESM.zip\]](#)

[\[40425\\_2019\\_787\\_MOESM4\\_ESM.tif\]](#)

[\[40425\\_2019\\_787\\_MOESM5\\_ESM.zip\]](#)

[\[40425\\_2019\\_787\\_MOESM6\\_ESM.tif\]](#)

[\[40425\\_2019\\_787\\_MOESM7\\_ESM.tif\]](#)

[\[40425\\_2019\\_787\\_MOESM8\\_ESM.docx\]](#)

combiDCs

combined CD1c<sup>+</sup> myeloid and plasmacytoid dendritic cells

CRPC

Castration-resistant prostate cancer

CTCAE

Common Terminology Criteria for Adverse Events

DCs  
Dendritic cells

dm<sup>+</sup>  
tetramer or dextramer-positive

DTH  
Delayed-type hypersensitivity

ECOG  
Eastern Cooperative Oncology Group

GMP  
Good Manufacturing Practices

IFN- $\gamma$ <sup>+</sup>  
IFN- $\gamma$ -producing

KLH  
Keyhole limpet hemocyan

mDCs  
CD1c<sup>+</sup> myeloid dendritic cells/cDC2

OS  
Overall survival

PBMCs  
Peripheral blood mononuclear cells

PCa  
Prostate cancer

PCWG2  
Prostate Cancer Clinical Trials Working Group 2

pDCs  
plasmacytoid dendritic cells

PSA  
Prostate-specific antigen

PSAdt  
Prostate-specific antigen doubling time

PSMA  
Prostate-specific membrane antigen

RECIST  
Response Evaluation Criteria In Solid Tumours

rPFS  
radiological progression-free survival

SKILs  
Skin-test infiltrating lymphocytes

TAA  
Tumor-associated antigen

Th1  
T helper 1; Th2: T helper 2

## References

1. 1. 

1. Siegel RL,
2. Miller KD,
3. Jemal A

. Cancer statistics, 2019. *CA Cancer J Clin.* 2019;**69**:17–34.

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

2. 2. 

1. Tannock IF,
2. de Wit R,
3. Berry WR,
4. Horti J,
5. Pluzanska A,
6. Chi KN, *et al*

. Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer. *N Engl J Med.* 2004;**351**:151502–1512. [doi:10.1056/NEJMoa040720](https://doi.org/10.1056/NEJMoa040720)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

3. 3.

1. Berthold DR,
2. Pond GR,
3. Soban F,
4. de Wit R,
5. Eisenberger M,
6. Tannock IF

. Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer: updated survival in the TAX 327 study. *J Clin Oncol.* 2008;**26**:2242–245. [doi:10.1200/JCO.2007.12.4008](https://doi.org/10.1200/JCO.2007.12.4008)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

4. 4. 

1. Petrylak DP,
2. Tangen CM,
3. Hussain MH,
4. Lara PN Jr.,
5. Jones JA,
6. Taplin ME, *et al*

. Docetaxel and estramustine compared with mitoxantrone and prednisone for advanced refractory prostate cancer. N Engl J Med. 2004;**351**:151513–1520. [doi:10.1056/NEJMoa041318](https://doi.org/10.1056/NEJMoa041318)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

5. 5. 

1. de Bono JS,
2. Logothetis CJ,
3. Molina A,
4. Fizazi K,
5. North S,
6. Chu L, *et al*

. Abiraterone and increased survival in metastatic prostate cancer. N Engl J Med. 2011;**364**:211995–2005. pmcid:3471149 [doi:10.1056/NEJMoa1014618](https://doi.org/10.1056/NEJMoa1014618)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

6. 6.

1. Scher HI,
2. Fizazi K,
3. Saad F,
4. Taplin ME,
5. Sternberg CN,
6. Miller K, *et al*

. Increased survival with enzalutamide in prostate cancer after chemotherapy. N Engl J Med. 2012;**367**:131187–1197. [doi:10.1056/NEJMoa1207506](https://doi.org/10.1056/NEJMoa1207506)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

7. 7.

1. Ryan CJ,
2. Smith MR,
3. Fizazi K,
4. Saad F,
5. Mulders PF,
6. Sternberg CN, *et al*

. Abiraterone acetate plus prednisone versus placebo plus prednisone in chemotherapy-naive men with metastatic castration-resistant prostate cancer (COU-AA-302): final overall survival analysis of a randomised, double-blind,



placebo-controlled phase 3 study. *Lancet Oncol.* 2015;**16**:2152–160. [doi:10.1016/S1470-2045\(14\)71205-7](https://doi.org/10.1016/S1470-2045(14)71205-7)

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

8. 8. 

1. Beer TM,
2. Armstrong AJ,
3. Rathkopf DE,
4. Lortot Y,
5. Sternberg CN,
6. Higano CS, *et al*

. Enzalutamide in metastatic prostate cancer before chemotherapy. *N Engl J Med.* 2014;**371**:5424–433. pmcid:4418931 [doi:10.1056/NEJMoa1405095](https://doi.org/10.1056/NEJMoa1405095)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

9. 9. 

1. Kantoff PW,
2. Higano CS,
3. Shore ND,
4. Berger ER,
5. Small EJ,
6. Penson DF, *et al*

. Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med.* 2010;**363**:5411–422. [doi:10.1056/NEJMoa1001294](https://doi.org/10.1056/NEJMoa1001294)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

10. 10. 

1. Parker C,
2. Nilsson S,
3. Heinrich D,
4. Helle SI,
5. O'Sullivan JM,
6. Fossa SD, *et al*

. Alpha emitter radium-223 and survival in metastatic prostate cancer. *N Engl J Med.* 2013;**369**:3213–223. [doi:10.1056/NEJMoa1213755](https://doi.org/10.1056/NEJMoa1213755)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

11. 11.↵  
1. de Bono JS,  
2. Oudard S,  
3. Ozguroglu M,  
4. Hansen S,  
5. Machiels JP,  
6. Kocak I, *et al*

. Prednisone plus cabazitaxel or mitoxantrone for metastatic castration-resistant prostate cancer progressing after docetaxel treatment: a randomised open-label trial. *Lancet*. 2010;**376**:97471147–1154. [doi:10.1016/S0140-6736\(10\)61389-X](https://doi.org/10.1016/S0140-6736(10)61389-X)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

12. 12.↵  
1. Eisenberger M,  
2. Hardy-Bessard AC,  
3. Kim CS,  
4. Geczi L,  
5. Ford D,  
6. Mourey L, *et al*

. Phase III study comparing a reduced dose of Cabazitaxel (20 mg/m<sup>2</sup>) and the currently approved dose (25 mg/m<sup>2</sup>) in Postdocetaxel patients with metastatic castration-resistant prostate Cancer-PROSELICA. *J Clin Oncol*. 2017;**35**:283198–3206. [doi:10.1200/JCO.2016.72.1076](https://doi.org/10.1200/JCO.2016.72.1076)

[Full Text @ Radboud UniversityGoogle Scholar](#)

13. 13.↵  
1. Fizazi K,  
2. Scher HI,  
3. Molina A,  
4. Logothetis CJ,  
5. Chi KN,  
6. Jones RJ, *et al*

. Abiraterone acetate for treatment of metastatic castration-resistant prostate cancer: final overall survival analysis of the COU-AA-301 randomised, double-blind, placebo-controlled phase 3 study. *Lancet oncol*. 2012;**13**:10983–992. [doi:10.1016/S1470-2045\(12\)70379-0](https://doi.org/10.1016/S1470-2045(12)70379-0)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

14. 14.↵  
1. Beer TM,  
2. Kwon ED,  
3. Drake CG,  
4. Fizazi K,  
5. Logothetis C,  
6. Gravis G, *et al*

. Randomized, double-blind, phase III trial of Ipilimumab versus placebo in asymptomatic or minimally symptomatic patients with metastatic chemotherapy-naive castration-resistant prostate Cancer. *J Clin Oncol.* 2017;**35**:140–47. [doi:10.1200/JCO.2016.69.1584](https://doi.org/10.1200/JCO.2016.69.1584)

[Full Text @ Radboud UniversityCrossRefGoogle Scholar](#)

15. 15.↵  
1. Kwon ED,  
2. Drake CG,  
3. Scher HI,  
4. Fizazi K,  
5. Bossi A,  
6. van den Eertwegh AJ, *et al*

. Ipilimumab versus placebo after radiotherapy in patients with metastatic castration-resistant prostate cancer that had progressed after docetaxel chemotherapy (CA184-043): a multicentre, randomised, double-blind, phase 3 trial. *Lancet Oncol.* 2014;**15**:7700–712. pmcid:4418935 [doi:10.1016/S1470-2045\(14\)70189-5](https://doi.org/10.1016/S1470-2045(14)70189-5)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

16. 16.↵  
EMA Human Medicine European Public Assessment Report (EPAR) of Provenge (last updated 19/05/2015); EMA website: <https://www.ema.europa.eu/en/medicines/human/EPAR/provenge>.

[Google Scholar](#)

17. 17.↵  
1. Sheikh NA,  
2. Petrylak D,  
3. Kantoff PW,  
4. Dela Rosa C,  
5. Stewart FP,

6. Kuan LY, *et al*

. Sipuleucel-T immune parameters correlate with survival: an analysis of the randomized phase 3 clinical trials in men with castration-resistant prostate cancer. *Cancer Immunol Immunother.* 2013;**62**:1137–147. [doi:10.1007/s00262-012-1317-2](https://doi.org/10.1007/s00262-012-1317-2)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

18. 18. 

1. Burch PA,
2. Breen JK,
3. Buckner JC,
4. Gastineau DA,
5. Kaur JA,
6. Laus RL, *et al*

. Priming tissue-specific cellular immunity in a phase I trial of autologous dendritic cells for prostate cancer. *Clin Cancer Res.* 2000;**6**:62175–2182.

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

19. 19. 

1. Banchereau J,
2. Steinman RM

. Dendritic cells and the control of immunity. *Nature.* 1998;**392**:6673245–252. [doi:10.1038/32588](https://doi.org/10.1038/32588)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

20. 20. 

1. Figdor CG,
2. de Vries IJ,
3. Lesterhuis WJ,
4. Melief CJ

. Dendritic cell immunotherapy: mapping the way. *Nat Med.* 2004;**10**:5475–480. [doi:10.1038/nm1039](https://doi.org/10.1038/nm1039)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

21. 21.

1. Palucka K,

2. Banchereau J

. Dendritic-cell-based therapeutic cancer vaccines. *Immunity*.. 2013;**39**:138–48. pmcid:3788678 [doi:10.1016/j.immuni.2013.07.004](https://doi.org/10.1016/j.immuni.2013.07.004)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

22. 22.

1. Bol KF,
2. Figdor CG,
3. Aarntzen EH,
4. Welzen ME,
5. van Rossum MM,
6. Blokx WA, *et al*

. Intranodal vaccination with mRNA-optimized dendritic cells in metastatic melanoma patients. *Oncoimmunology*.. 2015;**4**:84570143[doi:10.1080/2162402X.2015.1019197](https://doi.org/10.1080/2162402X.2015.1019197)

[Google Scholar](#)

23. 23. ↵

1. Tel J,
2. Aarntzen EH,
3. Baba T,
4. Schreibelt G,
5. Schulte BM,
6. Benitez-Ribas D, *et al*

. Natural human plasmacytoid dendritic cells induce antigen-specific T-cell responses in melanoma patients. *Cancer Res*. 2013;**73**:31063–1075. [doi:10.1158/0008-5472.CAN-12-2583](https://doi.org/10.1158/0008-5472.CAN-12-2583)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

24. 24. ↵

1. Schreibelt G,
2. Bol KF,
3. Westdorp H,
4. Wimmers F,
5. Aarntzen EH,
6. Duiveman-de Boer T, *et al*


. Effective clinical responses in metastatic melanoma patients after vaccination with primary myeloid dendritic cells. *Clin Cancer Res.* 2016;**22**:92155–2166. [doi:10.1158/1078-0432.CCR-15-2205](https://doi.org/10.1158/1078-0432.CCR-15-2205)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

25. 25. 
1. MacDonald KP,
  2. Munster DJ,
  3. Clark GJ,
  4. Dzionek A,
  5. Schmitz J,
  6. Hart DN


. Characterization of human blood dendritic cell subsets. *Blood.* 2002;**100**:134512–4520. [doi:10.1182/blood-2001-11-0097](https://doi.org/10.1182/blood-2001-11-0097)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

26. 26. 
1. Piccioli D,
  2. Sammicheli C,
  3. Tavarini S,
  4. Nuti S,
  5. Frigimelica E,
  6. Manetti AG, *et al*

. Human plasmacytoid dendritic cells are unresponsive to bacterial stimulation and require a novel type of cooperation with myeloid dendritic cells for maturation. *Blood.* 2009;**113**:184232–4239. [doi:10.1182/blood-2008-10-186890](https://doi.org/10.1182/blood-2008-10-186890)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

27. 27. 
1. Nizzoli G,
  2. Krietsch J,
  3. Weick A,
  4. Steinfeld S,
  5. Facciotti F,
  6. Guarini P, *et al*

. Human CD1c+ dendritic cells secrete high levels of IL-12 and potently prime cytotoxic T-cell responses. *Blood.* 2013;**122**:6932–942. [doi:10.1182/blood-2013-04-495424](https://doi.org/10.1182/blood-2013-04-495424)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

28. 28.↵  
1. Bakdash G,  
2. Schreurs I,  
3. Schreibelt G,  
4. Tel J

. Crosstalk between dendritic cell subsets and implications for dendritic cell-based anticancer immunotherapy. *Expert Rev Clin Immunol*. 2014;**10**:7915–926. [doi:10.1586/1744666X.2014.912561](https://doi.org/10.1586/1744666X.2014.912561)

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

29. 29.↵  
1. Dzionek A,  
2. Fuchs A,  
3. Schmidt P,  
4. Cremer S,  
5. Zysk M,  
6. Miltenyi S, *et al*

. BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood. *J Immunol*. 2000;**165**:116037–6046. [doi:10.4049/jimmunol.165.11.6037](https://doi.org/10.4049/jimmunol.165.11.6037)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

30. 30.↵  
1. Bol KF,  
2. Schreibelt G,  
3. Rabold K,  
4. Wculek SK,  
5. Schwarze JK,  
6. Dzionek A, *et al*

. The clinical application of cancer immunotherapy based on naturally circulating dendritic cells. *J Immunother Cancer*. 2019;**7**:1109. pmcid:6471787 [doi:10.1186/s40425-019-0580-6](https://doi.org/10.1186/s40425-019-0580-6)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

31. 31.↵  
1. van Beek JJ,  
2. Gorris MA,  
3. Skold AE,  
4. Hatipoglu I,  
5. Van Acker HH,

6. Smits EL, *et al*

. Human blood myeloid and plasmacytoid dendritic cells cross activate each other and synergize in inducing NK cell cytotoxicity. *Oncoimmunology*. 2016;**5**:105087293 [doi:10.1080/2162402X.2016.1227902](https://doi.org/10.1080/2162402X.2016.1227902)

[Google Scholar](#)

32. 32. 

1. Heidenreich A,
2. Bastian PJ,
3. Bellmunt J,
4. Bolla M,
5. Joniau S,
6. van der Kwast T, *et al*

. EAU guidelines on prostate cancer. Part II: treatment of advanced, relapsing, and castration-resistant prostate cancer. *Eur Urol*. 2014;**65**:2467–479. [doi:10.1016/j.eururo.2013.11.002](https://doi.org/10.1016/j.eururo.2013.11.002) pmcid:24321502

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

33. 33. 

1. Eiber M,
2. Maurer T,
3. Souvatzoglou M,
4. Beer AJ,
5. Ruffani A,
6. Haller B, *et al*

. Evaluation of hybrid (6)(8)Ga-PSMA ligand PET/CT in 248 patients with biochemical recurrence after radical prostatectomy. *J Nucl Med*. 2015;**56**:5668–674. [doi:10.2967/jnumed.115.154153](https://doi.org/10.2967/jnumed.115.154153) pmcid:25791990

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

34. 34. 

1. Heesakkers RA,
2. Hovels AM,
3. Jager GJ,
4. van den Bosch HC,
5. Witjes JA,
6. Raat HP, *et al*

. MRI with a lymph-node-specific contrast agent as an alternative to CT scan and lymph-node dissection in patients with prostate cancer: a prospective




multicohort study. *Lancet Oncol.* 2008;**9**:9850–856. [doi:10.1016/S1470-2045\(08\)70203-1](https://doi.org/10.1016/S1470-2045(08)70203-1) pmcid:18708295

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

35. 35. 
1. Fortuin Ansje S.,
  2. Brüggemann Roger,
  3. van der Linden Janine,
  4. Panfilov Ilia,
  5. Israël Bas,
  6. Scheenen Tom W.J.,
  7. Barentsz Jelle O.


. Ultra-small superparamagnetic iron oxides for metastatic lymph node detection: back on the block. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology.* 2017;**10**:1e1471

[Full Text @ Radboud UniversityGoogle Scholar](#)

36. 36. 
1. Eisenhauer EA,
  2. Therasse P,
  3. Bogaerts J,
  4. Schwartz LH,
  5. Sargent D,
  6. Ford R, *et al*

. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer.* 2009;**45**:2228–247. pmcid:19097774 [doi:10.1016/j.ejca.2008.10.026](https://doi.org/10.1016/j.ejca.2008.10.026)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

37. 37. 
1. Scher HI,
  2. Halabi S,
  3. Tannock I,
  4. Morris M,
  5. Sternberg CN,
  6. Carducci MA, *et al*

. Design and end points of clinical trials for patients with progressive prostate cancer and castrate levels of testosterone: recommendations of the prostate

Cancer clinical trials working group. J Clin Oncol. 2008;**26**:71148–1159. pmcid:18309951 [doi:10.1200/JCO.2007.12.4487](https://doi.org/10.1200/JCO.2007.12.4487)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

38. 38.↵
1. Hoos A,
  2. Parmiani G,
  3. Hege K,
  4. Sznol M,
  5. Loibner H,
  6. Eggermont A, *et al*

. A clinical development paradigm for cancer vaccines and related biologics. J Immunother. 2007;**30**:11–15. [doi:10.1097/01.cji.0000211341.88835.ae](https://doi.org/10.1097/01.cji.0000211341.88835.ae)

[Full Text @ Radboud UniversityCrossRefGoogle Scholar](#)

39. 39.
1. Hoos A,
  2. Eggermont AM,
  3. Janetzki S,
  4. Hodi FS,
  5. Ibrahim R,
  6. Anderson A, *et al*

. Improved endpoints for cancer immunotherapy trials. J Natl Cancer Inst. 2010;**102**:181388–1397. pmcid:20826737 [doi:10.1093/jnci/djq310](https://doi.org/10.1093/jnci/djq310)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

40. 40.
1. Hoos A

. Evolution of end points for cancer immunotherapy trials. Ann Oncol. 2012;**23**:Suppl 8viii47–viii52. [doi:10.1093/annonc/mds263](https://doi.org/10.1093/annonc/mds263)

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

41. 41.↵
1. Seymour L,
  2. Bogaerts J,
  3. Perrone A,
  4. Ford R,
  5. Schwartz LH,
  6. Mandrekar S, *et al*

. iRECIST: guidelines for response criteria for use in trials testing immunotherapeutics. *Lancet Oncol*. 2017;**18**:3e143–ee52. pmcid:5648544 [doi:10.1016/S1470-2045\(17\)30074-8](https://doi.org/10.1016/S1470-2045(17)30074-8)

[Full Text @ Radboud UniversityGoogle Scholar](#)

42. 42. 
1. de Vries IJ,
  2. Bernsen MR,
  3. Lesterhuis WJ,
  4. Scharenborg NM,
  5. Strijk SP,
  6. Gerritsen MJ, *et al*

. Immunomonitoring tumor-specific T cells in delayed-type hypersensitivity skin biopsies after dendritic cell vaccination correlates with clinical outcome. *J Clin Oncol*. 2005;**23**:245779–5787. [doi:10.1200/JCO.2005.06.478](https://doi.org/10.1200/JCO.2005.06.478)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

43. 43. 
1. Westdorp H,
  2. Skold AE,
  3. Snijer BA,
  4. Franik S,
  5. Mulder SF,
  6. Major PP, *et al*

. Immunotherapy for prostate cancer: lessons from responses to tumor-associated antigens. *Front Immunol*. 2014;**5**:191. pmcid:4018526 [doi:10.3389/fimmu.2014.00191](https://doi.org/10.3389/fimmu.2014.00191)

[Full Text @ Radboud UniversityPubMedGoogle Scholar](#)

44. 44. 
1. Skold AE,
  2. van Beek JJ,
  3. Sittig SP,
  4. Bakdash G,
  5. Tel J,
  6. Schreibelt G, *et al*

. Protamine-stabilized RNA as an ex vivo stimulant of primary human dendritic cell subsets. *Cancer Immunol Immunother*. 2015;**64**:111461–1473. pmcid:4612318 [doi:10.1007/s00262-015-1746-9](https://doi.org/10.1007/s00262-015-1746-9)

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

45. 45.↵  
1. Aarntzen EH,  
2. Bol K,  
3. Schreibelt G,  
4. Jacobs JF,  
5. Lesterhuis WJ,  
6. Van Rossum MM, *et al*

. Skin-test infiltrating lymphocytes early predict clinical outcome of dendritic cell-based vaccination in metastatic melanoma. *Cancer Res.* 2012;**72**:236102–6110. [doi:10.1158/0008-5472.CAN-12-2479](https://doi.org/10.1158/0008-5472.CAN-12-2479)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

46. 46.↵  
1. Benitez-Ribas D,  
2. Adema GJ,  
3. Winkels G,  
4. Klasen IS,  
5. Punt CJ,  
6. Figdor CG, *et al*

. Plasmacytoid dendritic cells of melanoma patients present exogenous proteins to CD4+ T cells after fc gamma RII-mediated uptake. *J Exp Med.* 2006;**203**:71629–1635. pmcid:2118356 [doi:10.1084/jem.20052364](https://doi.org/10.1084/jem.20052364)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

47. 47.↵  
1. Ryan CJ,  
2. Smith MR,  
3. de Bono JS,  
4. Molina A,  
5. Logothetis CJ,  
6. de Souza P, *et al*

. Abiraterone in metastatic prostate cancer without previous chemotherapy. *N Engl J Med.* 2013;**368**:2138–148. [doi:10.1056/NEJMoa1209096](https://doi.org/10.1056/NEJMoa1209096)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

48. 48.↵  
1. Beer TM,  
2. Armstrong AJ,  
3. Rathkopf D,

4. Loriot Y,
5. Sternberg CN,
6. Higano CS, *et al*

. Enzalutamide in men with chemotherapy-naive metastatic castration-resistant prostate Cancer: extended analysis of the phase 3 PREVAIL study. *Eur Urol.* 2017;**71**:2151–154. [doi:10.1016/j.eururo.2016.07.032](https://doi.org/10.1016/j.eururo.2016.07.032)

[Full Text @ Radboud UniversityGoogle Scholar](#)

49. 49. ↵
1. Giobbie-Hurder A,
  2. Gelber RD,
  3. Regan MM

. Challenges of guarantee-time bias. *J Clin Oncol.* 2013;**31**:232963–2969. pmcid:3732313 [doi:10.1200/JCO.2013.49.5283](https://doi.org/10.1200/JCO.2013.49.5283)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

50. 50. ↵
1. Boudewijns S,
  2. Westdorp H,
  3. Koornstra RH,
  4. Aarntzen EH,
  5. Schreibelt G,
  6. Creemers JH, *et al*

. Immune-related adverse events of dendritic cell vaccination correlate with Immunologic and clinical outcome in stage III and IV melanoma patients. *J Immunother.* 2016;**39**:6241–248. pmcid:4902323 [doi:10.1097/CJI.0000000000000127](https://doi.org/10.1097/CJI.0000000000000127)

[Full Text @ Radboud UniversityGoogle Scholar](#)

51. 51. ↵
1. Scher HI,
  2. Morris MJ,
  3. Stadler WM,
  4. Higano C,
  5. Basch E,
  6. Fizazi K, *et al*


. Trial design and objectives for castration-resistant prostate Cancer: updated recommendations from the prostate Cancer clinical trials working group 3. *J Clin Oncol.* 2016;**34**:121402–1418. pmcid:26903579 [doi:10.1200/JCO.2015.64.2702](https://doi.org/10.1200/JCO.2015.64.2702)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

52. 52. 
1. Scarpelli M,
  2. Zahm C,
  3. Perlman S,
  4. McNeel DG,
  5. Jeraj R,
  6. Liu G


. FLT PET/CT imaging of metastatic prostate cancer patients treated with pTVG-HP DNA vaccine and pembrolizumab. *J Immunother Cancer*.. 2019;**7**:123. pmcid:6354338 [doi:10.1186/s40425-019-0516-1](https://doi.org/10.1186/s40425-019-0516-1)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

53. 53. 
1. von Boehmer L,
  2. Keller L,
  3. Mortezaei A,
  4. Provenzano M,
  5. Sais G,
  6. Hermanns T, *et al*


. MAGE-C2/CT10 protein expression is an independent predictor of recurrence in prostate cancer. *PLoS One*. 2011;**6**:7 [doi:10.1371/journal.pone.0021366](https://doi.org/10.1371/journal.pone.0021366)

[Google Scholar](#)

54. 54. 
1. Fossa A,
  2. Berner A,
  3. Fossa SD,
  4. Hernes E,
  5. Gaudernack G,
  6. Smeland EB

. NY-ESO-1 protein expression and humoral immune responses in prostate cancer. *Prostate*.. 2004;**59**:4440–447. [doi:10.1002/pros.20025](https://doi.org/10.1002/pros.20025)

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

55. 55. 
1. Wong N,
  2. Major P,

3. Kapoor A,
4. Wei F,
5. Yan J,
6. Aziz T, *et al*

. Amplification of MUC1 in prostate cancer metastasis and CRPC development. *Oncotarget*. 2016;**7**:5083115–83133. pmcid:5347757 [doi:10.18632/oncotarget.13073](https://doi.org/10.18632/oncotarget.13073)

[Full Text @ Radboud UniversityGoogle Scholar](#)

56. 56. ↵
1. Prue RL,
  2. Vari F,
  3. Radford KJ,
  4. Tong H,
  5. Hardy MY,
  6. D'Rozario R, *et al*

. A phase I clinical trial of CD1c (BDCA-1)+ dendritic cells pulsed with HLA-A\*0201 peptides for immunotherapy of metastatic hormone refractory prostate cancer. *J Immunother*. 2015;**38**:271–76. [doi:10.1097/CJI.0000000000000063](https://doi.org/10.1097/CJI.0000000000000063)

[Full Text @ Radboud UniversityCrossRefGoogle Scholar](#)

57. 57. ↵
1. Aarntzen EH,
  2. Srinivas M,
  3. Bonetto F,
  4. Cruz LJ,
  5. Verdijk P,
  6. Schreibelt G, *et al*


. Targeting of 111In-labeled dendritic cell human vaccines improved by reducing number of cells. *Clin Cancer Res*. 2013;**19**:61525–1533. [doi:10.1158/1078-0432.CCR-12-1879](https://doi.org/10.1158/1078-0432.CCR-12-1879)

[Full Text @ Radboud UniversityAbstract/FREE Full TextGoogle Scholar](#)

58. 58. ↵
1. Graff JN,
  2. Alumkal JJ,
  3. Drake CG,
  4. Thomas GV,
  5. Redmond WL,
  6. Farhad M, *et al*

. Early evidence of anti-PD-1 activity in enzalutamide-resistant prostate cancer. *Oncotarget*. 2016;**7**:3352810–52817. pmcid:5288150 [doi:10.18632/oncotarget.10547](https://doi.org/10.18632/oncotarget.10547)

[Full Text @ Radboud University](#)[Google Scholar](#)

59. 59.   
1. Mehra N,  
2. Gerritsen W

. Now the dust has settled over immune checkpoint blockade in metastatic prostate cancer. *Ann Oncol*. 2018;**29**:81620–1622. [doi:10.1093/annonc/mdy239](https://doi.org/10.1093/annonc/mdy239)

[Full Text @ Radboud University](#)[Google Scholar](#)

60. 60. 

Higano CSF, Somer B, Curti B, Petrylak DP, Drake CG, et al. A phase III trial of GVAX immunotherapy for prostate cancer vs docetaxel plus prednisone in asymptomatic, castration-resistant prostate cancer (CRPC). . Genitourinary Cancer Symposium: Proc Am Soc Clin Oncol, 2009 (abstract #LBA150). 2009.

[Google Scholar](#)

61. 61. 

Small EJDT, Gerritsen WR et al. A phase III trial of GVAX immunotherapy for prostate cancer in combination with docetaxel versus docetaxel plus prednisone in symptomatic, castration-resistant prostate cancer (CRPC). Genitourinary Cancer Symposium: Proc Am Soc Clin Oncol, 2009 (abstract 7). 2009.

[Google Scholar](#)

62. 62.   
1. Gulley JL,  
2. Borre M,  
3. Vogelzang NJ,  
4. Ng S,  
5. Agarwal N,  
6. Parker CC, *et al*

. Phase III trial of PROSTVAC in asymptomatic or minimally symptomatic metastatic castration-resistant prostate Cancer. *J Clin Oncol*. 2019;**37**:131051–1061. [doi:10.1200/JCO.18.02031](https://doi.org/10.1200/JCO.18.02031)



[Full Text @ Radboud UniversityGoogle Scholar](#)

63. 63. ↵
1. Herbst RS,
  2. Soria JC,
  3. Kowanetz M,
  4. Fine GD,
  5. Hamid O,
  6. Gordon MS, *et al*

. Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. *Nature*.. 2014;**515**:7528563–567. pmcid:4836193 [doi:10.1038/nature14011](https://doi.org/10.1038/nature14011)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

64. 64. ↵
1. Huang AC,
  2. Postow MA,
  3. Orlovski RJ,
  4. Mick R,
  5. Bengsch B,
  6. Manne S, *et al*

. T-cell invigoration to tumour burden ratio associated with anti-PD-1 response. *Nature*.. 2017;**545**:765260–65. pmcid:5554367 [doi:10.1038/nature22079](https://doi.org/10.1038/nature22079)

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

65. 65.
1. Tumei PC,
  2. Harview CL,
  3. Yearley JH,
  4. Shintaku IP,
  5. Taylor EJ,
  6. Robert L, *et al*

. PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature*.. 2014;**515**:7528568–571. pmcid:4246418 [doi:10.1038/nature13954](https://doi.org/10.1038/nature13954)

[Full Text @ Radboud UniversityCrossRefPubMedWeb of ScienceGoogle Scholar](#)

66. 66. ↵

1. Veglia F,
2. Perego M,
3. Gabrilovich D

. Myeloid-derived suppressor cells coming of age. *Nat Immunol.* 2018;**19**:2108–119. pmcid:29348500 [doi:10.1038/s41590-017-0022-x](https://doi.org/10.1038/s41590-017-0022-x)

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

67. [67.](#)
1. Pauken KE,
  2. Wherry EJ

. Overcoming T cell exhaustion in infection and cancer. *Trends Immunol.* 2015;**36**:4265–276. pmcid:4393798 [doi:10.1016/j.it.2015.02.008](https://doi.org/10.1016/j.it.2015.02.008)

[Full Text @ Radboud UniversityCrossRefPubMedGoogle Scholar](#)

68. [68.](#)

Gulley JL, Madan RA, Pachynski R, Mulders P, Sheikh NA, Trager J, et al. Role of Antigen Spread and Distinctive Characteristics of Immunotherapy in Cancer Treatment. *J Natl Cancer Inst.* 2017;109(4).

[Google Scholar](#)

[View Abstract](#)

## Footnotes

- **Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.
- Jeroen H. A. Creemers, Inge M. van Oort, I. Jolanda M. de Vries and Winald R. Gerritsen contributed equally to this work.
- **Funding** This work was supported by Stichting Afweer Tegen Kanker, Dr. Paul A.J. Speth Stichting and H2020 EU Grant PROCROP (Grant Number 635122). Carl G. Figdor received ERC Adv Grant PATHFINDER (269019) and the NWO Spinoza grant. I. Jolanda M. de Vries received NWO-Vici grant (918.14.655).
- **Supplementary information** **Supplementary information** accompanies this paper at [10.1186/s40425-019-0787-6](https://doi.org/10.1186/s40425-019-0787-6).
- **Authors' contributions** HW, GS, CGF, IJMdV and WRG conceived the study and drafted the clinical protocol. HW and JHAC provided care for the patients under supervision of IMvO, NM and WRG. EHJG, RDMM and MG administered the dendritic cell vaccines intranodally. EHJG and MG assessed the <sup>68</sup>Ga-PSMA

PET/CT findings. RDMM and JOB assessed the MRI findings. MAJG and MS performed and supervised immune histochemistry experiments and analyses. HW, JHAC and MAJG collected and analyzed the data and wrote the manuscript under supervision of GS, IJMdV and WRG. All authors reviewed the manuscript. All authors read and approved the final manuscript.

- **Availability of data and materials** The data that support the findings of this clinical trial are available from the authors upon request.
  - **Ethics approval and consent to participate** The Dutch Central Committee on Research involving Human Subjects has approved the study (NL49143.000.14) and written informed consent was obtained from all individuals. [ClinicalTrials.gov](https://clinicaltrials.gov) identifier of the study is [NCT02692976](https://clinicaltrials.gov/ct2/show/study/NCT02692976). The study was conducted in accordance with the Good Clinical Practice guidelines and with the provisions of the Declaration of Helsinki (October 9th 2004).
  - **Consent for publication** The patients have signed consent for the use of anonymized patient data for research purposes.
  - **Competing interests** IMvO has received grants, personal fees and non-financial support from Astellas; grants and personal fees from Janssen-Cilag; grants and personal fees from Bayer and personal fees from Sanofi (all outside the submitted work). NM reports personal fees from Bayer, grants and personal fees from Jansen-Cilag, personal fees from Merck Sharp & Dohme, grants and personal fees from Roche, grants and personal fees from Astellas, grants and personal fees from Sanofi (all outside the submitted work). JAW reports personal fees from Sanofi, Merck Sharp & Dohme, Bristol-Myers Squibb and Roche (all outside the submitted work). MB is employed as a Project Manager R&D at Miltenyi Biotec GmbH. KP is employed as a Senior Clinical Project Manager at Miltenyi Biotec GmbH. MG has received grants from Astellas and Sanofi, personal fees from Boehringer Ingelheim, Sanofi and Sandoz (all outside the submitted work). JOB is an advisor for SPL Medical and Soteria Medical. WRG reports personal fees from Bristol-Myers Squibb, Merck Sharp & Dohme, Janssen-Cilag, Bayer, Astellas, Bavarian Nordic, Sanofi, Amgen, Curevac, IQVIA, Dendreon, Morphosys, ORCA Pharmaceuticals and Sotio (all outside the submitted work).
- 

## Request Permissions

If you wish to reuse any or all of this article please use the link below which will take you to the Copyright Clearance Center's RightsLink service. You will be able to get a quick price and instant permission to reuse the content in many different ways.

[Request permissions](#)

## Copyright information:

© The Author(s). **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

## We recommend

1. [AB0009 Genetic association of mitochondrial dna polymorphisms with behÇet's disease in a korean population](#)  
M.-H. Kwon et al., Ann Rheum Dis, 2018
  2. [Differences in hemodynamic characteristics under high packing density between the porous media model and finite element analysis in computational fluid dynamics of intracranial aneurysm virtual treatment](#)  
Yeqing Jiang et al., J Neurointerv Surg, 2019
  3. [Chief physicians' perceived need for improvement of morbidity and mortality conferences: the role of structural and procedural characteristics](#)  
David L B Schwappach et al., Leader, 2018
  4. [Recurrent fetal hydrops with maternal M alloimmunisation: not a benign condition](#)  
Michelle Yu et al., Case Rep, 2019
  5. [1 Learning to lead](#)  
Emma Davies et al., Leader, 2018
1. [Synthesis of RNA containing a methylated blocked 5' terminus by HeLa nuclear homogenates.](#)  
Y Groner et al., Proc Natl Acad Sci U S A, 1975
  2. [Separation of reticulocyte initiation factor M 2 activity into two components.](#)  
D A Shafritz et al., Proc Natl Acad Sci U S A, 1972

3. [Effect of Applied Potential on Pit Propagation in Copper as Function of Water Chemistry](#)

Hung Ha et al., Journal of The Electrochemical Society, 2011

4. [Transmission of Bursaphelenchus xylophilus \(Nematoda: Aphelenchoididae\) through feeding activity of Monochamus alternatus \(Coleoptera: Cerambycidae\)](#)

WANG Yang<sup>1</sup> et al., Journal of Nanjing Forestry University (Natural Sciences), 2019

5. [Current Efficiency of an Ion-Selective Membrane: Effect of Fixed Charge Distribution](#)

Jyh-Ping Hsu et al., Journal of The Electrochemical Society, 1998