Letter to the Editor

Comments to “Persistent problems in the construction of matrix population models”

The use of matrix population models (MPMs*) to summarize and analyze the empirically observed demography of both plant and animal populations has exhibited rapid growth in recent decades (Salguero-Gómez et al., 2015, 2016). Comparative analyses using independent collections of MPMs have generated valuable ecological insights (e.g., Franco and Silvertown, 1990; Franco and Silvertown, 2004; Ramula et al., 2008; Buckley et al., 2010; Burns et al., 2010; Salguero-Gómez and Casper, 2010; Stott et al., 2010; Crone et al., 2011; Zeigler et al., 2013), leading to the recognition that a comprehensive MPM dataset would be a valuable resource for the ecological research community. We therefore launched the COMPADRE Plant Matrix Database (Salguero-Gómez et al., 2015) and the COMADRE Animal Matrix Database (Salguero-Gómez et al., 2016; <http://compadre-db.com/>), which comprise 1204 plant and animal species and 10,949 MPMs as of October 2019.

Our initial goal with COMPADRE and COMADRE was to present MPMs as intended by the authors of the original publications, relying on the authors and the peer review process to ensure that the published population projection matrix correctly described the intended life history. Thus, our error-checking procedure to date has focused for the most part on eliminating typographical errors introduced during digitization, rather than errors of MPM construction made by the original authors. However, Kendall et al. (2019) have shown that a substantial fraction of animal MPMs in peer-reviewed papers contain errors in construction, such that model components are missing and/or the model does not accurately represent the described life history of the species. Plant MPMs are also prone to errors in construction, including dormant life stages (most commonly a seed bank) that are missing (Doak et al., 2002; Nguyen et al., 2019) or incorrectly incorporated, leading to a one-year delay in the life cycle (Caswell, 2001, p. 60). Although some construction errors may not dramatically affect estimates of demographic parameters in some cases, they can be crucial in others. Estimates of generation time and population growth rate, in particular, can be sensitive to missing components or life stages (Kendall et al., 2019; Nguyen et al., 2019). Construction errors may also lead to directional biases, such as underestimating the elasticities of population growth rate to survival when fertility coefficients do not include survival (Kendall et al., 2019), and these biases may affect results in comparative analyses. Moving forward, there is a clear need for better training materials to help ecologists and conservation scientists (many with little experience in mathematical modeling) construct MPMs from their data. But how can we ensure that previously published MPMs, including those already compiled in COMPADRE and COMADRE and those remaining to be digitized, correctly represent the intended life histories, and thus support meaningful synthetic analyses?

We believe that an important principle in big data analyses can be summed up as “*Just because you can, does not mean you should*”. That is, although large volumes of data and tools that enable their fast analysis are available, they should not be used without careful consideration

and evaluation. There are many possible sources of error whenever large datasets are compiled, and we have previously discussed some of these (including errors in matrix construction by authors) as a cautionary note to data users (Salguero-Gómez et al., 2016). We also maintain an open, frequently updated digitization protocol, and an extensive user guide (both available on our GitHub repository, <https://github.com/jonesor/compadreDB>). These open, transparent policies are essential for users to fully understand the data and choose the subsets of MPMs that are most appropriate for addressing their specific research questions.

In our collective cases, comparative analyses of demographic data from COMPADRE and COMADRE (e.g., Vindenes and Engen, 2017; Beckman et al., 2018; Che-Castaldo et al., 2018; Davison et al., 2019; Healy et al., 2019; Jones et al., 2019; Nguyen et al., 2019; Roper et al., 2019; Pistón et al., 2019) have each resulted in a separate sample size because the research questions required a different set of selection criteria. For instance, for some questions related to estimating various moments of longevity, the subcomponents of fertility are not necessary (if one assumes no trade-offs between reproduction and survival), and so MPMs that do not include those (indicated by the metadata field “MatrixFec”) can be used. Similarly, to calculate the population growth rate from an MPM, the subcomponents of survival, sexual reproduction and clonality do not need to be separated (indicated by the metadata field “MatrixSplit”), but to estimate life history traits and vital rate sensitivities/elasticities, it is fundamental that they are separated. Thus, the set of selection criteria used for a specific study must be carefully designed before the analysis begins. Moreover, the availability of digitized data should not be a substitute for examining the original sources, which may provide important details of interest for the user’s specific goals. For this reason (and to facilitate the citation of original sources), COMPADRE and COMADRE include bibliographic information for the source studies.

Nevertheless, we agree that addressing the newly-identified MPM construction errors in COMPADRE and COMADRE would be a useful service for the scientific community. In fact, in recent years we have started to include information on data quality in the metadata. For example, there is a field (“SurvivalIssue”) for noting when the summed survival for a given age/stage exceeds 1.0, which may indicate a typographical error or a sensitivity matrix being incorrectly identified as an MPM in the original source. Other issues, such as when GPS coordinates are approximate rather than accurate, or environmental conditions that, while naturally occurring, may not be representative of typical conditions for a given population (e.g., fires, herbivory, droughts), are noted in the unstructured “Observations” field. In the coming months, we will take further steps including updating the metadata to record (1) whether the MPM is based on a pre-breeding or post-breeding census whenever the information is available, and (2) whether the MPM in the database has been altered from the original published one and why [e.g., typo in the original source; inclusion of an

unnecessary seed stage where no permanent seed bank exists (Caswell, 2001, p. 60)], or whether the MPM was calculated from other published data along with reference to those data (e.g., a life table, integral projection model, individual-based model). We have updated the digitization protocol to include Kendall et al. (2019) as essential reading so that our team can avoid the identified common errors when constructing MPMs from published life cycle data. Future releases of the database will include a unique identification number for each existing and new MPM, so that users can look up a specific MPM and all of its associated observations and metadata. They can also share with the community any additional metadata they collect for their own analyses, which could be linked back to the full dataset via the unique ID.

In the longer term, our plans include identifying MPMs that are likely to have construction issues following the protocol presented in Appendix B of Kendall et al. (2019). If the publication contains sufficient data, we will attempt to correct the MPMs and indicate why and how we have done so. This way, users will be able to identify the modified matrices and determine whether those modifications are appropriate for their use. In the last year, we have transitioned to a relational database that will allow us to keep a record of changes to the MPM data, so that we can record the MPMs as presented by original authors and also update them to fix known errors such as those mentioned above. However, often the published paper and supplementary materials do not include the necessary information (e.g., a separate life table or mean development time for each life history stage) to help us contextualize the MPMs and determine specific construction methods. Our digitization team invests a significant amount of time contacting authors for clarification and additional information. Without the original researchers' input, there may be many MPMs that we cannot correct or even flag with potential issues. This highlights the need for authors to consider providing as much detail on the life cycles, data collection, and biogeographic context for their species and populations as possible to enable reproducible research.

Although finding errors like the ones reported by Kendall et al. (2019) is worrisome, we emphasize that an evaluation of data quality issues and their extent (not to mention large-scale assessments in general) would not be possible without the existence of a database for MPMs. A database also allows us to implement solutions systematically across the entire set of demographic studies represented in the database. As a centralized repository of MPM data, COMPADRE and COMADRE can: (1) serve as a point of reference for recording whether a particular publication may have miscalculated MPMs, (2) pool efforts from independent groups or individuals to identify and/or fix the miscalculated MPMs, and (3) enable users to access and benefit from previously corrected MPMs. The database will only improve as we continue to aggregate data, assess data quality, and make corrections. Moreover, the mission of the COMPADRE and COMADRE team goes beyond releasing large volumes of data, and includes teaching workshops around the globe about MPMs and engaging the demographic community. We can therefore use our existing platform, including educational materials on our GitHub repository (see link above) and our regular user engagement workshops, to help raise awareness and promote more careful construction of MPMs.

We encourage and welcome the community to help validate the MPM data in COMPADRE and COMADRE, as the databases are currently supported by limited funding and volunteered researcher time. Conducting a comprehensive check in the databases for construction errors will require substantial effort, and we are only beginning to understand the extent of these issues. We are developing website tools (such as the new function for reporting data errors and corrections) to gather input and facilitate collaborative data validation. Participation from users and original authors will be vital in this effort to help us

deliver reliable, open-access MPM data for demographic researchers and the broader scientific community.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was supported by National Science Foundation, USA, DBI-1661342-000.

References

- Buckley, Y.M., et al., 2010. Causes and consequences of variation in plant population growth rate: a synthesis of matrix population models in a phylogenetic context. *Ecol. Lett.* 13, 1182–1197.
- Burns, J.H., et al., 2010. Empirical tests of life-history evolution theory using phylogenetic analysis of plant demography. *J. Ecol.* 98, 334–344.
- Beckman, N., Bullock, J., Salguero-Gómez, R., 2018. High dispersal ability is related to fast life history strategies. *J. Ecol.* 4, 1349–1362.
- Caswell, H., 2001. *Matrix Population Models: Construction, Analysis, and Interpretation*, 2nd edn. Sinauer Associates, Inc., Sunderland, MA, USA.
- Che-Castaldo, J., Che-Castaldo, C., Neel, M.C., 2018. Predictability of demographic rates based on phylogeny and biological similarity. *Conserv. Biol.* 32, 1290–1300.
- Crone, E.E., et al., 2011. How do plant ecologists use matrix population models? *Ecol. Lett.* 14, 1–8.
- Davison, R., Stadman, M., Jongejans, E., 2019. Stochastic effects contribute to population fitness differences. *Ecol. Modell.* 408, 108760.
- Doak, D.F., Thomson, D., Jules, E.S., 2002. Population viability analysis for plants: understanding the demographic consequences of seed banks for population health. In: Beissinger, S.R., McCullough, D.R. (Eds.), *Population Viability Analysis*. The University of Chicago Press, Chicago, Illinois, USA.
- Franco, M., Silvertown, J., 1990. Plant demography: what do we know? *Evol. Trends Plants* 4, 74–76.
- Franco, M., Silvertown, J., 2004. A comparative demography of plants based upon elasticities of vital rates. *Ecology* 85, 531–538.
- Healy, K., Ezard, T., Jones, O., Salguero-Gómez, R., Buckley, Y., 2019. Animal life history is shaped by the pace of life and the distribution of age-specific mortality and reproduction. *Nat. Ecol. Evol.* 3, 1217–1224.
- Jones, O.R., Ezard, T.H.G., Dooley, C., Healy, K., Hodgson, D.J., Mueller, M., Townley, S., Salguero-Gómez, R., 2019. My family and other animals: human demography under a comparative cross-species lens. In: Burger, O., Lee, R., Sear, R. (Eds.), *Human Evolutionary Demography*. Open Book Publishers, Cambridge, UK.
- Kendall, B.E., Fujiwara, M., Diaz-Lopez, J., Schneider, S., Voigt, J., Wiesner, S., 2019. Persistent problems in the construction of matrix population models. *Ecol. Modell.* 406, 33–43. <https://doi.org/10.1016/j.ecolmodel.2019.03.011>.
- Nguyen, V., Buckley, Y.M., Salguero-Gómez, R., Wardle, G.M., 2019. Consequences of neglecting cryptic life stages from demographic models. *Ecol. Modell.* 408 108723.
- Pistón, N., de Bello, F., Dias, A., Götzenberger, L., de Mattos, E., Rosado, B., Salguero-Gómez, R., Carmona, C., 2019. Multidimensional ecological analyses demonstrate how interactions between functional traits shape fitness and life history strategies. *J. Ecol.* 107, 2317–2328.
- Ramula, S., Knight, T.M., Burns, J.H., Buckley, Y.M., 2008. General guidelines for invasive plant management based on comparative demography of invasive and native plant populations. *J. Appl. Ecol.* 45, 1124–1133.
- Roper, M., Capdevila, P., Salguero-Gómez, R., 2019. Senescence - still an unsolved problem of biology. *Biorxiv*. <https://doi.org/10.1101/739730>.
- Salguero-Gómez, R., Casper, B.B., 2010. Keeping plant shrinkage in the demographic loop. *J. Ecol.* 98, 312–323.
- Salguero-Gómez, R., et al., 2015. The COMPADRE Plant Matrix Database: an open online repository for plant demography. *J. Ecol.* 103, 202–218.
- Salguero-Gómez, R., et al., 2016. COMADRE: a global data base of animal demography. *J. Anim. Ecol.* 85, 371–384.
- Stott, I., Franco, M., Carslake, D., Townley, S., Hodgson, D., 2010. Boom or bust? A comparative analysis of transient population dynamics in plants. *J. Ecol.* 98, 302–311.
- Vindenes, Y., Engen, S., 2017. Demographic stochasticity and temporal autocorrelation in the dynamics of structured populations. *Oikos* 126, 462–471.
- Zeigler, S., Che-Castaldo, J., Neel, M., 2013. Actual and potential use of population viability analysis in recovery of plant species listed under the U.S. Endangered Species Act. *Conserv. Biol.* 27, 1265–1278.

Judy Che-Castaldo*

Alexander Center for Applied Population Biology, Conservation & Science
Department, Lincoln Park Zoo, Chicago, IL 60614-4712, USA
E-mail address: jchecastaldo@lpzoo.org.

Owen R. Jones

Interdisciplinary Center on Population Dynamics & Department of Biology,
Campusvej 55, University of Southern Denmark, 5230 Odense M, Denmark

Bruce E. Kendall

Bren School of Environmental Science & Management, University of
California Santa Barbara, Santa Barbara, CA 93106-5131, USA

Jean H. Burns

Department of Biology, Case Western Reserve University, Cleveland, OH
44106-7080, USA

Dylan Z. Childs

Department of Animal and Plant Sciences, University of Sheffield, Sheffield
S10 2TN, UK

Thomas H.G. Ezard

Ocean & Earth Science, National Oceanography Centre Southampton,
University of Southampton Waterfront Campus, Southampton SO14 3ZH,
UK

Haydee Hernandez-Yanez

Alexander Center for Applied Population Biology, Conservation & Science
Department, Lincoln Park Zoo, Chicago, IL 60614-4712, USA

David J. Hodgson

Centre for Ecology & Conservation, College of Life and Environmental
Sciences, University of Exeter, Cornwall, UK

Eelke Jongejans

Animal Ecology and Physiology Department, Radboud University, Nijmegen,
the Netherlands

Tiffany Knight^{a,b,c}

^a Institute of Biology, Martin Luther University Halle-Wittenberg, Am
Kirchtor 1, 06108 Halle (Saale), Germany

^b Department of Community Ecology, Helmholtz Centre for Environmental
Research-UFZ, Theodor-Lieser-Straße 4, 06120 Halle (Saale), Germany

^c German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-
Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

Cory Merow

Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT
06269, USA

Satu Ramula

Department of Biology, University of Turku, FI-20014 Turku, Finland

Iain Stott

School of Life Sciences, University of Lincoln, Brayford Pool, LN6 7TS
Lincoln, UK

Yngvild Vindenes

Centre for Ecological and Evolutionary Synthesis, Department of
Biosciences, University of Oslo, Oslo, Norway

Hiroyuki Yokomizo

Center for Health and Environmental Risk Research, National Institute for
Environmental Studies, Tsukuba 305-8506, Japan

Roberto Salguero-Gómez

Department of Zoology, 11a Mansfield Rd, University of Oxford, OX1 3SZ
Oxford, UK

* Corresponding author.