

Review

## Hyperhomocysteinaemia: a role in the accelerated atherogenesis of chronic renal failure?

M.J.F.M. Janssen <sup>a,\*</sup>, M. van den Berg <sup>b</sup>, C.D.A. Stehouwer <sup>a</sup>, G.H.J. Boers <sup>c</sup>

<sup>a</sup> *ICaR-VU, Department of Internal Medicine, University Hospital, Free University, De Boelelaan 1117, 1081 HV Amsterdam, Netherlands*

<sup>b</sup> *Department of Surgery, University Hospital, Free University, De Boelelaan 1117, 1081 HV Amsterdam, Netherlands*

<sup>c</sup> *Department of Internal Medicine, University Hospital, Nijmegen, Netherlands*

Received 21 July 1994; revision received 14 September 1994; accepted 15 September 1994

---

### Abstract

Moderate hyperhomocysteinaemia has recently been established as an independent risk factor for atherothrombotic disease. It might be caused by heterozygosity for cystathionine  $\beta$ -synthase deficiency, an enzyme involved in the conversion of methionine to cysteine through the transsulphuration pathway or by inherited thermolability of the enzyme which remethylates homocysteine into methionine. In chronic renal failure (CRF) homocysteine levels are significantly elevated at a relatively early stage. The normal kidney possibly plays an important role in homocysteine catabolism, which cannot be performed in CRF. Alternatively, decreased extrarenal catabolism can contribute to the hyperhomocysteinaemia in this disease state. Treatment with folic acid, 5 mg daily, significantly lowers homocysteine levels in chronic renal patients.

*Keywords:* Hyperhomocysteinaemia; Chronic renal failure; Haemodialysis; Folic acid

---

### 1. Introduction

Homocysteine is a sulphhydryl amino acid that does not occur in protein structures; it is formed as an intermediate by demethylation of methionine [1]. In many tissues, it can be converted to cystathionine by condensation with serine (Fig. 1). This irreversible reaction, part of the transsulphuration pathway, is catalyzed by the enzyme, cystathionine  $\beta$ -synthase (EC 4.2.1.22; CBS), and

requires pyridoxal 5'-phosphate (active form of vitamin B<sub>6</sub>) as co-factor. Alternatively, homocysteine can be remethylated to methionine by two different reactions. The first is catalyzed by 5-methyltetrahydrofolate-homocysteine methyltransferase (methionine synthase, EC 2.1.1.13; MH), with 5-methyltetrahydrofolate (5-mTHF, from folic acid) as the methyl-group donor and methylcobalamin (from vitamin B<sub>12</sub>) as coenzyme. The second pathway is catalyzed by betaine-homocysteine methyltransferase (EC 2.1.1.5; BH), with betaine (Fig. 2) serving as the methyl donor. In normal subjects about 70% of total homocysteine in plasma is protein-bound, for the largest

---

\* Corresponding author. Tel. (20) 5489111; Fax (20) 5484898.

part to albumin [2–4]. In freshly prepared plasma, the free homocysteine, i.e. the homocysteine that is not protein-bound, exists as a mixture of cysteine-homocysteine mixed disulphide (CHMD, Fig. 2) and trace amounts of homocystine and homocysteine [5].

Normal fasting concentrations of free and total homocysteine are about 3 and about 10  $\mu\text{mol/l}$ , respectively [4–6], premenopausal women having significantly lower values than postmenopausal women and men [7,8]. It has been suggested that efficient homocysteine metabolism is a protective factor against vascular disease in women during their reproductive years [9].

Homocystinuria is an inborn error of metabolism, usually associated with a defect of transsulphuration caused by CBS deficiency, inherited as an autosomal recessive trait. Heritable defects in methyltetrahydrofolate or methylcobalamin synthesis are very rare causes of homo-

cystinuria [1]. Untreated (homozygotic) CBS-deficient patients have high total homocysteine levels in the fasting state up to 200  $\mu\text{mol/l}$  [5]. Clinical characteristics are dislocation of the optic lens, osteoporosis, vertebral anomalies and mental retardation. Arterial and venous thromboembolism, occurring at a rate of 50% by the age of 30 years, constitutes the major cause of death.

The frequency of heterozygosity for homocystinuria is estimated to be 1:70 to 1:290 [1]. Carriers have slightly elevated fasting homocysteine values which overlap with those of normal controls, but they can be detected by l-methionine loading [10,11]. When given an oral dose of methionine of 0.1 g/kg body weight, heterozygotes, when compared with controls, reach higher homocysteine levels 6 h after loading, due to their partially deficient transsulphuration. On the basis of such testing also, up to 35% of patients presenting with symptoms of arterial occlusive dis-




Fig. 1. The main reactions in methionine and homocysteine metabolism by transsulphuration and remethylation. MH = 5,10-methyltetrahydrofolate-homocysteine methyltransferase (methionine synthase); BH = betaine-homocysteine methyltransferase; 5-meTHFR = 5-methylenetetrahydrofolate reductase; CBS = cystathionine  $\beta$ -synthase; 5-mTHF = 5-methyltetrahydrofolate; 5-meTHF = 5-methylenetetrahydrofolate; B<sub>6</sub> = pyridoxal 5'-phosphate (from vitamin B<sub>6</sub>).

ease before the age of 50 years could be shown to be hyperhomocysteinaemic, what identified mild hyperhomocysteinaemia as a risk factor for premature occlusive arterial disease [12–15]. In only a few of the reports has the cause of this hyperhomocysteinaemia in vascular patients been investigated. In the majority of these cases an intermediately deficient activity of CBS has been demonstrated in their cultured fibroblasts [12,16]. About 50% of adult patients with the homozygous enzyme defect respond to oral doses of pyridoxine of 750–1000 mg/day by normalization of the biochemical abnormalities [17]. There is evidence of the effectiveness of pyridoxine in preventing further thromboembolic events in these patients [17].

In the case of non-responsiveness to pyridoxine, folic acid and betaine can be given in order

to stimulate the remethylation of homocysteine [18,19].

Patients with premature arterial disease and mild hyperhomocysteinaemia comparable with that in carriers of CBS deficiency are currently being treated with pyridoxine 250 mg/day and folic acid 5 mg/day. The effectiveness of this medication in the prevention of further vascular damage is a matter of investigation.

Apart from heterozygosity for CBS deficiency, other enzyme defects may be the cause of moderate hyperhomocysteinaemia. A thermolabile form of 5,10-methylenetetrahydrofolate reductase (EC 1.1.1.68; 5-meTHFR, Fig. 1) has been demonstrated in 17% of patients with premature coronary artery disease, possibly causing a lack of 5-methyltetrahydrofolate (5-mTHF) and hampering homocysteine remethylation [20].




Fig. 2. Structural formulae of some molecules involved in homocysteine metabolism.

Acquired forms of hyperhomocysteinaemia have been found in severe liver disease [21], in vitamin B<sub>12</sub> [22] and folate deficiency [23], and in chronic renal failure (CRF).

## 2. The hyperhomocysteinaemia of chronic renal failure

Cardiovascular disease is the major cause of death in patients on maintenance dialysis [24] and is only partially accounted for by the high prevalence of hypertension, smoking, hyperlipidaemia and diabetes mellitus in this population [25].

### *Plasma homocysteine in CRF patients not requiring maintenance dialysis*

Wilcken et al. were the first to demonstrate that plasma homocysteine is increased in patients with chronic renal failure [26]. In 22 patients not requiring dialysis (mean serum creatinine  $\pm$ SD:  $680 \pm 300 \mu\text{mol/l}$ ) the fasting plasma cysteine-homocysteine mixed disulphide concentration (CHMD), reflecting unbound homocysteine, was increased significantly compared to the values of an equally large control group:  $8.2 \pm 3.4$  versus  $3.1 \pm 1.0 \mu\text{mol/l}$  ( $P < 0.001$ ). The increase was positively correlated with the loss of renal function, as assessed by the serum creatinine concentration ( $r = 0.62$ ;  $P < 0.01$ ).

Soria et al. studied total homocysteine levels (= free + protein-bound homocysteine) in 53 patients with different degrees of chronic renal failure [27]. Group I consisted of 25 patients with moderate renal failure (serum creatinine  $\leq 300 \mu\text{mol/l}$ ). Group II comprised 18 patients with serum creatinine levels between 300 and 600  $\mu\text{mol/l}$ . The remaining 10 patients constituted Group III and had serum creatinine values above 600  $\mu\text{mol/l}$ . Plasma homocysteine in Group I was substantially elevated:  $18.2 \pm 1.7 \mu\text{mol/l}$  versus  $8.0 \pm 0.3 \mu\text{mol/l}$  (mean  $\pm$  SEM) in 45 controls.

Group II had even higher homocysteine levels:  $27.3 \pm 3.2$  and group III had values comparable with group II, i.e.  $27.3 \pm 1.7 \mu\text{mol/l}$ . Again the homocysteine concentration was correlated with the serum creatinine concentration ( $r = 0.76$ ).

Recently, a third study was published on hyperhomocysteinaemia in patients with various degrees of chronic renal failure [28]. Again, three groups were distinguished. The first one ( $n = 28$ ), having a creatinine clearance ( $C_{\text{cr}}$ ) of 30–75 ml/min, had a mean ( $\pm$ SD) total plasma homocysteine concentration of  $16.2 \pm 8.1 \mu\text{mol/l}$ . In the second group ( $n = 29$ ), with  $C_{\text{cr}}$  ranging from 10–30 ml/min, homocysteine levels were  $23.3 \pm 14.7 \mu\text{mol/l}$ , while in the third group ( $n = 22$ ), with advanced renal failure ( $C_{\text{cr}} < 10 \text{ ml/min}$ ), homocysteine values of  $29.5 \pm 14.4 \mu\text{mol/l}$  were observed. All groups had significantly higher levels than the 45 controls ( $8.2 \pm 2.2 \mu\text{mol/l}$ ). Linear regression analysis showed a negative correlation between  $C_{\text{cr}}$  and total plasma homocysteine ( $r = 0.40$ ;  $P < 0.01$ ). All patients received moderate protein restriction (0.6–1.0 g/kg/day) according to the degree of renal failure. Folic acid supplementation was given in none of these three studies.

### *Plasma homocysteine in dialysis patients*

Wilcken et al. studied 19 chronic haemodialysis patients [29]. The cysteine-homocysteine mixed disulphide concentration (CHMD) before dialysis was  $6.3 \pm 3.7 \mu\text{mol/l}$ . After haemodialysis CHMD was  $2.7 \pm 1.2 \mu\text{mol/l}$ , even lower than the values obtained in 22 normal subjects ( $3.1 \pm 1.0 \mu\text{mol/l}$ ).

Smolin et al. reported elevated CHMD and protein-bound homocysteine concentrations in chronic haemodialysis patients taking 1 mg folic acid and 10 mg pyridoxine per day [30]. Haemodialysis substantially lowered the unbound fraction from  $14.7 \pm 6.4$  to  $7.9 \pm 4.9 \mu\text{mol/l}$ , leaving the protein-bound fraction largely intact. Pre-dialysis protein-bound homocysteine of  $349 \pm 77 \text{ nmol/g}$  protein in the renal patients remained high after dialysis, i.e.  $312 \pm 71 \text{ nmol/g}$  protein, versus  $153 \pm 50 \text{ nmol/g}$  protein in fasting controls.

Kang et al. determined the protein-bound homocysteine in 13 haemodialysis patients on 5 mg folic acid daily [31]. Before dialysis they measured  $197.7 \pm 19.6 \text{ nmol/g}$  protein, after dialysis  $153.9 \pm 19.1 \text{ nmol/g}$  protein, constituting a decrease of 23%. Protein-bound homocysteine in 13 controls was  $53.4 \pm 3.6 \text{ nmol/g}$  protein.

### *The mechanism of hyperhomocysteinaemia in CRF*

The renal handling of homocysteine is incompletely understood. Since about 30% of plasma homocysteine is not protein-bound, this fraction freely passes the glomerular basement membrane; the daily filtered load can be calculated to be about 550  $\mu\text{mol}$ . Thus, a large part of the circulating unbound homocysteine is delivered to the proximal tubular cells. The quantity of homocysteine in 24-h urine samples varies from 3.5 to 10  $\mu\text{mol}$  [3]; urinary homocysteine clearance is only 0.3% of creatinine clearance [7]. The low urinary excretion is accounted for by the avid reabsorption taking place in the cortical tubule: more than 99.5% of the filtered homocysteine is reabsorbed by the proximal tubular cells [32]. Reabsorption in the proximal tubule occurs via two saturable systems. The first, a high-affinity, low- $K_m$  (0.17 mM) system, is shared with cystine, arginine and lysine [32]. Urinary homocysteine excretion can be stimulated by infusion of these amino acids [33]. The second system with low-affinity, high- $K_m$  (7.65 mM) characteristics, seems to be shared with cystine only [32]. The fate of the reabsorbed homocysteine is unknown. In the human kidney cystathionine  $\beta$ -synthase activity is low [34], whereas the activity of both the homocysteine remethylation enzymes is high [35,36]. Assuming that the filtered and reabsorbed homocysteine is catabolized in the proximal tubule cell, the kidney may normally play a substantial role in homocysteine metabolism. In chronic renal failure, renal catabolism may be hampered because homocysteine, due to the decreased glomerular filtration, cannot reach its renal metabolic site, i.e. the proximal tubular cell. Moreover, tubular cell dysfunction itself in renal failure may contribute to the raised homocysteine levels due to the fact that homocysteine is passing the tubular cell into the peritubular capillaries without being catabolized.

In CRF, total body catabolism of homocysteine has been thought to be changed, mainly on the basis of the low serine levels measured in this condition. Serine is involved in homocysteine metabolism in two ways (Fig. 1). Firstly, it is converted to glycine by donating a methyl group in the restoration of 5,10-methylenetetrahydrofo-

late from tetrahydrofolate, and secondly, it condenses with homocysteine to form cystathionine in the transsulphuration route. The low serine values in CRF have been said to point to enhanced catabolism of homocysteine via these two reactions [37]. The normal kidney, however, produces rather large quantities of serine [38]. In CRF, there is a 80–90% reduction of this production, which itself can explain the low serine levels seen in CRF [38,39]. Consequently, low serine levels do not necessarily reflect an increased catabolism of homocysteine. On the contrary, a feedback inhibition of homocysteine metabolism in the liver and other organs by heightened levels of uraemic toxins has been suggested to contribute to the hyperhomocysteinaemia of CRF [37].

Could suboptimal levels of folic acid, vitamin B<sub>6</sub> or B<sub>12</sub> contribute to the raised homocysteine concentration in CRF patients?

All studies of hyperhomocysteinaemia in dialysis patients have been performed in patients on haemodialysis. Folate deficiency is unlikely to develop in such patients on a regular diet [40–43]. Moreover, Smolin et al. found high homocysteine levels in 24 haemodialysis patients on a daily multivitamin supplement providing 1 mg of folic acid and 10 mg of pyridoxine [30]. All the patients with stable CRF that were included in the published trials exploring the effect of folic acid on the homocysteine concentration had normal baseline serum- and red cell folate levels, as well as normal serum B<sub>12</sub> concentrations [44,45]. Furthermore, supplementation with B<sub>12</sub> and with high-dose pyridoxine was shown not to influence the homocysteine levels of these patients [44]. Therefore, it is unlikely that deficiencies of folic acid, vitamin B<sub>12</sub> or pyridoxine contribute significantly to the elevated homocysteine levels in CRF patients.

A reduction of the protein intake has been shown to retard the progression of CRF. Protein restriction may lower the plasma homocysteine concentration in advanced CRF. However, the reduced intake of methionine that is induced by protein restriction may only level off the homocysteine levels in advanced renal failure [27].

### 3. Treatment options for hyperhomocysteinaemia in CRF

There is a limited number of publications on the treatment of heightened homocysteine levels in renal patients. In 1981, Wilcken et al. tested the effect of administering the co-factors for homocysteine metabolism to CRF patients [44]. Eleven long-term renal transplant recipients, with a mean ( $\pm$ SD) serum creatinine concentration of  $210 \pm 90 \mu\text{mol/l}$  and a mean ( $\pm$ SD) cysteine homocysteine mixed disulphide (CHMD) of  $7.3 \pm 2.1 \mu\text{mol/l}$ , received folic acid in a daily dose of 5 mg. After 2 weeks CHMD had declined to  $5.1 \pm 0.7 \mu\text{mol/l}$ . After 4 weeks of treatment CHMD was  $4.3 \pm 0.8 \mu\text{mol/l}$ ; continuation of the treatment did not further lower CHMD. Separate or additive effects of pyridoxine (100 mg/day) or vitamin B<sub>12</sub> (1000  $\mu\text{g}$  im) on plasma CHMD were not detectable. Wilcken later confirmed the effectiveness of folic acid in 21 CRF patients with a mean ( $\pm$ SD) serum creatinine concentration of  $560 \pm 240 \mu\text{mol/l}$  [45]. Baseline free homocysteine was  $12.9 \pm 6.8 \mu\text{mol/l}$  for the patients and  $4.2 \pm 0.8 \mu\text{mol/l}$  in 24 normal controls. Administration of folic acid 5 mg/day to the patients for  $15 \pm 6$  days decreased the free homocysteine to  $6.8 \pm 2.8 \mu\text{mol/l}$ . This fall was linearly related to the pre-treatment level of free homocysteine ( $r = 0.92$ ). The decline of homocysteine concentrations in CRF patients treated with folic acid coincided with a further decrease of plasma serine and a rise in plasma glycine, indicating the stimulation of remethylation by folic acid [45]. In both studies by Wilcken on the treatment of hyperhomocysteinaemia in CRF the patients had normal pre-treatment levels of folic acid and vitamin B<sub>12</sub>; the concentration of vitamin B<sub>6</sub> had not been determined.

### 4. Conclusion

There is a significant increase in plasma homocysteine concentration in patients with CRF, hyperhomocysteinaemia being present from the incipient stage of renal failure and increasing further as renal function deteriorates. To date, the

origin of hyperhomocysteinaemia in CRF is now largely unclear and awaits further investigation. Decreased catabolism of homocysteine in the proximal tubular cells due to reduced filtration and to tubular dysfunction and decreased extrarenal homocysteine metabolism due to inhibition by retained metabolites may both contribute. Folic acid treatment can lower homocysteine concentrations in these patients. Further research is needed into the optimal dose of folic acid, not only in patients with advancing renal failure, but also in haemodialysis and peritoneal dialysis patients. The possible contribution of betaine, the other known stimulator of remethylation, should also be explored. Lowering the homocysteine concentration in CRF patients could aid in the prevention of cardiovascular disease in CRF patients, which constitutes a major problem in medical practice.

### Acknowledgement

We wish to thank Prof. Dr. A.J.M. Donker (Internist, Department of Internal Medicine, Free University Hospital, Amsterdam) for a critical reading of the manuscript.

### References

- [1] Mudd SH, Levy HL, Skovby F. Disorders of transsulfuration. In: Stanbury JB, ed. *Metabolic basis of inherited diseases*. New York, NY: McGraw-Hill Publications, 1989;693–734.
- [2] Kang SS, Wong PWK, Becker N. Protein-bound homocyste(i)ne in normal subjects and in patients with homocystinuria. *Pediatr Res* 1979;13:1141–1143.
- [3] Refsum H, Helland S, Ueland PM. Radioenzymic determination of homocysteine in plasma and urine. *Clin Chem* 1985;31:624–628.
- [4] Wiley VC, Dudman PB, Wilcken DEL. Interrelations between plasma free and protein-bound homocysteine and cysteine in homocystinuria. *Metabolism* 1988;37:191–195.
- [5] Ueland PM, Refsum H. Plasma homocysteine, a risk factor for vascular disease: plasma levels in health, disease and drug therapy. *J Lab Clin Med* 1989;114:473–501.
- [6] Araki A, Sako Y. Determination of free and total homocysteine in human plasma by high-performance liquid chromatography with fluorescence detection. *J Chromatogr* 1987;422: 43–52.

- [7] Stabler SP, Marcell PD, Podell ER, Allen RH. Quantitation of total homocysteine, total cysteine, and methionine in normal serum and urine using capillary gas chromatography–mass spectrometry. *Anal Biochem* 1987;162:185–196.
- [8] Kang SS, Wong PWK, Cook HY, Norusis M, Messer JV. Protein-bound homocysteine: a possible risk factor for coronary artery disease. *J Clin Invest* 1986;77:1482–1486.
- [9] Boers GHJ, Smals AG, Trijbels FJ, Leermakers AI, Kloppenburg PW. Unique efficiency of methionine metabolism in premenopausal women may protect against vascular disease in the reproductive years. *J Clin Invest* 1983;72: 1971–1976.
- [10] Boers GH, Fowler B, Smals AGH, et al. Improved identification of heterozygotes for homocystinuria due to cystathionine synthase deficiency by the combination of methionine loading and enzyme determination in cultured fibroblasts. *Hum Genet* 1985;69:164–169.
- [11] Sardharwalla IB, Fowler B, Robins AJ, Komrower GM. Detection of heterozygotes for homocystinuria. Study of sulphur-containing amino acids in plasma and urine after L-methionine loading. *Arch Dis Child* 1974;49:553–559.
- [12] Boers GHJ, Smals AGH, Trijbels FJM et al. Heterozygosity for homocystinuria in premature peripheral and cerebral occlusive arterial disease. *N Engl J Med* 1985;313:709–715.
- [13] Israelsson B, Brattstrom L, Hultberg B, et al. Homocysteine and myocardial infarction. *Atherosclerosis* 1988;71: 227–233.
- [14] Malinow MR, Kang SS, Taylor LM et al. Prevalence of hyperhomocyst(e)inaemia in patients with peripheral arterial occlusive disease. *Circulation* 1989;79:1180–1188.
- [15] Coull BM, Malinow MR, Beamer N, Sexton G, Nordt F, de Garmo P. Elevated plasma homocyst(e)ine concentration as a possible independent risk factor for stroke. *Stroke* 1990;21: 572–576.
- [16] Clarke R, Daly L, Robinson K, et al. Hyperhomocysteinemia: an independent risk factor for vascular disease. *N Engl J Med* 1991;324: 1149–1155.
- [17] Mudd SH, Skovby F, Levy HL et al. The natural history of homocystinuria due to cystathionine  $\beta$ -synthase deficiency. *Am J Hum Genet* 1985;37:1–31.
- [18] Brattström LE, Israelsson B, Norrving B et al. Impaired homocysteine metabolism in early-onset cerebral and peripheral occlusive disease. Effects of pyridoxine and folic acid treatment. *Atherosclerosis* 1990;81:51–60.
- [19] Wilcken DEL, Wilcken B, Dudman NPB et al. Homocystinuria – the effects of betaine in the treatment of patients not responsive to pyridoxine. *N Engl J Med* 1983;309: 448–453.
- [20] Kang SS, Zhou J, Wong PWK, Kowalysyn J, Strokosch G. Intermediate homocystineinemia: a thermolabile variant of methylenetetrahydrofolate reductase. *Am J Hum Genet* 1988;43: 414–421.
- [21] Horowitz JH, Rypins EB, Henderson JM et al. Evidence for impairment of transsulfuration pathway in cirrhosis. *Gastroenterology* 1981;81:668–675.
- [22] Stabler SP, Marcell PD, Podell ER, Allen RH, Savage DG, Lindenbaum J. Elevation of total homocysteine in the serum of patients with cobalamin or folate deficiency detected by gas chromatography–mass spectrometry. *J Clin Invest* 1988;81: 466–474.
- [23] Kang SS, Wong PWK, Norusis M. Homocysteinemia due to folate deficiency. *Metabolism* 1987;36:458–462.
- [24] Port FK. Mortality and causes of death in patients with end-stage renal failure. *Am J Kidney Dis* 1990;15: 215–217.
- [25] Ibels LS, Stewart JH, Mahogoney JF et al. Occlusive arterial disease in uraemic and haemodialysis patients and renal transplant recipients. A study of the incidence of arterial disease and of the prevalence of risk factors implicated in the pathogenesis of arteriosclerosis. *Q J Med* 1977; 46:197–214.
- [26] Wilcken DEL, Gupta VJ. Sulfur containing aminoacids in chronic renal failure with particular reference to homocystine and cysteine-homocysteine mixed disulphide. *Eur J Clin Invest* 1979;9:301–307.
- [27] Soria C, Chadeaux B, Coudé M, Gaillard O, Kamoun P. Concentrations of total homocysteine in plasma in chronic renal failure. *Clin Chem* 1990;36:2137–2138.
- [28] Chauveau P, Chadeaux B, Coudé M et al. Increased plasma homocysteine concentration in patients with chronic renal failure. *Min Electr Metab* 1992;18:196–198.
- [29] Wilcken DEL, Gupta VJ, Reddy SG. Accumulation of sulphur containing amino acids including cysteine – homocysteine in patients on maintenance haemodialysis. *Clin Sci* 1980; 58:427–430.
- [30] Smolin LA, Laidlaw SA, Kopple JD. Altered plasma free and protein-bound sulfur aminoacid levels in patients undergoing maintenance hemodialysis. *Am J Clin Nutr* 1987;45: 737–743.
- [31] Kang SS, Wong PWK, Bidani A, Milanez S. Plasma protein-bound homocyst(e)ine in patients requiring chronic haemodialysis. *Clin Sci* 1983;65:335–336.
- [32] Foreman JW, Wald H, Blumberg G, Pepe LM, Segal S. Homocystine uptake in isolated cortical tubules. *Metabolism* 1982;31: 613–619.
- [33] Cusworth DC, Gattreau A. Inhibition of renal tubular reabsorption of homocystine by lysine and arginine. *Lancet* 1968;ii: 916–917.
- [34] Sturman JA, Rassin DK, Gaull GE. Distribution of transsulfuration enzymes in various organs and species. *Int J Biochem* 1970;1:251–253.
- [35] Mudd SH, Levy HL, Morrow G III. Deranged B<sub>12</sub> metabolism: effects on sulfur amino acid metabolism. *Biochem Med* 1970;4: 193–214.
- [36] Gaull GE, von Berg W, Raiha NCR, Sturman JA. Development of methyltransferase activities of human fetal tissues. *Pediatr Res* 1973;7:527–533.
- [37] Hultberg B, Agardh E, Andersson A et al. Increased levels of plasma homocysteine are associated with nephropathy, but not severe retinopathy in type I diabetes mellitus. *Scand J Clin Lab Invest* 1991;51:277–282.
- [38] Tizianello A, De Ferrari G, Garibotto G, Gurreri G,

- Robaudo C. Renal metabolism of amino acids and ammonia in subjects with normal renal function and in patients with chronic renal insufficiency. *J Clin Invest* 1980;65:1162–1173.
- [39] Fukuda S, Kopple JD. Uptake and release of amino acids by the kidney of dogs made chronically uremic with uranyl nitrate. *Min Elect Metab* 1980;3:248–260.
- [40] Westhuyzen J, Matherson K, Tracey R, Fleming SJ. Effect of withdrawal of folic acid supplementation in maintenance hemodialysis patients. *Clin Nephrol* 1993;40:96–99.
- [41] Cunningham J, Sharman VL, Goodwin FJ, Marsh FP. Do patients receiving haemodialysis need folic acid supplements? *Br Med J* 1981;282:1582.
- [42] Swainson CP, Winney RJ. Do dialysis patients need extra folate? *Lancet* 1983;i:239.
- [43] Lash J, Smith EC. Is folate supplementation necessary in dialysis patients? *Int J Artif Org* 1990;13:785–786.
- [44] Wilcken DEL, Gupta VJ, Betts AK. Homocysteine in the plasma of renal transplant recipients: effects of cofactors for methionine metabolism. *Clin Science* 1981;61:743–749.
- [45] Wilcken DEL, Dudman NPB, Tyrrell PA, Robertson MR. Folic acid lowers elevated plasma homocysteine in chronic renal insufficiency: possible implications for prevention of vascular disease. *Metabolism* 1988;37:697–701.