



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Ten myths about Functional Neurological Disorder

**Citation for published version:**

Lidstone, SC, Araújo, R, Stone, J & Bloem, BR 2020, 'Ten myths about Functional Neurological Disorder', *European Journal of Neurology*. <https://doi.org/10.1111/ene.14310>

**Digital Object Identifier (DOI):**

[10.1111/ene.14310](https://doi.org/10.1111/ene.14310)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

European Journal of Neurology

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Ten myths about Functional Neurological Disorder

Journal:	<i>European Journal of Neurology</i>
Manuscript ID	EJoN-20-0838
Wiley - Manuscript type:	Letters to the Editor
Date Submitted by the Author:	27-Apr-2020
Complete List of Authors:	Lidstone, Sarah; Toronto Western Hospital Krembil Neuroscience Centre, Medicine, Division of Neurology; University Health Network, Movement Disorders Clinic and the Edmond J. Safra Program in Parkinson's Disease Araújo, Rui; Centro Hospitalar Universitário de São João, Neurology Stone, Jon; University of Edinburgh, Clinical Neurosciences Bloem, Bastiaan; Radboud university medical Centre; Donders Institute for Brain, Cognition and Behaviour, Department of Neurology
Keywords:	Functional disorders < Psychiatric disorders < NEUROLOGICAL DISORDERS, conversion disorder, functional movement disorder, psychogenic, myths

# Ten myths about Functional Neurological Disorder

Sarah C. Lidstone, MD, PhD, Rui Araújo MD,  
Jon Stone MD, PhD, Bastiaan R. Bloem, MD, PhD

Sarah C. Lidstone, MD, PhD, FRCPC  
Gloria and Morton Shulman Movement Disorders Clinic  
Toronto Western Hospital  
399 Bathurst Street, Toronto, Ontario  
Sarah.lidstone@uhnresearch.ca

Rui Araújo, MD, FEBN  
Department of Neurology, Centro Hospitalar Universitário de São João, E.P.E,  
Department of Clinical Neurosciences and Mental Health, Faculty of Medicine of  
University of Porto  
Porto, Portugal  
rmma22@gmail.com

Jon Stone, MB ChB PhD  
Centre for Clinical Brain Sciences,  
University of Edinburgh,  
Edinburgh, Scotland  
Jon.stone@ed.ac.uk

Bastiaan R. Bloem, MD, PhD, FRCPE (corresponding author)  
Radboud University Medical Centre  
Donders Institute for Brain, Cognition and Behaviour  
Department of Neurology  
Nijmegen, the Netherlands  
bas.bloem@radboudumc.nl

**Word count:** 1590

**Key words:** functional neurological disorder; functional movement disorder; myths;  
diagnosis; treatment; psychogenic; conversion disorder

**Figures:** 0

**Tables:** 0

**Running title:** Ten myths of FND

1  
2  
3 Functional neurological disorder (FND), is the second most common reason for  
4 referral to a neurologist. Despite being potentially treatable, FND remains  
5 underdiagnosed and associated with high levels of disability and distress. In this  
6 article, we address ten common myths about FND that continue to obstruct the  
7 diagnosis and treatment for these patients, and present them with accompanying  
8 lessons for the clinician.  
9  
10

### 11 ***Myth 1: FND is a diagnosis of exclusion***

12  
13  
14 The diagnosis of FND should be approached in the same way as other medical  
15 conditions and be diagnosed using positive features, rather than by ruling out other  
16 conditions.(1) Positive signs on examination include Hoover's sign of functional limb  
17 weakness, where hip extension weakness returns to normal when attention is  
18 diverted to the contralateral leg. Functional tremor may 'entrain' to the frequency of  
19 imposed rhythmic movements in another body part. Functional (dissociative)  
20 seizures can be recognised by a combination of typical features such as eyes tightly  
21 closed during an episode, long duration, or awareness of generalized shaking. Thus,  
22 the diagnosis of FND should be based primarily on typical features only seen in this  
23 disorder or internal inconsistency of the symptoms and signs.(2)

24  
25 Lesson: The diagnosis of FND should be "ruled in" based on the presence of positive  
26 signs.  
27  
28

### 29 ***Myth 2: Patients have either FND or another neurological disorder***

30  
31  
32 The presence of another neurological condition is a powerful risk factor for the  
33 development of FND and they commonly occur together. Physical injury, surgical  
34 procedures and migraine often precipitate functional movements in vulnerable  
35 individuals, and up to 20% of patients with functional (dissociative) seizures also  
36 have epilepsy.(3) Some conditions like Parkinson's disease appear particularly likely  
37 to present with comorbid FND, perhaps because of shared neurobiology, but  
38 symptoms and disability from any neurological condition may be sufficient. For this  
39 reason, there should be a low threshold for investigations.  
40  
41

42 Lesson: FND commonly co-occurs with other neurological disorders.  
43  
44

### 45 ***Myth 3: A bizarre presentation indicates FND***

46  
47 FND is frequently equated to bizarre or unrecognizable clinical neurological  
48 presentations, such as complex movement disorders. In fact, several genetic or  
49 acquired movement disorders present with strikingly bizarre phenomenology.(4–6)  
50 In such cases, the "bizarreness" is not the key to the diagnosis, but rather it is that  
51 the bizarre pattern remains consistently present, despite variations in task  
52 performance, or with distraction. Conversely, the key to recognizing FND is the  
53 variability of the symptoms and signs across different situations over time and within  
54 the physical examination itself. In addition, in the case of functional movement  
55 disorders, the movements are not inherently more bizarre, they just abide by their  
56 own rules, and therefore can also produce recognizable phenotypes. For example,  
57 functional facial dystonia is remarkably identical amongst patients.(7)  
58  
59  
60

Lesson: A bizarre presentation does not equate to a diagnosis of FND.

**Myth 4: Different phenotypes of FND indicate different disorders**

Various presentations of FND are commonly defined as the functional correlates of known neurological signs, e.g. a functional tremor, dystonia, seizures, etc. However, given the shared risk factors among patients with FND, it is more likely that these different phenotypes represent a variable expression of the same underlying disorder (FND), manifesting differently in different individuals. FND is often accompanied by the presence of pain, fatigue, cognitive symptoms, and/or other systemic functional symptoms. Recognizing this broader FND syndrome can be helpful diagnostically in challenging cases, particularly in complex presentations that do not neatly conform to known phenotypes.

Lesson: Functional symptoms are often part of a broader FND syndrome including pain, fatigue, and cognitive symptoms.

**Myth 5. FND symptoms are voluntary**

FND symptoms do arise from the voluntary nervous system which is one reason why concerns about exaggeration or malingering persist in many doctor's minds when thinking about FND. Multiple convergent evidence suggests that feigning is a highly improbable reason for these symptoms including studies showing dysfunction of brain regions involved in movement planning, attention, body monitoring and sense of agency in patients with FND, in contrast with observations in experimental feigning. (14,17,18) Neurophysiological studies, differential recovery in randomised trials, and consistent presentations and comorbidities across cultures and across time are all supportive of FND as a distinct clinical brain disorder, and not a result of voluntary feigning of symptoms.

Lesson: FND symptoms are involuntary; patients are not "putting them on" and feigning is rare.

**Myth 6: There is no role for investigations in the diagnosis of FND**

Having a low threshold for investigations is important because additional neurological disease, such as radiculopathy or demyelination, is such a strong risk factor for FND. Care must be taken in explaining to patients why investigations are being done and to prepare the patient for potentially incidental findings. Laboratory-supported criteria can also be helpful in difficult cases, such as the presence of a cortical *Bereitschaftspotential* in functional myoclonus which suggests use of self-initiated movement pathways. Electromyography can be useful in detecting positive signs such as coherence, distractibility or entrainment in functional tremor, which can then be demonstrated to the patient. Video electroencephalography is helpful in differentiating epileptic and functional (dissociative) seizures. Ideally, the FND diagnosis is presented before the outcome of the investigations to highlight that it is a 'rule-in' diagnosis.

1  
2  
3 Lesson: Investigations can be useful to identify co-morbid neurological conditions,  
4 diagnose phenotypically challenging cases, and reinforce positive signs to the  
5 patient.  
6  
7

8 ***Myth 7: There is less harm in missing a diagnosis of FND than missing another***  
9 ***neurological disease***  
10

11 There is often the perception that it is worse to miss a diagnosis of another  
12 neurological disease than a functional disorder. Published frequencies of  
13 misdiagnosing FND has been consistently around 4% from the early 1970s to 2005,  
14 similar to other neurological and psychiatric disorders, with even lower rates in more  
15 recent studies(8). To put this into perspective, the rate of misdiagnosis of Parkinson's  
16 disease in tertiary centres is approximately 25%, at least upon the initial  
17 contact.(9,10) It is appropriate to be concerned about missing any diagnosis,  
18 particularly in young and disabled patients, and especially when potentially treatable  
19 like FND. Misdiagnosis of patients with FND with diseases such as multiple sclerosis  
20 occurs just as commonly and can lead to just as much harm.(11,12)

21 Lesson: FND is not misdiagnosed more than other conditions. Erroneously  
22 diagnosing FND as another neurological condition can be as harmful as the reverse.  
23  
24  
25  
26

27 ***Myth 8: FND is exclusively a psychological problem caused by psychological factors***  
28  
29

30 For much of the 20<sup>th</sup> century Freud's conversion hypothesis – that FND arises from a  
31 psychological conflict converted into physical symptoms – has been dominant. More  
32 recent work shows that a history of adverse life experience and psychological  
33 comorbidities are commonly seen in this population, but they do not occur in all  
34 patients, and even when present may not be relevant.(13) This is now reflected in  
35 the DSM-5 diagnostic criteria which no longer requires the patient to have a  
36 psychological stressor.(14) In addition psychological/psychiatric comorbidities are  
37 also common in other neurological disorders.(15–17) Just as hypertension and  
38 smoking are risk factors for stroke – disease comorbidity, health anxiety and stress  
39 are risk factors for FND. FND is a complex and heterogeneous disorder, with multiple  
40 potential biological and psychological causes and mechanisms that vary hugely  
41 between patients and which challenges conventional dualistic assumptions about  
42 the brain and mind.  
43  
44

45 Lesson: Psychological factors are one of many possible risk factors for FND and  
46 should not be considered the sole etiological cause.  
47  
48  
49

50 ***Myth 9: The prognosis of FND is usually good***  
51

52 There tends to be a perception that in FND 'nothing is wrong' or that, with  
53 treatment, all patients ought to improve. As in any neurological condition, there is a  
54 spectrum of disease severity treatment response but many studies show the  
55 majority of patients being the same or worse at follow-up.(8) Patients with FND have  
56 levels of disability and impairment in quality of life similar to those with similar  
57 debilitating conditions such as Parkinson's disease or epilepsy(18) which commonly  
58 includes chronic pain, fatigue, cognitive problems and psychological comorbidity.  
59  
60

1  
2  
3 Lesson: Patients with FND are as disabled and have as impaired quality of life as  
4 patients with other neurological conditions. When left untreated, prognosis is  
5 unfavourable for most.  
6  
7

8 ***Myth 10: The treatment of FND is solely referral to a psychologist or psychiatrist***  
9

10 FND patients require individualized treatment, beginning with a transparent and  
11 carefully explained diagnosis. This should avoid simply telling the patient what they  
12 don't have or jumping to conclusions about aetiology. Demonstration of positive  
13 signs to the patient can alter fundamental views about the nature and potential  
14 reversibility of the condition. As in other neurological conditions a multidisciplinary  
15 approach is often required for more complex cases.(19) Physical therapy alone is  
16 effective in some patients with functional movement disorders and has promising  
17 early data from randomised trials, even in patients with long duration  
18 symptoms.(20) There is an evidence base for psychological therapy in functional  
19 (dissociative) seizures(21) which may also be essential when anxiety, mood, or  
20 personality disorders are comorbid. Identifying and triaging patients with dominant  
21 pain or fatigue syndromes is important, as these may need to be the initial focus of  
22 treatment. In some severe cases, the main treatment focus may be on support and  
23 prevention of iatrogenic harm from unnecessary medication or interventions.  
24 Lesson: FND treatment is individualized and involves careful explanation  
25 combinations of physical and psychological rehabilitation.  
26  
27  
28  
29  
30

31 **Conclusion**  
32

33 FND is a common condition, lying at the interface of neurology and psychiatry. A  
34 clear diagnosis delivered in a timely manner can have a strong positive impact on the  
35 patient's symptoms, prognosis, and quality of life. Patients can improve with  
36 individualized, multidisciplinary treatment. Attitudes and practices are changing but  
37 the many misconceptions that surround FND continue to obstruct good medical care  
38 for these patients.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

1. Espay AJ, Aybek S, Carson A, Edwards MJ, Goldstein LH, Hallett M, et al. Current Concepts in Diagnosis and Treatment of Functional Neurological Disorders. *JAMA Neurol.* 2018 Sep 1;75(9):1132–41.
2. Stone J, Reuber M, Carson A. Functional symptoms in neurology: mimics and chameleons. *Pract Neurol.* 2013 Apr 1;13(2):104–13.
3. Kutlubaev MA, Xu Y, Hackett ML, Stone J. Dual diagnosis of epilepsy and psychogenic nonepileptic seizures: Systematic review and meta-analysis of frequency, correlates, and outcomes. *Epilepsy Behav EB.* 2018;89:70–8.
4. Edwards MJ, Bhatia KP. Functional (psychogenic) movement disorders: merging mind and brain. *Lancet Neurol.* 2012 Mar 1;11(3):250–60.
5. Wilcox RA, Winkler S, Lohmann K, Klein C. Whispering dysphonia in an Australian family (DYT4): A clinical and genetic reappraisal. *Mov Disord.* 2011;26(13):2404–8.
6. Carecchio M, Mencacci NE, Iodice A, Pons R, Panteghini C, Zorzi G, et al. ADCY5-related movement disorders: Frequency, disease course and phenotypic variability in a cohort of paediatric patients. *Parkinsonism Relat Disord.* 2017 Aug;41:37–43.
7. Kaski D, Bronstein AM, Edwards MJ, Stone J. Cranial functional (psychogenic) movement disorders. *Lancet Neurol.* 2015 Dec 1;14(12):1196–205.
8. Gelauff J, Stone J. Prognosis of functional neurologic disorders. *Handb Clin Neurol.* 2016;139:523–41.
9. Tolosa E, Wenning G, Poewe W. The diagnosis of Parkinson's disease. *Lancet Neurol.* 2006 Jan 1;5(1):75–86.
10. Stone J, Carson A, Duncan R, Coleman R, Roberts R, Warlow C, et al. Symptoms 'unexplained by organic disease' in 1144 new neurology out-patients: how often does the diagnosis change at follow-up? *Brain.* 2009 Oct 1;132(10):2878–88.
11. Walzl D, Carson AJ, Stone J. The misdiagnosis of functional disorders as other neurological conditions. *J Neurol.* 2019;266(8):2018–26.
12. Coebergh JA, Wren DR, Mumford CJ (2014) 'Undiagnosing' neurological disease: how to do it, and when not to. *Pract Neurol* 14:436–439.
13. Ludwig L, Pasma JA, Nicholson T, Aybek S, David AS, Tuck S, et al. Stressful life events and maltreatment in conversion (functional neurological) disorder: systematic review and meta-analysis of case-control studies. *Lancet Psychiatry.* 2018 Apr;5(4):307–20.



14. American Psychiatric Association, DSM-5 Task Force. Diagnostic and statistical manual of mental disorders: DSM-5™ (5th ed.). 5th ed. Arlington, VA, US: American Psychiatric Publishing, Inc.; 2013.
15. Gendre T, Carle G, Mesrati F, Hubsch C, Mauras T, Roze E, et al. Quality of life in functional movement disorders is as altered as in organic movement disorders. *J Psychosom Res.* 2019 Jan;116:10–6.
16. van der Hoeven RM, Broersma M, Pijnenborg GHM, Koops EA, van Laar T, Stone J, et al. Functional (psychogenic) movement disorders associated with normal scores in psychological questionnaires: A case control study. *J Psychosom Res.* 2015 Sep;79(3):190–4.
17. Kranick S, Ekanayake V, Martinez V, Ameli R, Hallett M, Voon V. Psychopathology and psychogenic movement disorders. *Mov Disord.* 2011;26(10):1844–50.
18. Anderson KE, Gruber-Baldini AL, Vaughan CG, Reich SG, Fishman PS, Weiner WJ, et al. Impact of psychogenic movement disorders versus Parkinson's on disability, quality of life, and psychopathology. *Mov Disord.* 2007 Nov 15;22(0885-3185 (Print)):2204–9.
19. Espay AJ, Aybek S, Carson A, Edwards MJ, Goldstein LH, Hallett M, et al. Current Concepts in Diagnosis and Treatment of Functional Neurological Disorders. *JAMA Neurol.* 2018 Sep 1;75(9):1132.
20. Nielsen G, Buszewicz M, Stevenson F, Hunter R, Holt K, Dudzic M, et al. Randomised feasibility study of physiotherapy for patients with functional motor symptoms. *J Neurol Neurosurg Psychiatry.* 2017 Jun 30;88(6):484–90.
21. Carlson P, Nicholson Perry K. Psychological interventions for psychogenic non-epileptic seizures: A meta-analysis. *Seizure.* 2017;45:142–50.

## Ten myths about Functional Neurological Disorder

Sarah C. Lidstone, MD, PhD, Rui Araújo MD,  
Jon Stone MD, PhD, Bastiaan R. Bloem, MD, PhD

Sarah C. Lidstone, MD, PhD, FRCPC  
Gloria and Morton Shulman Movement Disorders Clinic  
Toronto Western Hospital  
399 Bathurst Street, Toronto, Ontario  
Sarah.lidstone@uhnresearch.ca

Rui Araújo, MD, FEBN  
Department of Neurology, Centro Hospitalar Universitário de São João, E.P.E,  
Department of Clinical Neurosciences and Mental Health, Faculty of Medicine of  
University of Porto  
Porto, Portugal  
rmma22@gmail.com

Jon Stone, MB ChB PhD  
Centre for Clinical Brain Sciences,  
University of Edinburgh,  
Edinburgh, Scotland  
Jon.stone@ed.ac.uk

Bastiaan R. Bloem, MD, PhD, FRCPE (corresponding author)  
Radboud University Medical Centre  
Donders Institute for Brain, Cognition and Behaviour  
Department of Neurology  
Nijmegen, the Netherlands  
bas.bloem@radboudumc.nl

**Word count:** ~~1602590~~583

**Key words:** functional neurological disorder; functional movement disorder; myths;  
diagnosis; treatment; [psychogenic](#); [conversion disorder](#)

**Figures:** 0

**Tables:** 0

**Running title:** Ten myths of FND

1  
2  
3  
4  
5  
6  
7  
8  
9  
10 Functional neurological disorder (FND), is the second most common reason for  
11 referral to a neurologist. Despite being potentially treatable, FND remains  
12 underdiagnosed and associated with high levels of disability and distress. In this  
13 article, we address ten common myths about FND that continue to obstruct the  
14 diagnosis and treatment for these patients, and present them with accompanying  
15 lessons for the clinician.

16 ***Myth 1: FND is a diagnosis of exclusion***

17  
18 The diagnosis of FND should be approached in the same way as other medical  
19 conditions and be diagnosed using positive features, rather than by ruling out other  
20 conditions.(1) Positive signs on examination include Hoover's sign ~~or abductor sign~~  
21 of functional limb weakness, where hip extension weakness returns to normal when  
22 attention is diverted to the contralateral leg. Functional tremor may 'entrain' to the  
23 frequency of imposed rhythmic movements in another body part. Functional  
24 (dissociative) seizures can be recognised by a combination of typical features such as  
25 eyes tightly closed during an episode, long duration, or awareness of generalized  
26 shaking. Thus, the diagnosis of FND should be based primarily on typical features  
27 only seen in this disorder or internal inconsistency of the symptoms and signs.(2)  
28 Lesson: The diagnosis of FND should be "ruled in" based on the presence of positive  
29 signs.

30 ***Myth 2: Patients have either FND or another neurological disorder***

31  
32 The presence of another neurological condition is a powerful risk factor for the  
33 development of FND and they commonly occur together. Physical injury, surgical  
34 procedures and migraine often precipitate functional movements in vulnerable  
35 individuals, and up to 20% of patients with functional (dissociative) seizures also  
36 have epilepsy.(3) Some conditions like Parkinson's disease appear particularly likely  
37 to present with comorbid FND, perhaps because of shared neurobiology, but  
38 symptoms and disability from any neurological condition may be sufficient. For this  
39 reason, there should be a low threshold for investigations.  
40 Lesson: FND commonly co-occurs with other neurological disorders.

41 ***Myth 3: A bizarre presentation indicates FND***

42  
43 FND is frequently equated to bizarre or unrecognizable clinical neurological  
44 presentations, such as complex movement disorders. In fact, several genetic or  
45 acquired movement disorders present with strikingly bizarre phenomenology.(4-6)  
46 In such cases, the "bizarreness" is not the key to the diagnosis, but rather it is that  
47 the bizarre pattern remains consistently present, despite variations in task  
48 performance, or with distraction. Conversely, the key to recognizing FND is the  
49 variability of the symptoms and signs across different situations over time and within  
50 the physical examination itself. In addition, in the case of functional movement  
51 disorders, the movements are not inherently more bizarre, they just abide by their  
52 own rules, and therefore can also produce recognizable phenotypes. For example,  
53 functional facial dystonia is remarkably identical amongst patients.(7)  
54  
55  
56  
57  
58  
59  
60

Lesson: A bizarre presentation does not equate to a diagnosis of FND.

**Myth 4: Different phenotypes of FND indicate different disorders**

Various presentations of FND are commonly defined as the functional correlates of known neurological signs, e.g. a functional tremor, dystonia, seizures, etc. However, given the shared risk factors among patients with FND, it is more likely that these different phenotypes represent a variable expression of the same underlying disorder (FND), manifesting differently in different individuals. FND is often accompanied by the presence of pain, fatigue, cognitive symptoms, and/or other systemic functional symptoms. Recognizing this broader FND syndrome can be helpful diagnostically in challenging cases, particularly in complex presentations that do not neatly conform to known phenotypes.

Lesson: Functional symptoms are often part of a broader FND syndrome including pain, fatigue, and cognitive symptoms.

**Myth 5. FND symptoms are voluntary**

FND symptoms ~~are perceived as involuntary but do arise from the voluntary nervous system which is one reason why~~ concerns about exaggeration or malingering persist in many doctor's minds when thinking about FND. Multiple convergent evidence suggests that feigning is a highly improbable reason for these symptoms including studies showing dysfunction of brain regions involved in movement planning, attention, body monitoring and sense of agency in patients with FND, ~~which in contrast vary with those seen observations~~ in experimental feigning. (14,17,18) Neurophysiological studies, differential recovery in randomised trials, and consistent presentations and comorbidities across cultures and across time are all supportive of FND as a distinct clinical brain disorder, and not a result of voluntary feigning of symptoms.

Lesson: FND symptoms are involuntary; patients are not "putting them on" and feigning is rare.

**Myth 6: There is no role for investigations in the diagnosis of FND**

~~Investigations can be useful in establishing a diagnosis of FND. Having a low threshold to do so for investigations is reasonable because as mentioned, additional neurological disease, such as radiculopathy or demyelination, is such a strong risk factor for FND. Imaging should be done primarily to look for the presence of additional neurological disease, such as radiculopathy or demyelination. Care must be taken in how investigations are framed to the patient, since they do not change the diagnosis of FND in explaining to patients why investigations are being done and to prepare the patient for potentially incidental findings, or in the event that incidental findings are discovered. such as the clinically silent multiple sclerosis or root lesion contributing to the clinical picture. Having a low threshold to do so is reasonable because as mentioned, neurological disease is such a strong risk factor for FND.~~ Laboratory-supported criteria can also be helpful in difficult cases, such as the presence of a cortical *Bereitschaftspotential* in functional myoclonus which

Commented [JS1]: Weve used this phrasing earlier 'another neurological condition' - rather than trying to spearate from disease

suggests use of self-initiated movement pathways. Electromyography can be useful in detecting positive signs such as coherence, distractibility or entrainment in functional tremor, which can then be demonstrated to the patient. Video electroencephalography is helpful in differentiating epileptic and functional (dissociative) seizures. Ideally, the FND diagnosis is presented before the outcome of the investigations to highlight that it is a 'rule-in' diagnosis.

Lesson: Investigations can be useful to identify co-morbid neurological conditions, diagnose phenotypically challenging cases, and reinforce positive signs to the patient.

**Myth 7: There is less harm in missing a diagnosis of FND than missing another neurological disease**

There is often the perception that it is worse to miss a diagnosis of another neurological disease than a functional disorder. Published frequencies of misdiagnosing FND has been consistently around 4% from the early 1970s to 2005, similar to other neurological and psychiatric disorders, with even lower rates in more recent studies(8). To put this into perspective, the rate of misdiagnosis of Parkinson's disease in tertiary centres is approximately 25%, at least upon the initial contact.(9,10) It is appropriate to be concerned about missing any diagnosis, particularly in young and disabled patients, and especially when potentially treatable like FND. Misdiagnosis of patients with FND with diseases such as multiple sclerosis occurs just as commonly and can lead to just as much harm.(11,12)

Lesson: FND is not misdiagnosed more than other conditions. Erroneously diagnosing FND as another neurological condition can be as harmful as the reverse.

**Myth 8: FND is exclusively a psychological problem caused by psychological factors**

For much of the 20<sup>th</sup> century Freud's conversion hypothesis – that FND arises from is a psychological conflict converted into physical symptoms – has been dominant. More recent work shows that a history of adverse life experience and psychological comorbidities are commonly seen in this population, but they do not occur in all patients, and even when present may not be relevant.(132) This is now reflected in the DSM-5 diagnostic criteria which no longer requires the patient to have a psychological stressor.(143) In addition psychological/psychiatric comorbidities are also common in other neurological disorders.(154–176) Just as hypertension and smoking are risk factors for stroke – disease comorbidity, health anxiety and stress are risk factors for FND. FND is a complex and heterogenous neuropsychiatric disorder, best approached from a biopsychosocial perspective with predisposing, precipitating and perpetuating factors with multiple potential biological and psychological causes and mechanisms that vary hugely between patients and which challenges conventional dualistic assumptions about the brain and mind.

Lesson: Psychological factors are one of many possible risk factors for FND and should not be considered the sole etiological cause.

**Myth 9: The prognosis of FND is usually good**

**Commented [JS2]:** Insert Coebergh reference - 1. Coebergh JA, Wren DR, Mumford CJ (2014) 'Undiagnosing' neurological disease: how to do it, and when not to. Pract Neurol 14:436–439.

**Commented [JS3]:** The word neuropsychiatric unfortunately causes a lot of confusion both with patients and neurologists who assume it means psychiatric Like psychosomatic I have no problem with the word in its true meaning, but this is about communicating the right message I think it would be best to leave out I have antibodies to biopsychosocial as well but for different reasons – I think a biopsychosocial formulation should be used for all conditions in my medical practice. When its highlighted specially for FND we are reinforcing a stereotype that we need to use it more in this condition than in others. Also when neurologists see that word they mentally glaze over in my experience!

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

There tends to be a perception that in FND ‘nothing is wrong’ or that, with treatment, all patients ought to improve. As in any neurological condition, there is a spectrum of disease severity treatment response but many studies show the majority of patients being the same or worse at follow-up.(8) Patients with FND have levels of disability and impairment in quality of life similar to those with similar debilitating conditions such as Parkinson’s disease or epilepsy(187) which commonly includes chronic pain, fatigue, cognitive problems and psychological comorbidity. Lesson: Patients with FND are as disabled and have as impaired quality of life as patients with other neurological conditions. When left untreated, prognosis is unfavourable for most.

**Myth 10: The treatment of FND is solely referral to a psychologist or psychiatrist**

FND patients require individualized treatment, beginning with a transparent and carefully explained diagnosis. This should avoid simply telling the patient what they don’t have or jumping to conclusions about aetiology. Demonstration of positive signs to the patient can alter fundamental views about the nature and potential reversibility of the condition. As in other neurological conditions a multidisciplinary approach is often required for more complex cases.(198) Physical therapy alone is effective in some patients with functional movement disorders and has promising early data from randomised trials, even in patients with long duration symptoms.(2019) There is an evidence base for psychological therapy in functional (dissociative) seizures(210) which may also be essential when anxiety, mood, or personality disorders are comorbid. Identifying and triaging patients with dominant pain or fatigue syndromes is important, as these may need to be the initial focus of treatment. In some severe cases, the main treatment focus may be on support and prevention of iatrogenic harm from unnecessary medication or interventions. Lesson: FND treatment is individualized and involves careful explanation combinations of physical and psychological rehabilitation.

**Conclusion**

FND is a common condition, lying at the interface of neurology and psychiatry and is best approached from a biopsychosocial perspective. -neurological condition encountered in any neurology practice. A clear diagnosis delivered in a timely manner can have a strong positive impact on the patient’s symptoms, prognosis, and quality of life. Patients can improve with individualized, multidisciplinary treatment. Attitudes and practices are changing but the many misconceptions that surround FND continue to obstruct good medical care for these patients.

## References

1. Espay AJ, Aybek S, Carson A, Edwards MJ, Goldstein LH, Hallett M, et al. Current Concepts in Diagnosis and Treatment of Functional Neurological Disorders. *JAMA Neurol.* 2018 Sep 1;75(9):1132–41.
2. Stone J, Reuber M, Carson A. Functional symptoms in neurology: mimics and chameleons. *Pract Neurol.* 2013 Apr 1;13(2):104–13.
3. Kutlubaev MA, Xu Y, Hackett ML, Stone J. Dual diagnosis of epilepsy and psychogenic nonepileptic seizures: Systematic review and meta-analysis of frequency, correlates, and outcomes. *Epilepsy Behav EB.* 2018;89:70–8.
4. Edwards MJ, Bhatia KP. Functional (psychogenic) movement disorders: merging mind and brain. *Lancet Neurol.* 2012 Mar 1;11(3):250–60.
5. Wilcox RA, Winkler S, Lohmann K, Klein C. Whispering dysphonia in an Australian family (DYT4): A clinical and genetic reappraisal. *Mov Disord.* 2011;26(13):2404–8.
6. Carecchio M, Mencacci NE, Iodice A, Pons R, Panteghini C, Zorzi G, et al. ADCY5-related movement disorders: Frequency, disease course and phenotypic variability in a cohort of paediatric patients. *Parkinsonism Relat Disord.* 2017 Aug;41:37–43.
7. Kaski D, Bronstein AM, Edwards MJ, Stone J. Cranial functional (psychogenic) movement disorders. *Lancet Neurol.* 2015 Dec 1;14(12):1196–205.
8. Gelauff J, Stone J. Prognosis of functional neurologic disorders. *Handb Clin Neurol.* 2016;139:523–41.
9. Tolosa E, Wenning G, Poewe W. The diagnosis of Parkinson's disease. *Lancet Neurol.* 2006 Jan 1;5(1):75–86.
10. Stone J, Carson A, Duncan R, Coleman R, Roberts R, Warlow C, et al. Symptoms 'unexplained by organic disease' in 1144 new neurology out-patients: how often does the diagnosis change at follow-up? *Brain.* 2009 Oct 1;132(10):2878–88.
11. Walzl D, Carson AJ, Stone J. The misdiagnosis of functional disorders as other neurological conditions. *J Neurol.* 2019;266(8):2018–26.
12. Coebergh JA, Wren DR, Mumford CJ (2014) 'Undiagnosing' neurological disease: how to do it, and when not to. *Pract Neurol* 14:436–439.
13. Ludwig L, Pasman JA, Nicholson T, Aybek S, David AS, Tuck S, et al. Stressful life events and maltreatment in conversion (functional neurological) disorder: systematic review and meta-analysis of case-control studies. *Lancet Psychiatry.* 2018 Apr;5(4):307–20.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10 | **143.** American Psychiatric Association, DSM-5 Task Force. Diagnostic and statistical  
11 | manual of mental disorders: DSM-5™ (5th ed.). 5th ed. Arlington, VA, US:  
12 | American Psychiatric Publishing, Inc.; 2013.
- 13 | **154.** Gendre T, Carle G, Mesrati F, Hubsch C, Mauras T, Roze E, et al. Quality of life in  
14 | functional movement disorders is as altered as in organic movement disorders.  
15 | J Psychosom Res. 2019 Jan;116:10–6.
- 16 | **165.** van der Hoeven RM, Broersma M, Pijnenborg GHM, Koops EA, van Laar T, Stone  
17 | J, et al. Functional (psychogenic) movement disorders associated with normal  
18 | scores in psychological questionnaires: A case control study. J Psychosom Res.  
19 | 2015 Sep;79(3):190–4.
- 20 | **176.** Kranick S, Ekanayake V, Martinez V, Ameli R, Hallett M, Voon V.  
21 | Psychopathology and psychogenic movement disorders. Mov Disord.  
22 | 2011;26(10):1844–50.
- 23 | **187.** Anderson KE, Gruber-Baldini AL, Vaughan CG, Reich SG, Fishman PS, Weiner WJ,  
24 | et al. Impact of psychogenic movement disorders versus Parkinson's on  
25 | disability, quality of life, and psychopathology. Mov Disord. 2007 Nov  
26 | 15;22(0885-3185 (Print)):2204–9.
- 27 | **198.** Espay AJ, Aybek S, Carson A, Edwards MJ, Goldstein LH, Hallett M, et al. Current  
28 | Concepts in Diagnosis and Treatment of Functional Neurological Disorders.  
29 | JAMA Neurol. 2018 Sep 1;75(9):1132.
- 30 | **2019.** Nielsen G, Buszewicz M, Stevenson F, Hunter R, Holt K, Dudzic M, et al.  
31 | Randomised feasibility study of physiotherapy for patients with functional  
32 | motor symptoms. J Neurol Neurosurg Psychiatry. 2017 Jun 30;88(6):484–90.
- 33 | **210.** Carlson P, Nicholson Perry K. Psychological interventions for psychogenic non-  
34 | epileptic seizures: A meta-analysis. Seizure. 2017;45:142–50.

**Commented [LS4]:** New reference 12  
Coebergh JA, Wren DR, Mumford CJ (2014)  
'Undiagnosing' neurological disease: how to do it, and  
when not to. Pract Neurol 14:436–439.



European Journal of Neurology

Title: Ten myths about Functional Neurological Disorder

Manuscript ID: EJoN-20-0591

Authors: Lidstone, Sarah; Araújo, Rui; Stone, Jon; Bloem, Bastiaan

## Response to reviewers

### Reviewer: 1

#### COMMENTS TO AUTHOR(S)

The article is a helpful summary of important myths in an easily readable format suitable for the general neurologist and will hopefully help challenge those myths.

I had some minor comments only:

This sentence does not quite flow easily; explaining Hoover's sign after hip abductor sign (Positive signs on examination include Hoover's sign or abductor sign of functional limb weakness, where hip extension weakness returns to normal when attention is diverted to the contralateral leg.

*This has now been corrected to:*

*Positive signs on examination include Hoover's sign of functional limb weakness, where hip extension weakness returns to normal when attention is diverted to the contralateral leg.*

For this reason, there should be a low threshold for investigations

As below I feel an addition is important; to have low threshold for investigations but explain beforehand that because FND can co-exist investigation findings do not necessarily change management; and it can be helpful to explain investigations (especially MRI) can have incidental findings; at times after discussion of the incidental findings investigations can be postponed to allow clinical monitoring, e.g. in functional cognitive symptoms and mild functional tremor a review after 6-12 months can replace investigations if decision regarding investigations are shared decisions with patients.

the clinically silent multiple sclerosis or root lesion contributing to the clinical picture.

Reviewer: I do not see how clinically silent MS contributes to a clinical picture. That is radiologically isolated syndrome. It can lead to a clinical picture but as above careful conversation before investigation is done can help tremendously in dealing with results of investigations.

*We thank the reviewer for these excellent points. We have changed the section to highlight this:*

*Having a low threshold for investigations is important because additional neurological disease, such as radiculopathy or demyelination, is such a strong risk factor for FND. Care must be taken in explaining to patients why investigations are being done and to prepare the patient for*

1  
2  
3 *potentially incidental findings.*

4  
5  
6 which vary with those seen in experimental feigning.

7 Reviewer: Is it meant to say: contrasts or differs to those seen?

8  
9 *Yes – thank you for picking this up, the sentence has been corrected.*

10  
11  
12 Lesson: FND is not misdiagnosed more than other conditions. Erroneously  
13 diagnosing FND as another neurological condition can be as harmful as the reverse.

14 Reviewer: possible relevant reference to: how to undiagnose neurological disease (coebergh et  
15 al Practical Neurology

16 *We thank the reviewer for this additional reference, which we have now added to the revision.*

17  
18  
19 This sentence does not quite flow:

20 More recent work shows that a history of adverse life experience and psychological  
21 comorbidities are

22 commonly seen in this population, (MISSING: BUT) they do not occur in all patients, and even  
23 when present may not be relevant.(12)

24  
25  
26 *Agreed – this has been added.*

27  
28  
29 Overall however with minor adjustments on how to decide/explain investigations a helpful, easy  
30 to read letter for the general neurologist.

31  
32 *We thank the reviewer for the favorable review and helpful suggestions which have improved the*  
33 *manuscript.*

## 34 35 36 **Reviewer: 2**

### 37 38 COMMENTS TO AUTHOR(S)

39 I have read this article with interest as the topic of FND is of great importance to the clinical  
40 neurosciences. This article is well written and I agree with well over 90% of what is being  
41 conveyed. However, there are some subtle yet very important themes that I do not completely  
42 agree with and I am hopeful the authors can consider in more nuanced fashion:

43  
44  
45  
46 1. Myth 5 (minor comment) - it may be helpful to note that abnormal movements in FND are  
47 perceived as involuntary - however clinical features such as tremor entrainment underscore that  
48 functional abnormal movements engage volitional motor pathways.

49  
50  
51 *We agree with this point and have altered the text as follows*

52 *“FND symptoms are perceived as involuntary but do arise from the voluntary nervous system which is*  
53 *one reason why concerns about exaggeration or malingering may persist in many doctors’ minds when*  
54 *thinking about FND”*

55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 2. Myth 8 (major comment) - is not entirely a myth in my opinion. In some patients, FND is a  
6 condition where psychological factors are the primary drivers of the disorder (some patients will  
7 tell us this themselves!). As examples, we can think about patients whose dissociative seizures  
8 overlap considerably with panic attacks with a heightened tremulousness and a dissociative  
9 component. We can think of others whose functional neurological symptoms are occurring in  
10 the context of traumatic re-experiencing events (flashbacks). I think for Myth 8 it is critically  
11 important to be more nuanced to allow for the importance of neurological and  
12 psychiatric/psychological viewpoints. The equation for developing FND is highly heterogenous.  
13 For some patients, psychological factors are the primary driver of the condition. For others,  
14 medical / neurological factors (TBI and dissociative seizures; peripheral limb injury and  
15 functional dystonia) may play a stronger role. Where the field has gone wrong previously is  
16 forcing onto patients certain etiological models when they are not relevant. I don't know the  
17 cause of FND in any patient, but I do find that patients can find this out for themselves in the  
18 course of treatment. Sometimes when patients improve they make the psychologically rich  
19 formulation for themselves and providers.  
20  
21  
22  
23

24 3. Related to point 2, the authors note that for some patients psychological factors and life  
25 events are not relevant for their clinical picture. Why that can be the case in some, for others life  
26 events are relevant and the correlation between adverse early life events and symptom severity  
27 offers an alternative perspective of importance (i.e. Roelofs et al AJP 2002; Selkirk et al 2008  
28 Epilepsia; Perez, Matin et al 2017 JNNP). In addition to edits to the body of this section, a quick  
29 fix to the Myth 8 statement is to add the word "exclusively" (FND is exclusively a psychological  
30 problem caused by psychological factors).  
31  
32  
33

34 *We thank the reviewer for these important comments. We completely agree this should have been*  
35 *more nuanced and that very often psychological factors are drivers for symptom generation*  
36 *and/or maintenance. Here we were addressing the myth about etiology of FND being exclusively*  
37 *"psychological". The reviewer is correct that we should have added the word 'exclusively' to the*  
38 *title, and this has been done.*  
39  
40

41 *We have also added the following text to the revised manuscript:*

42 "FND is a complex and heterogeneous disorder, with multiple potential biological and  
43 psychological causes and mechanisms that vary hugely between patients, and which challenges  
44 conventional dualistic assumptions about the brain and mind.

45 Lesson: Psychological factors are one of many possible risk factors for FND and should not be  
46 considered as the sole etiological cause."  
47  
48

49 4. Myth 10 could also be more nuanced. Adding in the qualifier "solely" or "exclusively" would  
50 more accurately convey the point I think. In some instances, the primary treatment can be  
51 referral to a psychotherapist with experience in the management of FND. What you are trying to  
52 convey here (which is so important as well) is that the neurologist plays an important role in  
53 treatment, including catalyzing treatment engagement. Also - while I've heard many talk about  
54 how motor FND can improve with physiotherapy alone, this may be true of patients with modest  
55  
56  
57  
58  
59  
60

1  
2  
3 symptom burden, acute onset and/or low psychiatric comorbidities. The converse is much more  
4 familiar to me - physical and occupational therapists raising concerns that  
5 psychiatric/psychological factors are under addressed and impeding treatment  
6  
7

8 *We agree with the reviewer and have now amended the myth title. Unfortunately, the format of*  
9 *this piece requires a simplified discussion of this complex topic. In our experience many patients*  
10 *with motor FMD who improve or partially improve with PT, it is not the PT itself that is the helpful*  
11 *component, rather the education around FND, the disease model, and self-management strategies*  
12 *which are not elements of traditional neurorehabilitation. Thus, the specificity of what is being*  
13 *targeted in physical rehab for FND is not yet well understood.*  
14  
15

16  
17 5. The conclusion: I think that statements such as "FND is a common neurological condition"  
18 unqualified does not send the appropriately nuanced message. Not also pointing out that FND  
19 "is a common (neuro)psychiatric condition as well" disenfranchises our  
20 psychiatry/psychology/social work colleagues who are so needed in the front lines of FND  
21 clinical care (and research). I wonder if the authors can consider rephrasing to note that FND is a  
22 common condition at the interface of neurology and psychiatry.  
23  
24

25 *We agree, note that we had initially targeted this piece to Neurologists, reflecting the word choice*  
26 *here. We take the reviewer's point and do not want to underemphasize the importance of*  
27 *psychiatry, while also asserting the responsibility of the neurologist in daily practice not to over-*  
28 *estimate the role of psychiatric pathology in these patients. It is a fine line and we have attempted*  
29 *to soften this here:*  
30  
31

32 *FND is a common condition, lying at the interface of neurology and psychiatry.*  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60