

β -Adrenergic Relaxation in Mesenteric Resistance Arteries of Spontaneously Hypertensive and Wistar-Kyoto Rats: The Role of Precontraction and Intracellular Ca^{2+}

W. Matthijs Blankesteyn, *Nicolaas J. H. Raat, †Peter H. G. M. Willems, and Theo Thien

*Department of Medicine, Division of General Internal Medicine, University Hospital Nijmegen, and Departments of *Physiology and †Biochemistry, University of Nijmegen, The Netherlands*

Summary: An attenuated β -adrenergic vasodilation of small arteries may help explain the increased peripheral resistance in hypertension. To investigate this, we compared the isoprenaline-induced relaxation of mesenteric resistance arteries of spontaneously hypertensive rats (SHR) and Wistar-Kyoto rats (WKY) using a small vessel myograph. The arteries had similar diameters, but the contractile force induced by cumulative addition of K^+ (10–130 mM) was 1.3-fold higher for the SHR. The β -adrenoceptor-mediated relaxation of arteries, precontracted with 40 mM K^+ , was significantly less in SHR ($41 \pm 3\%$, $n = 11$) than in WKY ($56 \pm 3\%$, $n = 15$, $p = 0.003$), and the pD_2 value for isoprenaline was significantly lower in SHR (7.13 ± 0.09 vs. 7.41 ± 0.07 , $p = 0.02$). In contrast, when precontracted with phenyleph-

rine (PE, α_1 -adrenoceptor agonist, 3–10 μM), isoprenaline relaxation was almost complete in both SHR and WKY, and the pD_2 value for isoprenaline did not differ between strains. Forskolin induced complete relaxation of both precontractions. Because the β -adrenergic relaxation of the mesenteric resistance arteries was attenuated only after K^+ -precontraction, we conclude that alterations in this precontracting mechanism in SHR rather than a defect in the β -adrenoceptor system may provide an explanation for the decreased relaxation in these vessels. Intracellular Ca^{2+} measurements and a review of the literature support this conclusion. **Key Words:** β -Adrenergic relaxation—Mesenteric resistance arteries—Isoprenaline—Forskolin—Spontaneously hypertensive rats—Intracellular Ca^{2+} concentration—Precontraction.

Increased peripheral vascular resistance (PVR) is common in hypertension (1). Because stimulation of β -adrenoceptors on the smooth muscle cells (SMC) of the vascular wall results in vasodilation, a blunted β -adrenergic response may help explain the increased peripheral resistance of the blood vessels in hypertension (2,3). Alterations in the β -adrenoceptor system in hypertension have been investigated quite extensively. For most of these studies, animal models of hypertension, such as the spontaneously hypertensive rat (SHR), have been used. In studies of conduit arteries of SHR, such as aorta and femoral artery, Cohen and Berkowitz (4) and Asano and co-workers (5–10) showed that the β -adrenergic responsiveness was decreased. The main

part of the vascular resistance, however, is generated in the smaller arteries or resistance vessels (11). Therefore, these vessels appear to be more appropriate for investigation of alterations in β -adrenergic responsiveness in hypertension. In the present study, we examined the β -adrenoceptor-mediated relaxation in mesenteric resistance arteries from SHR and Wistar-Kyoto rats (WKY) with the β -adrenoceptor agonist isoprenaline.

The contraction and relaxation of vascular SMC (VSMC) is the result of a change in the intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) or a modulation of the sensitivity of the contractile apparatus to Ca^{2+} (12). The contraction of mesenteric resistance arteries with K^+ has been shown to be mainly the result of

Received March 15, 1995; revision accepted August 28, 1995.
Address correspondence and reprint requests to Professor Dr. Th. Thien at Department of Medicine, Division of General Internal Medicine, University Hospital Nijmegen St. Radboud,

P.O. Box 9101, 6500 HB Nijmegen, The Netherlands.

The present address of Drs. Blankesteyn and Raat is Department of Pharmacology, University of Limburg, Maastricht, The Netherlands.

an increase in $[Ca^{2+}]_i$, whereas phenylephrine (PE, α_1 -adrenoceptor agonist) mainly induces a sensitization to Ca^{2+} (13). To determine the influence of these different mechanisms of contraction on the β -adrenoceptor-mediated relaxation, we contracted the arteries with either K^+ or PE and then exposed them to isoprenaline. The effect of K^+ , PE, isoprenaline and forskolin on the intracellular Ca^{2+} concentration of VSMC muscle cells was investigated by digital imaging microscopy of Fura-2-loaded mesenteric resistance arteries from WKY.

MATERIALS AND METHODS

Materials

Isoprenaline was purchased from Lansberg BV (Uden, The Netherlands), PE from Genfarm BV (Maarssen, The Netherlands), and forskolin and dibutyryl cyclic AMP from Sigma (St. Louis, MO, U.S.A.). Fura-2 AM and pluronic F127 were obtained from Molecular Probes (Eugene, OR, U.S.A.). All other reagents were of analytical quality.

Preparation of the mesenteric arteries

Male SHR and WKY were obtained from Harlan Olac, Bicester, U.K. At the age of 12–18 weeks, the rats were killed by cervical dislocation. The intestine was removed and stored for <24 h at 4°C in phosphate-buffered saline solution (PSS: NaCl 120 mM, KCl 4.7 mM, $CaCl_2$ 2.5 mM, $NaHCO_3$ 25 mM, KH_2PO_4 1.2 mM, $MgCl_2$ 1.17 mM, EDTA 2.7 μ M, and glucose 5.5 mM). Storage of the intestine for \leq 24 h did not affect any of the properties determined in this study. Parts of the small intestine, 7–15 cm from the stomach, were fixed on a wax plate and submerged in PSS. Side branches of the superior mesenteric artery were dissected and mounted on a small vessel myograph as described by Mulvany and Halpern (11). Two stainless-steel wires (diameter 40 μ m) were inserted into the lumen of the resistance arteries. The wires were mounted between an isometric force transducer (Kistler Morse DSC 6, Seattle, WA, U.S.A.) and a displacement device (Mitutoyo, Tokyo, Japan) in a 10-ml organ chamber. The organ chamber was filled with PSS solution maintained at 37°C and continuously aerated with 95% O_2 /5% CO_2 (pH 7.4). After equilibration for 1 h at 37°C, the arteries were set to a normalized internal circumference (L_1), estimated to be 0.9 times the circumference they would maintain if relaxed and exposed to a transmural pressure of 100 mm Hg (L_{100}), a procedure described in detail by Mulvany and Halpern (11). The mean internal diameter at 100 mm Hg was 238 ± 39 μ m ($n = 11$, mean \pm SD) and 246 ± 67 μ m ($n = 15$) for SHR and WKY, respectively, which is characteristic for resistance vessels (11). The arteries mounted in the myograph were 1.8–2.0 mm long.

Contraction and relaxation of resistance arteries

After normalization, the resistance arteries were contracted for 1 min by exchanging the PSS for K-PSS, in which all NaCl was replaced by KCl (total K^+ concentration 125 mM), followed by rinsing twice with PSS and repeated after 15 min. A cumulative concentration-response curve was recorded by increasing the K^+ -concentration stepwise from 10 to 125 mM K^+ . After two

washes in 15 min, the arteries were contracted with a mixture of 30% K-PSS and 70% PSS (total K^+ concentration 40 mM), which gave precontractions of 70–90% of the maximal contractions observed in the concentration-response curves. The relaxing effect of isoprenaline was then studied by a stepwise increase in their concentration (10 nM to 3.2 μ M); after the washing procedure, a concentration-response curve for forskolin was recorded (10 nM to 3.2 μ M). The effect of PE precontraction on the isoprenaline relaxation was studied by contracting the arteries with 3 μ M PE; when no stable contraction was obtained, the PE concentration was increased to 10 μ M. When the response continued to be unstable, the artery was not used further ($n = 3$ in both SHR and WKY). The relaxing effect of isoprenaline was studied as already described. All experiments were executed in the same order because pilot experiments showed no effect of the sequence of precontractions on the parameters determined in this study.

$[Ca^{2+}]_i$ measurements with mesenteric resistance arteries

Mesenteric arteries for the $[Ca^{2+}]_i$ measurements were prepared in the same way as for the myograph studies. After a mesenteric resistance artery was mounted in the myograph, a glass capillary (diameter \sim 100 μ m) was gently inserted in the lumen to stretch the artery. Inserting a glass capillary into the arterial lumen appeared to be crucial because no changes in $[Ca^{2+}]_i$ could be observed without it. The artery was then removed from the myograph and placed in a small plastic tube containing PSS with 10 μ M Fura-2-AM and 1 μ M pluronic F127 (total volume 500 μ l) followed by incubations at room temperature for 2 h in the dark. The incubation solution was then changed, followed by a second incubation for 2 h, in which the artery was transferred to a thermostated (30°C) perfusion chamber (Leiden chamber). The incubation volume was \sim 300 μ l. The tissue was superfused at a flow rate of 1 ml/min with PSS and thoroughly gassed with 95% O_2 /5% CO_2 . The test substances were dissolved in the medium and delivered to the tissue by means of superfusion. To remove nonhydrolyzed Fura-2-AM, the tissue was superfused with PSS for 15 min. After this equilibration period, fluorescence measurements were started as previously described (14). The chamber was placed on the stage of an inverted microscope (Nikon Diaphot). The light from a 100-W xenon lamp was directed through a quartz neutral density filter (Ealing Electro-Optics, Holliston, MA, U.S.A.) with a density \geq 1.5 through excitation filters, mounted in a motor-driven rotating filter wheel, with transmission maxima at 340 and 380 nm (\pm 12 nm) (Ealing Electro-Optics). Dynamic video imaging of part of the artery was performed with a charge-coupled device (CCD) camera (Photonics Sciences) with MagiCal hardware and TARDIS software provided by Joyce Loebel (Dukesway, Team Valley, Gateshead, Tyne & Wear, U.K.) as described in detail by Neylon and colleagues (15). The fluorescence emission ratio at 492 nm, corrected for autofluorescence, was monitored as a measure of $[Ca^{2+}]_i$ after excitation at 340 and 380 nm (16).

Data analysis

The concentration-response curves were analyzed by GraphPad, a nonlinear curve fitting program (GraphPad, Institute for Scientific Information, San Diego, CA,




FIG. 1. Concentration-response curve for the contraction of mesenteric resistance arteries from spontaneously hypertensive rats (SHR, solid line, $n = 11$) and Wistar-Kyoto rats (WKY, dashed line, $n = 15$) to cumulative addition of K^+ . Curves were constructed by fitting the averaged data from the individual experiments. Error bars indicate SE.

U.S.A.). Sensitivities to drugs, expressed as pD_2 values, and maximal effects were calculated on the basis of data from individual arteries. The responses to relaxing agents were expressed as the absolute decrease in tension (mN) and as a percentage of the precontraction. The data are mean \pm SE. The results were analyzed statistically with a two-sample Wilcoxon rank sum test for comparison of the SHR and WKY arteries, and a paired Wilcoxon signed-rank test for comparison of the different precontractions. A p -value < 0.05 (two-sided) was considered statistically significant, insignificant differences are indicated by NS in the text.

RESULTS

Concentration-response curves for K^+

A cumulative increase in K^+ concentration induced a concentration-dependent contraction of the mesenteric resistance arteries, as shown in Fig. 1. The maximal contractile force was 7.1 ± 0.5 mN ($n = 11$) for SHR arteries and 5.5 ± 0.7 mN ($n = 15$) for WKY arteries, a difference which was statistically significant ($p = 0.03$). The pD_2 values for K^+ were 1.33 ± 0.04 and 1.35 ± 0.03 (NS) for SHR and WKY, respectively.

Relaxation of the arteries with isoprenaline

Isoprenaline-mediated relaxations were studied after precontraction of the arteries with 40 mM K^+ ,

a precontraction that was stable for at least 40 min. As shown in Fig. 2A, the precontraction was slightly but insignificantly greater in arteries from SHR. The absolute isoprenaline-mediated relaxation (in mN) was not significantly different between the two groups. However, when expressed as a percentage of the precontraction with 40 mM K^+ (Fig. 2B and Table 1), a significant attenuation of the relaxing effect of isoprenaline was observed in SHR as compared with WKY. The pD_2 value for isoprenaline was significantly lower in the SHR arteries, precontracted with 40 mM K^+ (Table 1).

To study the influence of the nature of precontraction on isoprenaline-mediated relaxations, we also used the α_1 -selective adrenergic agonist phenylephrine (17) as a precontracting agent. The contraction induced by PE (3 – 10 μ M in PSS) was significantly stronger than the response to K^+ in WKY arteries, but significantly weaker in SHR arteries (Table 2). PE induced an oscillating contraction in all arteries tested, a phenomenon which was also observed with oxymethazoline, norepinephrine, and serotonin (data not shown). In some of the preparations (3 of 11 SHR, 3 of 15 WKY) the PE response was transient and did not reach sufficient stability to allow study of the isoprenaline-mediated relaxation. In contrast to K^+ precontraction, the β -adrenoceptor-mediated relaxation was almost complete in PE-precontracted arteries (Fig. 3A). The relative relaxation shown in Fig. 3B was similar in SHR and WKY, although the concentration-response curve for WKY was shifted to the right. This resulted in a tendency toward a higher pD_2 value for isoprenaline in SHR (Table 1), which just failed to reach statistical significance.

Relaxation of the mesenteric resistance arteries with forskolin

Cumulative addition of forskolin induced an almost complete relaxation of mesenteric resistance arteries precontracted with 40 mM K^+ . As shown in Table 1, no marked differences in forskolin-mediated relaxation were evident between SHR and WKY arteries. Forskolin also completely relaxed




FIG. 2. Relaxation of mesenteric resistance arteries, precontracted with 40 mM K^+ , by cumulative addition of isoprenaline. Spontaneously hypertensive rats (SHR, solid line, $n = 11$), Wistar-Kyoto rats (WKY, dashed line, $n = 15$). **A:** Absolute relaxation. **B:** Relaxation expressed as a fraction of the precontraction. C, Force in the absence of isoprenaline. Error bars represent SEM.

TABLE 1. Comparison of the relaxations induced by isoprenaline and forskolin in SHR and WKY mesenteric resistance arteries

Relaxation	SHR	WKY	p-Value
Isoprenaline			
K⁺ precontraction			
Precontraction (mN)	5.7 ± 0.5	5.2 ± 0.3	NS
Maximum relaxation (mN)	2.3 ± 0.2	2.9 ± 0.5	NS
Maximum percentage of relaxation	41 ± 3	56 ± 3	0.003
pD ₂ (M)	7.13 ± 0.09	7.41 ± 0.07	0.02
n	11	15	
PE precontraction			
Precontraction (mN)	4.7 ± 0.6	6.5 ± 1.0	NS
Maximum relaxation (mN)	4.4 ± 0.7	6.2 ± 1.1	NS
Maximum percentage of relaxation	93 ± 2	91 ± 3	NS
pD ₂ (M)	7.05 ± 0.11	6.76 ± 0.11	0.07
n	8	12	
Forskolin			
K⁺ precontraction			
Precontraction (mN)	4.8 ± 0.4	4.3 ± 0.4	NS
Maximum relaxation (mN)	4.0 ± 0.4	3.7 ± 0.8	NS
Maximum percentage of relaxation	81 ± 3	84 ± 2	NS
pD ₂ (M)	7.08 ± 0.12	7.01 ± 0.13	NS
n	11	15	

SHR, spontaneously hypertensive rats; WKY, Wistar-Kyoto rats; PE, phenylephrine.

Data are mean ± SE calculated on the basis of the results of curve fits from individual arteries, with statistical analysis by Wilcoxon rank-sum test for unpaired samples.

arteries precontracted with PE (data not shown).

[Ca²⁺]_i measurements

Measurements of the intracellular Ca²⁺ concentration were performed under conditions similar to those in the myograph. Loading of the arteries with Fura-2 resulted in fluorescent staining of the SMC spanning the arterial lumen. A tracing of a [Ca²⁺]_i measurement is shown in Fig. 4. Superfusion with 40 mM K⁺ induced an increase in the 340/380 ratio from 0.87 ± 0.10 to 0.98 ± 0.12 (n = 4); this increase was rapidly reversible by superfusion with PSS. Superfusion with PE (10 μM) resulted in an average increase in the basal 340/380 ratio of 0.05 in three experiments, whereas in a fourth experiment no increase in ratio was detected. In three experiments, isoprenaline (10 μM) was added to the superfusate while the 340/380 ratio was increased with 40 mM K⁺, but this did not induce a decrease in the

ratio. Similar results were obtained with forskolin (10 μM) and dibutyryl-cyclicAMP (100 μM).

DISCUSSION

In the present study, the vasorelaxant properties of the β-adrenoceptor system of mesenteric resistance arteries of SHR and WKY rats were compared. To our knowledge, this is the first time that this receptor system has been studied in resistance vessels, which contribute strongly to vascular resistance and are therefore of primary interest in hypertension research.

The maximal contraction of the mesenteric resistance arteries, induced by K⁺, was significantly higher in SHR as compared with WKY. This finding is in agreement with results of other studies (8–10) in which strips of conducting arteries, mainly femoral arteries, from SHR and WKY were used.

TABLE 2. Effect of different precontractions on the relaxing properties of isoprenaline

Parameter	K ⁺	PE	p-Value
SHR (n = 8)			
Precontraction (mN)	5.8 ± 0.5	4.7 ± 0.6	0.005
Maximum relaxation (mN)	2.4 ± 0.2	4.4 ± 0.7	0.001
Maximum percentage of relaxation	41 ± 3	93 ± 2	0.007
pD ₂ (M)	7.14 ± 0.07	7.05 ± 0.11	NS
WKY (n = 12)			
Precontraction (mN)	5.6 ± 0.8	6.5 ± 1.0	0.003
Relaxation (mN)	3.1 ± 0.5	6.2 ± 1.1	0.001
Relaxation (%)	56 ± 3	91 ± 3	0.001
pD ₂ (M)	7.37 ± 0.05	6.76 ± 0.11	0.001

Abbreviations as in Table 1.

Statistical analysis by Wilcoxon signed-rank test for paired observations.




FIG. 3. Relaxation of mesenteric resistance arteries precontracted with 3–10 μ M phenylephrine by cumulative addition of isoprenaline. Spontaneously hypertensive rats (SHR, solid line, $n = 8$); Wistar-Kyoto rats (WKY, dashed line, $n = 12$). Details as in legend to Fig. 2.

These results indicate that both in conducting arteries and in resistance arteries of SHR, an abnormality exists, causing an increased K^+ -induced contractile force. Besides an increased ratio of wall to lumen in SHR, Kojima and associates (18) described a malfunction of the sarcoplasmic reticulum (SR) in femoral arteries from SHR as compared with WKY that leads to an increased intracellular Ca^{2+} concentration on contractile stimulus.

In tissues from SHR and WKY precontracted with 40 mM K^+ , cumulative addition of isoprenaline induced a partial relaxation of the mesenteric resistance arteries. In a previous study, we showed that this relaxation was not the result of a release of endothelium-derived relaxing factor (EDRF) and therefore is probably the result of a direct stimula-




FIG. 4. Tracing of a measurement of the intracellular Ca^{2+} concentration in smooth muscle cells of a mesenteric resistance artery from Wistar-Kyoto rats (WKY). The artery was loaded with Fura-2-AM for 4 h. On the Y-axis, the ratio between the emission of the Ca^{2+} -sensitive dye after excitation with light from 340 and 380 nm (340/380 ratio), a measure for the intracellular Ca^{2+} concentration, is shown. Unless indicated otherwise (top) the artery was superfused with phosphate-buffered saline solution 1 ml/min aerated with 95% O_2 /5% CO_2 . Superfusion with K^+ induced an increase in 340/380 ratio ($K^+ 40 = 40$ mM K^+ ; $K^+ 125 = 125$ mM K^+) that could not be reversed with isoprenaline (ISO, 10 μ M) or forskolin (FORSK, 10 μ M). Phenylephrine (10 μ M) induced only a slight increase in 340/380 ratio as compared with K^+ .

tion of β -adrenoceptors on VSMC (19). When expressed as a fraction of the precontraction, the isoprenaline-induced relaxation was significantly less in SHR arteries. Moreover, the pD_2 value for isoprenaline was significantly lower in SHR as compared with WKY. These results are similar to those of Cohen and Berkowitz (4) and Asano and co-workers (5–10) in studies of conducting arteries of SHR, in which the β -adrenoceptor mediated relaxation was expressed as a fraction of the papaverine-induced relaxation.

To investigate the effect of the type of precontraction on the β -adrenergic relaxation, the α_1 -adrenoceptor agonist PE was also used as a precontracting agent. The compound induces contraction through inositol phospholipid metabolism (20), without the α_2 -adrenoceptor-mediated inhibition of cyclicAMP production (21). Because the latter would interfere with the isoprenaline-mediated stimulation of cyclicAMP production, it would hamper the interpretation of the experiments. When the mesenteric resistance arteries from either SHR or WKY were precontracted with PE, isoprenaline could almost completely relax them, and no difference in relaxation was observed between the two groups. Moreover, the pD_2 value for isoprenaline was no longer decreased in SHR. These results indicate that the β -adrenoceptor-mediated relaxation is dependent on the type of precontraction used. These results are in agreement with the data of Abe and Karaki (22), who showed that dibutyryl cyclic AMP could relax rat aorta strips more effectively after α -adrenoceptor-mediated precontraction than after depolarization with K^+ .

The stronger isoprenaline-mediated relaxation of PE-precontracted arteries as compared with K^+ -precontracted arteries may be explained by the mechanisms by which these compounds induce contraction. Jensen and colleagues (13) showed that K^+ induces contraction of mesenteric resistance arteries mainly by increasing the intracellular Ca^{2+} concentration, whereas norepinephrine induces contraction mainly by increasing the sensitivity of the contractile apparatus for Ca^{2+} . Relaxation by cyclicAMP-generating agents can result from a desensitization of the contractile apparatus to Ca^{2+} (12); in the present study, this finding was supported by the measurements of the intracellular Ca^{2+} concentration in WKY. Therefore, the attenuated isoprenaline-mediated relaxation observed in SHR arteries precontracted with K^+ is more likely to result from an increased contraction of these arteries to K^+ than from a defect in the β -adrenergic system. At 40 mM K^+ , the SHR arteries tend to contract more strongly than those of WKY, which can be explained by a malfunction of the SR (18). Due to this malfunction, K^+ induces a stronger increase in the intracellular Ca^{2+} concentration in SHR arteries,

which can less effectively be counteracted by isoprenaline, leading to an attenuated isoprenaline-induced relaxation of SHR arteries with the compound. Although one of the limitations of our study is that the $[Ca^{2+}]_i$ measurements were made only in WKY arteries, we suggest that the attenuated β -adrenergic relaxation in SHR arteries precontracted with K^+ is better explained by an alteration in intracellular Ca^{2+} handling in SHR arteries than by a defect in the β -adrenergic system itself. This finding is in agreement with recent reports concerning β -adrenoceptor density and function in human primary hypertension (23,24), in which no alterations in this receptor system were noted.

The β -adrenergic relaxation of mesenteric resistance arteries from SHR was attenuated after precontraction with K^+ . This finding could not be reproduced with a PE precontraction and can probably be explained by the reported malfunction of the SR in SHR arteries. The β -adrenergic relaxation itself is likely to be unaltered in mesenteric resistance arteries of SHR.

Acknowledgment: We thank Dr. J. F. Rodrigues de Miranda and E. Nagtegaal for helpful discussions.

REFERENCES

1. Folkow B. Physiological aspects of primary hypertension. *Physiol Rev* 1982;62:347-504.
2. Feldman RD. β -Adrenoceptor alterations in hypertension-physiological and molecular correlates. *Can J Physiol Pharmacol* 1987;65:1666-72.
3. Michel MC, Brodde O-E, Insel PA. Peripheral adrenoceptors in hypertension. *Hypertension* 1990;16:107-20.
4. Cohen ML, Berkowitz BA. Decreased vascular relaxation in hypertension. *J Pharmacol Exp Ther* 1976;196:396-406.
5. Asano M, Aoki K, Matsuda T. Reduced beta adrenoceptor interactions of norepinephrine enhance contraction in the femoral artery from spontaneously hypertensive rats. *J Pharmacol Exp Ther* 1982;233:207-14.
6. Fujimoto S, Dohi Y, Aoki K, Asano M, Matsuda T. Diminished β -adrenoceptor-mediated relaxation of arteries from spontaneously hypertensive rats before and during development of hypertension. *Eur J Pharmacol* 1987;136:179-87.
7. Asano M, Masuzawa K, Matsuda T. Evidence for reduced β -adrenoceptor coupling to adenylate cyclase in femoral arteries from spontaneously hypertensive rats. *Br J Pharmacol* 1988;94:73-86.
8. Asano M, Masuzawa K, Matsuda T, Asano T. Reduced function of the stimulatory GTP-binding protein in beta adrenoceptor-adenylate cyclase system of femoral arteries isolated from spontaneously hypertensive rats. *J Pharmacol Exp Ther* 1988;245:709-18.
9. Masuzawa K, Matsuda T, Asano M. Decreased arterial responsiveness to multiple cyclic AMP-generating receptor agonists in spontaneously hypertensive rats. *Br J Pharmacol* 1989;96:227-35.
10. Asano M, Masuzawa K, Matsuda T, Asano T. Decreased responsiveness to β -adrenoceptor agonists in arterial strips from spontaneously hypertensive rats is not associated with alterations in β -adrenoceptors. *J Hypertens* 1991;9:607-13.
11. Mulvany MJ, Halpern W. Contractile properties of small arterial resistance vessels in spontaneously hypertensive and normotensive rats. *Circ Res* 1977;41:19-26.
12. Rembold CM. Regulation of contraction and relaxation in arterial smooth muscle. *Hypertension* 1992;20:129-37.
13. Jensen PE, Mulvany MJ, Aalkjær C. Endogenous and exogenous agonist-induced changes in the coupling between $[Ca^{2+}]_i$ and force in rat resistance arteries. *Pflugers Arch* 1992;420:536-43.
14. Willems PHGM, van Emst-de Vries SE, van Os CH, de Pont JJHMM. Dose-dependent recruitment of pancreatic acinar cells during receptor-mediated calcium mobilization. *Cell Calcium* 1993;14:145-59.
15. Neylon CB, Hoyland J, Mason WT, Irvine RF. Spatial dynamics of intracellular calcium in agonist-stimulated vascular smooth muscle cells. *Am J Physiol* 1990;259:C675-86.
16. Nitschke R, Fröbe U, Greger R. Antidiuretic hormone acts via V_1 receptors on intracellular calcium in the isolated perfused rabbit cortical thick ascending limb. *Pflugers Arch* 1991;417:622-32.
17. Van Zwieten PA. Antihypertensive drugs interacting with α - and β -adrenoceptors. A review of basic pharmacology. *Drugs* 1988;35(suppl 6):6-19.
18. Kojima M, Aoki K, Asano M, Fujimoto S, Matsuda T. Malfunction of arterial sarcoplasmic reticulum leading to faster and greater contraction induced by high-potassium depolarization in young spontaneously hypertensive rats. *J Hypertens* 1991;9:783-8.
19. Blankesteijn WM, Thien Th. Effect of N^G -monomethyl-L-arginine on the β -adrenoceptor-mediated relaxation of rat mesenteric resistance arteries. *Life Sci* 1993;52:PL135-9.
20. Minneman KP. α_1 -Adrenoceptor subtypes, inositol phosphates and sources of cell Ca^{2+} . *Pharmacol Rev* 1988;40:87-115.
21. Timmermans PBMWM. α -Adrenoceptors. In: Williams M, Glennon RA, Timmermans PBMWM, eds. *Receptor pharmacology and function*. New York, Marcel Dekker, 1989: 173-205.
22. Abe A, Karaki H. Mechanisms underlying the inhibitory effect of dibutyryl cyclic AMP in vascular smooth muscle. *Eur J Pharmacol* 1992;211:305-11.
23. Blankesteijn WM, Graafsma SJ, Van Tits LJH, Hectors MPC, Thien Th. Adrenoceptors on blood cells in patients with primary hypertension: correlations with blood pressure and some related variables. *J Hypertens* 1993;11:995-1002.
24. Blankesteijn WM, Willems PHGM, Graafsma SJ, de Pont JJHMM, Thien Th. Forskolin stimulated adenylyl cyclase activity is decreased in primary hypertension. *J Cardiovasc Pharmacol* 1993;22:695-701.