

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/24181>

Please be advised that this information was generated on 2019-02-21 and may be subject to change.

# Acquired ichthyosiform erythroderma and sarcoidosis

Annette G. Feind-Koopmans, MD, George P. H. Lucker, MD, and Peter C. M. van de Kerkhof, MD, PhD *Nijmegen, The Netherlands*

We describe a patient with ichthyosiform erythroderma as a manifestation of sarcoidosis. This is the first report of the simultaneous occurrence of erythroderma and ichthyosis in sarcoidosis. (*J Am Acad Dermatol* 1996;35:826-8.)

Cutaneous manifestations occur in 10% to 35% of the patients with systemic sarcoidosis.<sup>1</sup> The most common of these manifestations are papular, nodular, plaque lesions and involvement of scars.<sup>2</sup> Rare cutaneous expressions of sarcoidosis are erythroderma<sup>3,4</sup> and acquired ichthyosis.<sup>5-8</sup> Erythroderma with a fine, silvery and lamellar scaling has also been described.<sup>9-10</sup>

## CASE REPORT

A 60-year-old man had an erythroderma. Two years previously he had complained of malaise and weight loss (10 kg in 1 year). Examination revealed generalized lymphadenopathy and hepatosplenomegaly. His white blood cell count was  $17.5 \times 10^9/L$ ; differential determinations were neutrophils, 38%; lymphocytes, 24%; monocytes, 3%; and eosinophils, 35%. Examination of liver and lymph node biopsy specimens revealed epithelioid cell granulomas with extensive eosinophilia. Low-grade lymphoma, the hypereosinophilic syndrome, and an unspecified viral infection were considered as diagnoses. Two years later hypercalcemia, increased urinary calcium excretion, progressive hepatosplenomegaly, and a non-pruritic generalized eruption developed.

Examination revealed erythroderma on the extensor surfaces of the extremities (Fig. 1). On diascopy, apple jelly lesions were observed. The extensor aspects of the extremities were covered with large lamellar scales (Fig. 2). The flexures, palms and soles, were relatively free of scales. The patient also had enlarged, painless lymph




Fig. 1. Generalized erythroderma.



This article is made possible through an educational grant from the Dermatological Division, Ortho Pharmaceutical Corporation.

From the Department of Dermatology, University Hospital Nijmegen, The Netherlands.

Reprint requests: Dr. Annette Feind-Koopmans, Department of Dermatology, University Hospital Nijmegen, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands.

Copyright © 1996 by the American Academy of Dermatology, Inc. 0190-9622/96 \$5.00 + 0 16/4/75749




Fig. 2. Lamellar scaling and erythema of leg.



Fig. 3. Biopsy specimen showing dense infiltrate of epithelioid cells, mononuclear cells, and eosinophils. (Hematoxylin-eosin stain;  $\times 400$ .)

nodes of the head and neck and axillary and inguinal regions. The liver and spleen had a normal consistency and were palpated 12 cm and 6 cm, respectively, below the costal margin.

Several biopsy specimens revealed multiple noncaseating epithelioid cell granulomas surrounded by a mixed infiltrate of lymphocytes and eosinophils (Fig. 3). The epidermis was slightly acanthotic, without a granular layer and with hyperkeratosis, focal acanthosis, and focal parakeratosis.<sup>1</sup>

Laboratory studies revealed a white blood cell count of  $17.5 \times 10^9/L$ , 35% eosinophils, a platelet count of  $106 \times 10^9/L$ , and serum levels of creatinine, 200  $\mu\text{mol/L}$ ; calcium, 3.88  $\text{mmol/L}$ ; alkaline phosphatase, 315  $\text{U/L}$ ; and angiotensin-converting enzyme, 29  $\text{U/ml}$  (normal, less than 20  $\text{U/L}$ ). Serum levels of parathyroid hormone and vitamin  $\text{D}_3$  were normal. Results of serum protein electrophoresis were normal. No abnormalities of the

lungs or eyes were found. A gallium scan revealed no skeletal abnormalities, and the results of the Mantoux test were negative.

The results of clinical, histologic, and laboratory investigations indicated that the patient had systemic sarcoidosis with an acquired ichthyosiform erythroderma.

The patient was treated with oral prednisone and topical betamethasone valerate cream. After 6 months of treatment the hepatomegaly, splenomegaly, and lymphadenopathy had improved markedly, and erythroderma had faded. Apart from a generalized fine, silvery scaling, the skin had a normal appearance.

## DISCUSSION

Acquired ichthyosis and erythroderma are rare cutaneous manifestations of sarcoidosis.<sup>4-8</sup> Acquired ichthyosis usually occurs as diffuse, marked scaling

overlying erythematous papules and plaques on the legs.<sup>5,11</sup> In one patient a generalized hyperpigmented ichthyosiform scaling was present.<sup>6</sup> Although in most patients the distribution and the histologic picture are compatible with ichthyosis vulgaris, it should be emphasized that in all cases the ichthyosis was acquired.<sup>11</sup> However, the absence of the stratum granulosum in the sarcoid ichthyosis, as in the report of Griffith et al.<sup>5</sup> and in the present case, is inconsistent with ichthyosis vulgaris. In the present case, parakeratosis was present. Lamellar ichthyosis is usually characterized by hyperkeratosis,<sup>12</sup> but parakeratosis has been described in some forms of lamellar ichthyosis.<sup>13</sup>

In the previously reported cases, both erythroderma and acquired ichthyosis were associated with systemic involvement.<sup>14-16</sup> In our patient the eosinophilia and lack of pulmonary involvement were unusual findings.

#### REFERENCES

1. Kerdel FA, Moschella SL. Sarcoidosis: an updated review. *J Am Acad Dermatol* 1984;11:1-19.
2. Veien NK, Stahl D, Brodthagen H. Cutaneous sarcoidosis in Caucasians. *J Am Acad Dermatol* 1987;16:534-40.
3. Mittag H, Rupec M, Kalbfleisch H. Zur Frage der erythrodermischen Sarkoidose. *Z Hautkr Geschlechtskr* 1985;61:673-87.
4. Kenneth E, Greer L, Harman JR, et al. Unusual cutaneous manifestations of sarcoidosis. *South Med J* 1977;70:666-9.
5. Griffith CEM, Leonard JN, Walker MM. Acquired ichthyosis and sarcoidosis. *Clin Exp Dermatol* 1986;11:296-8.
6. Banse-Kupin L, Pelachyk M. Ichthyosiform sarcoidosis. *J Am Acad Dermatol* 1987;17:616-20.
7. Kelley AP. Ichthyosiforme sarcoid. *Arch Dermatol* 1978;114:1551-2.
8. Matsuoka LY, Levine M, Glasser S, et al. Ichthyosiform sarcoid. *Cutis* 1980;25:188-9.
9. Wigley JEM, Musso LA. A case of sarcoidosis with erythrodermic lesions. Treatment with calciferol. *Br J Dermatol* 1951;63:398.
10. Lever WF, Freiman DG. Sarcoid. *Arch Dermatol Syph* 1948;57:629.
11. Kauh YC, Goody HE, Luscombe HA. Ichthyosiforme sarcoidosis. *Arch Dermatol* 1978;114:100-1.
12. Lever WF, Schaumburg-Lever. *Histopathology of the skin*. 7th ed. Philadelphia: WB Saunders, 1990:67-8.
13. Traup H. *The ichthyosis. A guide to dermal diagnosis, genetic counseling and therapy*. Berlin: Springer-Verlag, 1989:119-22.
14. Samtsov AV. Review of cutaneous sarcoidosis. *Int J Dermatol* 1992;31:385-93.
15. Elgart ML. Cutaneous sarcoidosis: definitions and types of lesions. *Clin Dermatol* 1986;4:35-45.
16. Hanno R, Needleman A, Eiferman RA. Cutaneous sarcoid granulomas and the development of systemic sarcoidosis. *Arch Dermatol* 1981;117:203-7.