

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/26252>

Please be advised that this information was generated on 2019-04-26 and may be subject to change.



ELSEVIER

6 November 1997

PHYSICS LETTERS B

Physics Letters B 413 (1997) 159–166

Hard-photon production at $\sqrt{s} = 161$ and 172 GeV at LEP

L3 Collaboration

M. Acciarri^{ac}, O. Adriani^r, M. Aguilar-Benitez^{ab}, S. Ahlen^l, J. Alcaraz^{ab},
G. Alemanni^x, J. Allaby^s, A. Aloisio^{ae}, G. Alverson^m, M.G. Alviggi^{ae},
G. Ambrosi^u, H. Anderhub^{ay}, V.P. Andreev^{g,an}, T. Angelescuⁿ, F. Anselmo^j,
A. Arefiev^{ad}, T. Azemoon^c, T. Aziz^k, P. Bagnaia^{am}, L. Baksay^{at}, S. Banerjee^k,
Sw. Banerjee^k, K. Banicz^{av}, A. Barczyk^{ay,aw}, R. Barillère^s, L. Barone^{am},
P. Bartalini^{aj}, A. Baschirotto^{ac}, M. Basile^j, R. Battiston^{aj}, A. Bay^x, F. Becattini^r,
U. Becker^q, F. Behner^{ay}, J. Berdugo^{ab}, P. Berges^q, B. Bertucci^{aj}, B.L. Betev^{ay},
S. Bhattacharya^k, M. Biasini^s, A. Biland^{ay}, G.M. Bilei^{aj}, J.J. Blaising^d,
S.C. Blyth^{ak}, G.J. Bobbink^b, R. Bock^a, A. Böhm^a, L. Boldizar^o, B. Borgia^{am},
D. Bourilkov^{ay}, M. Bourquin^u, S. Braccini^u, J.G. Branson^{ap}, V. Brigljevic^{ay},
I.C. Brock^{ak}, A. Buffini^r, A. Buijs^{au}, J.D. Burger^q, W.J. Burger^u, J. Busenitz^{at},
A. Button^c, X.D. Cai^q, M. Campanelli^{ay}, M. Capell^q, G. Cara Romeo^j,
G. Carlino^{ae}, A.M. Cartacci^r, J. Casaus^{ab}, G. Castellini^r, F. Cavallari^{am},
N. Cavallo^{ae}, C. Cecchi^u, M. Cerrada^{ab}, F. Cesaroni^y, M. Chamizo^{ab},
Y.H. Chang^{ba}, U.K. Chaturvedi^t, S.V. Chekanov^{ag}, M. Chemarin^{aa}, A. Chen^{ba},
G. Chen^h, G.M. Chen^h, H.F. Chen^v, H.S. Chen^h, X. Chereau^d, G. Chiefari^{ae},
C.Y. Chien^e, L. Cifarelli^{ao}, F. Cindolo^j, C. Civinini^r, I. Clare^q, R. Clare^q,
H.O. Cohn^{ah}, G. Coignet^d, A.P. Colijn^b, N. Colino^{ab}, V. Commichau^a,
S. Costantiniⁱ, F. Cotorobaiⁿ, B. de la Cruz^{ab}, A. Csilling^o, T.S. Dai^q,
R. D'Alessandro^r, R. de Asmundis^{ae}, A. Degré^d, K. Deiters^{aw}, D. della Volpe^{ae},
P. Denes^{al}, F. DeNotaristefani^{am}, D. DiBitonto^{at}, M. Diemoz^{am},
D. van Dierendonck^b, F. Di Lodovico^{ay}, C. Dionisi^{am}, M. Dittmar^{ay},
A. Dominguez^{ap}, A. Doria^{ae}, M.T. Dova^{t,l}, D. Duchesneau^d, P. Duinker^b,
I. Duran^{aq}, S. Dutta^k, S. Easo^{aj}, Yu. Efremenko^{ah}, H. El Mamouni^{aa}, A. Engler^{ak},
F.J. Eppling^q, F.C. Erné^b, J.P. Ernenwein^{aa}, P. Extermann^u, M. Fabre^{aw},
R. Faccini^{am}, S. Falciano^{am}, A. Favara^r, J. Fay^{aa}, O. Fedin^{an}, M. Felcini^{ay},
B. Fenyi^{at}, T. Ferguson^{ak}, F. Ferroni^{am}, H. Fesefeldt^a, E. Fiandrini^{aj}, J.H. Field^u,

F. Filthaut ^{ak}, P.H. Fisher ^q, I. Fisk ^{ap}, G. Forconi ^q, L. Fredj ^u, K. Freudenreich ^{ay},
 C. Furetta ^{ac}, Yu. Galaktionov ^{ad,q}, S.N. Ganguli ^k, P. Garcia-Abia ^{ax}, S.S. Gau ^m,
 S. Gentile ^{am}, N. Gheordanescu ⁿ, S. Giagu ^{am}, S. Goldfarb ^x, J. Goldstein ^l,
 Z.F. Gong ^v, A. Gougas ^e, G. Gratta ^{ai}, M.W. Gruenewald ⁱ, V.K. Gupta ^{al}, A. Gurtu ^k,
 L.J. Gutay ^{av}, B. Hartmann ^a, A. Hasan ^{af}, D. Hatzifotiadou ^j, T. Hebbeker ⁱ,
 A. Hervé ^s, W.C. van Hoek ^{ag}, H. Hofer ^{ay}, S.J. Hong ^{as}, H. Hoorani ^{ak}, S.R. Hou ^{ba},
 G. Hu ^e, V. Innocente ^s, K. Jenkes ^a, B.N. Jin ^h, L.W. Jones ^c, P. de Jong ^s,
 I. Josa-Mutuberria ^{ab}, A. Kasser ^x, R.A. Khan ^t, D. Kamrad ^{ax}, Yu. Kamyshkov ^{ah},
 J.S. Kapustinsky ^z, Y. Karyotakis ^d, M. Kaur ^{t,2}, M.N. Kienzle-Focacci ^u, D. Kim ^{am},
 D.H. Kim ^{as}, J.K. Kim ^{as}, S.C. Kim ^{as}, Y.G. Kim ^{as}, W.W. Kinnison ^z, A. Kirkby ^{ai},
 D. Kirkby ^{ai}, J. Kirkby ^s, D. Kiss ^o, W. Kittel ^{ag}, A. Klimentov ^{q,ad}, A.C. König ^{ag},
 A. Kopp ^{ax}, I. Korolko ^{ad}, V. Koutsenko ^{q,ad}, R.W. Kraemer ^{ak}, W. Krenz ^a,
 A. Kunin ^{q,ad}, P. Ladron de Guevara ^{ab}, I. Laktineh ^{aa}, G. Landi ^r, C. Lapoint ^q,
 K. Lassila-Perini ^{ay}, P. Laurikainen ^w, M. Lebeau ^s, A. Lebedev ^q, P. Lebrun ^{aa},
 P. Lecomte ^{ay}, P. Lecoq ^s, P. Le Coultre ^{ay}, J.M. Le Goff ^s, R. Leiste ^{ax},
 E. Leonardi ^{am}, P. Levchenko ^{an}, C. Li ^v, C.H. Lin ^{ba}, W.T. Lin ^{ba}, F.L. Linde ^{b,s},
 L. Lista ^{ae}, Z.A. Liu ^h, W. Lohmann ^{ax}, E. Longo ^{am}, W. Lu ^{ai}, Y.S. Lu ^h,
 K. Lübelmeyer ^a, C. Luci ^{am}, D. Luckey ^q, L. Luminari ^{am}, W. Lustermann ^{aw},
 W.G. Ma ^v, M. Maity ^k, G. Majumder ^k, L. Malgeri ^{am}, A. Malinin ^{ad}, C. Mañá ^{ab},
 D. Mangeol ^{ag}, S. Mangla ^k, P. Marchesini ^{ay}, A. Marin ^l, J.P. Martin ^{aa},
 F. Marzano ^{am}, G.G.G. Massaro ^b, D. McNally ^s, R.R. McNeil ^g, S. Mele ^{ae},
 L. Merola ^{ae}, M. Meschini ^r, W.J. Metzger ^{ag}, M. von der Mey ^a, Y. Mi ^x, A. Mihul ⁿ,
 A.J.W. van Mil ^{ag}, G. Mirabelli ^{am}, J. Mnich ^s, P. Molnar ⁱ, B. Monteleoni ^r,
 R. Moore ^c, S. Morganti ^{am}, T. Moulík ^k, R. Mount ^{ai}, S. Müller ^a, F. Muheim ^u,
 A.J.M. Muijs ^b, S. Nahn ^q, M. Napolitano ^{ae}, F. Nessi-Tedaldi ^{ay}, H. Newman ^{ai},
 T. Niessen ^a, A. Nippe ^a, A. Nisati ^{am}, H. Nowak ^{ax}, Y.D. Oh ^{as}, H. Opitz ^a,
 G. Organtini ^{am}, R. Ostonen ^w, C. Palomares ^{ab}, D. Pandoulas ^a, S. Paoletti ^{am},
 P. Paolucci ^{ae}, H.K. Park ^{ak}, I.H. Park ^{as}, G. Pascale ^{am}, G. Passaleva ^r, S. Patricelli ^{ae},
 T. Paul ^m, M. Pauluzzi ^{aj}, C. Paus ^a, F. Pauss ^{ay}, D. Peach ^s, Y.J. Pei ^a, S. Pensotti ^{ac},
 D. Perret-Gallix ^d, B. Petersen ^{ag}, S. Petrak ⁱ, A. Pevsner ^e, D. Piccolo ^{ae},
 M. Pieri ^r, J.C. Pinto ^{ak}, P.A. Piroué ^{al}, E. Pistoiesi ^{ac}, V. Plyaskin ^{ad}, M. Pohl ^{ay},
 V. Pojidaev ^{ad,r}, H. Postema ^q, N. Produit ^u, D. Prokofiev ^{an}, G. Rahal-Callot ^{ay},
 N. Raja ^k, P.G. Rancoita ^{ac}, M. Rattaggi ^{ac}, G. Raven ^{ap}, P. Razis ^{af}, K. Read ^{ah},
 D. Ren ^{ay}, M. Rescigno ^{am}, S. Reucroft ^m, T. van Rhee ^{au}, S. Riemann ^{ax}, K. Riles ^c,
 A. Robohm ^{ay}, J. Rodin ^q, B.P. Roe ^c, L. Romero ^{ab}, S. Rosier-Lees ^d, Ph. Rosselet ^x,
 W. van Rossum ^{au}, S. Roth ^a, J.A. Rubio ^s, D. Ruschmeier ⁱ, H. Rykaczewski ^{ay},
 J. Salicio ^s, E. Sanchez ^{ab}, M.P. Sanders ^{ag}, M.E. Sarakinos ^w, S. Sarkar ^k,

M. Sassowsky ^a, C. Schäfer ^a, V. Schegelsky ^{an}, S. Schmidt-Kaerst ^a, D. Schmitz ^a,
 P. Schmitz ^a, N. Scholz ^{ay}, H. Schopper ^{az}, D.J. Schotanus ^{ag}, J. Schwenke ^a,
 G. Schwering ^a, C. Sciacca ^{ae}, D. Sciarrino ^u, L. Servoli ^{aj}, S. Shevchenko ^{ai},
 N. Shivarov ^{ar}, V. Shoutko ^{ad}, J. Shukla ^z, E. Shumilov ^{ad}, A. Shvorob ^{ai},
 T. Siedenburger ^a, D. Son ^{as}, A. Sopczak ^{ax}, B. Smith ^q, P. Spillantini ^r, M. Steuer ^q,
 D.P. Stickland ^{al}, A. Stone ^g, H. Stone ^{al}, B. Stoyanov ^{ar}, A. Straessner ^a,
 K. Strauch ^p, K. Sudhakar ^k, G. Sultanov ^l, L.Z. Sun ^v, G.F. Susinno ^u, H. Suter ^{ay},
 J.D. Swain ^t, X.W. Tang ^h, L. Tauscher ^f, L. Taylor ^m, Samuel C.C. Ting ^q,
 S.M. Ting ^q, M. Tonutti ^a, S.C. Tonwar ^k, J. Tóth ^o, C. Tully ^{al}, H. Tuchscherer ^{at},
 K.L. Tung ^h, Y. Uchida ^q, J. Ulbricht ^{ay}, U. Uwer ^s, E. Valente ^{am},
 R.T. Van de Walle ^{ag}, G. Vesztegombi ^o, I. Vetlitsky ^{ad}, G. Viertel ^{ay},
 M. Vivargent ^d, R. Völkert ^{ax}, H. Vogel ^{ak}, H. Vogt ^{ax}, I. Vorobiev ^{ad},
 A.A. Vorobyov ^{an}, A. Vorvolakos ^{af}, M. Wadhwa ^f, W. Wallraff ^a, J.C. Wang ^q,
 X.L. Wang ^v, Z.M. Wang ^v, A. Weber ^a, F. Wittgenstein ^s, S.X. Wu ^t, S. Wynhoff ^a,
 J. Xu ^l, Z.Z. Xu ^v, B.Z. Yang ^v, C.G. Yang ^h, X.Y. Yao ^h, J.B. Ye ^v, S.C. Yeh ^{ba},
 J.M. You ^{ak}, An. Zalite ^{an}, Yu. Zalite ^{an}, P. Zemp ^{ay}, Y. Zeng ^a, Z. Zhang ^h,
 Z.P. Zhang ^v, B. Zhou ^l, G.Y. Zhu ^h, R.Y. Zhu ^{ai}, A. Zichichi ^{j,s,t}, F. Ziegler ^{ax}

^a I. Physikalisches Institut, RWTH, D-52056 Aachen, FRG ³

III. Physikalisches Institut, RWTH, D-52056 Aachen, FRG ³

^b National Institute for High Energy Physics, NIKHEF, and University of Amsterdam, NL-1009 DB Amsterdam, The Netherlands

^c University of Michigan, Ann Arbor, MI 48109, USA

^d Laboratoire d'Annecy-le-Vieux de Physique des Particules, LAPP, IN2P3-CNRS, BP 110, F-74941 Annecy-le-Vieux CEDEX, France

^e Johns Hopkins University, Baltimore, MD 21218, USA

^f Institute of Physics, University of Basel, CH-4056 Basel, Switzerland

^g Louisiana State University, Baton Rouge, LA 70803, USA

^h Institute of High Energy Physics, IHEP, 100039 Beijing, China ⁴

ⁱ Humboldt University, D-10099 Berlin, FRG ³

^j University of Bologna and INFN-Sezione di Bologna, I-40126 Bologna, Italy

^k Tata Institute of Fundamental Research, Bombay 400 005, India

^l Boston University, Boston, MA 02215, USA

^m Northeastern University, Boston, MA 02115, USA

ⁿ Institute of Atomic Physics and University of Bucharest, R-76900 Bucharest, Romania

^o Central Research Institute for Physics of the Hungarian Academy of Sciences, H-1525 Budapest 114, Hungary ⁵

^p Harvard University, Cambridge, MA 02139, USA

^q Massachusetts Institute of Technology, Cambridge, MA 02139, USA

^r INFN Sezione di Firenze and University of Florence, I-50125 Florence, Italy

^s European Laboratory for Particle Physics, CERN, CH-1211 Geneva 23, Switzerland

^t World Laboratory, FBLJA Project, CH-1211 Geneva 23, Switzerland

^u University of Geneva, CH-1211 Geneva 4, Switzerland

^v Chinese University of Science and Technology, USTC, Hefei, Anhui 230 029, China ⁴

^w SEFT, Research Institute for High Energy Physics, P.O. Box 9, SF-00014 Helsinki, Finland

^x University of Lausanne, CH-1015 Lausanne, Switzerland

^y INFN-Sezione di Lecce and Università Degli Studi di Lecce, I-73100 Lecce, Italy

^z Los Alamos National Laboratory, Los Alamos, NM 87544, USA

^{aa} Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, F-69622 Villeurbanne, France

^{ab} Centro de Investigaciones Energeticas, Medioambientales y Tecnológicas, CIEMAT, E-28040 Madrid, Spain ⁶

^{ac} INFN-Sezione di Milano, I-20133 Milan, Italy

- ^{ad} Institute of Theoretical and Experimental Physics, ITEP, Moscow, Russia
^{ae} INFN-Sezione di Napoli and University of Naples, I-80125 Naples, Italy
^{af} Department of Natural Sciences, University of Cyprus, Nicosia, Cyprus
^{ag} University of Nijmegen and NIKHEF, NL-6525 ED Nijmegen, The Netherlands
^{ah} Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA
^{ai} California Institute of Technology, Pasadena, CA 91125, USA
^{aj} INFN-Sezione di Perugia and Università Degli Studi di Perugia, I-06100 Perugia, Italy
^{ak} Carnegie Mellon University, Pittsburgh, PA 15213, USA
^{al} Princeton University, Princeton, NJ 08544, USA
^{am} INFN-Sezione di Roma and University of Rome, "La Sapienza", I-00185 Rome, Italy
^{an} Nuclear Physics Institute, St. Petersburg, Russia
^{ao} University and INFN, Salerno, I-84100 Salerno, Italy
^{ap} University of California, San Diego, CA 92093, USA
^{aq} Dept. de Física de Partículas Elementales, Univ. de Santiago, E-15706 Santiago de Compostela, Spain
^{ar} Bulgarian Academy of Sciences, Central Lab. of Mechatronics and Instrumentation, BU-1113 Sofia, Bulgaria
^{as} Center for High Energy Physics, Korea Adv. Inst. of Sciences and Technology, 305-701 Taejon, South Korea
^{at} University of Alabama, Tuscaloosa, AL 35486, USA
^{au} Utrecht University and NIKHEF, NL-3584 CB Utrecht, The Netherlands
^{av} Purdue University, West Lafayette, IN 47907, USA
^{aw} Paul Scherrer Institut, PSI, CH-5232 Villigen, Switzerland
^{ax} DESY-Institut für Hochenergiephysik, D-15738 Zeuthen, FRG
^{ay} Eidgenössische Technische Hochschule, ETH Zürich, CH-8093 Zürich, Switzerland
^{az} University of Hamburg, D-22761 Hamburg, FRG
^{ba} High Energy Physics Group, Taiwan, ROC

Received 1 July 1997

Editor: K. Winter

Abstract

We have studied the process $e^+e^- \rightarrow n\gamma$ ($n \geq 2$) at centre-of-mass energies of 161.3 GeV and 172.1 GeV. The analysis is based on a sample of events collected by the L3 detector in 1996 corresponding to total integrated luminosities of 10.7 pb⁻¹ and 10.1 pb⁻¹ respectively. The observed rates of events with two and more photons and the characteristic distributions are in good agreement with the Standard Model expectations. This is used to set lower limits on contact interaction energy scale parameters, on the QED cut-off parameters and on the mass of excited electrons. © 1997 Elsevier Science B.V.

1. Introduction

During 1996 LEP increased the centre-of-mass energy above 160 GeV providing a unique opportunity to search for new physics beyond the Standard Model. The process $e^+e^- \rightarrow n\gamma$ ($n \geq 2$) is well suited

for this purpose. On one hand it is a clean process with negligible background and with small non-QED radiative corrections. On the other hand it may be influenced by new phenomena, like compositeness or effective contact interactions, and its sensitivity increases with the centre-of-mass energy.

¹ Also supported by CONICET and Universidad Nacional de La Plata, CC 67, 1900 La Plata, Argentina.

² Also supported by Panjab University, Chandigarh-160014, India.

³ Supported by the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie.

⁴ Supported by the National Natural Science Foundation of China.

⁵ Supported by the Hungarian OTKA fund under contract numbers T14459 and T24011.

⁶ Supported also by the Comisión Interministerial de Ciencia y Tecnología.

In this paper we present the results on the search for new physics based on the process $e^+e^- \rightarrow n\gamma$ ($n \geq 2$). The analysis is performed with a sample of events collected by the L3 experiment in 1996 which corresponds to a total integrated luminosity of 10.69 pb^{-1} at the centre-of-mass energy of 161.3 GeV and 10.09 pb^{-1} at 172.1 GeV . Previous results have been published at lower centre-of-mass energies [1–3].

The L3 detector and its performance is described in detail in [4]. In 1996 a lead scintillator fibre calorimeter [5] was installed in the gap between the electromagnetic calorimeter barrel region and the end-caps to measure more precisely the energy of the particles which go into this region.

2. Event selection

To obtain a clean sample of $e^+e^- \rightarrow n\gamma$ ($n \geq 2$) events different selection criteria are applied. They are based on "photon candidates" defined as:

- i) A shower in the electromagnetic calorimeter with an energy above 1 GeV or in the lead scintillator fibre calorimeter with an energy above 10 GeV . Spurious signals are rejected by requiring a shower profile consistent with that of a photon. For the lead scintillator fibre calorimeter region, where no shower profile is available, we require a scintillator signal in time within a cone of 14° half-opening angle. A scintillator signal is always observed for electromagnetic particles of energy above 10 GeV , due to the leakage of charged particles from the shower;
- ii) The number of signals in the vertex chamber within an azimuthal angle of $\pm 8^\circ$ around the path of any photon candidate must be less than 40% of that expected for a charged particle. This requirement makes the selection insensitive to the presence of noise at low polar angles, where only few signals are expected. No change in the number of selected events is observed when the occupancy cut is varied in the range 20%–40%.

To ensure a good identification a fiducial cut is applied requiring that the events have:

- At least two photon candidates with a polar angle θ_γ between 16° and 164° and an angular separation of more than 15° .

The main sources of background come from $e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$ and cosmic rays. To reduce their contribution we require that:

- The sum of the energies of the photon candidates must be larger than $\sqrt{s}/2$.

With these selection cuts the contamination from other processes, estimated from Monte Carlo simulations, is negligible. In order to determine the acceptance, the same analysis is applied to a sample of $e^+e^- \rightarrow \gamma\gamma(\gamma)$ Monte Carlo generated events passed through the L3 simulation and reconstruction programs. The overall selection efficiency is found to be $79.3 \pm 0.2\%$ for θ_γ between 16° and 164° and the trigger efficiency is estimated to be above 99.7%.

3. Analysis of $e^+e^- \rightarrow n\gamma$ ($n \geq 2$) events

After applying these selection cuts the number of observed events, classified according to the number of isolated photons within the range $16^\circ < \theta_\gamma < 164^\circ$, is given in Table 1 together with the number of expected events from the process $e^+e^- \rightarrow n\gamma$ ($n = 2,3,4$) for the two different centre-of-mass energies [6]. No events with 5 or more photons have been observed.

For the two most energetic photons of the $n \geq 2 \gamma$ events the distribution of the acollinearity is shown




Fig. 1. Distribution of the acollinearity angle between the two most energetic photons in the $e^+e^- \rightarrow \gamma\gamma(\gamma)$ process. Data samples at $\sqrt{s} = 161.3$ and $\sqrt{s} = 172.1 \text{ GeV}$ have been combined. The points are data and the histogram is the Monte Carlo prediction.




Fig. 2. Distribution of the invariant mass of the two most energetic photons of the process $e^+e^- \rightarrow n\gamma (n \geq 2)$ for $\sqrt{s} = 161$ GeV (a) and 172 GeV (b). The points are data and the histogram is the Monte Carlo prediction.

in Fig. 1 and of the invariant mass in Fig. 2 together with the Monte Carlo expected distributions.

The distribution of the $\cos\theta^*$ of the event⁷ is shown in Fig. 3 compared with the Monte Carlo prediction. The data shows good agreement with QED.

The 137 and 112 observed events at $\sqrt{s} = 161.3$

⁷The polar angle θ^* of the event is defined as $\cos\theta^* = |\sin(\frac{\theta_1 - \theta_2}{2}) / \sin(\frac{\theta_1 + \theta_2}{2})|$, where θ_1 and θ_2 are the polar angles of the two most energetic photons in the event.




Fig. 3. Distribution of the polar angle of the event for the selected $e^+e^- \rightarrow \gamma\gamma(\gamma)$ sample. Data samples at $\sqrt{s} = 161.3$ and $\sqrt{s} = 172.1$ GeV have been combined. The points are data and the histogram is the Monte Carlo prediction.

GeV and $\sqrt{s} = 172.1$ GeV with $n \leq 3$ γ correspond to values of the total measured cross-sections of:

$$\sigma_{\gamma\gamma(\gamma)}(\sqrt{s} = 161.3 \text{ GeV}) = 16.2 \pm 1.4 \text{ pb}$$

and

$$\sigma_{\gamma\gamma(\gamma)}(\sqrt{s} = 172.1 \text{ GeV}) = 13.9 \pm 1.3 \text{ pb}$$

when at least two photons are in the range $16^\circ < \theta_\gamma < 164^\circ$. The quoted error is purely statistical. The possible systematic effects have been found to be much smaller than the statistical errors and are neglected. The same holds for the error on the measured luminosity and for the error associated to the contribution of the different sources of background. The predicted cross-sections for the process $e^+e^- \rightarrow \gamma\gamma(\gamma)$ at the two centre-of-mass energies are 16.40 ± 0.09 pb and 14.25 ± 0.09 pb [6] respectively, in good agreement with the observed values.

The two measured cross-sections are shown in Fig. 4 as a function of the centre-of-mass energy

Table 1

Observed and expected number of events with 2, 3 and 4 photons

Event	$\sqrt{s} = 161.3$ GeV		$\sqrt{s} = 172.1$ GeV	
	Observed	Expected	Observed	Expected
2 γ	131	130.6	109	108.7
3 γ	6	7.9	3	5.8
4 γ	0	0.3	2	0.2




Fig. 4. Measured cross-sections as function of the centre-of-mass energy for θ_γ between 16° and 164° compared with the QED prediction. The value at $\sqrt{s} = 90$ GeV has been extrapolated to the aforementioned angular range from the one given in [1].

together with the prediction of QED and our previously determined values at $\sqrt{s} = 91.2$ GeV [1] and $\sqrt{s} = 133.3$ GeV [2].

4. Limits on deviations from QED

The possible deviations from QED are parametrised in terms of effective Lagrangians, and their effect on the observables can be expressed as a multiplicative correction term to the QED differential cross-section. Depending on the type of parametrisation two general forms are considered:

$$\frac{d\sigma}{d\Omega} = \left(\frac{d\sigma}{d\Omega} \right)_{\text{QED}} \left(1 + \frac{s^2}{\alpha} \frac{1}{\Lambda^4} \sin^2\theta \right) \quad (1)$$

and

$$\frac{d\sigma}{d\Omega} = \left(\frac{d\sigma}{d\Omega} \right)_{\text{QED}} \left(1 + \frac{s^3}{32\pi\alpha^2} \frac{1}{\Lambda'^6} \frac{\sin^2\theta}{1 + \cos^2\theta} \right) \quad (2)$$

which depend on the centre-of-mass energy, the polar angle θ and the scale parameter Λ which has dimension of energy. A simpler and more standard way of parametrisation the deviations from QED is the introduction of the cut-off parameters Λ_\pm [7]. The differential cross-section can be obtained from Eq. 1 by replacing Λ^4 by $\pm(2/\alpha)\Lambda_\pm^4$.

Limits on the different scale parameters have

already been set in our previous publication [2]. However, since the sensitivity to possible deviations from QED increases rapidly with the centre-of-mass energy they are improved with the present data. In order to quantify the possible deviations from QED we define, for each sample at a given centre-of-mass energy, a likelihood for the different hypotheses of Λ in terms of the observed polar angle of the event (θ_i) and the total number of observed events (N_o) as:

$$L(\lambda_p) = \frac{1}{\sqrt{2\pi}\sigma(\lambda_p)} \exp\left(-\frac{(N_o - N_t(\lambda_p))^2}{2\sigma^2(\lambda_p)}\right) \times \prod_{i=1}^{N_o} f(\cos\theta_i|\lambda_p) \quad (3)$$

In this expression λ_p stands for the parameter under consideration ($1/\Lambda^4$ or $1/\Lambda'^6$); $N_t(\lambda_p)$ is the total number of expected events, $\sigma(\lambda_p)$ the statistical error on the number of expected events and $f(\cos\theta_i|\lambda_p)$ the probability density function of the polar angle θ . The choice of λ_p as a parameter has the advantage of giving, to a good approximation, a parabolic shaped log-likelihood around the maximum. The estimated parameters from the combined data samples at the two centre-of-mass energies are:

$$\frac{1}{\Lambda^4} = (-0.03^{+0.11}_{-0.10}) 10^{-11} \text{ GeV}^{-4},$$

$$\frac{1}{\Lambda'^6} = (-0.11^{+0.32}_{-0.30}) 10^{-16} \text{ GeV}^{-6}$$

consistent with no deviations from QED. To determine the confidence levels the probability distribution is normalised over the physically allowed range of the parameters. At the 95% C.L. the following limits are obtained:

$$\Lambda > 844 \text{ GeV}, \quad \Lambda_+ > 207 \text{ GeV},$$

$$\Lambda_- > 205 \text{ GeV}, \quad \Lambda' > 507 \text{ GeV}$$

Another way to study possible deviations from QED is to postulate the existence of an excited electron (e^*) of mass m_{e^*} which couples to the electron and the photon via magnetic interactions. To describe this interaction two different phenomenological Lagrangians are used; one with a magnetic interaction [8]:

$$\mathcal{L} = \frac{e}{2\Lambda_{e^*}} \bar{\Psi}_{e^*} \cdot \sigma^{\mu\nu} \Psi_e F_{\mu\nu} + \text{h.c.} \quad (4)$$

and another one with a magnetic interaction where only left-handed or right-handed fermions are involved [9]:

$$\mathcal{L} = \frac{e}{2\Lambda_e} \bar{\Psi}_e \cdot \sigma^{\mu\nu} (1 \pm \gamma^5) \Psi_e F_{\mu\nu} + \text{h.c.} \quad (5)$$

In both cases Λ_e is related to the effective scale of the interaction and m_e is the additional mass parameter. Fixing the interaction scale Λ_e to m_e we obtain

$$\frac{1}{m_e^4} = (-0.10^{+0.28}_{-0.28}) 10^{-9} \text{ GeV}^{-4}$$

for the first case and

$$\frac{1}{m_e^4} = (-0.25^{+0.78}_{-1.04}) 10^{-9} \text{ GeV}^{-4}$$

for the second one. From them we derive the 95% C.L. lower limits of:

$$m_e > 210 \text{ GeV}$$

and

$$m_e > 157 \text{ GeV}$$

respectively.

Acknowledgements

We wish to express our gratitude to the CERN accelerator divisions for the excellent performance of the LEP machine. We acknowledge the effort of all

the engineers and technicians who have participated in the construction and maintenance of the experiment.

References

- [1] L3 Collab., M. Acciarri et al., Phys. Lett. B 353 (1995) 136.
- [2] L3 Collab., O. Adriani et al., Phys. Lett. B 384 (1996) 323.
- [3] ALEPH Collab., D. Buskulic et al., Phys. Lett. B 384 (1996) 333;
DELPHI Collab., P. Abreu et al., Phys. Lett. B 327 (1994) 242;
OPAL Collab., G. Alexander et al., Phys. Lett. B 377 (1996) 222.
- [4] The L3 Collaboration, B. Adeva et al., Nucl. Instr. and Meth. A 289 (1990) 35;
M. Chemarin et al., Nucl. Instr. and Meth. A 349 (1994) 345;
M. Acciarri et al., Nucl. Instr. and Meth. A 351 (1994) 300;
I.C. Brock et al., Nucl. Instr. and Meth. A 381 (1996) 236;
A. Adam et al., Nucl. Instr. and Meth. A 383 (1996) 342.
- [5] G. Basti et al., Nucl. Instr. and Meth. A 374 (1996) 293.
- [6] F.A. Berends, R. Kleiss, Nucl. Phys. B 186 (1981) 22;
CALKUL Collaboration, F.A. Berends et al., Nucl. Phys. B 239 (1984) 395.
- [7] F.E. Low, Phys. Rev. Lett. 14 (1965) 238;
R.P. Feynman, Phys. Rev. Lett. 74 (1948) 939;
F.M. Renard, Phys. Lett. B 116 (1982) 264;
S. Drell, Ann. Phys. (NY) 4 (1958) 75; Z. Phys. C 29 (1985) 115.
- [8] A. Litke, Harvard Univ., Ph.D. Thesis, 1970, unpublished.
- [9] K. Hagiwara, S. Komamyia, D. Zeppenfeld, Z. Phys. C 29 (1985) 115;
N. Cabibbo, L. Maiani, Y. Srivastava, Phys. Lett. B 139 (1984) 459;
F.M. Renard, Nucl. Phys. B 196 (1982) 93.