

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF NIJMEGEN The Netherlands

**A Class of Counterexamples to the Biregular
Cancellation Problem**

Arno van den Essen Peter van Rossum

Report No. 9826 (December 1998)

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF NIJMEGEN
Toernooiveld
6525 ED Nijmegen
The Netherlands

A Class of Counterexamples to the Biregular Cancellation Problem

Arno van den Essen Peter van Rossum

Abstract

If V and W are algebraic varieties over \mathbb{C} , then the Biregular Cancellation Problem asks whether or not $V \times \mathbb{C}^k \cong W \times \mathbb{C}^k$ implies $V \cong W$. This paper presents a class of examples where the answer is negative, using non-free stably free modules and locally nilpotent derivations.

1 Introduction

The Biregular Cancellation Problem for algebraic varieties asks the following question.

Problem 1 (Biregular Cancellation Problem, geometric formulation). *Let V and W be algebraic varieties over a field k and $n \in \mathbb{N}$. Does $V \times k^n \cong W \times k^n$ imply that $V \cong W$?*

This question can be reformulated as follows.

Problem 2 (Biregular Cancellation Problem, algebraic formulation). *Let A and B be affine domains over a field k and $n \in \mathbb{N}$. Consider the polynomial rings $A[X_1, \dots, X_n]$ and $B[X_1, \dots, X_n]$. Does $A[X_1, \dots, X_n] \cong B[X_1, \dots, X_n]$ imply that $A \cong B$ over k ?*

A special case of the Biregular Cancellation Problem is the Cancellation Problem.

Problem 3 (Cancellation Problem, geometric formulation). *Let V be an algebraic variety over a field k and $n \in \mathbb{N}^*$. Does $V \times k \cong k^n$ imply that $V \cong k^{n-1}$?*

First results on the Biregular Cancellation Problem were obtained by Abhyankar, Heinzer and Eakin ([AHE72]) in 1972, who proved the claim in case $\text{trdeg}_k Q(A) \leq 1$. In the same year, Hochster gave a counterexample for the case $k = \mathbb{R}$ ([Hoc72]). In 1989, Danielewski gave a counterexample to the Biregular Cancellation Problem ([Dan89], unpublished) over \mathbb{C} . He showed that for the algebraic varieties $Y_n := V(x^n y + z^2 - 1) \subseteq \mathbb{C}^3$, for $n \in \mathbb{N}^*$, the stabilizations $Y_n \times \mathbb{C}$ are all isomorphic. It was, however, shown by Fieseler that all Y_n are of different homotopy type. Recently, Makar-Limanov has studied algebraic varieties of the form $V(x^n y + p(z))$, for $n \in \mathbb{N}^*$ and $p(z) \in \mathbb{C}[z]$, and has characterized when such varieties are isomorphic ([Mak98],

preprint). In particular, he has obtained a purely algebraic proof of the fact that the varieties Y_n are non-isomorphic.

See also the paper by Kraft ([Kra89]) and the survey papers by Kang ([Kan87, Kan92]) for background information on these and other cancellation problems in algebraic geometry.

This paper shows how to construct a whole class of counterexamples to the Biregular Cancellation Problem.

The construction in Section 3 has two ingredients. On the one hand, it uses the existence of commutative rings A with a unimodular row (a_1, \dots, a_n) over A that cannot be completed to an invertible square matrix. In other words, it uses the existence of commutative rings A for which there exists a stably free module of type 1 that is not free. This was in fact also a basic ingredient in Hochster's paper, who considered the ring $\mathbb{R}[x, y, z]/(x^2 + y^2 + z^2 - 1)$ and the unimodular row $(\bar{x}, \bar{y}, \bar{z})$. On the other hand, our construction uses the notion of locally nilpotent derivations. Section 2 contains a brief overview of the required facts about these derivations.

2 Derivations

Let k be a field of characteristic zero and let A be a commutative k -algebra. A k -derivation on A is a k -linear map $D: A \rightarrow A$ satisfying the Leibniz-rule, $D(ab) = a(Db) + (Da)b$ for all $a, b \in A$. It is said to be *locally nilpotent* if for all $a \in A$ there is an $n \in \mathbb{N}$ such that $D^n(a) = 0$. The kernel of such a derivation D is denoted by A^D . A *slice* of D is an element $s \in A$ such that $D(s) = 1$.

If D is locally nilpotent and $t \in A$, then we can define a map $\phi_t: A \rightarrow A$ by $\phi_t(a) := \sum_{i=0}^{\infty} \frac{1}{i!} D^i(a)t^i$. If D also has a slice s , this map can be used to easily describe the kernel of D .

Proposition 4 ([Ess93], Proposition 2.1). *Let D be a locally nilpotent derivation on a finitely generated commutative k -algebra $A = k[a_1, \dots, a_n]$. Assume that D has a slice $s \in A$. Then*

$$A^D = \phi_{-s}(A) = k[\phi_{-s}(a_1), \dots, \phi_{-s}(a_n)].$$

Proposition 5 ([Wri81], Proposition 2.1). *Let D be a locally nilpotent derivation on a commutative k -algebra A and assume that D has a slice $s \in A$. Then*

1. $A = A^D[s]$;
2. s is algebraically independent over A [and therefore $A = A^D[s]$ is a polynomial ring in one variable over A^D];
3. $D = d/ds$.

Remark 6. Note that if, in the above situation, A is a domain and $\text{trdeg}_k Q(A)$ is finite, it follows that $\text{trdeg}_k Q(A^D) = \text{trdeg}_k Q(A) - 1$. In particular, if A is of the form $A = B[X_1, \dots, X_n]$ for some domain B whose quotient field is of finite transcendence degree over k and $A^D = B[F_1, \dots, F_{n-1}]$ for certain polynomials $F_1, \dots, F_{n-1} \in A$, then the F_i are algebraically independent over k .

For more information on locally nilpotent derivations see for instance [Ess98], Chapter 1 or [Now94].

3 Counterexamples

Let k be a field of characteristic zero and let A be a finitely generated commutative k -algebra without zero divisors. Let (a_1, \dots, a_n) be a unimodular row over A , say $b_1, \dots, b_n \in A$ with $b_1 a_1 + \dots + b_n a_n = 1$, and assume that it cannot be completed to an invertible square matrix.

Let $A[X]$ denote the polynomial ring $A[X_1, \dots, X_n]$ over A in n variables. Now define a k -derivation $D: A[X] \rightarrow A[X]$ by

$$D := b_1 \frac{\partial}{\partial X_1} + \dots + b_n \frac{\partial}{\partial X_n}.$$

This derivation is locally nilpotent and has a slice, namely $s := a_1 X_1 + \dots + a_n X_n$. Letting $B := A[X]^D$ be the kernel of the derivation, it follows from Proposition 5 that $A[X] = B[s]$, a polynomial ring over B in s , and from Proposition 4 that $B = A[X_1 - b_1 s, \dots, X_n - b_n s]$.

Notation 7. For $F \in A[X]$ and $j \in \mathbb{N}$ we denote by $F_{(j)}$ the homogeneous part of F of degree j . So $F = F_{(0)} + F_{(1)} + \dots + F_{(d)}$ where $d := \deg F$.

Lemma 8. Let $F_1, \dots, F_{n-1} \in A[X]$ and assume that $B = A[F_1, \dots, F_{n-1}]$. Take $f_i := F_{i(1)}$ (i.e. the linear part of F_i). Then $B = A[f_1, \dots, f_{n-1}]$.

Proof. \subseteq : We may assume, without loss of generality, that the polynomials F_1, \dots, F_{n-1} do not have a constant term.

Now consider $X_i - b_i s \in B = A[F_1, \dots, F_{n-1}]$. Then there is a polynomial $p(T_1, \dots, T_{n-1}) \in A[T_1, \dots, T_{n-1}]$ such that $X_i - b_i s = p(F_1, \dots, F_{n-1})$. Then

$$\begin{aligned} X_i - b_i s &= (p(F_1, \dots, F_{n-1}))_{(1)} && \text{(because } X_i - b_i s \text{ is linear)} \\ &= (p_{(1)}(F_1, \dots, F_{n-1}))_{(1)} && \text{(because } F_1, \dots, F_{n-1} \text{ have} \\ & && \text{no constant term)} \\ &= p_{(1)}(F_{1(1)}, \dots, F_{n-1(1)}) && \text{(because } p_{(1)} \text{ is linear)} \\ &= p_{(1)}(f_1, \dots, f_{n-1}) \in A[f_1, \dots, f_{n-1}] \end{aligned}$$

\supseteq : Because $F_i \in B = A[X]^D$, every homogeneous part $F_{i(j)}$ of F_i is also in B . In particular $f_i \in B$. \square

Lemma 9. Let $f_1, \dots, f_m \in A[X]$ be linear polynomials. Then

$$A[f_1, \dots, f_m] \cap AX_1 \oplus \dots \oplus AX_n = Af_1 + \dots + Af_m.$$

[i.e. every polynomial expression $p(f_1, \dots, f_m)$ in the f_i which is linear in the X_i is in fact an A -linear combination of the f_i .]

Proof. \subseteq : Take $p(T_1, \dots, T_m) \in A[T_1, \dots, T_m]$ and let $g := p(f_1, \dots, f_m)$ be a polynomial expression in the f_i . Assume that g is in fact linear in the X_i . Then, using essentially the same argument as in the proof of the previous lemma,

$$\begin{aligned} g &= (p(f_1, \dots, f_m))_{(1)} \\ &= (p_{(1)}(f_1, \dots, f_m))_{(1)} \\ &= p_{(1)}(f_1, \dots, f_m) \in Af_1 + \dots + Af_n. \end{aligned}$$

\supseteq : This is obvious. \square

Lemma 10. *Let $f_1, \dots, f_{n-1} \in A[X]$ be linear polynomials and assume that $B = A[f_1, \dots, f_{n-1}]$. Then*

$$As \oplus Af_1 \oplus \dots \oplus Af_{n-1} = AX_1 \oplus \dots \oplus AX_n.$$

[i.e. every linear polynomial in $A[X]$ can be written in a unique way as an A -linear combination of s, f_1, \dots, f_{n-1} .]

Proof. We first show that $As + Af_1 + \dots + Af_{n-1} = AX_1 \oplus \dots \oplus AX_n$.

\subseteq : This is obvious.

\supseteq : Take $g \in AX_1 \oplus \dots \oplus AX_n$. Then $Dg \in A$ and therefore $D(g - (Dg)s) = Dg - (D^2g)s - (Dg)(Ds) = Dg - Dg = 0$. So

$$\begin{aligned} g - (Dg)s &\in B \cap AX_1 \oplus \dots \oplus AX_n \\ &= A[f_1, \dots, f_{n-1}] \cap AX_1 \oplus \dots \oplus AX_n \\ &= Af_1 + \dots + Af_{n-1} \quad (\text{by Lemma 9}) \end{aligned}$$

and hence $g \in As + Af_1 + \dots + Af_{n-1}$.

To see that $As + Af_1 + \dots + Af_n$ is in fact a direct sum, take $\mu, \lambda_1, \dots, \lambda_{n-1} \in A$ and assume that $\mu s + \lambda_1 f_1 + \dots + \lambda_{n-1} f_{n-1} = 0$. Applying D to both sides yields $\mu = 0$, so $\lambda_1 f_1 + \dots + \lambda_{n-1} f_{n-1} = 0$. The f_i , however, are even algebraically independent (by Remark 6) and therefore $\lambda_1 = \dots = \lambda_{n-1} = 0$. \square

Theorem 11. $B \not\cong A[X_1, \dots, X_{n-1}]$, even though $B[s] = A[X_1, \dots, X_{n-1}][X_n]$.

Proof. Assume that $B \cong A[X_1, \dots, X_{n-1}]$. Then $B = A[F_1, \dots, F_{n-1}]$ for certain polynomials $F_1, \dots, F_{n-1} \in A[X]$ and by Lemma 8 even $B = A[f_1, \dots, f_{n-1}]$ for certain linear polynomials $f_1, \dots, f_{n-1} \in A[X]$. Now Lemma 10 implies that

$$As \oplus Af_1 \oplus \dots \oplus Af_{n-1} = AX_1 \oplus \dots \oplus AX_n,$$

say $f_i = \lambda_{i1}X_1 + \dots + \lambda_{in}X_n$ (and $s = a_1X_1 + \dots + a_nX_n$). This is an equality between free A -modules of rank n and the base transformation matrix is

$$\begin{pmatrix} a_1 & \dots & a_n \\ \lambda_{11} & \dots & \lambda_{1n} \\ \vdots & & \vdots \\ \lambda_{n-11} & \dots & \lambda_{n-1n} \end{pmatrix}.$$

This is an invertible matrix and hence the unimodular row (a_1, \dots, a_n) has been completed to an invertible matrix, which contradicts the assumption. \square

So, every coordinate ring A of an affine variety that has a unimodular row that cannot be completed to an invertible matrix, gives rise to a counterexample to the Biregular Cancellation Problem.

Over the real numbers, we recover Hochster's example mentioned in the Introduction. Over the complex numbers, one can consider the "generic" example $A = \mathbb{C}[a, b, c, x, y, z]/(ax + by + cz - 1)$. The unimodular row $(\bar{x}, \bar{y}, \bar{z})$ cannot be completed to an invertible square matrix. This was shown by Raynaud in [Ray68] using homological methods and, in a more general setting, by Suslin in [Sus82] (Theorem 2.8) using K -theory.

4 Acknowledgments

We would like to thank Wilberd van der Kallen for pointing out the references to the results of Raynaud and Suslin.

References

- [AHE72] Shreeram S. Abhyankar, William Heinzer, and Paul Eakin, *On the uniqueness of the coefficient ring in a polynomial ring*, J. Alg. **23** (1972), 310–342.
- [Dan89] W. Danielewski, *On the cancellation problem and automorphism group of affine algebraic varieties*, preprint, 1989.
- [Ess93] Arno van den Essen, *An algorithm to compute the invariant ring of a G_a -action on an affine variety*, J. Symb. Comp. **16** (1993), 551–555.
- [Ess98] Arno van den Essen, *Invertible polynomial maps and the Jacobian conjecture*, manuscript, 1998.
- [Hoc72] M. Hochster, *Nonuniqueness of coefficient rings in a polynomial ring*, Proc. Am. Math. Soc. **34** (1972), no. 1, 81–82.
- [Kan87] Ming-chang Kang, *The biregular cancellation problem*, J. Pure Appl. Alg. **45** (1987), 241–253.
- [Kan92] Ming-chang Kang, *A note on the birational cancellation problem*, J. Pure Appl. Alg. **77** (1992), 141–154.
- [Kra89] Hanspeter Kraft, *Algebraic automorphisms of affine space*, Topological Methods in Algebraic Transformation Groups (Boston-Basel-Berlin) (H. Kraft, T. Petrie, and G. Schwarz, eds.), Birkhäuser-Verlag, 1989.
- [Mak98] Makar-Limanov, *Locally nilpotent derivations, a new ring invariant and applications*, preprint, 1998.

- [Now94] A. Nowicki, *Commutative basis of derivations in polynomial and power series rings*, Univ. of Toruń, 1994.
- [Ray68] Michèle Raynaud, *Modules projectifs universels*, Inven. Math. **6** (1968), 1–26.
- [Sus82] A.A. Suslin, *Mennicke symbols and their applications in the K-theory of fields*, Algebraic K-Theory, Proceedings of a Conference Held at Oberwolfach, June 1980, Part I (Berlin) (R. Keith Dennis, ed.), Lecture Notes in Mathematics, vol. 966, Springer-Verlag, 1982, pp. 344–356.
- [Wri81] David Wright, *On the Jacobian conjecture*, Illinois J. Math. **25** (1981), no. 3, 423–440.