

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/28959>

Please be advised that this information was generated on 2019-05-23 and may be subject to change.



ELSEVIER

4 May 1995

PHYSICS LETTERS B

Physics Letters B 350 (1995) 109–119

Search for neutralinos in Z decays

L3 Collaboration

M. Acciarri^z, A. Adam^{aq}, O. Adriani^p, M. Aguilar-Benitez^y, S. Ahlen^j,
B. Alpat^{ag}, J. Alcaraz^y, J. Allaby^q, A. Aloisio^{ab}, G. Alverson^k, M.G. Alviggi^{ab},
G. Ambrosi^{ah}, Q. An^r, H. Anderhub^{at}, A.L. Anderson^o, V.P. Andreev^{ak},
T. Angelescu^l, D. Antreasyan^h, G. Alkhazov^{ak}, A. Arefiev^{aa}, T. Azemoon^c,
T. Azizⁱ, P.V.K.S. Baba^r, P. Bagnaia^{aj,q}, J.A. Bakken^{ai}, L. Baksay^{ap}, R.C. Ball^c,
S. Banerjeeⁱ, K. Banicz^{aq}, R. Barillère^q, L. Barone^{aj}, P. Bartalini^{ag},
A. Baschirotto^z, M. Basile^h, R. Battiston^{ag}, A. Bay^v, F. Becattini^p, U. Becker^o,
F. Behner^{at}, Gy.L. Bencze^m, J. Berdugo^y, P. Berges^o, B. Bertucci^q,
B.L. Betev^{at}, M. Biasini^{ag}, A. Biland^{at}, G.M. Bilei^{ag}, R. Bizzarri^{aj}, J.J. Blaising^q,
G.J. Bobbink^b, R. Bock^a, A. Böhm^a, B. Borgia^{aj}, A. Boucham^d, D. Bourilkov^{at},
M. Bourquin^s, D. Boutigny^d, B. Bouwens^b, E. Brambilla^o, J.G. Branson^{al},
V. Brigljevic^{at}, I.C. Brock^{ah}, A. Bujak^{aq}, J.D. Burger^o, W.J. Burger^s,
C. Burgos^u, J. Busenitz^{ap}, A. Buytenhuijs^{ad}, A. Bykov^{ak}, X.D. Cai^r, M. Capell^o,
G. Cara Romeo^h, M. Caria^g, G. Carlino^{ab}, A.M. Cartacci^p, J. Casaus^y,
G. Castellini^p, R. Castello^z, N. Cavallo^{ab}, C. Cecchi^s, M. Cerrada^y, F. Cesaroni^{aj},
M. Chamizo^y, Y.H. Chang^{au}, U.K. Chaturvedi^r, M. Chemarin^x, A. Chen^{au},
C. Chen^f, G. Chen^f, G.M. Chen^f, H.F. Chen^t, H.S. Chen^f, M. Chen^o,
G. Chiefari^{ab}, C.Y. Chien^c, M.T. Choi^{ao}, S. Chung^o, L. Cifarelli^h, F. Cindolo^h,
C. Civinini^p, I. Clare^o, R. Clare^o, T.E. Coan^w, H.O. Cohn^{ae}, G. Coignet^d,
N. Colino^q, V. Commichau^a, S. Costantini^{aj}, F. Cotorobai^l, B. de la Cruz^y,
X.T. Cui^r, X.Y. Cui^r, T.S. Dai^o, R. D'Alessandro^p, R. de Asmundis^{ab},
H. De Boeck^{ad}, A. Degré^d, K. Deiters^{ar}, E. Dénes^m, P. Denes^{ai},
F. DeNotaristefani^{aj}, D. DiBitonto^{ap}, M. Diemoz^{aj}, C. Dionisi^{aj}, M. Dittmar^{at},
A. Doria^{ab}, I. Dorne^d, M.T. Dova^{r,4}, E. Drago^{ab}, D. Duchesneau^q, F. Duhem^d,
P. Duinker^b, I. Duran^{am}, S. Duttaⁱ, S. Easo^{ag}, H. El Mamouni^x, A. Engler^{ah},
F.J. Eppling^o, F.C. Erné^b, J.P. Ernenwein^x, P. Extermann^s, R. Fabbretti^{ar},
M. Fabre^{ar}, R. Faccini^{aj}, S. Falciano^{aj}, A. Favara^p, J. Fay^x, M. Felcini^{at},
T. Ferguson^{ah}, D. Fernandez^y, G. Fernandez^y, F. Ferroni^{aj}, H. Fesefeldt^a,

E. Fiandrini ^{ag}, J.H. Field ^s, F. Filthaut ^{ah}, P.H. Fisher ^e, G. Forconi ^o, L. Fredj ^s,
 K. Freudenreich ^{at}, M. Gailloud ^v, Yu. Galaktionov ^{aa,o}, E. Gallo ^p, S.N. Ganguli ⁱ,
 P. Garcia-Abia ^y, S.S. Gau ^k, S. Gentile ^{aj}, J. Gerald ^e, N. Gheordanescu ^l,
 S. Giagu ^{aj}, S. Goldfarb ^v, J. Goldstein ^j, Z.F. Gong ^t, E. Gonzalez ^y, A. Gougas ^e,
 D. Goujon ^s, G. Gratta ^{af}, M.W. Gruenewald ^g, C. Gu ^r, M. Guanziroli ^r,
 V.K. Gupta ^{ai}, A. Gurtu ⁱ, H.R. Gustafson ^c, L.J. Gutay ^{aq}, B. Hartmann ^a,
 A. Hasan ^{ac}, J.T. He ^f, T. Hebbeker ^g, M. Hebert ^{al}, A. Hervé ^q, K. Hilgers ^a,
 W.C. van Hoek ^{ad}, H. Hofer ^{at}, H. Hoorani ^s, S.R. Hou ^{au}, G. Hu ^r, B. Ille ^x,
 M.M. Ilyas ^r, V. Innocente ^q, H. Janssen ^d, B.N. Jin ^f, L.W. Jones ^c, P. de Jong ^o,
 I. Josa-Mutuberria ^y, A. Kasser ^v, R.A. Khan ^r, Yu. Kamyshkov ^{ac}, P. Kapinos ^{as},
 J.S. Kapustinsky ^w, Y. Karyotakis ^d, M. Kaur ^r, S. Khokhar ^r,
 M.N. Kienzle-Focacci ^s, D. Kim ^e, J.K. Kim ^{ao}, S.C. Kim ^{ao}, Y.G. Kim ^{ao},
 W.W. Kinnison ^w, A. Kirkby ^{af}, D. Kirkby ^{af}, J. Kirkby ^q, S. Kirsch ^{as}, W. Kittel ^{ad},
 A. Klimentov ^{o,aa}, A.C. König ^{ad}, E. Koffeman ^b, O. Kornadt ^a, V. Koutsenko ^{o,aa},
 A. Koulbardis ^{ak}, R.W. Kraemer ^{ah}, T. Kramer ^o, W. Krenz ^a, H. Kuijten ^{ad},
 A. Kunin ^{o,aa}, P. Ladron de Guevara ^y, G. Landi ^p, M. Lanfranchi ^{ag}, S. Lanzano ^{ab,5},
 P. Laurikainen ^u, M. Lebeau ^q, A. Lebedev ^o, P. Lebrun ^x, P. Lecomte ^{at}, P. Lecoq ^q,
 P. Le Coultre ^{at}, J.S. Lee ^{ao}, K.Y. Lee ^{ao}, I. Leedom ^k, C. Leggett ^c, J.M. Le Goff ^q,
 R. Leiste ^{as}, M. Lenti ^p, E. Leonardi ^{aj}, P. Levtchenko ^{ak}, C. Li ^{t,r}, E. Lieb ^{as},
 W.T. Lin ^{au}, F.L. Linde ^b, B. Lindemann ^a, L. Lista ^{ab}, Y. Liu ^r, W. Lohmann ^{as},
 E. Longo ^{aj}, W. Lu ^{af}, Y.S. Lu ^f, K. Lübelmeyer ^a, C. Luci ^{aj}, D. Luckey ^o,
 L. Ludovici ^{aj}, L. Luminari ^{aj}, W. Lustermann ^{ar}, W.G. Ma ^t, M. MacDermott ^{at},
 M. Maity ⁱ, L. Malgeri ^{aj}, R. Malik ^r, A. Malinin ^{aa}, C. Maña ^y, S. Mangla ⁱ,
 M. Maolinbay ^{at}, P. Marchesini ^{at}, A. Marin ^j, J.P. Martin ^x, F. Marzano ^{aj},
 G.G.G. Massaro ^b, K. Mazumdar ⁱ, T. McMahon ^{aq}, D. McNally ^q, S. Mele ^{ab},
 M. Merk ^{ah}, L. Merola ^{ab}, M. Meschini ^p, W.J. Metzger ^{ad}, Y. Mi ^v, A. Mihul ^l,
 A.J.W. van Mil ^{ad}, Y. Mir ^r, G. Mirabelli ^{aj}, J. Mnich ^q, M. Möller ^a, V. Monaco ^{aj},
 B. Monteleoni ^p, R. Moore ^c, R. Morand ^d, S. Morganti ^{aj}, N.E. Moulai ^r,
 R. Mount ^{af}, S. Müller ^a, E. Nagy ^m, M. Napolitano ^{ab}, F. Nessi-Tedaldi ^{at},
 H. Newman ^{af}, M.A. Niaz ^r, A. Nippe ^a, H. Nowak ^{as}, G. Organtini ^{aj}, R. Ostonen ^u,
 D. Pandoulas ^a, S. Paoletti ^{aj}, P. Paolucci ^{ab}, G. Pascale ^{aj}, G. Passaleva ^p,
 S. Patricelli ^{ab}, T. Paul ^{ag}, M. Pauluzzi ^{ag}, C. Paus ^a, F. Pauss ^{at}, Y.J. Pei ^a,
 S. Pensotti ^z, D. Perret-Gallix ^d, A. Pevsner ^e, D. Piccolo ^{ab}, M. Pieri ^p, J.C. Pinto ^{ah},
 P.A. Piroué ^{ai}, E. Pistolesi ^p, F. Plasil ^{ac}, V. Plyaskin ^{aa}, M. Pohl ^{at}, V. Pojidaev ^{aa,p},
 H. Postema ^o, N. Produit ^s, K.N. Qureshi ^r, R. Raghavan ⁱ, G. Rahal-Callot ^{at},
 P.G. Rancoita ^z, M. Rattaggi ^z, G. Raven ^b, P. Razis ^{ac}, K. Read ^{ae}, M. Redaelli ^z,
 D. Ren ^{at}, Z. Ren ^r, M. Rescigno ^{aj}, S. Reucroft ^k, A. Ricker ^a, S. Riemann ^{as},
 B.C. Riemers ^{aq}, K. Riles ^c, O. Rind ^c, H.A. Rizvi ^r, S. Ro ^{ao}, A. Robohm ^{at},

F.J. Rodriguez ^y, B.P. Roe ^c, M. Röhner ^a, S. Röhner ^a, L. Romero ^y,
 S. Rosier-Lees ^d, Ph. Rosselet ^v, W. van Rossum ^b, S. Roth ^a, J.A. Rubio ^q,
 H. Rykaczewski ^{at}, J. Salicio ^q, J.M. Salicio ^y, E. Sanchez ^y, A. Santocchia ^{ag},
 M.E. Sarakinos ^{aq}, S. Sarkar ⁱ, G. Sartorelli ^r, M. Sassowsky ^a, G. Sauvage ^d,
 C. Schäfer ^a, V. Schegelsky ^{ak}, D. Schmitz ^a, P. Schmitz ^a, M. Schneegans ^d,
 B. Schoeneich ^{as}, N. Scholz ^{at}, H. Schopper ^{au}, D.J. Schotanus ^{ad}, R. Schulte ^a,
 K. Schultze ^a, J. Schwenke ^a, G. Schwering ^a, C. Sciacca ^{ab}, R. Sehgal ^r,
 P.G. Seiler ^{ar}, J.C. Sens ^{au}, L. Servoli ^{ag}, I. Sheer ^{al}, S. Shevchenko ^{af}, N. Shivarov ^{an},
 S. Shotkin ^o, V. Shoutko ^{aa}, J. Shukla ^v, E. Shumilov ^{aa}, D. Son ^{ao}, A. Sopczak ^q,
 V. Soulimov ^{ab}, C. Spartiotis ^u, T. Spickermann ^a, P. Spillantini ^p, M. Steuer ^o,
 D.P. Stickland ^{ai}, F. Sticozzi ^o, H. Stone ^{ai}, B. Stoyanov ^{an}, K. Strauch ⁿ,
 K. Sudhakar ⁱ, G. Sultanov ^r, L.Z. Sun ^{ts}, G.F. Susinno ^s, H. Suter ^{at}, J.D. Swain ^r,
 A.A. Syed ^{ad}, X.W. Tang ^f, L. Taylor ^k, R. Timellini ^h, Samuel C.C. Ting ^o,
 S.M. Ting ^o, O. Toker ^{ag}, M. Tonutti ^a, S.C. Tonwar ⁱ, J. Tóth ^m,
 A. Tsaregorodtsev ^{ak}, G. Tsipolitis ^{ah}, C. Tully ^{ai}, H. Tuchscherer ^{ap}, J. Ulbricht ^{at},
 L. Urbän ^m, U. Uwer ^a, E. Valente ^{aj}, R.T. Van de Walle ^{ad}, I. Vetlitsky ^{aa},
 G. Viertel ^{at}, P. Vikas ^r, U. Vikas ^r, M. Vivargent ^d, R. Voelkert ^{as}, H. Vogel ^{ah},
 H. Vogt ^{as}, I. Vorobiev ^{aa}, A.A. Vorobyov ^{ak}, An.A. Vorobyov ^{ak}, L. Vuilleumier ^v,
 M. Wadhwa ^y, W. Wallraff ^a, J.C. Wang ^o, X.L. Wang ^t, Y.F. Wang ^o,
 Z.M. Wang ^{tr}, A. Weber ^a, R. Weill ^v, C. Willmott ^y, F. Wittgenstein ^q, D. Wright ^{ai},
 S.X. Wu ^r, S. Wynhoff ^a, Z.Z. Xu ^t, B.Z. Yang ^t, C.G. Yang ^f, G. Yang ^r,
 X.Y. Yao ^f, C.H. Ye ^r, J.B. Ye ^t, Q. Ye ^r, S.C. Yeh ^{au}, J.M. You ^{ah}, N. Yunus ^r,
 M. Yzerman ^b, C. Zaccardelli ^{af}, P. Zemp ^{at}, M. Zeng ^r, Y. Zeng ^a,
 Z. Zhang ^f, Z.P. Zhang ^{tr}, B. Zhou ^j, G.J. Zhou ^f, J.F. Zhou ^a, Y. Zhou ^c,
 R.Y. Zhu ^{af}, A. Zichichi ^{h,q,r}, B.C.C. van der Zwaan ^b

^a I. Physikalisches Institut, RWTH, D-52056 Aachen, FRG ¹

III. Physikalisches Institut, RWTH, D-52056 Aachen, FRG ¹

^b National Institute for High Energy Physics, NIKHEF, NL-1009 DB Amsterdam, The Netherlands

^c University of Michigan, Ann Arbor, MI 48109, USA

^d Laboratoire d'Annecy-le-Vieux de Physique des Particules, LAPP, IN2P3-CNRS, BP 110, F-74941 Annecy-le-Vieux CEDEX, France

^e Johns Hopkins University, Baltimore, MD 21218, USA

^f Institute of High Energy Physics, IHEP, 100039 Beijing, China

^g Humboldt University, D-10099 Berlin, FRG

^h INFN-Sezione di Bologna, I-40126 Bologna, Italy

ⁱ Tata Institute of Fundamental Research, Bombay 400 005, India

^j Boston University, Boston, MA 02215, USA

^k Northeastern University, Boston, MA 02115, USA

^l Institute of Atomic Physics and University of Bucharest, R-76900 Bucharest, Romania

^m Central Research Institute for Physics of the Hungarian Academy of Sciences, H-1525 Budapest 114, Hungary ²

ⁿ Harvard University, Cambridge, MA 02139, USA

^o Massachusetts Institute of Technology, Cambridge, MA 02139, USA

^p INFN Sezione di Firenze and University of Florence, I-50125 Florence, Italy

^q European Laboratory for Particle Physics, CERN, CH-1211 Geneva 23, Switzerland

^r World Laboratory, FBLJA Project, CH-1211 Geneva 23, Switzerland

- ^s University of Geneva, CH-1211 Geneva 4, Switzerland
^t Chinese University of Science and Technology, USTC, Hefei, Anhui 230 029, China
^u SEFT, Research Institute for High Energy Physics, P.O. Box 9, SF-00014 Helsinki, Finland
^v University of Lausanne, CH-1015 Lausanne, Switzerland
^w Los Alamos National Laboratory, Los Alamos, NM 87544, USA
^x Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, F-69622 Villeurbanne Cedex, France
^y Centro de Investigaciones Energeticas, Medioambientales y Tecnológicas, CIEMAT, E-28040 Madrid, Spain ³
^z INFN-Sezione di Milano, I-20133 Milan, Italy
^{aa} Institute of Theoretical and Experimental Physics, ITEP, Moscow, Russia
^{ab} INFN-Sezione di Napoli and University of Naples, I-80125 Naples, Italy
^{ac} Department of Natural Sciences, University of Cyprus, Nicosia, Cyprus
^{ad} University of Nymegen and NIKHEF, NL-6525 ED Nymegen, The Netherlands
^{ae} Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA
^{af} California Institute of Technology, Pasadena, CA 91125, USA
^{ag} INFN-Sezione di Perugia and Università Degli Studi di Perugia, I-06100 Perugia, Italy
^{ah} Carnegie Mellon University, Pittsburgh, PA 15213, USA
^{ai} Princeton University, Princeton, NJ 08544, USA
^{aj} INFN-Sezione di Roma and University of Rome, "La Sapienza", I-00185 Rome, Italy
^{ak} Nuclear Physics Institute, St. Petersburg, Russia
^{al} University of California, San Diego, CA 92093, USA
^{am} Dept. de Física de Partículas Elementales, Univ. de Santiago, E-15706 Santiago de Compostela, Spain
^{an} Bulgarian Academy of Sciences, Central Laboratory of Mechatronics and Instrumentation, BU-1113 Sofia, Bulgaria
^{ao} Center for High Energy Physics, Korea Advanced Inst. of Sciences and Technology, 305-701 Taejon, South Korea
^{ap} University of Alabama, Tuscaloosa, AL 35486, USA
^{aq} Purdue University, West Lafayette, IN 47907, USA
^{ar} Paul Scherrer Institut, PSI, CH-5232 Villigen, Switzerland
^{as} DESY-Institut für Hochenergiephysik, D-15738 Zeuthen, FRG
^{at} Eidgenössische Technische Hochschule, ETH Zürich, CH-8093 Zürich, Switzerland
^{au} University of Hamburg, D-22761 Hamburg, FRG
^{av} High Energy Physics Group, Taiwan, ROC

Received 9 February 1995

Editor: K. Winter

Abstract

We have searched for neutralinos produced via the reactions $e^+e^- \rightarrow \chi\chi'$ and $e^+e^- \rightarrow \chi'\chi'$, where the next-to lightest neutralino, χ' , decays into the lightest neutralino, χ , and either a photon or a fermion pair. Based on 1.8×10^6 hadronic Z decays collected with the L3 detector at LEP, no signal has been observed. We present upper limits of a few times 10^{-5} on the branching ratios $Z \rightarrow \chi\chi'$ and $Z \rightarrow \chi'\chi'$. In the framework of the Minimal Supersymmetric Standard Model, we exclude a lightest χ with m_χ less than 18 GeV, if either $\tan \beta > 2$ or the gluino mass $m_{\tilde{g}} > 100$ GeV.

1. Introduction

The Standard Model [1] has been very successful in describing data concerning electroweak interactions. However, it leaves many fundamental parameters unexplained such as the electroweak mixing parameter $\sin^2\theta_w$. The quadratic divergences of scalar masses at the one-loop level and the large difference between the electroweak and grand unification scales (hierarchy problem) are further prob-

¹ Supported by the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie.

² Supported by the Hungarian OTKA fund under contract number 2970.

³ Supported also by the Comisión Interministerial de Ciencia y Tecnología.

⁴ Also supported by CONICET and Universidad Nacional de La Plata, CC 67, 1900 La Plata, Argentina.

⁵ Deceased.

lems of the Standard Model. Supersymmetry (SUSY) addresses some of these questions. For every particle it predicts the existence of a partner particle with spin differing by half a unit. SUSY models in general require at least two higgs doublets. The partners of the W^\pm and H^\pm mix to form two mass eigenstates, the charginos $\tilde{\chi}_{1,2}^\pm$. The partners of the γ , Z and the neutral higgs mix to form at least four mass eigenstates, the neutralinos [2] χ , χ' , χ'' and χ''' , in order of increasing mass.

In the Minimal Supersymmetric Standard Model (MSSM) [3], the Lagrangian at the unification scale is globally supersymmetric, except for a set of “soft breaking” mass terms. Among these are the gaugino masses M_1 , M_2 and M_3 associated with the $U(1)_Y$, $SU(2)_L$ and $SU(3)_C$ gauge groups, respectively. These mass terms are assumed to be equal at the unification scale, leading to $M_1 = \frac{5}{3}M_2 \tan^2\theta_w$ at the electroweak scale [4]. From naturalness arguments [5] it is expected that the gaugino mass parameter, $M \equiv M_2$, is in the range $0 \leq M \leq 250$ GeV, the higgsino mass parameters, μ , is bounded to $0 \leq |\mu| \leq 200$ GeV and the ratio of the vacuum expectation values of the two higgs doublets, $\tan\beta = v_2/v_1$, is constrained to $1 \leq \tan\beta \leq m_t/m_b \sim 50$, where v_2 gives mass to quarks of charge $2/3e$ and v_1 gives mass to charged leptons and quarks of charge $-1/3e$. In the MSSM, the masses and interactions of the neutralinos and charginos are entirely described in terms of $\tan\beta$ and the two mass parameters M and μ . We will make the usual assumption that χ is the lightest supersymmetric particle (LSP), which is stable by R-parity conservation and escapes detection due to its weakly interacting nature.

Many experiments have looked for supersymmetric particles, but so far no signature has been found. There exist mass limits up to 45 GeV for most of the supersymmetric particles [6] except for neutralinos. The other LEP experiments derived lower mass limits of 20 GeV and 40 GeV for the two lightest neutralinos if $\tan\beta \geq 2$ and no limit for $\tan\beta < 1.6$ [7]. The production cross section of neutralinos not only depends on their masses but also on their particle content and varies from almost 0, if one of the produced neutralinos is a photino, to several hundred picobarns. The analysis of a high statistics event sample, as presented here, is needed to derive conclusions.

We present the event selection and results of our search, which are based on 1.8×10^6 hadronic Z decays collected in 1991–1993. These results are interpreted in the MSSM context as well as in a more general way.

2. The L3 detector

The L3 detector [8,9] covers 99% of the 4π solid angle. It consists of a central tracking chamber (TEC), a forward–backward tracking chamber, a high resolution electromagnetic calorimeter (ECAL) composed of Bismuth Germanium Oxide (BGO) crystals, a ring of scintillation counters, a uranium brass hadron calorimeter with proportional wire chambers (HCAL), and a high-precision muon chamber spectrometer (MUCH). These detectors are located in a 12 m diameter magnet providing a uniform field of 0.5 T along the beam direction. Forward BGO arrays (LUMI) on each side of the detector measure the luminosity by detecting the energy deposit of small angle Bhabha events.

3. Event selection

We searched for neutralinos produced in the reaction

$$e^+ e^- \longrightarrow Z \longrightarrow \chi\chi' \text{ or } \chi'\chi'$$

with χ' decaying via

$$\chi' \longrightarrow \chi Z^* \longrightarrow \chi f\bar{f} \text{ or } \chi' \longrightarrow \chi\gamma.$$

The signature is missing energy due to the undetected χ and one or two photons, two or four acollinear and acoplanar leptons, or one to four hadronic jets from the primary quarks.

We assume that the masses of the lightest higgs, h^0 or A^0 , are charginos, $\tilde{\chi}_1^\pm$, are sufficiently large [10] so that the decays $\chi' \rightarrow \chi h^0$, χA^0 and $\chi' \rightarrow \tilde{\chi}_1^\pm f\bar{f}'$, are kinematically forbidden. However, the signature of such decays would be very similar to $\chi' \rightarrow \chi Z^* \rightarrow \chi q\bar{q}$ and could lead to even higher detection efficiencies. Because the SUSY partner of the electron, the selectron, is assumed to be heavier than 45 GeV [6], we neglect neutralino production via t-channel exchange, which is of the order of

10^{-3} relative to the s-channel contribution in the MSSM, if the neutralino is not a pure photino.

We used Monte Carlo generators to estimate the background arising from all Standard Model reactions. The main background sources for the fermionic neutralino decays are fermion pair production $e^+e^- \rightarrow q\bar{q}(\gamma)$ [12], $e^+e^- \rightarrow \tau^+\tau^-(\gamma)$ [13] and four-fermion processes $e^+e^- \rightarrow e^+e^-\bar{f}f$ [14]. The backgrounds for the radiative neutralino decays are $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ [15] and $e^+e^- \rightarrow \gamma\gamma(\gamma)$ [16]. The detector response of the final state particles is simulated with the GEANT package using the GHEISHA program for hadronic interactions [17]. The simulated events are reconstructed and analyzed in the same way as the real data.

To determine the efficiency for neutralino detection, we have simulated the neutralino production and decay based on the formulae given in Ref. [11]. The detection efficiency for neutralinos depends on their masses, their relative CP -sign and their decay mode. A finite set of neutralino mass combinations was fully simulated as described above. To interpolate between these points in the $m_\chi - m_{\chi'}$ mass plane, a fast simulation has been developed. Based on particle momenta, the detector response is evaluated taking detector (including trigger) efficiencies and resolutions into account. The agreement between the fast and full simulation is found to be better than 3%.

We have looked for the electroweak neutralino decay $\chi' \rightarrow \chi\bar{f}$, where $f = q, \mu, e$ and for the radiative decay $\chi' \rightarrow \chi\gamma$. Due to the low visible energy and the small branching fraction, we did not investigate decays involving τ leptons.

3.1. Hadronic final state

We select events with one or two jets in the final state, which may be interpreted as $Z \rightarrow \chi\chi' \rightarrow \chi\chi q\bar{q}$, $Z \rightarrow \chi'\chi' \rightarrow \chi q\bar{q} \chi\nu\bar{\nu}$, or $Z \rightarrow \chi'\chi' \rightarrow \chi q\bar{q} \chi q'\bar{q}'$ events. Calorimetric jets are reconstructed using the Durham clustering algorithm [19] with $y_{\text{cut}} = 0.04$ and are required to have at least one associated TEC track. Depending on the kinematics of the reaction, the two initial decay quarks may be constructed as one or two jets. We did not study the three and four-jet topology, because of the large background arising from Standard Model $q\bar{q}(g)$ events. Also, their relative fraction compared to one and two-jet




Fig. 1. Acoplanarity distribution of the hadronic two-jet events. The points correspond to the data, while the solid lines indicate the results of the Standard Model Monte Carlo simulation (MC). The dotted line shows a possible signal for $\chi - \chi'$ production with neutralinos of mass $m_\chi = 0$ GeV and $m_{\chi'} = 85$ GeV and a 2.5 nb cross section of production, while the dashed line corresponds to $m_\chi = 35$ GeV, $m_{\chi'} = 50$ GeV and a 0.5 nb cross section.

events was found to be small. For hadronic one-jet events we require:

- (1) The jet to have at least 10 GeV transverse momentum in order to reject four-fermion background.
- (2) The event to contain at least 4 TEC tracks to remove single jets originating from one or three single prong τ decays, where the other τ remains undetected.
- (3) There to be no TEC track in the $r-\phi$ plane in the hemisphere opposite to the jet direction, and the total calorimeter energy deposited in a cone of half opening angle of 30° around the missing energy direction to be less than 500 MeV. The contribution of most standard Z decays is therefore eliminated.

For hadronic two-jet events we require:

- (1) The acoplanarity and acollinearity angle of the two jets to exceed 40° to remove most standard Z decays (Fig. 1).
- (2) The total calorimetric energy deposited in a cone of half opening angle of 30° around the missing energy direction to be less than 500 MeV. There must be no TEC track within 30° of the missing energy direction in the $r-\phi$ plane. These requirements remove any remaining $q\bar{q}$ or $\tau^+\tau^-$ events.
- (3) We reduce the four-fermion background by requiring that the direction of the missing energy points more than 25° away from the beam axis, that

each jet has at least 3 GeV transverse momentum and that no more than 10 GeV is deposited in the luminosity monitors.

No events survive the one-jet selection, while three events are left after applying the two-jet selection cuts consistent with an expected background from Standard Model four-fermion processes of 0.9 ± 0.4 events.⁶

3.2. Muon final state

In order to select candidates of the type $Z \rightarrow \chi\chi' \rightarrow \chi\chi\mu^+\mu^-$, one of the two muons must be identified in the muon chambers and the other one either in the muon chambers or as an isolated TEC track. We reject cosmic rays as described in Ref. [20]. Further selection criteria are:

(1) The acoplanarity between the two muons has to exceed 40° , and the TEC track multiplicity has to be at most 2 to suppress the four-fermion background.

(2) The most energetic ECAL or LUMI cluster should not exceed 2 GeV in order to remove radiative dimuons and four-fermion events respectively. This requirement also ensures that a muon candidate which is only identified by a TEC track corresponds to a real muon.

(3) If both muons have been identified in the MUCH, the most energetic one has to have momentum $p > 6$ GeV. The missing transverse momentum of the event also has to exceed 6 GeV. In the case where one muon has been identified in the TEC, it should have a transverse momentum $p_T > 3$ GeV, while the other one (identified in the MUCH) should have $p > 10$ GeV. These cuts further reduce the four-fermion contribution.

No candidate events are observed after applying these cuts.

3.3. Electron final state

$Z \rightarrow \chi\chi' \rightarrow \chi\chi e^-e^+$ candidate events are selected by requiring at most three clusters in the

ECAL. The two most energetic clusters are associated with a TEC track and must have more than 3 GeV and 2 GeV of energy. The energy of a possible third cluster has to be below 0.5 GeV. Furthermore:

(1) The acoplanarity and acollinearity angle of the two most energetic clusters has to exceed 15° to reduce the four-fermion and Bhabha background.

(2) The missing transverse momentum has to exceed 6 GeV and point more than 12° away from the beam axis and more than 5° away from the closest ECAL cluster to further reduce the $\tau^+\tau^-$ and four-fermion contamination.

(3) The sum of the visible energy and missing momentum should not exceed the center-of-mass energy minus 5 GeV in order to suppress three-body final states with an undetected particle.

No events are left after applying these cuts.

3.4. Photon final states

Photonic final states may result from the $\chi' \rightarrow \chi\gamma$ decay. They are selected by allowing up to three electromagnetic clusters in the ECAL and no other significant detector activity. The selection criteria are [21]:

(1) The most energetic cluster in the LUMI should not exceed 5 GeV, while the most energetic cluster in the HCAL has to be below 3 GeV. This reduces the background from radiative Bhabha scattering.




Fig. 2. Transverse momentum distribution of the single-photon events. The points correspond to the data, while the solid lines indicates the results of a Standard Model Monte Carlo simulation (MC). The dotted line shows a possible signal for $\chi - \chi'$ production with neutralinos of mass $m_\chi = 0$ GeV and $m_{\chi'} = 85$ GeV and 1.2 pb production cross section, while the dashed line corresponds to $m_\chi = 35$ GeV, $m_{\chi'} = 50$ GeV and a 1.6 pb cross section.

⁶ The cross section of the four-fermion Monte Carlo events has been normalized to fit the observed distributions in hadronic two-jet events.

Table 1

The number of the observed events versus the number of expected background events for different neutralino decay channels

Signature	Events	
	Observed	Expected
one jet	0	< 0.5
two jets	3	0.9 ± 0.4
two muons	0	< 0.5
two electrons	0	< 0.5
one photon	13	15.7 ± 1.5
two photons	0	0.7 ± 0.5

(2) Radiative cosmic muon and four-fermion backgrounds are removed by requiring no tracks in the TEC or MUCH.

(3) Single-photon events are selected starting 20° away from the beam line. The transverse momentum of the photon has to exceed 10 GeV to suppress events from radiative neutrino production $\nu\bar{\nu}(\gamma)$ (Fig. 2).

(4) Events with two photons are selected allowing for a third cluster. To suppress the Standard Model $\gamma\gamma(\gamma)$ final states, we require that the two most energetic photons have less than 40 GeV of energy each and an acoplanarity in excess of 3.5° . The missing transverse momentum has to be above 6 GeV.

No events survive the two-photon selection. 13 events survive in the one-photon sample, in good

agreement with the Standard Model expectation from $\nu\bar{\nu}\gamma$ and $\gamma\gamma(\gamma)$ events of 15.7 ± 1.5 .

4. Results

No excess over the Standard Model expectation has been observed (Table 1). The efficiency to detect neutralinos is high in the case where the $\chi-\chi'$ mass difference is greater than 10 GeV (see Table 2 for details). It reaches 70% for the single-photon selection and up to 30%–60% for signatures with hadronic jets, muons, electrons or two photons. The trigger efficiency for events passing all our cuts is more than 90% for most of the final state configurations.

In the absence of a neutralino signal, we set limits on the branching fractions $\text{BR}(Z \rightarrow \chi\chi')$ and $\text{BR}(Z \rightarrow \chi'\chi')$, which do not depend on the neutralino coupling constants. We subsequently interpret these results in the MSSM framework, excluding regions of the parameter space as well as establishing limits on the neutralino masses.

We present upper limits on the branching ratios $\text{BR}(Z \rightarrow \chi\chi', \chi'\chi')$ as a function of the neutralino masses, using the relative neutralino CP -sign that leads to the lowest detection efficiency. The branching ratios $\chi' \rightarrow \chi Z^* \rightarrow \chi(e^+e^-, \mu^+\mu^-, q\bar{q}, \nu\bar{\nu})$ are calculated according to the Z partial width of these channels, while the neutralino decay width to pho-

Table 2

Total neutralino detection efficiencies in percent

Particle mass		Efficiency (%)					
$m_{\chi'}$ (GeV)	m_{χ} (GeV)	$\chi\chi q\bar{q}$	$\chi\chi e^+e^-$	$\chi\chi\mu^+\mu^-$	$\chi\chi\gamma$	$\chi q\bar{q}\chi$ ($q' \bar{q}, \nu\bar{\nu}$)	$\chi\gamma\chi\gamma$
5	0	25	50	42	50	9	37
10	5	35	50	40	43	12	43
30	0	36	51	50	60	15	61
30	25	8	24	17	9	4	14
45	0	36	53	50	71	10	67
45	35	9	30	23	20	7	53
55	0	32	51	47	73		
55	35	23	49	38	64		
75	0	27	45	41	76		
75	15	26	54	41	63		
90	0	27	26	43	76		

The quoted values correspond to the relative neutralino CP -sign with the lowest efficiency.




Fig. 3. Contour plot of the 95% C.L. upper limits on $\text{Br}(Z \rightarrow \chi\chi')$ versus the χ and χ' masses. The solid lines separate the regions, where the limit on the branching ratio is smaller than the value shown, while the dashed lines show the kinematical limits of the channel.

tons is assumed to be unknown. We then vary the relative photonic branching ratio between 0 and 1 and quote the highest, most conservative, limit obtained. The different channels are combined to calculate limits on the $Z \rightarrow \chi\chi'$ and $Z \rightarrow \chi'\chi'$ decay modes taking the different branching ratios, detection efficiencies, numbers of candidates and expected background events properly into account using Poisson statistics in the Bayesian approach [22]. The 95% confidence level (C.L.) upper limits on $\text{BR}(Z \rightarrow \chi\chi')$ and $\text{BR}(Z \rightarrow \chi'\chi')$ are shown in Figs. 3 and 4.




Fig. 4. Contour plot of the 95% C.L. upper limits on $\text{Br}(Z \rightarrow \chi\chi')$ versus the χ and χ' masses. The solid lines separate the regions, where the limit on the branching ratio is smaller than the value shown, while the dashed lines show the kinematical limits of the channel.




Fig. 5. The excluded regions of the MSSM parameter space at the 95% C.L. as a function of the parameters M and μ . The kinematical limit corresponds to the sum of the lightest and next to lightest neutralino masses being equal to the center-of-mass energy. The exclusion coming from the lineshape measurement goes beyond the kinematic limit of the direct search.

4.1. Limits within the MSSM

In addition to the limits on $\text{BR}(Z \rightarrow \chi\chi')$ and $\text{BR}(Z \rightarrow \chi'\chi')$, we use the constraints coming from the precise LEP Z lineshape measurements⁷

$$\Delta\Gamma_Z < 23.1 \text{ MeV (95\% C.L.)}$$

$$\Delta\Gamma_{\text{inv}} < 8.4 \text{ MeV (95\% C.L.)}$$

to restrict the MSSM parameter space using

$$\begin{aligned} \Delta\Gamma_Z = & \Gamma(Z \rightarrow \chi\chi) + \Gamma(Z \rightarrow \chi\chi') + \Gamma(Z \rightarrow \chi\chi'') \\ & + \Gamma(Z \rightarrow \chi\chi''') + \Gamma(Z \rightarrow \chi'\chi') \\ & + \Gamma(Z \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^-) \end{aligned}$$

$$\Delta\Gamma_{\text{inv}} = \Gamma(Z \rightarrow \chi\chi).$$

⁷ The limits on $\Delta\Gamma_Z$ and $\Delta\Gamma_{\text{inv}}$ are obtained with the method described in Ref. [9] using the results given by Ref. [23]. We have used $m_{H^0} = 1000 \text{ GeV}$, $m_t = 131 \text{ GeV}$, $\alpha_s = 0.117$, $m_Z = 91.180 \text{ GeV}$, $\Gamma_Z = 2490 \pm 7 \text{ MeV}$ and $\Gamma_{\text{inv}} = 498.2 \pm 4.2 \text{ MeV}$, which are the values that give the most conservative constraints.

Table 3
Lower neutralino mass limits (95% C.L.) in GeV

Particle	tan β			$m_{\tilde{g}} > 100$ GeV all tan β
	> 1	> 2	> 3	
χ	0	20	23	18
χ'	0	46	52	20
χ''	60	78	84	60
χ'''	90	115	127	98

Changing the limits on $\Delta\Gamma_Z$ and $\Delta\Gamma_{\text{inv}}$ by a factor of two does not change the excluded MSSM parameter space significantly. The excluded regions for different values of tan β are shown in Fig. 5. For moderate or high values of tan β , a significant part of the accessible parameter space is excluded.

All neutralino masses are functions of the parameters M , μ and tan β . Therefore, constraints on the MSSM parameter space translate into limits on these masses, summarized in Table 3. The dependence on tan β is illustrated in Fig. 6.

These neutralino mass limits can be further improved by using a limit on the gluino mass $m_{\tilde{g}} > 100$ GeV [24], as suggested by Hidaka [25], which limits the parameter M via

$$m_{\tilde{g}} = \frac{\alpha_s}{\alpha} \sin^2 \theta_w M \quad \text{thus} \quad M \approx 0.3 m_{\tilde{g}} > 30 \text{ GeV.}$$

This further restriction leads to mass limits for all tan β values, which are also shown in Table 3.




Fig. 6. Neutralino lower mass limits (95% C.L.). The lines correspond to the lower mass limit for the different neutralinos. The regions below the lines are excluded.

5. Conclusion

Using the 1991–1993 data of the L3 experiment, we have searched for neutralinos with a large variety of event signatures. No evidence for neutralinos was found and upper limits of a few times 10^{-5} have been set on the branching ratio for Z decaying to $\chi\chi'$ or $\chi'\chi'$. A significant part of the MSSM parameter space accessible at LEP has been excluded. In this paper the branching ratio limits are significantly improved compared to earlier results [7] and the dependence on assumptions on the neutralino decay modes has been minimized. In the MSSM the lightest neutralino is found to be heavier than 18 GeV, if either tan $\beta > 2$ or $m_{\tilde{g}} > 100$ GeV.

Acknowledgments

We wish to express our gratitude to the CERN accelerator division for the excellent performance of the LEP machine. We acknowledge the efforts of all engineers and technicians who have participated in the construction and maintenance of this experiment.

References

- [1] S.L. Glashow, Nucl. Phys. 22 (1961) 579; S. Weinberg, Phys. Rev. Lett. 19 (1967) 1264; A. Salam, Elementary Particle Theory, ed. N. Svartholm (Almquist and Wiksell, Stockholm, 1968) p. 367.
- [2] R. Barbieri et al., Z Physics at LEP1, CERN 89-08 (1989) Vol. 2, 121; J.M. Frere and G.L. Kane, Nucl. Phys. B 223 (1983) 331; J. Ellis et al., Phys. Lett. B 127 (1983) 233; G. Gamberini, Z. Phys. C 30 (1986) 605; J. Ellis et al., Phys. Lett. B 123 (1983) 436.
- [3] Y.A. Golfand and E.P. Likhthman, JETP Lett. 13 (1971) 323; D.V. Volkhov and V.P. Akulov, Phys. Lett. B 46 (1973) 109; J. Wess and B. Zumino, Nucl. Phys. B 70 (1974) 39; P. Fayet and S. Ferrara, Phys. Rep. 32 (1977) 249; A. Salam and J. Strathdee, Fortschr. Phys. 26 (1978) 57.
- [4] For reviews, see H. Nilles, Phys. Rep. 110 (1984) 1; R. Barbieri, Riv. Nuovo Cimento 11, No. 4 (1988); R. Barbieri et al., Nucl. Phys. B 296 (1988) 75.
- [5] J. Ellis et al., Mod. Phys. Lett. A 1 (1986) 57; R. Barbieri and G.F. Giudice, Nucl. Phys. B 306 (1988) 63; Z. Kunszt and F. Zwirner, CERN-TH.6150/91, ETH-TH/91-7 (December 1991).

- [6] J. Ellis et al., CERN-PPE/92-180.
- [7] ALEPH Collab., D. Decamp et al., Phys. Rev. Lett. B 244 (1990) 541; Phys. Rep. 216 (1992) 253; DELPHI Collab., P. Abreu et al., Phys. Lett. B 247 (1990) 157; OPAL Collab., K. Ahmet et al., Phys. Rev. Lett. B 248 (1990) 211.
- [8] L3 Collab., B. Adeva et al., Nucl. Instrum. Methods A 289 (1990) 35.
- [9] L3 Collab., O. Adriani et al., Phys. Rep. 236 (1993) 1.
- [10] ALEPH Collab., D. Decamp et al., Phys. Lett. B 236 (1990) 86; B 237 (1990) 291.
- [11] A. Bartl, H. Fraas and W. Majerotto, Nucl. Phys. B 278 (1986) 1; H.E. Haber and D. Wyler, Nucl. Phys. B 323 (1989) 267.
- [12] T. Sjöstrand and M. Bengtsson, JETSET 7.3, Comput. Phys. Commun. 43 (1987) 367; M. Böhm, A. Denner and W. Hollik, Nucl. Phys. B 304 (1988) 687.
- [13] S. Jadach et al., KORALZ, Z Physics at LEP 1, eds. G. Altarelli et al., CERN Report CERN-89-08, Vol. 3 CERN, Geneva (1989).
- [14] F.A. Berends et al., DIAG 36; Nucl. Phys. B 253 (1985) 441.
- [15] R. Miquel et al., Z. Phys. C 48 (1990) 309; F.A. Berends et al., Nucl. Phys. B 301 (1988) 583.
- [16] F.A. Berends and R. Kleiss, Nucl. Phys. B 186 (1981) 22.
- [17] R. Brun et al., GEANT 3.15, CERN-DD/78-2; H. Fesefeld, GHEISHA, RWTH Aachen Preprint PITHA 85/02 (1985).
- [18] P. Béné et al., Nucl. Instrum. Methods A 306 (1991) 150; R. Bizzarri et al., A 317 (1992) 463; P. Bagnaia et al., A 324 (1993) 101.
- [19] S. Bethke et al., Nucl. Phys. B 370 (1992) 310.
- [20] L3 Collab., O. Adriani et al., Z. Phys. C 62 (1994) 551.
- [21] A. Weber, Thesis, RWTH Aachen Preprint PITHA 94/18 (1994), for further details.
- [22] A.G. Frodesen, O. Skjeggstad and H. Töfte, Probability and statistics in Particle Physics (Universitetsforlaget, Bergen, 1979).
- [23] L. Montanet et al., Particle Oala Group, Review of Particle Properties, Phys. Rev. D 50 (1994) 1173.
- [24] CDF Collab., F. Abe et al., Phys. Rev. Lett. 69 (1992) 3439; H. Baer, X. Tata and J. Woodside, Phys. Rev. D 44 (1991) 207.
- [25] K. Hidaka, Phys. Rev. D 44 (1991) 927; I. Antoniadis, J. Ellis and D.V. Nanopoulos, Phys. Lett. B 262 (1991) 109.