

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/29134>

Please be advised that this information was generated on 2019-04-22 and may be subject to change.

Note

# Synthesis and X-ray analysis of 1-((1S)-phenylethyl)-azetidine-(2R)-piperidinamide

R. de Gelder,<sup>(1)\*</sup> J. M. M. Smits,<sup>(1)</sup> W. A. J. Starman,<sup>(2)</sup> L. Thijs,<sup>(2)</sup> and B. Zwanenburg<sup>(2)</sup>

Received February 2, 1996


The crystal and molecular structure of a new azetidine-2-carboxylic amide derivative is described. The structure was solved by direct methods and refined by least squares methods to  $R1 = 0.0393$  for 4264 reflections (with  $I > 2\sigma(I)$ ). The structure consists of two independent molecules which are chemically the same with slight differences in geometry. Crystal data:  $C_{17}H_{24}N_2O$ , monoclinic, space group  $P2_1$ ,  $a = 8.3782(4)$ ,  $b = 20.0342(13)$ ,  $c = 9.7769(8)\text{\AA}$ ,  $\beta = 109.687(6)^\circ$ ,  $V = 1545.1(2)\text{\AA}^3$ ,  $Z = 4$ .

**KEY WORDS:** Crystal structure; chiral auxiliaries.

## Introduction

Optically active azetidine-2-carboxylic acid is a versatile starting material for chiral auxiliaries<sup>1a,b</sup> and was first synthesized in enantiomerically pure form by

tion.<sup>3b</sup> We now describe an improved route towards two stereoisomers of azetidine-2-carboxylic ester derivatives derived from both enantiomers of the parent acid using optically active (S)-( $\alpha$ )-methyl-benzylamine in stoichiometric amounts (Scheme 1).



Miyoshi *et al.*<sup>2</sup> An easier route by Rodebaugh *et al.*<sup>3a</sup> afforded racemic azetidine-2-carboxylic acid which still had to be separated into its enantiomers by resolu-

Methyl 2,4-dibromobutyrate 1 was ringclosed with 1.1 equivalent (S)-( $\alpha$ )-methyl-benzylamine and 2 equivalents  $K_2CO_3$  in acetonitrile/water heated at reflux temperature. A 1:1 mixture of diastereomeric methyl 1-((1S)-phenylethyl)azetidine-2-carboxylate 2 was obtained which was separated by flash-column chromatography. The configuration of thus obtained diastereomers could not be determined by spectroscopic methods and therefore one of the diastereomers was transformed into the crystalline piperidinamide 3.<sup>4</sup> An X-ray diffraction analysis was undertaken to

<sup>(1)</sup> Department of Inorganic Chemistry, Nijmegen SON Research Center, University of Nijmegen, Toernooiveld, 6525 ED Nijmegen, The Netherlands.

<sup>(2)</sup> Department of Organic Chemistry, Nijmegen SON Research Center, University of Nijmegen, Toernooiveld, 6525 ED Nijmegen, The Netherlands.

\* To whom correspondence should be addressed.

**Table 1.** Crystal data and summary of intensity data collection, structure solution, and refinement

Crystal data	
Compound	C <sub>17</sub> H <sub>24</sub> N <sub>2</sub> O
Color/shape	Colorless/irregular
Crystallization	Hexane
Formula weight	272.38
Crystal system	Monoclinic
Space group	P 2 <sub>1</sub>
Temperature, K	208(2)
Cell constants <sup>a</sup>	
<i>a</i> , Å	8.3782(4)
<i>b</i> , Å	20.0342(13)
<i>c</i> , Å	9.7769(8)
β, °	109.687(6)
Cell volume; Å <sup>3</sup>	1545.1(2)
Formula units/unit cell	4
<i>D</i> -calc, g·cm <sup>-3</sup>	1.171
Mu-calc, cm <sup>-1</sup>	0.682
<i>F</i> (000), electrons	592
Intensity data collection	
Diffractometer/scan	Enraf-Nonius CAD-4/ω-scan
Radiation, graphite monochromator	MoKα (λ = 0.71073 Å)
Crystal dimensions, mm	0.65 × 0.51 × 0.18
Scan width, °	1.5
Standard reflections	3, every 7200 seconds exposure time
Decay of standards	1.000 – 0.979
Reflections measured	6473
2θ-range, °	up to 49.94
Range of <i>h, k, l</i>	–9 ≤ <i>h</i> ≤ 1, –23 ≤ <i>k</i> ≤ 23, –11 ≤ <i>l</i> ≤ 11
Corrections	
Decay correction	
Lorentz-polarization	
EMPABS <sup>7</sup> correction	0.988 – 1.013
Reflns (obs., <i>I</i> <sub>0</sub> > 2σ( <i>I</i> <sub>0</sub> ))	5410(4264)
Rmerge <sup>b</sup>	0.0185
Computer programs <sup>c</sup>	Local programs
Structure solution and refinement	
Structure solution	Direct methods/automatic Fourier extension
Computer programs	CRUNCH <sup>5</sup>
Structure refinement	Full matrix least squares on <i>F</i> <sup>2</sup>
Non-H-atoms	anisotropic
H-atoms	isotropic
Computer program	SHELXL <sup>8</sup>
Weights <sup>8</sup>	0.0286 0.1150
Shift/esd	less than 0.001
No. of restraints/parameters	1/553
Goodness-of-fit on <i>F</i> <sup>2</sup>	1.103
<i>R</i> index [ <i>I</i> > 2σ( <i>I</i> )]	<i>R</i> 1 = 0.0393
<i>R</i> indices (all data)	<i>R</i> 1 = 0.0612, <i>wR</i> 2 = 0.0798,
Largest diff. peak and hole, e·Å <sup>-3</sup>	0.12 and –0.14

**Table 2.** Atomic coordinates (× 10<sup>4</sup>) and equivalent isotropic displacement parameters (Å<sup>2</sup> × 10<sup>3</sup>) for 3<sup>a</sup>

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> <sub>(eq)</sub>
Molecule A				
N(1A)	–6335(2)	599(1)	–12213(2)	41(1)
C(2A)	–4557(3)	828(1)	–11920(3)	43(1)
C(3A)	–4642(5)	545(2)	–13402(3)	77(1)
C(4A)	–6576(4)	568(2)	–13784(3)	65(1)
C(5A)	–7467(3)	1063(1)	–11831(3)	40(1)
C(6A)	–9263(4)	783(2)	–12369(4)	59(1)
C(7A)	–6873(3)	1183(1)	–10207(2)	35(1)
C(8A)	–6283(3)	667(1)	–9218(3)	44(1)
C(9A)	–5830(3)	786(2)	–7745(3)	53(1)
C(10A)	–5957(3)	1418(2)	–7238(3)	55(1)
C(11A)	–6524(3)	1931(2)	–8210(3)	52(1)
C(12A)	–6972(3)	1818(1)	–9686(3)	41(1)
C(13A)	–3246(3)	496(1)	–10642(3)	43(1)
O(14A)	–3312(2)	–108(1)	–10481(2)	73(1)
N(15A)	–1963(2)	857(1)	–9767(2)	38(1)
C(16A)	–636(4)	525(1)	–8596(3)	51(1)
C(17A)	–703(4)	740(2)	–7136(3)	54(1)
C(18A)	–523(4)	1487(1)	–6972(3)	51(1)
C(19A)	–1859(4)	1828(1)	–8227(3)	49(1)
C(20A)	–1819(4)	1584(1)	–9677(3)	43(1)
Molecule B				
N(1B)	–7511(2)	–1608(1)	–9573(2)	31(1)
C(2B)	–5706(3)	–1406(1)	–8987(2)	34(1)
C(3B)	–5685(3)	–1395(2)	–10556(3)	46(1)
C(4B)	–7593(3)	–1323(1)	–10997(2)	38(1)
C(5B)	–8626(3)	–1320(1)	–8862(2)	37(1)
C(6B)	–10457(3)	–1499(2)	–9739(3)	50(1)
C(7B)	–8099(3)	–1565(1)	–7312(2)	38(1)
C(8B)	–8038(3)	–2239(1)	–7005(3)	47(1)
C(9B)	–7499(4)	–2463(2)	–5587(3)	56(1)
C(10B)	–7046(3)	–2018(2)	–4459(3)	55(1)
C(11B)	–7108(4)	–1348(2)	–4733(3)	55(1)
C(12B)	–7628(3)	–1122(1)	–6154(3)	44(1)
C(13B)	–4523(3)	–1882(1)	–7912(2)	33(1)
O(14B)	–4854(2)	–2478(1)	–7940(2)	49(1)
N(15B)	–3077(2)	–1631(1)	–6971(2)	39(1)
C(16B)	–2017(4)	–2041(1)	–5761(3)	48(1)
C(17B)	–2160(4)	–1785(1)	–4351(3)	50(1)
C(18B)	–1720(4)	–1045(1)	–4136(3)	51(1)
C(19B)	–2757(3)	–642(1)	–5458(3)	42(1)
C(20B)	–2565(3)	–929(1)	–6820(3)	38(1)

<sup>a</sup> *U*<sub>(eq)</sub> is defined as one third of the trace of the orthogonalized *U*<sub>*ij*</sub> tensor.

<sup>a</sup> Least-squares refinement for 25 reflections 11.527 < θ < 17.844°.

<sup>b</sup> Rmerge = Σ[*F*<sub>0</sub> – ⟨*F*<sub>0</sub>⟩]/Σ*F*<sub>0</sub>.

<sup>c</sup> Using neutral scattering factors and anomalous dispersion corrections.<sup>10</sup>

Table 3. Bond lengths (Å) and angles (°) for 3

Molecule A		Molecule B	
N(1A)–C(2A)	1.491(3)	N(1B)–C(2B)	1.481(3)
N(1A)–C(4A)	1.481(3)	N(1B)–C(4B)	1.485(3)
N(1A)–C(5A)	1.463(3)	N(1B)–C(5B)	1.458(3)
C(2A)–C(3A)	1.534(4)	C(2B)–C(3B)	1.540(3)
C(2A)–C(13A)	1.512(3)	C(2B)–C(13B)	1.515(3)
C(3A)–C(4A)	1.536(5)	C(3B)–C(4B)	1.516(3)
C(5A)–C(6A)	1.523(4)	C(5B)–C(6B)	1.527(4)
C(5A)–C(7A)	1.514(3)	C(5B)–C(7B)	1.510(3)
C(7A)–C(8A)	1.388(3)	C(7B)–C(8B)	1.382(3)
C(7A)–C(12A)	1.384(3)	C(7B)–C(12B)	1.387(3)
C(8A)–C(9A)	1.381(4)	C(8B)–C(9B)	1.379(3)
C(9A)–C(10A)	1.377(4)	C(9B)–C(10B)	1.369(4)
C(10A)–C(11A)	1.372(4)	C(10B)–C(11B)	1.365(4)
C(11A)–C(12A)	1.382(4)	C(11B)–C(12B)	1.385(4)
O(13A)–C(14A)	1.225(3)	O(13B)–C(14B)	1.225(3)
N(13A)–C(15A)	1.338(3)	N(13B)–C(15B)	1.348(3)
N(15A)–C(16A)	1.461(3)	N(15B)–C(16B)	1.468(3)
N(15A)–C(20A)	1.461(3)	N(15B)–C(20B)	1.462(3)
C(16A)–C(17A)	1.510(4)	C(16B)–C(17B)	1.513(4)
C(17A)–C(18A)	1.507(4)	C(17B)–C(18B)	1.525(4)
C(18A)–C(19A)	1.516(4)	C(18B)–C(19B)	1.523(3)
C(19A)–C(20A)	1.510(4)	C(19B)–C(20B)	1.508(3)
N(1A)–C(2A)–C(3A)	88.9(2)	N(1B)–C(2B)–C(3B)	88.3(2)
N(1A)–C(2A)–C(13A)	115.1(2)	N(1B)–C(2B)–C(13B)	116.1(2)
N(1A)–C(4A)–C(3A)	89.1(2)	N(1B)–C(4B)–C(3B)	89.1(2)
N(1A)–C(5A)–C(6A)	109.2(2)	N(1B)–C(5B)–C(6B)	108.9(2)
N(1A)–C(5A)–C(7A)	110.9(2)	N(1B)–C(5B)–C(7B)	109.4(2)
C(2A)–N(1A)–C(4A)	89.5(2)	C(2B)–N(1B)–C(4B)	88.7(2)
C(2A)–N(1A)–C(5A)	116.6(2)	C(2B)–N(1B)–C(5B)	116.2(2)
C(2A)–C(3A)–C(4A)	85.9(2)	C(2B)–C(3B)–C(4B)	85.4(2)
C(2A)–C(13A)–O(14A)	119.2(2)	C(2B)–C(13B)–O(14B)	120.6(2)
C(2A)–C(13A)–N(15A)	119.6(2)	C(2B)–C(13B)–N(15B)	117.8(2)
C(3A)–C(2A)–C(13A)	114.1(2)	C(3B)–C(2B)–C(13B)	118.4(2)
C(4A)–N(1A)–C(5A)	114.7(2)	C(4B)–N(1B)–C(5B)	117.4(2)
C(5A)–C(7A)–C(8A)	121.7(2)	C(5B)–C(7B)–C(8B)	120.8(2)
C(5A)–C(7A)–C(12A)	119.5(2)	C(5B)–C(7B)–C(12B)	121.2(2)
C(6A)–C(5A)–C(7A)	111.0(2)	C(6B)–C(5B)–C(7B)	112.7(2)
C(7A)–C(8A)–C(9A)	120.4(3)	C(7B)–C(8B)–C(9B)	120.8(3)
C(7A)–C(12A)–C(11A)	120.4(2)	C(7B)–C(12B)–C(11B)	121.1(3)
C(8A)–C(7A)–C(12A)	118.7(2)	C(8B)–C(7B)–C(12B)	117.9(2)
C(8A)–C(9A)–C(10A)	120.5(3)	C(8B)–C(9B)–C(10B)	120.4(3)
C(9A)–C(10A)–C(11A)	119.4(3)	C(9B)–C(10B)–C(11B)	120.0(3)
C(10A)–C(11A)–C(12A)	120.6(3)	C(10B)–C(11B)–C(12B)	119.8(3)
C(13A)–N(15A)–C(16A)	119.3(2)	C(13B)–N(15B)–C(16B)	120.2(2)
C(13A)–N(15A)–C(20A)	127.5(2)	C(13B)–N(15B)–C(20B)	126.4(2)
O(14A)–C(13A)–N(15A)	120.9(2)	O(14B)–C(13B)–N(15B)	121.6(2)
N(15A)–C(16A)–C(17A)	110.5(2)	N(15B)–C(16B)–C(17B)	109.5(2)
N(15A)–C(20A)–C(19A)	110.6(2)	N(15B)–C(20B)–C(19B)	109.9(2)
C(16A)–N(15A)–C(20A)	112.5(2)	C(16B)–N(15B)–C(20B)	111.4(2)
C(16A)–C(17A)–C(18A)	110.5(2)	C(16B)–C(17B)–C(18B)	111.4(2)
C(17A)–C(18A)–C(19A)	109.9(2)	C(17B)–C(18B)–C(19B)	110.8(2)
C(18A)–C(19A)–C(20A)	111.7(2)	C(18B)–C(19B)–C(20B)	110.3(2)




Fig. 1. ORTEP<sup>11</sup> drawing of the molecule with atomic numbering for the two independent molecules. Thermal ellipsoids are at 50% probability.

establish the configuration at C-2 relative to the (1*S*)-phenylethyl group. It was found that the analyzed compound had the *R*-configuration at C-2.

## Experimental

The crystal data and a summary of the data collection, the structure solution and refinement are given in Table 1. The atomic positional and vibrational parameters are given in Table 2. The structure was solved by the program CRUNCH.<sup>5</sup> The hydrogen atoms were placed at calculated positions and were freely refined subsequently.

## Discussion

The structure and atomic numbering are presented in Fig. 1. Bond distances and angles are given in Table

Table 4. Selected torsion angles (°)

	Molecule A	Molecule B
C(3)–C(2)–C(13)–O(14)	–57.04(0.38)	–76.91(0.29)
C(2)–C(13)–N(15)–C(20)	14.58(0.39)	3.81(0.33)
O(14)–C(13)–N(15)–C(16)	–0.72(0.38)	–12.25(0.34)
C(2)–N(1)–C(5)–C(6)	173.24(0.22)	171.72(0.20)
C(2)–N(1)–C(5)–C(7)	–64.17(0.25)	–64.66(0.23)
N(1)–C(5)–C(7)–C(8)	–41.88(0.29)	–57.46(0.28)

3. The crystal structure consists of two independent molecules which are chemically the same with slight differences in geometry. In Table 4 selected torsion angles are given which illustrate the geometrical differences between both molecules. Geometrical calculations revealed neither unusual geometric features, nor unusual short intermolecular contacts. Calculations with PLATON<sup>6</sup> revealed no higher symmetry and no solvent accessible areas.

**Supplementary Material.** Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication CCDC-1003/5050. Copies of available material can be obtained, free of charge, on application to the Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK, (fax: +44-(0)1223-336033 or e-mail: teched@chemcrs.cam.ac.uk).

## References

- (a) Behnen, W.; Dauelsberg, Ch.; Wallbaum, S.; Martens, J. *Synth. Commun.* **1992**, *22*, 2143. b) Behnen, W.; Mehler, Th.; Martens, J. *Tetrahedron Asymm.* **1993**, *4*, 1413.
- Miyoshi, M.; Sugano, H.; Fujii, T.; Ishihara, T.; Yoneda, N. *Chem. Lett.* **1973**, 5.
- (a) Rodebaugh, R.M.; Cromwell, N.H. *J. Heterocycl. Chem.* **1969**, *6*, 435. (b) Rodebaugh, R.M.; Cromwell, N.H. *J. Heterocycl. Chem.* **1969**, *6*, 993.
- Matsumoto, K.; Hashimoto, S.; Uchida, T.; Okamoto, T.; Otani, S. *Chem. Ber.* **1989**, *122*, 1357.
- Gelder, R. de; Graaff, R.A.G. de; Schenk, H. *Acta Crystallogr.* **1993**, *A49*, 287.
- Spek, A.L. (1995) PLATON, Vakgroep Kristal en Structuurchemie, University of Utrecht.
- North, A.C.T.; Philips, D.C.; Mathews, F.S. *Acta Crystallogr.* **1968**, *A24*, 351.
- Sheldrick, G.M.; (1993) *SHELXL-93, Program for the Refinement of Crystal Structures*; Univ. of Goettingen: Germany.
- Walker, N.; Stuart, D. *Acta Crystallogr.* **1983**, *A39*, 158.
- International Tables for X-ray Crystallography*, Vol. IV; Birmingham: Kynoch Press, 1974.
- Spek, A.L. ORTEP, part of PLATON.<sup>6</sup>