

PAPER • OPEN ACCESS

Is the present acceleration of the Universe caused by merging with other universes?

To cite this article: J. Ambjørn and Y. Watabiki JCAP12(2023)011

View the [article online](#) for updates and enhancements.

You may also like

- [Self-Purification and Entanglement Revival in Lambda Matter](#)
Dongni Chen, Stefano Chesi and Mahn-Soo Choi
- [New Active Jupiter Family Comet 2008 QZ44: a Discovery with Citizen Science](#)
Colin Orion Chandler, William J. Oldroyd, Chadwick A. Trujillo et al.
- [Phononic band calculations and experimental imaging of topological boundary modes in a hexagonal flexural wave machine](#)
Hayato Takeda, Ryoya Minami, Osamu Matsuda et al.

Is the present acceleration of the Universe caused by merging with other universes?

J. Ambjørn^{a,b} and Y. Watabiki^c

^aThe Niels Bohr Institute, Copenhagen University,
Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark

^bInstitute for Mathematics, Astrophysics and Particle Physics (IMAPP),
Radboud University Nijmegen,
Heyendaalseweg 135, 6525 AJ, Nijmegen, The Netherlands

^cTokyo Institute of Technology, Dept. of Physics, High Energy Theory Group,
2-12-1 Oh-okayama, Meguro-ku, Tokyo 152-8551, Japan

E-mail: ambjorn@nbi.dk, watabiki@th.phys.titech.ac.jp

Received August 30, 2023

Revised October 25, 2023

Accepted October 26, 2023

Published December 12, 2023

Abstract. We show that by allowing our Universe to merge with other universes one is lead to modified Friedmann equations that explain the present accelerated expansion of our Universe without the need of a cosmological constant.

Keywords: quantum cosmology, quantum gravity phenomenology

ArXiv ePrint: [2308.10924](https://arxiv.org/abs/2308.10924)



Contents

1	Introduction	1
2	Expansion by merging with other universes	2
3	Including matter	5
4	Comparison of the different models	8
5	Discussion	13
A	Analytic results for the three models	15
A.1	The f_{ds} -model	15
A.2	The f_{gcdt} -model	16
A.3	The f_{mod} -model	16
A.4	Duality in the f_{mod} -model	17

1 Introduction

We have proposed a modified Friedmann equation [1–3] which changes the late time cosmology, such that one does not need a cosmological constant to explain the present day acceleration of our Universe. While it is our hope that this modified Friedmann equation can eventually be derived from an underlying microscopic theory [4–7], we will here treat it as a phenomenological model that we can obtain in a simple way from the standard Friedmann equation.¹

Our starting point is the Hartle-Hawking minisuperspace action, which after the rotation of the conformal factor can be written as

$$S = \int dt \left(\frac{\beta \dot{v}^2}{2Nv} + \lambda Nv \right). \quad (1.1)$$

Before rotating to Euclidean spacetime and the rotation of the conformal factor, this action is just the standard minisuperspace action

$$S = \int dt \left(-\frac{\beta \dot{v}^2}{2Nv} - \lambda Nv \right), \quad (1.2)$$

written using the metric

$$ds^2 = -N^2(t)dt^2 + a^2(t)d\Omega_d, \quad d\Omega_d = \sum_{i=1}^d dx_i^2. \quad (1.3)$$

In (1.1) and (1.2) we are using units where $c = \hbar = 1$ and the constant $\beta = (d-1)/d$, where d is the dimension of space. We have used

$$v(t) = \frac{1}{\kappa} a^d(t), \quad \kappa = 8\pi G, \quad G = \text{the gravitational constant}. \quad (1.4)$$

¹It should be noted that ideas somewhat related to the ones presented here and in [1–7] have also been advocated in [8].

as variable rather than the scale factor $a(t)$. $v(t)$ is proportional to the spatial d -volume at time t and below we will just call it the d -volume. For the Hubble parameter $H(t)$ we then have

$$H(t) \equiv \frac{\dot{a}}{a} = \frac{1}{d} \frac{\dot{v}}{v}. \quad (1.5)$$

The reason we prefer to use $v(t)$ instead of $a(t)$ is that the minisuperspace action (1.1)–(1.2) is then valid in all space dimensions $d > 1$. Finally λ denotes the cosmological constant. Presently we will ignore matter, only assume a cosmological constant term. Below we will include matter in our considerations.

It should be noted that (1.1) also appears as the leading term in an effective action in a model of four-dimensional quantum gravity known as Causal Dynamical Triangulations (CDT). In that case one is not assuming a minisuperspace reduction, but performs the path integral over all degrees of freedom except $v(t)$. Thus (1.1) might be more general than suggested by the minisuperspace reduction. This CDT result is a numerical result, obtained via Monte Carlo Simulations of the CDT lattice gravity model where one identifies G/ε^2 and $\lambda\varepsilon^4$ with the corresponding dimensionless lattice coupling constants, ε denoting the length of the lattice links, i.e. the UV lattice cut-off (see [9, 10], and [11, 12] for reviews). Quite remarkably, for CDT in $d = 1$, (1.1) can be derived analytically (with $\beta \neq 0$), and it is an effective action coming entirely from the path integral measure [13], since the classical Einstein action is a topological invariant in the case of $d = 1$.

The classical Hamiltonian corresponding to (1.2) is²

$$\mathcal{H}(v, p) = Nv \left(-\frac{p^2}{2\beta} + \lambda \right), \quad (1.6)$$

where p denotes the momentum conjugate to v . In the following we will be interested in $d = 3$, i.e. $2\beta = 4/3$.

A classical solution corresponding to the action (1.2) is the de Sitter spacetime³

$$v(t) = v(t_0) e^{\sqrt{3\lambda}(t-t_0)}. \quad (1.7)$$

We now want to go beyond this classical picture, but staying as close as possible to the minisuperspace picture, by trying to include the possibility that our Universe can absorb other universes, which we denote baby universes even if they are not necessarily small.

2 Expansion by merging with other universes

Since we will allow for other universes to merge with our Universe, we are really discussing a multi-universe theory and like in a many-particle theory it is natural to introduce creation and annihilation operators $\Psi^\dagger(v)$ and $\Psi(v)$ for single universes of spatial volume v . In a full theory of four-dimensional quantum gravity the spatial volume alone will of course not

²Since there is no time derivative of $N(t)$ in (1.2), we should strictly speaking treat it as a constraint system and we can choose as a Hamiltonian $\mathcal{H}_u = \mathcal{H} + u(t)P_N$, where P_N is the momentum conjugate to N , and impose the constraint $P_N = 0$ in phase space. This leads to $\mathcal{H} = 0$ on the constraint surface as a consistent secondary constraint and $\dot{N} = u(t)$, expressing the invariance with respect to time reparametrization. In the following we will use this invariance to choose N constant ($N = 1$), and then consider “on shell” solutions $\mathcal{H} = 0$.

³Note that the solution of $\mathcal{H}(v, p) = 0$ corresponding to an expanding universe has $p = -\sqrt{\lambda} < 0$. The reason for the somewhat counter-intuitive negative values of p is the negative sign of the “kinetic” terms in (1.2) and (1.6).

provide a complete characterization of a state at a given time t . As mentioned we will here make the drastic simplification to work in a minisuperspace approximation where the spatial universe *is* characterized entirely by the spatial volume v . Thus, denote the quantum state of a spatial universe with volume v by $|v\rangle$. We consider now the multi-universe Fock space constructed from such single universe states and denote the Fock vacuum state by $|0\rangle$. Then

$$[\Psi(v), \Psi^\dagger(v')] = \delta(v - v'), \quad \Psi^\dagger(v)|0\rangle = |v\rangle, \quad \Psi(v)|0\rangle = 0. \quad (2.1)$$

In this way the (minisuperspace) quantum Hamiltonian that includes the creation and destruction of universes can be written as [14]

$$\begin{aligned} \hat{H} = \hat{H}^{(0)} - g \int dv_1 \int dv_2 \Psi^\dagger(v_1) \Psi^\dagger(v_2) (v_1 + v_2) \Psi(v_1 + v_2) \\ - g \int dv_1 \int dv_2 \Psi^\dagger(v_1 + v_2) v_2 \Psi(v_2) v_1 \Psi(v_1) - \int \frac{dv}{v} \rho(v) \Psi^\dagger(v), \quad \rho(v) = \delta(v) \end{aligned} \quad (2.2)$$

$$\hat{H}^{(0)} = \int_0^\infty \frac{dv}{v} \Psi^\dagger(v) \hat{\mathcal{H}}^{(0)} v \Psi(v), \quad \hat{\mathcal{H}}^{(0)} = v \left(-\frac{3}{4} \frac{d^2}{dv^2} + \lambda \right) \quad (2.3)$$

$\hat{H}^{(0)}$ describes the quantum Hamiltonian corresponding to the action (1.1) (with $\beta = 2/3$) and describes the propagation of a single universe, while the two cubic terms describe the splitting of a universe into two and the merging of two universes into one, respectively. Finally, the last term implies that a universe can be created from the Fock vacuum $|0\rangle$ provided the spatial volume is zero. If it was not for this term $\hat{H}|0\rangle = 0$, and the Fock vacuum would be stable. Again, in our minisuperspace approximation we do not attempt to describe how such a merging or splitting realistically takes place, the only thing which has our interest is how the volume of space can be influenced by such merging or splitting processes, and for this the minisuperspace model might give us some interesting hints.

Even the minisuperspace Hamiltonian \hat{H} is too complicated to be solved in general. A universe can successively split in many, be joined by many and a part that splits off can later rejoin, thereby changing the topology of spacetime. Since the Hamiltonian is essentially dimension independent (all dimension dependence is absorbed in the coupling constants κ , λ and g), what we are describing is the so-called string field theory of two-dimensional CDT [14]. For this string field theory there exists a truncation that can be solved analytically [15],⁴ called generalized CDT (GDCT), and that at the same time has our main interest from a cosmological point of view. It follows the evolution of a universe (let us call it “our” Universe) in time. During this evolution it can merge with other universes (denoted baby universes), created at times t_i with spatial volumes $v_i(t_i) = 0$ (see figure 1). We will assume these universes have the same coupling constants as our Universe. The effective Hamiltonian, the so-called *inclusive* Hamiltonian, first introduced in [18], governing the evolution of the Universe, is obtained from the path integral by integrating over the times t_i and summing over the number of baby universes merging with our Universe. In the path integral, the various baby universes, characterized at a given time t by spatial volumes $v_i(t)$, can themselves be merged with other baby universes. One integrates over all possible $v_i(t)$, not necessarily related to solutions of any classical equations.

Let us describe in some detail how such an inclusive Hamiltonian can be obtained from \hat{H} . The simplest way to incorporate an absorption of baby universes into the propagation of our

⁴Surprisingly, some CDT string field amplitudes can actually be calculated non-perturbatively to all order in the genus expansion, see [16, 17].




Figure 1. Our Universe (blue), represented as one-dimensional circles, propagating in time, with baby universes (red) merging and increasing the spatial volume.

Universe is to replace the quantum field $\Psi(v)$, representing the disappearance of a universe of spatial volume v when it is absorbed by our Universe, by a classical field $\psi(v)$. Thus we make the following replacement in the third term on the r.h.s. of eq. (2.2)

$$\Psi^\dagger(v_1 + v_2)\Psi(v_1)\Psi(v_2) \rightarrow \Psi^\dagger(v_1 + v_2)\psi(v_1)\Psi(v_2) + \Psi^\dagger(v_1 + v_2)\Psi(v_1)\psi(v_2) \quad (2.4)$$

The terms on the r.h.s. of eq. (2.4) contribute to the propagation of our Universe since they are of the form $\Psi^\dagger(v_1)(\cdot)\Psi(v_2)$, but contrary to the terms in \hat{H}_0 they are non-local in v . By Taylor expanding $\Psi^\dagger(v + w)$ around v , this non-locality can be expanded as a power series in the non-local operator $(d/dv)^{-1}$. After some algebra this leads to

$$\int dv dw \Psi^\dagger(v + w) w \psi(w) v \Psi(v) = \int dv \Psi^\dagger(v) F\left(\frac{d}{dv}\right) v \Psi(v), \quad (2.5)$$

where $F(p)$ is the Laplace transform of $\phi(v) = v\psi(v)$:

$$\phi(v) = \phi_0 + \phi_1 v + \phi_2 v^2 + \dots, \quad F(p) = \int_0^\infty dv e^{-pv} \phi(v) = \frac{\Gamma(1)\phi_0}{p} + \frac{\Gamma(2)\phi_1}{p^2} + \dots \quad (2.6)$$

Thus $\hat{\mathcal{H}}_0$ from (2.3) will be replaced by the inclusive Hamiltonian

$$\hat{\mathcal{H}}_{\text{inc}} = \hat{\mathcal{H}}_0 - 2gF\left(\frac{d}{dv}\right) v \quad (2.7)$$

In GCDT $F(p)$ is determined by the self-consistency requirement that our Universe, modified by the impact $\phi(v)$ of baby universes, should be identical to the baby universes it absorbs. We refer to [15] (or to the Lecture Notes [19, 20]) for the details of this determination.⁵

⁵The only difference compared the derivation in [15] is that the sign of g appearing in (2.9) and (2.8) will be different from the sign appearing in [15], the reason being that we consider the *absorption* rather than the emission of baby universes. In earlier articles discussing a modified Friedmann equation ([1, 2]) we used the notation from [15], but with negative g , which effectively meant that we were considering absorption rather than emission of baby universes, like here.

Finally, rotating back from the Hartle-Hawking metric to Lorentzian signature, as when going from (1.1) to (1.2) and replacing⁶ the $-id/dv$ by the classical momentum p conjugate to v we obtain the following semiclassical Hamiltonian

$$\mathcal{H} = v \left(-\frac{3}{4} \left(p^2 + \lambda - 2gF(p) \right) \right) = \frac{3}{4} v \left((p + \alpha) \sqrt{(p - \alpha)^2 + \frac{2g}{\alpha}} \right), \quad (2.8)$$

where α satisfies the equation⁷

$$\alpha^3 - \frac{4}{3} \lambda \alpha - g = 0. \quad (2.9)$$

It is seen that if $g = 0$ one obtains precisely the H in (1.6) (with $N = 1$). A solution to the “on shell” Hamiltonian equations is

$$v(t) = v(t_0) e^{\frac{3}{2} \Sigma(t-t_0)}, \quad \Sigma = \sqrt{\alpha^2 + \frac{g}{2\alpha}}. \quad (2.10)$$

Again, if we choose $g = 0$ we obtain the de Sitter solution (1.7). Increasing g will increase the expansion rate of the universe, the intuition behind this being illustrated in figure 1, and we can actually take the cosmological constant $\lambda = 0$ and still obtain an expanding universe:

$$v(t) = v(t_0) e^{\sqrt{\frac{27}{8}} g^{1/3} (t-t_0)}. \quad (2.11)$$

Thus we have the same classical de Sitter solution as (1.7) provided

$$g^{2/3} = \frac{8}{9} \lambda, \quad (2.12)$$

but the origin of this exponential expansion is now not a cosmological constant but instead the “bombardment” of our Universe by baby universes.

3 Including matter

Until now we have considered our Universe, but without matter. We will now include matter in the most simple way, as a matter density $\rho_m(v)$ in the Hamiltonian (2.8).⁸ In addition we will only consider relatively late times in the evolution of our universe, namely the period after the time of last scattering (t_{LS}), where matter to a good approximation can be considered as dust that exerts zero pressure. Thus the Hamiltonian has the form

$$\mathcal{H}[v, p] = v \left(-f(p) + \kappa \rho_m(v) \right), \quad v \rho_m(v) = v_0 \rho_m(v_0), \quad (3.1)$$

where v_0 and $\rho_m(v_0)$ denote the values at the present time t_0 and where we will later consider three different choices of $f(p)$. Before that, let us discuss the solution of the eoms for arbitrary

⁶The rotation from the Hartle-Hawking metric to the Lorentzian metric involves a double analytic continuation, namely both in time and also a conformal factor rotation, which in the minisuperspace metric parametrization becomes $v \rightarrow -iv$.

⁷In the case of $g = 0$ we choose the solution $\alpha = \sqrt{\lambda} > 0$. In this case, as remarked in footnote 3, $p = -\alpha < 0$ and $\sqrt{(p - \alpha)^2} = \alpha - p$. Since we consider $g \geq 0$, the solution $\alpha > 0$ will only increase with increasing g and $\mathcal{H} = 0$ leads to $p = -\alpha < 0$.

⁸The inclusion of matter in this way introduces an asymmetry between the baby universes and our Universe, in the sense that we do not include such a matter term in the evolution of the baby universes.

$f(p)$ in (3.1). The eoms simplify since $v\rho_m(v)$ is constant.

$$\dot{v} = \frac{\partial \mathcal{H}}{\partial p} = -v f'(p), \quad \text{i.e.} \quad 3 \frac{\dot{a}}{a} = \frac{\dot{v}}{v} = -f'(p) \quad (3.2)$$

$$\dot{p} = -\frac{\partial \mathcal{H}}{\partial v} = f(p), \quad \text{i.e.} \quad t - t_{\text{LS}} = \int_{p_{\text{LS}}}^p \frac{dp}{f(p)} \quad (3.3)$$

By construction any solution to (3.2)–(3.3) will satisfy $\mathcal{H} = \text{const}$, and we are interested in the “on-shell” solutions $\mathcal{H} = 0$, which by (3.1) implies that

$$f(p) = \kappa\rho_m(v) = \kappa\rho_m(v_0) \frac{v_0}{v} = f(p_0) \frac{v_0}{v} = f(p_0)(1+z)^3, \quad (3.4)$$

where p_0 denotes the value of p at present time t_0 and z denotes the redshift at time t , i.e.

$$z(t) + 1 = \frac{a(t_0)}{a(t)} = \left(\frac{v(t_0)}{v(t)} \right)^{1/3}. \quad (3.5)$$

Eq. (3.4) is the generalized Friedmann when eq. (3.2) is used to express p in terms of \dot{a}/a .

Using p as a parameter instead of t (the relation between the two parameters is defined by eq. (3.3)) we can immediately write parametric expressions for a number of relevant functions expressed in terms of $f(p)$. Let us define these. The redshift $z(t)$ or $z(p)$ is

$$z = \frac{a(t_0)}{a(t)} - 1 = \left(\frac{f(p)}{f(p_0)} \right)^{1/3} - 1. \quad (3.6)$$

The Hubble parameter $H(t)$ or $H(p)$ is defined as (see also eq. (1.5))

$$H = \frac{\dot{a}(t)}{a(t)} = -\frac{1}{3} f'(p). \quad (3.7)$$

The *formal density* $\rho_f(t)$ or $\rho_f(p)$ related to the function $f(p)$ is obtained by writing the generalized Friedmann equation (3.4) as

$$\left(\frac{\dot{a}(t)}{a(t)} \right)^2 = \frac{\kappa\rho_m(v)}{3} + \frac{\kappa\rho_f(v)}{3} \quad (3.8)$$

from which we deduce (using the eoms)

$$\kappa\rho_f(p) = \frac{1}{3} (f'(p))^2 - f(p). \quad (3.9)$$

$$\kappa \frac{d\rho_f}{dt} = f(p) f'(p) \left(\frac{2}{3} f''(p) - 1 \right). \quad (3.10)$$

We define the *formal pressure* P_f related to ρ_f by the energy conservation equation

$$\frac{d}{dt}(v\rho_f) + P_f \frac{d}{dt}v = 0. \quad (3.11)$$

This leads to

$$P_f = f(p) \left(\frac{2}{3} f''(p) - 1 \right) - \rho_f(v) \quad (3.12)$$

and the *formal equation of state parameter* w_f is defined (for $\rho_f \neq 0$) by

$$w_f = \frac{P_f}{\rho_f} = \frac{f(p) \left(\frac{2}{3} f''(p) - 1 \right)}{\frac{1}{3} (f'(p))^2 - f(p)} - 1 \quad (3.13)$$

The definitions of ρ_f and P_f ensure that our eoms can be written in the standard GR form, expressed in terms of a , ρ_m , ρ_f and P .

Finally, we will need the standard definitions of some cosmological distances. D_H is defined as the inverse Hubble parameter

$$D_H = \frac{1}{H} = -\frac{3}{f'(p)} \quad (3.14)$$

while

$$D_M = \int_0^z \frac{dz'}{H(z')} = \int_t^{t_0} dt' \frac{a(t_0)}{a(t')} = \int_{p_0}^p \frac{dp}{f(p_0)} \left(\frac{f(p_0)}{f(p)} \right)^{2/3} \quad (3.15)$$

and

$$D_V = \sqrt[3]{z D_H D_M^2}, \quad (3.16)$$

represents a kind of average of the various distances and the so-called angular diameter is often defined as

$$\theta = \frac{r_s}{D_V}, \quad (3.17)$$

where r_s is the co-moving sound horizon at t_{drag} . θ is an important variable, e.g. in the observations of baryon acoustic oscillations (BAO). Since the physics associated with these oscillations takes place before t_{LS} and thus $t_{\text{drag}} < t_{\text{LS}}$, r_s will be independent of the functions f we consider, for reasons to be discussed below.

In these formulas the coupling constants κ , λ and g are so far arbitrary, and so are t_0 and t_{LS} (or equivalently, via (3.3)) p_0 and p_{LS} . If we from observations are given H_0 , the Hubble parameter at present time t_0 , and z_{LS} ,⁹ the redshift at the time of last scattering t_{LS} , we can determine p_0 and p_{LS} :

$$f'(p_0) = -3H_0, \quad f(p_{\text{LS}}) = f(p_0)(1 + z_{\text{LS}})^3. \quad (3.18)$$

This finally leads to the determination of $t_0 - t_{\text{LS}}$ by eq. (3.3). We need more (experimental) input to determine the coupling constants that enter in $f(p)$. We will discuss this in the next section.

Let us now consider the three choices of $f(p)$ in (3.1), where $f(p)$ is given by (2.8). The first choice corresponds to $g = 0$, $\lambda > 0$:

$$f_{\text{ds}}(p) = \frac{3}{4} p^2 - \lambda. \quad (3.19)$$

This is of course just the $f(p)$ associated with the standard de Sitter Hamiltonian (1.6). We include this choice in order to compare the late time cosmology of the standard Λ CDM model

⁹Of course, what is given by observations is the temperature $T(t_0)$ of the CMB. $T(t_{\text{LS}})$ can be calculated by atomic physics and is to a large extent independent of the cosmological model, as is also the statement that $T(t_{\text{LS}})/T(t_0) = a(t_0)/a(t_{\text{LS}}) = 1 + z_{\text{LS}}$.

with the results from our new cosmological models. The second choice corresponds to $\lambda = 0$ and $g > 0$. Thus

$$f_{\text{gcdt}}(p) = \frac{3}{4} (p + \alpha) \sqrt{(p - \alpha)^2 + 2\alpha^2}, \quad \alpha = g^{1/3}. \quad (3.20)$$

This is our model where baby universes of any size v can merge with our Universe. The third choice is motivated by expanding $f_{\text{gcdt}}(p)$ in powers of α/p :

$$f_{\text{gcdt}}(p) = \frac{3}{4} \left(p^2 + \alpha^2 \left(\frac{2\alpha}{p} + O\left(\frac{\alpha^2}{p^2}\right) \right) \right) \quad (3.21)$$

At early times, $v(t)$ becomes small and according to (3.1) $\rho(v(t))$ will be large. Thus $\mathcal{H}(v, p) = 0$ implies that $|p|$ is large at early times (i.e. times larger than but close to t_{LS}). In addition we expect that in our Universe g is small according to (2.12). Thus we expect that keeping only the first term in the expansion (3.21) is a good approximation except at quite late times where the exponential expansion will dominate and where $|p|$ is close to $g^{1/3}$, and this was one of the reasons we in earlier work considered this approximation, which when inserted in (3.4) led to what we denoted the *modified Friedmann equation*. We thus define this $f(p)$ as follows

$$f_{\text{mod}}(p) = \frac{3}{4} \left(p^2 + \frac{2g}{p} \right). \quad (3.22)$$

The last term in (3.22) is effectively a time dependent cosmological constant. At very late time, when the matter term $\rho(v) \propto 1/v$ plays no role, the modified Friedmann equation implies that $p = -(2g)^{1/3}$ and the term acts precisely as a cosmological constant, as already shown in (2.11), and in analogy with the $-\lambda$ term in (3.19). However, for smaller t , $|p|$ will increase, as discussed above, and the last term in (3.22) will be less important.

The three functions f_{ds} , f_{mod} and f_{gcdt} are so simple that even the integrals appearing in (3.3) and (3.15) can be expressed as known analytic functions and in this sense the models are fully solvable. In an appendix we present some details and point out a curious symmetry of the f_{mod} -model.

Returning to the expansion (2.6) that defined the inclusive Hamiltonian, it is seen that the approximation (3.22) corresponds to $\phi(v) = \phi_0 + O(v)$, i.e. to the absorption of baby universes of infinitesimal size (which is one reason we introduced the word “baby universe” for the universes absorbed by our Universe). The absorption of such small baby universes can only result in a change of the spatial volume if infinitely many are absorbed per unit time and unit spatial volume.

4 Comparison of the different models

In the three models we introduced above there are two coupling constants. The gravitational coupling constant κ is fixed by local experiments and we will not discuss it any further. $f_{\text{ds}}(p)$ has the cosmological coupling constant λ as a parameter, while $f_{\text{mod}}(p)$ and $f_{\text{gcdt}}(p)$ have the universe-merging coupling constant g as parameter. λ or g are actually determined already from the values H_0 and z_{LS} mentioned above provided we can assume they play no role for $t < t_{\text{LS}}$. We know this is true for λ in the Λ CDM model and from (2.12), which is of course not exactly true when we add matter to the model, we nevertheless expect that it is also true for g in the other models. This implies that

$$f(p) \approx \frac{3}{4} p^2 \quad \text{for } 0 < t < t_{\text{LS}}. \quad (4.1)$$

That being the case, we can calculate t_{LS} as well as the relation between t_{LS} and p_{LS} without reference to the specific models. For the purpose of illustration, assume (incorrectly) that we can ignore the radiation density $\rho_r(t)$ all the way down to $t \approx 0$. Eq. (3.3) then leads to

$$t_{\text{LS}} = \int_{-\infty}^{p_{\text{LS}}} \frac{dp}{\frac{3}{4}p^2} = -\frac{4}{3} \frac{1}{p_{\text{LS}}}. \quad (4.2)$$

Eqs. (3.18) determine our coupling constant since the first equation determines p_0 and the last equation (the generalized Friedmann equation at t_{LS}) can be written as

$$f(p_0) = \frac{3p_{\text{LS}}^2}{4(1+z_{\text{LS}})^3} \quad (4.3)$$

So given t_{LS} , and thus p_{LS} , we can determine either λ from $f_{\text{ds}}(p)$, or g from $f_{\text{mod}}(p)$ or $f_{\text{gcdt}}(p)$.

The t_{LS} used in cosmology is obtained by also including the radiation density. In this case we obtain an additional term in eq. (3.3) from $\partial H/\partial v$ and

$$t_{\text{LS}} = \int_{-\infty}^{p_{\text{LS}}} \frac{dp}{\frac{3}{4}p^2 + \frac{1}{3}\kappa\rho_r(v(p))}, \quad \frac{3}{4}p^2 = \kappa\rho_m(v) + \kappa\rho_r(v). \quad (4.4)$$

This t_{LS} is smaller than the one given in (4.2) and leads to a somewhat different p_{LS} , which we should use in (4.3) to determine the coupling constants. We can effectively compensate for the smaller t_{LS} by shifting the origin of time.

Having determined the coupling constants we can now compare the different models for given H_0 and z_{LS} . While z_{LS} is fixed (almost) model independent to 1090, there are presently two values of H_0 that do not agree within 5σ (a fact that is denoted the H_0 tension). One value is deduced from ‘‘local’’ measurements, using various space candle techniques [21]. We denote it H_0^{SC} . This value ($H_0^{\text{SC}} = 73.04 \pm 1.04$ km/s/Mpc) is almost independent of cosmological models. The other value is deduced from the CMB data created at t_{LS} . It is using cosmological models in a number of ways, among those to extrapolate to present time t_0 . This value [22] ($H_0^{\text{CMB}} = 67.4 \pm 0.5$ km/s/Mpc), which we denote H_0^{CMB} , is model dependent and the value usually referred to is based on the Λ CDM model.

When comparing our two models with the Λ CDM model we will only use the CMB data to determine T_0 , the temperature of the microwave background radiation at the present time. We have not attempted to use the wealth of information contained in the CMB temperature fluctuations since it requires the use of metric perturbations and it is presently not clear how to incorporate such perturbations in our extended models. We will simply present some of the variables we listed in the previous section, namely $H(z)$, $D_V(z)$ and $w_f(z)$, and t_0 , calculated for the three cosmological models, in two cases, namely for $(H_0^{\text{SC}}, z_{\text{LS}})$ (case A) and for $(H_0^{\text{CMB}}, z_{\text{LS}})$ (case B). For $H(z)$ and $D_V(z)$ we can also compare with measurements from which one can extract these observables for relatively low values of z .

In figure 2 we have shown $a(z)H(z) = H(z)/(1+z)$ in case A and B for the three models. It is seen that in case A the f_{gcdt} and the f_{mod} models fit the data better than the f_{ds} model (the Λ CDM model), while in case B the opposite is the case. Of course we do not know yet whether H_0^{SC} or H_0^{CMB} is the correct H_0 value, but *if* H_0^{SC} turns out to be the correct value the f_{gcdt} and the f_{mod} models seems a better choice for the data presented here. Of course there are many other data to take into account if one wants to make a multi-parameter fit, but as already remarked most these data should be matched to the cosmological model using perturbation expansions not yet available for the f_{gcdt} and f_{mod} models. In table 1 we have




Figure 2. $a(z)H(z) = H(z)/(1+z)$ shown for the three models for $H_0 = 73.0$ (case A). We have included in red the error bars around this point, corresponding to $z = 0$, but the curves all start at 73.0. The orange curve corresponds to f_{ds} , the red curve to f_{gcdt} and the blue curve to f_{mod} . In case B the three models all start at $H_0 = 67.4$, the value suggested by the Planck Collaboration. The green curve corresponds to f_{ds} , the dashed red curve to f_{gcdt} and the dashed blue curve to f_{mod} . Inserted in red are data from $0 < z < 3$: the first three points from the baryon acoustic oscillation data [24], the next from quasars [25] and the last two data points from Ly- α measurements [26, 27].

	$\chi_{\text{red}}^2(\text{ds})$	$\chi_{\text{red}}^2(\text{gcdt})$	$\chi_{\text{red}}^2(\text{mod})$
case A	3.5	1.5	1.8
case B	1.2	1.7	5.6

Table 1. The reduced χ^2 values obtained from figure 2.

	$\chi_{\text{red}}^2(\text{ds})$	$\chi_{\text{red}}^2(\text{gcdt})$	$\chi_{\text{red}}^2(\text{mod})$
case A	4.7	2.1	1.0
case B	1.6	4.1	13.7

Table 2. The reduced χ^2 values obtained from figure 3.

listed the reduced χ^2 values just calculated from the data and the corresponding error bars shown in figure 2 in Case A and B.

Figure 3 shows $D_V(z)/r_s$ for the f_{ds} , f_{gcdt} and f_{mod} models, normalized by $(D_V(z)/r_s)^{\text{CMB}}$, i.e. the $D_V(z)/r_s$ for the f_{ds} model in case B. We see here the same trend as in figure 2: the f_{gcdt} and the f_{mod} models fit the data better in case A, while the f_{ds} -model fits better in case B. In table 2 we list the reduced χ^2 values obtained from figure 3.

In figure 4 we display $w_f(z)$ for the three models. Of course $w_{f_{\text{ds}}}(z) = -1$. The two other models have negative $w_f(z) < -1$. In standard cosmology this is a sign that some unphysical




Figure 3. The three curves starting at value 1 for $z = 0$ are curves of $(D_V^f(z)/r_s)/(D_V^{f_{\text{ds}}}(z)/r_s)^{\text{CMB}}$ for the models with $f = f_{\text{gcdt}}$ (red, dotted), f_{mod} (blue, dotted) and f_{ds} (green), all for the chosen value $H_0 = H_0^{\text{CMB}}$ (case B). The last curve is of course 1, except for the error bars reflecting the uncertainty of H_0^{CMB} . The three curves starting at value 0.94 for $z = 0$ are the curves for $f = f_{\text{gcdt}}$ (red), f_{mod} (blue) and f_{ds} (orange), all for $H_0 = H_0^{\text{SC}}$ (case A), and again normalized by $(D_V^{f_{\text{ds}}}(z)/r_s)^{\text{CMB}}$. The data points and error bars are obtained from various BAO surveys (see [3] for a detailed table).

degrees of freedom have been added to the system. However, here one cannot conclude that, since merging with baby universe seems more like having a time dependent cosmological constant, but without the problem that a time dependent cosmological constant will break the invariance of the model under time-reparametrization. Allowing a time dependence of the cosmological “constant”: $\lambda \rightarrow \lambda(t)$, changes $w_{f_{\text{ds}}}$:

$$w_{f_{\text{ds}}}(z) = -1 \rightarrow w_{\tilde{f}_{\text{ds}}} = -1 - \frac{1}{3H(t)} \frac{\dot{\lambda}(t)}{\lambda(t)}. \quad (4.5)$$

Thus, if $\lambda(t)$ is growing with time, assuming the universe is expanding (i.e. $H(t) > 0$), it follows that $w_{\tilde{f}_{\text{ds}}}(t) < -1$. If we consider the f_{mod} model then $\lambda(t)$ is replaced by $-3g/2p$, and effectively it acts in the same way: for small $t \rightarrow 0$ $p(t) \rightarrow -\infty$ while for $t \rightarrow \infty$ $p(t) \rightarrow -(2g)^{1/3}$. In fact using (3.13) it follows immediately that $w_{f_{\text{mod}}}(z)$ goes monotonically from $-3/2$ at $z = \infty$ ($t = 0$), to -1 for $z = -1$ ($t = \infty$), as also illustrated in figure 4.

The Planck data, combined with BAO as well as supernovae type Ia data, is consistent with the Λ CDM model with a value $H_0 = 67.4 \pm 0.5$. The consistency is corroborated by treating w in the dark energy equation of state $p = w\rho$ as a parameter, the value of which is then determined to be $w = -1.03 \pm 0.03$ [22]. However, w is not well determined by the Planck data alone ($w = -1.56 \pm 0.5$, ([22], table 4)). If one allows for a more general time dependent $w(z)$ extension of the Λ CDM model, the picture is the same, but even including BAO as well as supernovae type Ia data the best fit will produce a curve which is both consistent with the $w(z)$ curve for our gcdt-model as well as consistent with the pure Λ CDM model within error bars (figure 5, [23]). One trend to observe in the analysis presented in [23] (see figure 3, 4




Figure 4. $w_f(z)$ for the three models. The orange curve is $w_{f_{\text{ds}}} = -1$, which is true for any value of H_0 . The red curves are for the f_{gcdt} -model, dotted curve for H_0^{SC} (case A) and full curve for H_0^{CMB} (case B). The blue curves are for the f_{mod} -model, dotted curve for H_0^{SC} (case A) and full curve for H_0^{CMB} (case B). For $z \rightarrow -1$ ($t \rightarrow \infty$) $w_f(z) \rightarrow -1$.

	$t_0(\text{ds})$	$t_0(\text{gcdt})$	$t_0(\text{mod})$
case A	13.3 Gyr	13.5 Gyr	13.9 Gyr
case B	13.8 Gyr	14.0 Gyr	14.4 Gyr

Table 3. The values of t_0 for the three models in cases A and B.

and 6) is that larger values of H_0 correlate with smaller values of $w(z)$. If H_0 is in the range 72–74 km/s/Mpc, as is the case for the best fits to our models, one obtains in the Planck collaboration analysis of the extended Λ CDM models, values of $w(z)$ that are typically in the range predicted by our models. In this sense our $w(z)$ is compatible with the Planck $w(z)$.

Finally, in table 3 we list the various values of t_0 in the six cases discussed. In case A $t_0(\text{ds}) = 13.3$ Gyr and $t_0(\text{gcdt}) = 13.5$ Gyr seem uncomfortable short and in case B $t_0(\text{mod}) = 14.4$ Gyr is probably too long. However, had we started out with a H_0 in between H_0^{SC} and H_0^{CMB} we can obtain values of both $t_0(\text{mod})$ and $t_0(\text{gcdt})$ that are comfortable in agreement with the age of the universe as determined from the oldest stars. In particular for the f_{gcdt} -model, just looking at the data in figures 2 and 3 it is clear that if we tried to determine the best value of H_0 for the model from the data (including now the observed H_0^{SC} as a data point), the optimal value of H_0 would precisely be between H_0^{SC} and H_0^{CMB} . In table 4 we have listed values of H_0 obtained for the three model by minimizing the χ^2 for the data shown in figure 2, together with the corresponding (minimal) value of the reduced χ^2 , gt_0^3 and $g/\Lambda^{3/2}$, where we for Λ have used the value corresponding to the f_{ds} -model with the H_0 listed in table 4. The f_{gcdt} - and f_{mod} -models are doing slightly better than the f_{ds} -model, but the error bars in figure 2 are too large to obtain any precision determination of H_0 in this way. The same is even more true if one had used the data in figure 3 to determine the “best value” of H_0 for the three models. In figure 5 we have shown the equivalent of figure 3, using the values of H_0 from table 4.

While we have some choice for different models for absorption of baby universes, once H_0 is fixed the values of observables like $H(z)$, $D_V(z)$ and $f_m(z)\sigma_8(z)$ are fixed for each such model. If, for some reason we decide that a specific value of H_0 is trustworthy the values

	$f_{\text{ds}}\text{-model}$	$f_{\text{gcdt}}\text{-model}$	$f_{\text{mod}}\text{-model}$
H_0	71.2	72.2	73.9
χ_{red}^2	2.4	1.2	1.4
t_0	13.5 Gyr	13.6 Gyr	13.8 Gyr
gt_0^3		2.983	1.224
$g/\Lambda^{3/2}$		0.971	0.380

Table 4. The best values of H_0 for the three models together with the corresponding parameters.




Figure 5. The graphs for $H(z)/(1+z)$ for the three models using the values of H_0 from table 4. Orange graph corresponds to $f_{\text{ds}}\text{-model}$, blue graph to the $f_{\text{mod}}\text{-model}$ and red graph to the $f_{\text{gcdt}}\text{-model}$.

of these observables can then be viewed as predictions for a given model. It was in this spirit we presented the data from the observed $H(z)$ and $D_V(z)$ for case A and B in figure 2 and figure 3 and the corresponding tables 1 and 2. Another “local” variable is the density fluctuations of matter for $z < 2$. Presently the error bars on these data are too large for being used in a simple analysis like done above in figure 2 and figure 3 (see [3] for an attempt). However, this will most likely change dramatically with the new observations to be obtained by the Euclid satellite.

5 Discussion

Let us start by emphasizing the extreme simplicity of the models considered. It is based on the Hamiltonian

$$\mathcal{H} = Nv(-f(p) + \kappa\rho_m(v)), \quad \rho_m(v)v = \rho_m(v_0)v_0, \quad (5.1)$$

and the corresponding eoms are

$$f(p) = \kappa\rho_m(v), \quad \frac{\dot{v}}{v} = -f'(p), \quad \dot{p} = f(p). \quad (5.2)$$

Thus, (for suitable f), if the universe starts out with a small spatial volume $v(t)$ for small t , $\rho_m(v(t))$ is large and the first equation implies that $-p$ is large. The second equation then implies that the expansion is fast, and the last equation that $-p$ increases, i.e. $|p|$ decreases, which again implies that the rate of expansion per unit volume \dot{v}/v decreases. Eventually, if $v(t)$ continues to grow to infinity, $\rho_m(v)$ goes to zero. Thus $f(p)$ goes to zero, i.e. p goes to a constant. The middle equation then implies that $v(t)$ approaches an pure exponential expansion. As we have argued, assuming the absorption of baby universes this exponential expansion is very natural if the chance of such an absorption per unit time is proportional to the spatial volume. Therefore, there is no need to introduce a cosmological constant (“negative gravity”) by hand: the gravitational force indeed wants to limit the expansion of the universe, but this is counteracted by the “bombardment” of our Universe by baby universes.

The model we have suggested is of course quite primitive and unrealistic, but taking two limits, one where only universes with infinitesimal spatial volumes are absorbed, the f_{mod} model, and one where universes of any size can be absorbed and where our Universe and the other universes are on equal footing, the f_{gcdt} model, show that the models are reasonable insensitive to the detailed distribution of baby universe sizes. One can therefore hope that they reflect the results one would obtain from more realistic models.

We have presented our model as a “late time” cosmological model and in the present formulation it has nothing to say about the universe for $t < t_{\text{LS}}$. Viewed as such a late time cosmological model it is interesting that it favors the local measured value H_0^{SC} somewhat compared to the value H_0^{CMB} . However, in view of the simplicity of the model, we will not press this as an important point. Let us rather discuss if this multi-universe scenario has a chance to answer some of the early-time questions in cosmology.

We have nothing to add about inflation as it is presented in various models. However, the fact that the universe has expanded from, say, a Planckian size to 10^{-5}m in a very short time, invites the suggestion that this expansion was caused by a collision with a larger universe, i.e. that it was really our Universe which was absorbed in another “parent” universe. Since we have presently no detailed description of the absorption process, it is difficult to judge if such a scenario could take place in a way that would actually solve the problems inflation was designed to solve, but one interesting aspect of such a scenario is that there is no need for an inflaton field.

While a continuous absorption of microscopic baby universes probably can be accommodated in a non-disruptive way in our Universe, it is less clear what happens if the “baby” universe is not small, since we have not suggested an actual mechanism for such absorption. Maybe the least disruptive situation would be one where the absorption happened inside a black hole. The unknown mechanism of absorption could maybe favor such a scenario when the sizes of the baby universes are not infinitesimal. Recall that a Reissner-Nordström black hole actually connects to different universes. We are not seriously suggesting such a black hole scenario, but we mention it to point out that there is room for a lot of interesting considerations. Ultimately, any realistic model should be specific about how the absorption occurs.

The present value of the cosmological constant λ in the ΛCDM model is according to some viewpoints embarrassingly small. According to (2.12), λ being small when expressed in Planck units ($\kappa = 1$) also implies that g is small. Historically, before observations pointed to a small λ in the ΛCDM model, many people favored $\lambda = 0$ as a result of some underlying mechanism not yet fully understood, e.g. Coleman’s mechanism [31]. We might still need such a mechanism to explain why a λ coming from the zero-point fluctuations of quantum

fields will not create a large λ .¹⁰ However, it might be easier to explain why λ is exactly zero than to explain why it is unnaturally small. If $\lambda = 0$ can be proven, we could then still have an exponentially expanding universe caused by baby universe absorption.

Like λ , g appears in our model just as a coupling constant, reflecting in some way the “density” of the baby universes “surrounding” our Universe. By comparing our model to observations g has to be small. However, it is a coupling constant in a larger multiverse theory that we have not yet been able to solve. This leaves the hope that a consistent solution of this larger theory might determine g . We have started the program to unveil this theory [4–7], but it is still work in progress.

Acknowledgments

JA thanks the Perimeter Institute for hospitality while this work was completed and the research was supported in part by the Perimeter Institute for Theoretical Physics. Research at the Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development and by the Province of Ontario through the Ministry of Colleges and Universities.

YW acknowledges the support from JSPS KAKENHI Grant Number JP18K03612.

A Analytic results for the three models

In this appendix we list some of the analytical results for the three models defined by f_{ds} , f_{gcdt} and f_{mod} , respectively. As remarked above we have already analytic parametric representations of all the observables considered, using κp as parameter. Here we will provide the explicit expressions and also provide some of them as analytic functions of time t .

A.1 The f_{ds} -model

$f_{\text{ds}}(p)$ is given (3.19). From (3.3) we can find t as a function of p :

$$t = \int_{-\infty}^p \frac{dp'}{f_{\text{ds}}(p')} = -\frac{2}{\sqrt{3\lambda}} \tanh^{-1} \left(\frac{2\sqrt{\lambda}}{\sqrt{3}} \frac{1}{p} \right) \quad (\text{A.1})$$

$$p = -\frac{2\sqrt{\lambda}}{\sqrt{3}} \coth \left(\frac{\sqrt{3\lambda}}{2} t \right). \quad (\text{A.2})$$

$$f_{\text{ds}}(p) = \frac{\lambda}{\sinh^2 \left(\frac{\sqrt{3\lambda}}{2} t \right)}, \quad \text{i.e.} \quad v(t) = \frac{\kappa \rho_{\text{m}0} v_0}{\lambda} \sinh^2 \left(\frac{\sqrt{3\lambda}}{2} t \right) \quad (\text{A.3})$$

The only non-trivial function is D_M defined by eq. (3.15). The integral can be expressed by hypergeometric functions and one representation is

$$D_M = \frac{6}{\sqrt{3\lambda}} \left(\frac{\lambda}{\kappa \rho_{\text{m}0}} \right)^{1/3} \sinh^{1/3} \left(\frac{\sqrt{3\lambda}}{2} t' \right) {}_2F_1 \left[\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; -\sinh^2 \left(\frac{\sqrt{3\lambda}}{2} t' \right) \right] \Big|_t^{t_0} \quad (\text{A.4})$$

$$= \frac{6}{\sqrt{3\lambda}} \left(\frac{\lambda}{\kappa \rho_{\text{m}0}} \right)^{1/3} \left(\frac{\lambda}{f_{\text{ds}}(p')} \right)^{1/6} {}_2F_1 \left[\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; -\frac{\lambda}{f_{\text{ds}}(p')} \right] \Big|_p^{p_0} \quad (\text{A.5})$$

¹⁰There are other viewpoints where the value of λ is claimed to be natural (or at least not embarrassingly small) in a quantum field context when one uses renormalization group arguments [28, 29] (see [30] for a review).

where

$$\frac{\lambda}{\kappa\rho_{m0}} = \frac{\lambda}{f_{ds}(p_0)} = \sinh^2\left(\frac{\sqrt{3\lambda}}{2}t_0\right). \quad (\text{A.6})$$

A.2 The f_{gcdt} -model

$f_{\text{gcdt}}(p)$ is given by (3.20). From (3.3) we can find t as a function of p :

$$t = \int_{-\infty}^p \frac{dp'}{f_{\text{gcdt}}(p')} = \frac{4}{3\sqrt{6}\alpha} \left(\tanh^{-1}\left(\frac{\sqrt{\frac{2}{3}}(-p+2\alpha)}{\sqrt{(p-\alpha)^2+2\alpha^2}}\right) - \tanh^{-1}\left(\sqrt{\frac{2}{3}}\right) \right) \quad (\text{A.7})$$

Defining t_c by

$$\frac{3\sqrt{6}\alpha}{4}t_c = \tanh^{-1}\left(\sqrt{\frac{2}{3}}\right) = \cosh^{-1}\sqrt{3} = \sinh^{-1}\sqrt{2}, \quad (\text{A.8})$$

we can invert (A.7) to find:

$$p = -\alpha \frac{\sinh\frac{3\sqrt{6}\alpha}{4}(t+t_c) + 2\sqrt{2}}{\sinh\frac{3\sqrt{6}\alpha}{4}(t+t_c) - \sqrt{2}} \quad (\text{A.9})$$

$$f_{\text{gcdt}}(p) = \frac{9\sqrt{3}\alpha^2}{2} \frac{\cosh\frac{3\sqrt{6}\alpha}{4}(t+t_c)}{\left(\sinh\frac{3\sqrt{6}\alpha}{4}(t+t_c) - \sqrt{2}\right)^2} \quad (\text{A.10})$$

Again the only non-trivial function is D_M defined by eq. (3.15). In this case the integral in eq. (3.15) is a generalized hypergeometric functions, a so-called Appell- F_1 function:

$$D_M = \frac{1}{f_{\text{gcdt}}(p_0)^{1/3}} \left(\frac{p'+\alpha}{6\alpha^2}\right)^{1/3} F_1\left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{4}{3}; x^+(p'), x^-(p')\right) \Big|_p^{p_0} \quad (\text{A.11})$$

where

$$f_{\text{gcdt}}(p_0) = \kappa\rho_{m0}, \quad x^\pm(p) = \frac{2 \pm i\sqrt{2}}{3\alpha}(p+\alpha). \quad (\text{A.12})$$

A.3 The f_{mod} -model

$f_{\text{mod}}(p)$ is given by (3.21). From (3.3) we can find t as a function of p :

$$t = \int_{-\infty}^p \frac{dp'}{f_{\text{mod}}(p')} = \frac{4}{3\sqrt{3}a} \left(\arctan\frac{2p-a}{\sqrt{3}a} + \frac{1}{2\sqrt{3}} \log\frac{(p-a)^2+ap}{(p+a)^2} + \frac{\pi}{2} \right) \quad (\text{A.13})$$

where

$$a = 2^{1/3}\alpha, \quad t \rightarrow 0 \text{ for } p \rightarrow -\infty, \quad t \rightarrow \infty \text{ for } p \rightarrow -a. \quad (\text{A.14})$$

Again we can express D_M , defined by the integrals in eq. (3.15), in terms of hypergeometric functions. One representation is:

$$D_M = \frac{3}{5} \left(\frac{2}{3}\right)^{2/3} \frac{1}{\alpha^2 f(p_0)^{1/3}} (p')^{5/3} {}_2F_1\left(\frac{5}{9}, \frac{2}{3}; \frac{14}{9}; -\frac{(p')^3}{2\alpha^3}\right) \Big|_p^{p_0}. \quad (\text{A.15})$$

A.4 Duality in the f_{mod} -model

We end by observing that we have a kind of duality in the f_{mod} -model. Define a (modified) Hubble parameter by $h(t) = \dot{v}/v = 3\dot{a}/a = 3H(t)$, where $H(t)$ is the usual Hubble parameter. We then have the equations:

$$\frac{3}{4} \left(p^2 + \frac{2\alpha^3}{p} \right) = \kappa\rho_{\text{m}}(v), \quad h(v) = -\frac{3}{4} \left(2p - \frac{2\alpha^3}{p^2} \right). \quad (\text{A.16})$$

These equations are invariant under the replacement

$$p \longleftrightarrow -\frac{\gamma}{\alpha} \frac{\alpha^3}{p}, \quad \kappa\rho_{\text{m}} \longleftrightarrow \frac{\alpha}{\gamma} h, \quad \gamma = 2^{1/3}. \quad (\text{A.17})$$

The mapping $p \rightarrow -\gamma\alpha^2/p$ has some similarity with a T -duality map in string theory.

References

- [1] J. Ambjørn and Y. Watabiki, *A modified Friedmann equation*, *Mod. Phys. Lett. A* **32** (2017) 1750224 [[arXiv:1709.06497](#)] [[INSPIRE](#)].
- [2] J. Ambjørn and Y. Watabiki, *Easing the Hubble constant tension*, *Mod. Phys. Lett. A* **37** (2022) 2250041 [[arXiv:2111.05087](#)] [[INSPIRE](#)].
- [3] J. Ambjørn and Y. Watabiki, *The large-scale structure of the universe from a modified Friedmann equation*, *Mod. Phys. Lett. A* **38** (2023) 2350068 [[arXiv:2208.02607](#)] [[INSPIRE](#)].
- [4] J. Ambjørn and Y. Watabiki, *A model for emergence of space and time*, *Phys. Lett. B* **749** (2015) 149 [[arXiv:1505.04353](#)] [[INSPIRE](#)].
- [5] J. Ambjørn and Y. Watabiki, *Creating 3, 4, 6 and 10-dimensional spacetime from W_3 symmetry*, *Phys. Lett. B* **770** (2017) 252 [[arXiv:1703.04402](#)] [[INSPIRE](#)].
- [6] J. Ambjørn and Y. Watabiki, *CDT and the Big Bang*, *Acta Phys. Polon. Supp.* **10** (2017) 299 [[arXiv:1704.02905](#)] [[INSPIRE](#)].
- [7] J. Ambjørn and Y. Watabiki, *Models of the Universe based on Jordan algebras*, *Nucl. Phys. B* **955** (2020) 115044 [[arXiv:2003.13527](#)] [[INSPIRE](#)].
- [8] Y. Hamada, H. Kawai and K. Kawana, *Baby universes in 2d and 4d theories of quantum gravity*, *JHEP* **12** (2022) 100 [[arXiv:2210.05134](#)] [[INSPIRE](#)].
- [9] J. Ambjørn, A. Gorlich, J. Jurkiewicz and R. Loll, *Planckian Birth of the Quantum de Sitter Universe*, *Phys. Rev. Lett.* **100** (2008) 091304 [[arXiv:0712.2485](#)] [[INSPIRE](#)].
- [10] J. Ambjørn, A. Gorlich, J. Jurkiewicz and R. Loll, *The Nonperturbative Quantum de Sitter Universe*, *Phys. Rev. D* **78** (2008) 063544 [[arXiv:0807.4481](#)] [[INSPIRE](#)].
- [11] J. Ambjørn, A. Goerlich, J. Jurkiewicz and R. Loll, *Nonperturbative Quantum Gravity*, *Phys. Rept.* **519** (2012) 127 [[arXiv:1203.3591](#)] [[INSPIRE](#)].
- [12] R. Loll, *Quantum Gravity from Causal Dynamical Triangulations: A Review*, *Class. Quant. Grav.* **37** (2020) 013002 [[arXiv:1905.08669](#)] [[INSPIRE](#)].
- [13] J. Ambjørn and R. Loll, *Nonperturbative Lorentzian quantum gravity, causality and topology change*, *Nucl. Phys. B* **536** (1998) 407 [[hep-th/9805108](#)] [[INSPIRE](#)].
- [14] J. Ambjørn et al., *A String Field Theory based on Causal Dynamical Triangulations*, *JHEP* **05** (2008) 032 [[arXiv:0802.0719](#)] [[INSPIRE](#)].
- [15] J. Ambjørn, R. Loll, W. Westra and S. Zohren, *Putting a cap on causality violations in CDT*, *JHEP* **12** (2007) 017 [[arXiv:0709.2784](#)] [[INSPIRE](#)].

- [16] J. Ambjørn, R. Loll, W. Westra and S. Zohren, *Summing over all Topologies in CDT String Field Theory*, *Phys. Lett. B* **678** (2009) 227 [[arXiv:0905.2108](#)] [[INSPIRE](#)].
- [17] J. Ambjørn et al., *A new continuum limit of matrix models*, *Phys. Lett. B* **670** (2008) 224 [[arXiv:0810.2408](#)] [[INSPIRE](#)].
- [18] H. Kawai, N. Kawamoto, T. Mogami and Y. Watabiki, *Transfer matrix formalism for two-dimensional quantum gravity and fractal structures of space-time*, *Phys. Lett. B* **306** (1993) 19 [[hep-th/9302133](#)] [[INSPIRE](#)].
- [19] J. Ambjørn, *Elementary Quantum Geometry*, [arXiv:2204.00859](#) [[INSPIRE](#)].
- [20] J. Ambjørn, *Elementary Introduction to Quantum Geometry*, Taylor & Francis (2022) [[INSPIRE](#)].
- [21] A.G. Riess et al., *A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 km s⁻¹ Mpc⁻¹ Uncertainty from the Hubble Space Telescope and the SH0ES Team*, *Astrophys. J. Lett.* **934** (2022) L7 [[arXiv:2112.04510](#)] [[INSPIRE](#)].
- [22] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [*Erratum ibid.* **652** (2021) C4] [[arXiv:1807.06209](#)] [[INSPIRE](#)].
- [23] PLANCK collaboration, *Planck 2015 results. XIV. Dark energy and modified gravity*, *Astron. Astrophys.* **594** (2016) A14 [[arXiv:1502.01590](#)] [[INSPIRE](#)].
- [24] BOSS collaboration, *The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample*, *Mon. Not. Roy. Astron. Soc.* **470** (2017) 2617 [[arXiv:1607.03155](#)] [[INSPIRE](#)].
- [25] P. Zarrouk et al., *The clustering of the SDSS-IV extended Baryon Oscillation Spectroscopic Survey DR14 quasar sample: measurement of the growth rate of structure from the anisotropic correlation function between redshift 0.8 and 2.2*, *Mon. Not. Roy. Astron. Soc.* **477** (2018) 1639 [[arXiv:1801.03062](#)] [[INSPIRE](#)].
- [26] J.E. Bautista et al., *Measurement of baryon acoustic oscillation correlations at $z = 2.3$ with SDSS DR12 Ly α -Forests*, *Astron. Astrophys.* **603** (2017) A12 [[arXiv:1702.00176](#)] [[INSPIRE](#)].
- [27] BOSS collaboration, *Quasar-Lyman α Forest Cross-Correlation from BOSS DR11: Baryon Acoustic Oscillations*, *JCAP* **05** (2014) 027 [[arXiv:1311.1767](#)] [[INSPIRE](#)].
- [28] C. Moreno-Pulido and J. Sola Peracaula, *Renormalizing the vacuum energy in cosmological spacetime: implications for the cosmological constant problem*, *Eur. Phys. J. C* **82** (2022) 551 [[arXiv:2201.05827](#)] [[INSPIRE](#)].
- [29] C. Moreno-Pulido and J. Sola, *Running vacuum in quantum field theory in curved spacetime: renormalizing ρ_{vac} without $\sim m^4$ terms*, *Eur. Phys. J. C* **80** (2020) 692 [[arXiv:2005.03164](#)] [[INSPIRE](#)].
- [30] J. Sola Peracaula, *The cosmological constant problem and running vacuum in the expanding universe*, *Phil. Trans. Roy. Soc. Lond. A* **380** (2022) 20210182 [[arXiv:2203.13757](#)] [[INSPIRE](#)].
- [31] S.R. Coleman, *Why There Is Nothing Rather Than Something: A Theory of the Cosmological Constant*, *Nucl. Phys. B* **310** (1988) 643 [[INSPIRE](#)].