



Consistent differences in tissue oxygen levels across 15 insect species reflect a balance between oxygen supply and demand and highlight a hitherto unknown adaptation for extracting sufficient oxygen from water

Jackson H. Birrell^{a,1,*}, Wilco C.E.P. Verberk^b, H. Arthur Woods^a

^a Department of Biology, University of Montana, Missoula, MT 59812 USA

^b Department of Animal Ecology and Physiology, Radboud Institute for Biological and Environmental Sciences, Radboud University, AJ, Nijmegen 6525 The Netherlands

ARTICLE INFO

Keywords:

Aquatic insects
Hypoxia
Lifespan
Metabolism
Reactive oxygen species
Oxygen uptake
Oxygen availability

ABSTRACT

Animals, including insects, need oxygen for aerobic respiration and eventually asphyxiate without it. Aerobic respiration, however, produces reactive oxygen species (ROS), which contribute to dysfunction and aging. Animals appear to balance risks of asphyxiation and ROS by regulating internal oxygen relatively low and stable, but sufficient levels. How much do levels vary among species, and how does variation depend on environment and life history? We predicted that lower internal oxygen levels occur in insects with either limited access to environmental oxygen (i.e., insects dependent on aquatic respiration, where low internal levels facilitate diffusive oxygen uptake, and reduce asphyxiation risks) or consistently low metabolic rates (i.e., inactive insects, requiring limited internal oxygen stores). Alternatively, we predicted insects with long life-stage durations would have internal oxygen levels > 1 kPa (preventing high ROS levels that are believed to occur under tissue hypoxia). We tested these predictions by measuring partial pressures of oxygen (PO₂) in tissues from juvenile and adult stages across 15 species comprising nine insect orders. Tissue PO₂ varied greatly (from 0 to 18.8 kPa) and variation across species and life stages was significantly related to differences in habitat, activity level, and life stage duration. Individuals with aquatic respiration sustained remarkably low PO₂ (mean = 0.88 kPa) across all species from Ephemeroptera (mayflies), Plecoptera (stoneflies), Trichoptera (caddisflies), and Diptera (true flies), possibly reflecting a widespread, but hitherto unknown, adaptation for extracting sufficient oxygen from water. For Odonata (dragonflies), aquatic juveniles had higher PO₂ levels (mean = 6.12 kPa), but these were still lower compared to terrestrial adults (mean = 13.3 kPa). Follow-up tests in juvenile stoneflies showed that tissue PO₂ remained low even when exposed to hyperoxia, suggesting that levels were down-regulated. This was further corroborated since levels could be modulated by ambient oxygen levels in dead individuals. In addition, tissue PO₂ was positively related to activity levels of insect life stages across all species and was highest in stages with short durations. Combined, our results support the idea that internal PO₂ is an evolutionarily labile trait that reflects the balance between oxygen supply and demand within the context of the environment and life-history of an insect.

Introduction

For animals, oxygen is a double-edged sword (Harrison et al., 2012). During oxidative phosphorylation of cellular respiration, electrons are transported in the electron transport chain and eventually transferred onto an oxygen molecule, the electron acceptor. Because oxygen is such an effective acceptor, it can release more energy from substrates than is possible with other electron acceptors. The majority of animals have

come to rely on oxygen for respiration and asphyxiate when starved of it. Oxygen-based energy production was central to the evolution and diversification of modern eukaryotes and supports key aspects of modern ecological communities (Knoll 1992), many of which include large-bodied animals with energy-intensive lifestyles. Indeed, current interest in the mechanisms by which organisms sense and adapt to oxygen availability remains high, and work on this topic was recently awarded the 2019 Nobel Prize in Physiology or Medicine (Zhang et al.

* Corresponding author.

E-mail addresses: jackson.birrell@umontana.edu (J.H. Birrell), wilco.verberk@ru.nl (W.C.E.P. Verberk), art.woods@umontana.edu (H.A. Woods).

¹ Lead contact

2019). At the same time, oxygen can be toxic. While respiring aerobically, mitochondria produce reactive oxygen species (ROS) (e.g., superoxide anion radical, hydroxyl radical, hydrogen peroxide), which damage cell components, including lipid bilayers and DNA, and whose effects contribute to disease and aging (Brieger et al. 2012; Davalli et al. 2016).

Animals thus need to balance risks of toxicity with those of asphyxiation (Verberk and Atkinson 2013), regulating the uptake of oxygen and its transport down the oxygen cascade so that enough is supplied to meet the oxygen demand of the mitochondria without triggering excess production of ROS. Thus, the partial pressure of oxygen, PO_2 , tends to decrease across the steps of the oxygen transport cascade from ambient (~21 kPa at sea-level) via a convective distribution system (e.g., blood, hemolymph, or tracheal system) before the final diffusion-based step into the mitochondria. Holding internal PO_2 at low levels is likely beneficial because it increases the PO_2 gradient from atmosphere to mitochondria, which elevates rates of diffusive oxygen transport (Harrison et al. 2012).

Internal PO_2 also has consequences for rates of ROS production, although the relationship between ROS production and PO_2 is not straightforward (Zorov et al. 2014). While ROS production approaches zero when there is no oxygen, somewhat paradoxically, ROS production tends to increase under cellular hypoxia (e.g., < ~1 kPa) (Chandel et al. 1998; Guzy and Schumacker 2006; Murphy 2009; Semenza 2011; Zorov et al. 2014), despite being relatively stable across most other oxygen concentrations (Starkov 2008). In addition, ROS production tends to increase sharply during reoxygenation, leading to widespread cellular damage (Li et al. 2002; Millar et al. 2007; Wang et al. 2018). This suggests that organisms should hold internal PO_2 stable and low enough to support oxygen delivery but not so low that high levels of ROS are generated.

Data on mammals show that tissue PO_2 is indeed moderately low, in the range of 2.3 - 4.5 kPa in mammalian resting red muscle, with values somewhat lower around mitochondria (Wittenberg and Wittenberg 1989; Poole et al. 2020). Values in other mammalian tissues and organs (measured by a variety of techniques) are typically lower than 7 kPa, and sometimes as low as in red muscle (Vanderkooi et al. 1991). Fewer data have been obtained on insects, but those available suggest that hemolymph and intratracheal PO_2 typically lie between 5 and 18 kPa O_2 (Komai 1998; Kirkton 2007; Matthews and White 2011; Lehmann et al. 2019; Rowe et al. 2022), with substantial variation across body compartments (e.g., abdomen versus femur in grasshoppers; Kirkton 2007), between modes of breathing (e.g., continuous versus discontinuous; Rowe et al. 2022), or between levels of activity (e.g., preflight versus flight in hawkmoths; Komai 1998). Internal PO_2 tends to decrease with progressive atmospheric hypoxia and there is an attempt to minimize such decreases by increased ventilation rates once PO_2 of the meta-thoracic ganglion reaches 3 kPa (Harrison et al. 2020). Hetz & Bradley (2005) also reported active regulation of internal PO_2 levels. They measured intratracheal PO_2 in pupae of the silk moth *Attacus atlas* subjected to experimentally manipulated levels of external oxygen. Via spiracular opening and closing, pupae closely regulated intratracheal PO_2 to ~ 4 kPa, which the authors proposed was a mechanism for avoiding oxygen toxicity.

To date, published data on internal levels of PO_2 in insects have come from terrestrial species, which have ready access to high levels of oxygen in air. A significant proportion of insects, however, have aquatic juveniles, which extract oxygen from the surrounding water for respiration. These insects face fundamentally different oxygen challenges stemming from low concentrations of oxygen in water and the difficulty of extracting it (Lancaster and Downes 2013; Verberk and Bilton 2013). Many aquatic juveniles, moreover, have closed tracheal systems and thus cannot regulate oxygen levels via spiracular control (Buchwalter et al. 2020). We hypothesized that aquatic juvenile insects would sustain lower overall internal oxygenation for two reasons. First, for any given external PO_2 in water, sustaining the lowest possible internal PO_2

maximizes the driving gradient (ΔPO_2) for oxygen transport from environment to tissues. Second, high internal PO_2 is likely unnecessary for aquatic juveniles because they typically have stable, cool body temperatures and low activity levels that may be supported by small oxygen reservoirs and low metabolic rates. We thus also predicted that more active species would maintain higher levels of tissue PO_2 to provide larger reservoirs of oxygen for immediate use during rapid locomotion. Finally, we hypothesized that internal PO_2 levels reflect the need to avoid damage from ROS, something which should be a high priority for aquatic juvenile life stages which are typically longer lived, compared to either the adults they become or compared to terrestrial juveniles of other species. We thus predicted that life stages with long durations should have internal PO_2 above the threshold that is associated with cellular hypoxia (i.e. > 1 kPa), to avoid cellular damage, or perhaps alternative mechanisms for minimizing the effects of ROS production (e.g., enhanced antioxidant defenses).

Materials and methods

Insect collection and rearing

We measured the partial pressures of oxygen (PO_2) in juveniles and adults from nine orders of insects, five orders with terrestrial juveniles and adults (Coleoptera, Phasmatodea, Orthoptera, Diptera, Lepidoptera) and five orders with aquatic juveniles (Ephemeroptera, Plecoptera, Trichoptera, Odonata, Diptera). Obtaining juveniles and adults of the same species was often difficult, and more than one species was occasionally used from each order, for a total of fifteen species (Supplemental 1). Aquatic insects were all 'water-breathers' as juveniles, with closed tracheal systems and cutaneous gas-exchange. Aquatic insects that relied on exchange of aerial oxygen, e.g. by surfacing regularly to replenish air stores, on compressible physical gills, or on respiratory siphons at any point in development were not used. Adults of aquatic insects were air-breathers with functional spiracle systems. Terrestrial insects were air-breathers with open-spiracle systems both as juveniles and adults.

Insects were either collected near Missoula, Montana, USA from summer 2021 to spring 2022 or obtained from collaborators or commercially from a variety of sources (Supplemental 1). Field-collected insects were maintained in the laboratory at the University of Montana for up to two days until PO_2 measurements were made. Aquatic insects were held in buckets (10 L) with de-chlorinated tap water with an air stone to provide oxygen and to stir the water. Insects were held at roughly the mean temperature of the water source from which they were collected at the time of collection. Temperature was controlled by either placing the bucket of insects inside a refrigerator (5 °C) or cooling the water in the bucket with a recirculating water bath (10 - 20 °C) (ActiveAqua AACH10HP). Field-collected terrestrial insects were held in a plastic container (2 L) with a mesh lid until measurements were made. PO_2 of most insects from rearing sources were measured immediately upon arrival at the laboratory. Terrestrial Diptera and Lepidoptera were received from rearing sources as larvae and then reared to adulthood in the lab. Larvae of *Calliphora vicina* (Diptera) were held in a plastic container (2 L) with a mesh lid filled with food (wood shavings) at room temperature (~ 20 °C) until they began to pupate (~ 1 week). Once pupated, individuals were transferred to a refrigerator and held at 5 °C to prevent them from drying out in the lab. Caterpillars of *Manduca sexta* (Lepidoptera) were reared on commercial artificial diet until wandering, at which point they were transferred to wooden blocks with pre-drilled pupation chambers. Pupae were held at room temperature until eclosion.

After PO_2 measurements were taken, field-collected individuals were identified in the laboratory using a dissecting microscope and dichotomous keys (Tripplehorn and Johnson, 2004; Merritt et al. 2008). Data on the duration of each stage of species (juvenile or adult) were retrieved from a variety of peer-reviewed papers, non-peer-reviewed but

reputable online sources, and personal communications with experts who have unpublished data on study species' lifespans from laboratory rearing programs. Peer-reviewed sources could not be used in every case because of limited published data. For some species, no stage duration data were available, in which case stage durations of closely related species were substituted (Supplemental 1). No data were available on the temperature-sensitivity of stage duration, and so no adjustments could be made for rearing temperature. In addition, activity levels of 1 or 2 were assigned to each species based on our knowledge of their lifestyles and after carefully observing each in the laboratory under controlled temperatures, with a value of 1 being slow and inactive (e.g., stick insect) and a value of 2 being fast and active (e.g., flesh fly adult) (Supplemental 1). Subjecting individuals to activity level assays to collect more quantitative data was impractical as major, categorical differences in activity level were always apparent and because measurements would have overly stressed individuals prior to internal PO₂ measurements, necessitating time-intensive collections of many more individuals from the field.

Because differences in activity levels may be associated with variation in metabolism, metabolic rates were estimated for each species and stage. Resting metabolic rate data were retrieved from peer-reviewed papers or published, online datasets. For several species or stages, no metabolic data were available, and we then substituted data from the closest related species for which data were available (i.e., usually the same genus or family, sometimes order). When no metabolic data were available for species within the same order, a metabolic rate value was not assigned (Supplemental 1). Mean mass-specific metabolic rate was then calculated for each species and stage. All values were then converted to units of Watts g⁻¹. Because values reported in the literature were measured on different individuals, differing in mass and measurement temperatures, we derived temperature and mass corrected metabolic rates by fitting a linear model (R function: *lm*) of log₁₀(metabolic rate) ~ log₁₀(mean mass) + temperature. This model was used to derive mass specific metabolic rate at 20 °C, using the mass scaling coefficient and thermal dependency from the model (1.032 and 0.268 eV, respectively).

Oxygen measurements

PO₂ was measured using either Clark-style oxygen microelectrodes or an oxygen-sensitive optode. Oxygen microelectrodes (Unisense, 100 µm tip, with guard electrode; optimal measuring range 0 to 21 kPa) were connected to a picoammeter (Unisense, PA-2000) and calibrated at room temperature (~20 °C) using a custom glass apparatus holding tap water into which we bubbled pure N₂ gas (0 kPa O₂) or room air (19 kPa O₂, since Missoula is situated at an elevation of 978 m above sea level). Analog outputs from the picoammeter were sent to a Sable Systems acquisition system (UI-2, Expedata software) and visualized in real time during measurements. The optode was housed in a needle (Pyroscience, OXR430, 430 µm tip; optimal measuring range 0 to 50 kPa) and was connected to a meter (Pyroscience, FireSting O₂). The meter sent the outputs to a computer, where they were processed and visualized via the FireSting recording software. The optode was calibrated at room temperature using a plastic cup holding tap water into which we bubbled pure N₂ gas (0 kPa O₂) or room air (19 kPa O₂). Readings for both the electrodes and optodes were reliable, providing similar readings during calibration on different days with each method, and we performed the calibration procedure every day that measurements were taken.

PO₂ was measured inside the thorax and abdomen of individuals ($n = 118$) while they were still alive: Ephemeroptera ($n = 8$ and $n = 3$ for juveniles and adults, respectively), Plecoptera ($n = 8$, $n = 6$), Trichoptera ($n = 8$, $n = 4$), Odonata ($n = 4$, $n = 3$), Diptera ($n = 9$, $n = 8$); terrestrial orders: Coleoptera ($n = 8$, $n = 8$), Phasmatodea ($n = 5$, $n = 5$), Orthoptera ($n = 7$, $n = 7$), Diptera ($n = 7$, $n = 3$), Lepidoptera ($n = 8$, $n = 7$). Measurements were taken by securing each individual to a plastic Petri dish with rubber bands fed through holes cut in the dish. A small

hole was made in the cuticle of the thorax and abdomen of each individual, the exact location being haphazardly chosen within each body segment. An electrode or optode was mounted to a micro-manipulator secured to the lab bench-top, which was used to carefully insert the sensor tip into the hole cut in the insect as viewed through a stereomicroscope (Nikon SMZ 1500). Electrodes were used to measure internal PO₂ in most species, which were small or weakly sclerotized. Optodes were used to measure PO₂ in two larger, more heavily sclerotized species (*Trypoxylus* beetles and *Phyllium* stick insects). Optodes were used in these instances because the moving limbs of the insects would easily break the microelectrodes. Because juveniles of aquatic fly larvae (*Tipula* sp.) repeatedly wiggled out of the rubber band mount, individuals were instead cut in half transversely. The abdomen and thorax were then inserted onto the tip of the electrode mounted to the micro-manipulator. PO₂ was recorded based on the values shown on the acquisition system or computer program as soon as the reading stabilized (i.e., was constant $\pm \sim 2\%$) for ~ 2 s. Measurements were made soon after initial incisions were made in the body cavity, usually < 1 min. Based on the rapidity of measurements, changes to internal PO₂ due to diffusion of atmospheric air through holes or body openings, or due to reduced rates of ventilation post-injury, were considered to be negligible.

For each method, measurement locations and depths varied from individual to individual in an attempt to avoid biasing measurements. This is because little was known about the spatial variation of structures likely to support different internal PO₂ levels among species – i.e., the location of air sacs and large tracheae (with high PO₂), areas of high muscle fiber density (with low PO₂), etc. In addition, taking measurements in the same location or structure type (e.g., tissue or hemolymph) was problematic because most individuals were small (median body mass: 0.3 g), inhibiting our ability to take measurements in specific locations with certainty. However, for all individuals, measurements were made “deep” inside the body cavity (roughly 1/8 to 1/2 of the depth of each segment) and away from the center of the segment to avoid the gut lumen. Notably, analyses showed that internal PO₂ was no different in the thorax and abdomen for all species, suggesting that measurements were indeed representative of internal (tissue-hemolymph) oxygen levels, rather than specific body compartments.

Different methods and equipment used for measuring PO₂ were considered equivalent, based on preliminary PO₂ measurements on giant salmonfly nymphs (Plecoptera: *Pteronarcys californica*), in which measurements were made by inserting an optode through a hole cut in the cuticle (nine individuals, 16 measurements) or by inserting it into the abdomen and thorax after it has been cut transversely (one individual, two measurements). Mean PO₂ values from both methods using the optode were 0.14 kPa and were similar to measurements on the same species using the electrode in the main dataset (all were 0.0 kPa). Optodes and electrodes both showed nearly identical values when measuring the PO₂ of air-bubbled water near saturation and nitrogen-bubbled water near anoxia. Validity of our methods are also corroborated by similar mean internal PO₂ measurements made by Lee & Matthews (2024) on Odonata nymphs (6.2 and 6.3 kPa, respectively).

In a separate experiment, aquatic juveniles of the stonefly, *Pteronarcys californica*, were exposed to either normoxic (19 kPa O₂; $n = 7$) or hyperoxic (60 kPa; $n = 7$) water at 10 °C for 24 h. Internal PO₂ was measured twice inside the thorax of each individual. For the first measurement, a small hole was cut in the thorax of a haphazardly chosen location. The tip of an electrode was inserted into the hole using a micromanipulator and dissecting microscope. For the second measurement, the thorax was cut from the abdomen, and the thorax was inserted onto the tip of the electrode. For both measurements, we recorded the value shown on the acquisition system as soon as the reading stabilized ($\pm \sim 2\%$) for ~ 2 s, as above.

As an additional control, we measured internal PO₂ in the thorax of *P. californica* juveniles preserved in vials of 70 % ethanol at room temperature for at least one year. Here, the expectation was that biological processes affecting internal PO₂ would be negligible and should cause

internal PO₂ to track that of the external environment. Any differences in PO₂-change between exposures of live versus dead nymphs should thus reflect an effect of an unmeasured biological process (e.g., regulation of PO₂ levels). Before measurements were made, individuals were separated into unique vials with 5 ml of 70 % 'used' ethanol from the vial in which each juvenile was originally stored. Individuals were then assigned to either a normoxic (19 kPa O₂; $n = 7$) or hyperoxic (60 kPa; $n = 7$) treatment group. However, individuals from the same vial of origin were assigned to different treatments to avoid biasing the results based on slight differences in the percentage of ethanol, and therefore the solubility of oxygen, in each original storage vial (i.e., percentages of ethanol in preserved samples can change after storage as water from specimen bodies is drawn out into the solution) (Shchukarev and Tolmacheva 1968). For the normoxic treatment, thoracic PO₂ was measured for each individual at room temperature. Manipulating the PO₂ of the ethanol by bubbling in air was unnecessary because ethanol PO₂ was near saturation (~ 19 kPa O₂) in each vial. For the hyperoxic treatment, O₂ gas was bubbled into each vial until it reached 60 kPa, at which point the vial was tightly closed and stored for 24 h at room temperature. After exposure, we measured the PO₂ in both the ethanol and the thorax of each individual. For both treatments, one thoracic measurement was made for each nymph, in which a small hole was cut in the thorax of a haphazardly chosen location. The tip of an electrode was then inserted into the hole using a micromanipulator and dissecting microscope and the value shown on the acquisition system was recorded as soon as the reading stabilized for ~ 2 s, as above.

Statistical analysis

Tissue PO₂ data were zero-inflated, and we added 1 kPa to each measurement value and subsequently performed a log₁₀ transformation to improve normality. Although data transformation improved the distribution without yielding a truly normal distribution, we proceeded to use linear mixed-effects models (function: lmer; package: lme4) to analyze the effects of covariates on tissue PO₂ (Bates et al. 2015). To assess the performance of our models, we analyzed residuals and then checked the robustness of our linear mixed-effects models by reanalyzing our models using censored regression, which is designed for zero-inflated, semi-continuous data (function: censReg; package: censReg). Additionally, because strong correlations existed among some explanatory variables (Supplemental 2), results of each model were checked with residual regression – a common method for analyzing collinear data – following recommendations of Graham (2003). See additional details below.

In each linear mixed effects model, we included species as a random effect to account for effects of species identity and to isolate effects of habitat, life stage, activity and lifespan on PO₂. In preliminary analyses, we also tested for differences in location (i.e., on the thorax and abdomen), and body mass. Because both location and body mass did not contribute to the model fit and were non-significant as main effects nor in interaction with stage and habitat, we dropped them from further analyses and included measurements of both thoracic and abdominal PO₂ in our models. Because explanatory variables (habitat, life stage, activity and lifespan) could not be included in a single model without causing the fitting procedure to fail (likely due to strong collinearity among variables), we tested six more simplified candidate models, which, in total, included all possible combinations of covariates (Supplemental 3). Interactions between explanatory variables were only included when collinearity was low ($r < 0.30$) and models with explanatory variables that were highly correlated ($r > 0.30$) were checked with residual regressions (see additional details below). We calculated R² and AIC for use in model selection via the R functions, r.squaredGLMM (package: MuMIn) (Barton 2009) and aictab (package: AICcmodavg) (Mazerolle 2017).

Although R² and AIC did not agree on the single best model (Supplemental 3), model 1 (log₁₀(PO₂) ~ stage × habitat + activity level)

was strongly supported by both selection criteria and was presented in the results. Residuals of model 1 were calculated using the R function, residuals (R Core Team, 2022), and visual analysis suggested that they were approximately normally distributed (Supplemental 3). However, because model 1 did not include stage duration, which had significant effect on tissue PO₂ in several of the candidate models, we ran an additional linear mixed effect model (i.e., model 7) with stage duration as the sole predictor (log₁₀(PO₂) + stage duration). Although model 7 had a considerably poorer fit than the other models (Supplemental 3), the effect of stage duration was significant, suggesting that patterns of internal PO₂ in insects can be explained by a single, alternative explanatory variable. In addition to model 1, we therefore also present the results from model 7 in the results. As indicated above, outputs of all candidate models were checked with supplemental censored regression (Supplemental 2) and residual regression models (Supplemental 4). Outputs of linear mixed effects models were similar to those from the supplemental models and were therefore deemed reliable.

Because many of the explanatory mechanisms are rooted in metabolic rate, we also ran an additional analysis in which we tested whether mass-specific metabolic rate affected tissue PO₂ using linear mixed-effects models (R function: lmer). Mass-specific metabolic rate had no effect on tissue PO₂ and was thus dropped from subsequent analyses (Supplemental 5). In addition, we used linear mixed effects models, as above, to analyze the effect of insect order and clade (i.e. Paleoptera, Polyneoptera, and Holometabola) on internal PO₂, and no significant effects were found (Supplemental 6). We thus deemed phylogenetic effects to be negligible and further phylogenetic analyses unnecessary.

To test whether internal PO₂ levels varied with external oxygenation levels (i.e., normoxia vs. hyperoxia) and internal oxygen consumption we measured internal PO₂ of live and dead salmonfly (Plecoptera: *Pteronarcys californica*) nymphs. For live salmonfly nymphs, we used a linear mixed effects model (function: lmer; package: lme4). Before analysis, we removed a single, extreme outlier ($> Q3 + 3 * IQR$) from the dataset. As above, measurement values were zero-inflated and strongly right-skewed, and we added 1 kPa to each value and transformed each using log₁₀. Data transformation improved the distribution but did not yield a truly normal distribution. We nevertheless proceeded to use linear mixed-effects models (function: lmer; package: lme4) to analyze the effect of water oxygenation, and later checked the results using censored regression (Tobit) models with mixed effects, as above (function: censReg; R-package: censReg). The method used (i.e., cutting a small hole in the insect vs. cutting in half) and mass had no effect on internal PO₂, and these covariates were dropped from the analysis. Supplementary censored regression models yielded similar results to the linear mixed effects models, and outputs of linear mixed effects models were, therefore, considered valid. For the analysis of the effect of external oxygenation levels (i.e., normoxia vs. hyperoxia) on internal PO₂ of dead nymphs stored in ethanol, linear mixed effects models were used (R function: lme; package: lme4). Vial of origin was included as a random effect to account for any differences in the percentage of ethanol in which nymphs were stored. Mass has no effect in tissue PO₂ and was dropped from subsequent analyses.

Results

Across the 15 insect species and 9 orders examined, internal PO₂ ranged between 0 and 18.8 kPa (0.0–16.0 kPa, 5th and 95th percentile), with substantial variation among species and stages (Table 1). Tissue PO₂ of live insects was best explained by a model that included interactions between life stage and habitat (model 1: $P < 0.0001$) and activity level (model 1: $P < 0.0001$) (Fig. 1A) (Table 2). The interaction was driven by the remarkably low internal PO₂ values in aquatic juveniles (mean = 0.88 kPa), which were several times lower than in juveniles or adults of fully terrestrial species (mean = 3.85 and 5.33, respectively) and an order of magnitude lower compared to the air-breathing adults of the same or similar species (mean = 11.83 kPa).

Table 1Table of the number of replicates and the mean and standard deviation of internal PO₂ (kPa) for each species-stage, with the order and habitat to which they belong.

| Summary of internal PO ₂ for each species-stage | | | | | | |
|--|----------------------------------|----------|-------------|---|----------------------|--------------------|
| Order | Taxon | Stage | Habitat | N | Mean PO ₂ | Sd PO ₂ |
| Ephemeroptera | <i>Drunella grandis</i> | immature | aquatic | 8 | 0.00 | 0.00 |
| Ephemeroptera | <i>Drunella grandis</i> | adult | aquatic | 2 | 11.95 | 1.20 |
| Ephemeroptera | <i>Rhithrogena</i> sp. | adult | aquatic | 1 | 17.73 | 0.43 |
| Diptera | <i>Tipula</i> sp. | immature | aquatic | 9 | 0.43 | 1.60 |
| Diptera | <i>Tipula</i> sp. | adult | aquatic | 8 | 7.76 | 6.74 |
| Plecoptera | <i>Pteronarcys californica</i> | immature | aquatic | 8 | 0.04 | 0.11 |
| Plecoptera | <i>Pteronarcys californica</i> | adult | aquatic | 6 | 14.17 | 1.99 |
| Odonata | <i>Aeshna palmata</i> | immature | aquatic | 4 | 6.13 | 4.88 |
| Odonata | <i>Aeshna palmata</i> | adult | aquatic | 2 | 14.71 | 1.38 |
| Odonata | <i>Sympetrum danae</i> | adult | aquatic | 1 | 7.84 | NA |
| Trichoptera | <i>Parapsyche</i> sp. | immature | aquatic | 8 | 0.52 | 0.67 |
| Trichoptera | <i>Brachycentrus</i> sp. | adult | aquatic | 4 | 13.72 | 4.97 |
| Coleoptera | <i>Trypoxylus dichotomus</i> | immature | terrestrial | 8 | 1.34 | 0.73 |
| Coleoptera | <i>Trypoxylus dichotomus</i> | adult | terrestrial | 8 | 0.92 | 0.24 |
| Phasmatodea | <i>Phyllium philippinicum</i> | immature | terrestrial | 5 | 1.44 | 0.61 |
| Phasmatodea | <i>Phyllium philippinicum</i> | adult | terrestrial | 5 | 1.17 | 0.32 |
| Orthoptera | Acrididae | immature | terrestrial | 7 | 8.21 | 4.27 |
| Orthoptera | <i>Melanoplus bivittatus</i> | adult | terrestrial | 3 | 14.25 | 2.30 |
| Orthoptera | <i>Trimerotropis verruculata</i> | adult | terrestrial | 4 | 7.93 | 4.51 |
| Diptera | <i>Calliphora vicina</i> | immature | terrestrial | 7 | 2.64 | 4.35 |
| Diptera | <i>Calliphora vicina</i> | adult | terrestrial | 3 | 9.60 | 3.98 |
| Lepidoptera | <i>Manduca sexta</i> | immature | terrestrial | 8 | 5.12 | 5.84 |
| Lepidoptera | <i>Manduca sexta</i> | adult | terrestrial | 7 | 6.19 | 4.28 |

Aquatic juveniles also generally displayed low activity levels (mean = 1.1), contrasting with adults of aquatic insects which displayed high activity levels (mean = 2.0) (Fig. 1B). Internal PO₂ was also significantly affected by the duration of life stages (model 7: $P < 0.0001$) (Table 2), with aquatic juveniles exhibiting the longest stage durations (> 300 days) (Fig. 1C), again contrasting with adults that exhibited the shortest stage durations (mean = 10 days).

In addition, internal PO₂ of live salmonfly nymphs was not significantly affected by the PO₂ of the water (19 kPa vs. 60 kPa) during a 24 h exposure to hyperoxia ($P = 0.89$) (Fig. 2A) (Table 3). However, dead stoneflies preserved in ethanol had ~ 2 times higher tissue PO₂ after being exposed to hyperoxic ethanol (60 kPa) for 24 h than those exposed to normoxic ethanol (19 kPa) ($P < 0.0001$) (Fig. 2B) (Table 3).

Discussion

We observed impressive variation in internal levels of oxygen across the 15 insect species and 9 orders examined. Most notably, aquatic juveniles had remarkably low internal levels of oxygen, far lower than air-breathing adult insects or fully terrestrial species. Variation in PO₂ could be explained from differences in life stage, habitat, activity, and life-stage duration, indicating that insects likely actively regulate internal oxygen levels to balance competing functions (Fig. 1). While active regulation of PO₂ has long been known to occur in holometabolous, terrestrial insects (Hetz and Bradley 2005; Chown et al. 2006), it has been little-explored in aquatic species. Nevertheless, active regulation in aquatic insects is further supported by our experimental results, in which internal oxygen levels of juvenile stoneflies (*Pteronarcys californica*) remained low even when individuals were exposed to strong hyperoxia (~ 60 kPa) (Fig. 2). In contrast, internal oxygen levels of dead juvenile stoneflies had elevated internal oxygen levels (~ 18 kPa) in ambient normoxia and these levels were further elevated (~ 32 kPa) in hyperoxia, further supporting our conclusion that live aquatic juveniles actively regulate internal PO₂ to low levels.

What about life in water explains the low tissue oxygen levels in aquatic juveniles? One possibility is that low internal PO₂ evolved to facilitate oxygen uptake from water by maximizing the environment-to-mitochondria PO₂ gradient, which ultimately drives oxygen transport. In water (compared to in air), rates of oxygen flux are orders of magnitude lower because water imposes much lower oxygen diffusion coefficients (Woods, 1999; Verberk et al., 2011). In addition, water

holds 20–30-fold less oxygen than air and has a higher (dynamic) viscosity, which generates thicker boundary layers around insect respiratory surfaces and greatly increases the energetic costs of ventilation (Denny 1993; Verberk and Atkinson 2013; Woods and Moran 2020). Indeed, aquatic insects have evolved a suite of morphological (gills), physiological (oxygen-carrying proteins), and behavioral adaptations (exploiting high flow microhabitats) to help mitigate this biophysical problem (Hynes 1970; Harrison et al. 2012; Buchwalter et al. 2020; Frakes et al., 2021; Birrell and Woods, 2023). Our finding of low PO₂ in juveniles across four orders of aquatic insects nevertheless suggests that holding PO₂ to low levels is a widely used mechanism for increasing oxygen uptake. This innovation is likely important for aquatic juveniles to sustain sufficiently high basal metabolic rates, which we show are no different than those of air-breathers, despite much lower levels of oxygen availability in water than in air (Supplemental 5). It also suggests that oxygen fluxes in many but not all aquatic insects are well-approximated by indices of oxygen availability derived from external oxygen conditions, which often assume internal PO₂ to be zero (such as the OSI in Verberk et al. 2011). Exceptions in our data set include dragonfly nymphs, which, despite being aquatic, sustained relatively high mean PO₂ of 6.2 kPa. These results agree with those of Lee and Matthews (2024), who recently showed that nymphs of a dragonfly sustained hemolymph PO₂ of 6.3 kPa in normoxic water.

Maintaining low internal PO₂ may also require aquatic insects to have low activity levels. Across all stages and species, estimated activity level was positively correlated with tissue PO₂, with immature stages of aquatic insects generally displaying the lowest tissue PO₂ and activity levels. If insects are to regulate internal oxygen levels, rates of uptake and use must be matched, at least over timescales of minutes or longer. Indeed, many species of terrestrial, holometabolous insects regulate internal PO₂ by opening and closing their spiracles (Hetz and Bradley 2005; Chown et al. 2006), though to much higher levels than those observed here for aquatic juveniles. Thus, high internal PO₂ may support generally higher levels of activity by terrestrial insects, or greater variation in levels of activity. However, inactive insects may not require large reserves of oxygen for use during sustained bursts of locomotion – e.g., winged flight – and thus, hold tissue PO₂ at lower levels. Living life in the slow lane should thus allow insects to maintain sufficiently low internal PO₂ to support gas exchange, and may also help prevent accumulation of ROS (see below) and increase energy savings (Hetz and Bradley 2005; Speers-Roesch 2018). Indeed, energy savings could be




Fig. 1. Boxplot of tissue PO₂ of live insects from different life stages (adult vs. immature), habitats (aquatic vs. terrestrial) (A) and with different activity levels (B), and stage durations (C). Raw data are shown instead of log₁₀ + 1 transformed data used in final analyses for easier interpretation. The regression line in panel C was derived from a linear model (R function: lm), which showed a similar result to model 7, with a significant effect of life stage duration ($P < 0.001$).

paramount in long-lived aquatic stages that have access to only intermittent food (e.g., leaf shredders over an annual cycle). Nevertheless, the causal relationship between internal PO₂ and activity levels in

Table 2

Linear mixed effects models of the effects of stage, habitat, activity level and stage duration on tissue PO₂ of live insects. For models with categorical predictors, reference level is shown in brackets.

| PO ₂ in live insects – model 1 | | | | | |
|---|----------|------------|---------|---------|---------|
| Predictor | Estimate | Std. Error | Df | T-value | P-value |
| Intercept [reference: aquatic, adult] | 0.511 | 0.079 | 22.911 | 6.503 | < 0.001 |
| Stage [Immature] | -0.423 | 0.076 | 68.025 | -5.596 | < 0.001 |
| Habitat [Terrestrial] | -0.174 | 0.070 | 23.228 | -2.486 | 0.021 |
| Activity level [2] | 0.541 | 0.060 | 24.466 | 9.025 | < 0.001 |
| Stage [Immature] * Habitat [Terrestrial] | 0.503 | 0.079 | 189.453 | 6.400 | < 0.001 |
| PO ₂ in live insects – model 7 | | | | | |
| Predictor | Estimate | Std. Error | Df | T-value | P-value |
| Stage duration | -0.001 | < 0.001 | 245.400 | -12.189 | < 0.001 |




Fig. 2. Boxplots of tissue PO₂ of both live and dead juvenile salmonflies. Measurements on live individuals made after 24hr exposure to normoxia (19 kPa) and hyperoxia (60 kPa). Measurements on dead individuals in ethanol at normoxia (19 kPa) and 24 h exposure to hyperoxia (60 kPa).

Table 3

Linear mixed effects models of the effect of treatment (hyperoxia vs. normoxia) on tissue PO₂ of live salmonflies in water and dead salmonflies in ethanol.

| PO ₂ in live salmonflies in water | | | | | |
|--|----------|------------|--------|---------|---------|
| Predictor | Estimate | Std. Error | Df | T-value | P-value |
| Treatment | -0.005 | 0.038 | 23.000 | -0.141 | 0.889 |
| PO ₂ in dead salmonflies in ethanol | | | | | |
| Predictor | Estimate | Std. Error | Df | T-value | P-value |
| Treatment | 24.434 | 2.635 | 13.000 | 9.272 | < 0.001 |

aquatic juveniles is uncertain, as low levels of oxygen supply in water may instead *constrain* activity levels by requiring low tissue PO₂ for gas exchange, as discussed above.

We found no support for the alternative hypothesis that hypoxia-induced production of reactive oxygen species (ROS) prevents aquatic insects from having extremely low internal PO₂ levels (< 1 kPa). ROS, produced in mitochondria as a by-product of cellular respiration, are a major driver of cellular damage, disease, and aging (Boveris et al. 1973; Costa et al. 1993; Zorov et al. 2014; Shields et al. 2021). Because rates of

ROS production rise under both cellular hypoxia and reoxygenation (Murphy 2009; Wang et al. 2018), high levels of ROS damage under hypoxia could be expected to prevent aquatic juveniles from maintaining extremely low internal PO₂, especially in those with longer stage durations. We found, however, that insects with longer stage durations had significantly lower tissue PO₂, including many PO₂ values below 1 kPa, and this pattern was partly driven by aquatic juveniles, which had both the longest life stages and the lowest PO₂. This result, along with the taxonomic breadth of low internal PO₂ in aquatic juveniles, suggests that the benefits of improved gas exchange outweigh the costs of increased ROS production under cellular hypoxia and that aquatic juveniles employ a variety of mechanisms to minimize either the production or deleterious effects of ROS. Indeed, aquatic insects employ a host of antioxidant systems (Felton and Summers 1995; Xie et al. 2009; Sanz et al. 2017), yet their capacities to minimize ROS damage remains understudied compared to work on vertebrate or holometabolous insect model species. One possibility, however, is that aquatic juveniles reduce activity levels to consistently low levels, as shown by our data, to avoid excessive ROS production during activity-related reoxygenation (e.g., Wang et al. 2018).

In conclusion, using the first comparative dataset of tissue PO₂ from juvenile and adult insects from both aquatic and terrestrial environments, we show that aquatic juveniles across four orders have unexpectedly low internal oxygen levels. Because of the centrality of oxygen to both metabolic energy generation and oxidative damage causing senescence, this outcome likely reflects that competing priorities play out in fundamentally different ways in aquatic versus terrestrial environments. The mechanisms by which aquatic insects regulate internal oxygen levels provide fruitful direction for follow-up research. Potential mechanisms likely include shifts in tracheal morphology and the density and conductance of tracheoles (Wigglesworth 1983). In addition, we encourage additional studies on how the mitochondria of aquatic juveniles function under low internal oxygen levels and whether and how they avoid ROS production and damage during cellular hypoxia. More broadly, our data suggest that the problem of understanding patterns and mechanisms of oxygen regulation across insects would benefit from both focused and comparative studies utilizing species from a greater diversity of phylogenetic and ecological backgrounds.

CRediT authorship contribution statement

Jackson H. Birrell: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing. **Wilco C.E.P. Verberk:** Formal analysis, Writing – review & editing. **H. Arthur Woods:** Conceptualization, Formal analysis, Investigation, Methodology, Project administration, Resources, Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data and R scripts used in this study have been made publicly available on the Zenodo repository and can be accessed at: <https://doi.org/10.5281/zenodo.12745828>.

Acknowledgements

We thank Jon Harrison for helpful discussions on internal PO₂ of insects during initial preparation of the manuscript. We thank Romain Boisseau and Doug Emlen for donating rhinoceros beetles and stick

insects from their rearing programs to our experiments. We also thank Erin McCullough for providing raw metabolic rate data from O'Brien et al. (2019) and for helpful correspondence on how to interpret the data. We appreciate the University of Montana for general support and laboratory space for the project. W.C.E.P.V. gratefully acknowledges support from the Netherlands Organization for Scientific Research (NWO-VIDI Grant 016.161.321). The authors have no conflicts of interest to report.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.cris.2024.100095](https://doi.org/10.1016/j.cris.2024.100095).

References

- Barton, K., 2009. MuMIn: multi-model inference. R package version 1, 0. 0.
- Bates, D., Mächler, M., Bolker, B.M., Walker, S.C., 2015. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67. <https://doi.org/10.18637/jss.v067.i01>.
- Birrell, J.H., Woods, H.A., 2023. Going with the flow – how a stream insect, *Pteronarcys californica*, exploits local flows to increase oxygen availability. *J. Exp. Biol.* 226 (3), 1–10. <https://doi.org/10.1242/jeb.244609>.
- Boveris, A., Chance, B., 1973. The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem. J.* 134, 707–716. <https://doi.org/10.1042/bj1340707>.
- Brieger, K., Schiavone, S., Miller, F.J., Krause, K.H., 2012. Reactive oxygen species: from health to disease. *Swiss Med. Wkly.* 142. <https://doi.org/10.4414/smww.2012.13659>.
- Buchwalter, D.B., Resh, V.H., Lamberti, G.A., Verberk, W.C.E.P., 2020. Chapter 4: aquatic insect respiration. In: Merritt, R.W., Cummins, K.W., Berg, M.B. (Eds.), *In an introduction to the aquatic insects of North America*. Kendall Hunt Publishing.
- Chandel, N.S., Maltepe, E., Goldwasser, E., Mathieu, C.E., Simon, M.C., Schumacker, P. T., 1998. Mitochondrial reactive oxygen species trigger hypoxia-induced transcription. *Proc. Nat. Acad. Sci.* 95 (20), 11715–11720.
- Chown, S.L., Gibbs, A.G., Hetz, S.K., Klok, C.J., Lighton, J.R.B., Marais, E., 2006. Discontinuous gas exchange in insects: a clarification of hypotheses and approaches. *Physiol. Biochem. Zool.* 79, 333–343. <https://doi.org/10.1086/499992>.
- Costa, L.E., Llesuy, S., Boveris, A., 1993. Active oxygen species in the liver of rats submitted to chronic hypobaric hypoxia. *Am. J. Physiol. - Cell Physiol.* 264. <https://doi.org/10.1152/ajpcell.1993.264.6.c1395>.
- Davalli, P., Mitic, T., Caporali, T., Lauriola, A., D'Arca, D., 2016. ROS, cell senescence, and novel molecular mechanisms in aging and age-related diseases. *Oxid. Med. Cell. Longev.* 2016, 1–18. <https://doi.org/10.1155/2016/3565127>.
- Denny, M.W., 1993. *Air and water: the biology and physics of life's media*. Princeton University Press.
- Felton, G.W., Summers, C.B., 1995. Antioxidant systems in insects. *Arch. Insect Biochem. Physiol.* 29, 187–197. <https://doi.org/10.1002/arch.940290208>.
- Frakes, J.I., Birrell, J.H., Shah, A.A., Woods, H.A., 2021. Flow increases tolerance of heat and hypoxia of an aquatic insect. *Biol. Lett.* 17. <https://doi.org/10.1098/rsbl.2021.0004>.
- Guzy, R.D., Schumacker, P.T., 2006. Oxygen sensing by mitochondria at complex III: the paradox of increased reactive oxygen species during hypoxia. *Exp. Physiol.* 91 (5), 807–819.
- Harrison, J.F., Woods, H.A., Roberts, S.P., 2012. *Ecological and environmental physiology of insects*. Oxford University Press.
- Harrison, J.F., Wolfgang, W., Hetz, S.K., 2020. PO₂ of the metathoracic ganglion in response to progressive hypoxia in an insect. *Biol. Lett.*, 1620200548. <https://doi.org/10.1098/rsbl.2020.0548>.
- Hetz, S.K., Bradley, T.J., 2005. Insects breathe discontinuously to avoid oxygen toxicity. *Nature* 433, 516–519. <https://doi.org/10.1038/nature03106>.
- Hynes, H.B.N., 1970. *The ecology of running waters*. Liverpool University Press.
- Kirkton, S.D., 2007. Effects of insect body size on tracheal structure and function. In: Roach, R.C., Wagner, P.D., Hackett, P.H. (Eds.), *Hypoxia and the circulation. Advances in experimental medicine and biology*. Springer, Berlin, pp. 221–228.
- Knoll, A.H., 1992. The early evolution of eukaryotes: a geological perspective. *Science* 256, 622–627. <https://doi.org/10.1126/science.1585174>.
- Komai, Y., 1998. Augmented respiration in a flying insect. *J. Exp. Biol.* 201, 2359–2366. <https://doi.org/10.1242/jeb.201.16.2359>.
- Lancaster, J., Downes, Barbara, J., 2013. *Aquatic entomology*. Oxford University Press.
- Lee, D.J., Matthews, P.G.D., 2024. Mechanisms of sustaining oxygen extraction efficiency in dragonfly nymphs during aquatic hypoxia. *bioRxiv*. <https://doi.org/10.1101/2024.06.12.598728>.
- Lehmann, P., Javal, M., Terblanche, J.S., 2019. Oxygen limitation is not the cause of death during lethal heat exposure in an insect. *Biol. Lett.* 15, 20180701. <https://doi.org/10.1098/rsbl.2018.0701>.
- Li, C., Jackson, R.M., 2002. Reactive species mechanisms of cellular hypoxia-reoxygenation injury. *Am. J. Physiol.-Cell Physiol.* 282 (2), C227–C241.
- Matthews, P.G.D., White, C.R., 2011. Regulation of gas exchange and haemolymph pH in the cockroach *Nauphoeta cinerea*. *J. Exp. Biol.* 214, 3062–3073. <https://doi.org/10.1242/jeb.053991>.
- Mazerolle, M.J., Mazerolle, M.M.J., 2017. Package 'AICcmodavg'. R package, p. 281.

- Merritt, R.W., Cummins, K.W., Berg, M.B., 2008. An introduction to the aquatic insects of North America. Kendall Hunt, Kendall, 1996.
- Millar, T.M., Phan, V., Tibbles, L.A., 2007. ROS generation in endothelial hypoxia and reoxygenation stimulates MAP kinase signaling and kinase-dependent neutrophil recruitment. *Free Rad. Biol. Med.* 42 (8), 1165–1177.
- Murphy, M.P., 2009. How mitochondria produce reactive oxygen species. *Biochem. J.* 417, 1–13. <https://doi.org/10.1042/BJ20081386>.
- Poole, D.C., Pittman, R.N., Musch, T.I., Østergaard, L., 2020. August Krogh's theory of muscle microvascular control and oxygen delivery: a paradigm shift based on new data. *J. Physiol.* 598, 4473–4507. <https://doi.org/10.1113/JP279223>.
- R Core Team, 2022. R: A language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Rowe, T.T.C., Gutbrod, M.S., Matthews, P.G.D., 2022. Discontinuous gas exchange in Madagascar hissing cockroaches is not a consequence of hysteresis around a fixed PCO₂ threshold. *J. Exp. Biol.* 225. <https://doi.org/10.1242/jeb.242860>.
- Sanz, A., López-Rodríguez, M.J., García-Mesa, S., Trenzado, C.E., Ferrer, R.M., Tierno De Figueroa, J.M., 2017. Are antioxidant capacity and oxidative damage related to biological and autecological characteristics in aquatic insects? *J. Limnol.* 76, 170–181. <https://doi.org/10.4081/jlimnol.2016.1581>.
- Semenza, G.L., 2011. Hypoxia-inducible factor 1: regulator of mitochondrial metabolism and mediator of ischemic preconditioning. *Biochimica et Biophysica Acta (BBA) - Mole. Cell Res.* 1813 (7), 1263–1268.
- Shchukarev, S.A., Tolmacheva, T.A., 1968. Solubility of oxygen in ethanol – water mixtures. *J. Struct. Chem.* 9, 16–21.
- Shields, H.J., Traa, A., Van Raamsdonk, J.M., 2021. Beneficial and detrimental effects of reactive oxygen species on lifespan: a comprehensive review of comparative and experimental studies. *Front. Cell Dev. Biol.* 9, 1–27. <https://doi.org/10.3389/fcell.2021.628157>.
- Starkov, A.A., 2008. The role of mitochondria in reactive oxygen species metabolism and signaling. *Ann. N. Y. Acad. Sci.* 1147, 37–52. <https://doi.org/10.1196/annals.1427.015>.
- Speers-Roesch, B., Norin, T., Driedzic, W.R., 2018. The benefit of being still: energy savings during winter dormancy in fish come from inactivity and the cold, not from metabolic rate depression. *Proc. R. Soc. B Biol. Sci.* 285, 20181593. [10.1098/rspb.2018.1593](https://doi.org/10.1098/rspb.2018.1593).
- Tripplehorn, C.A., Johnson, N.F., 2004. Borror and DeLong's introduction to the study of insects. Thomson Brooks/Cole, Belmont, California.
- Vanderkooi, J.M., Erecinska, M., Silver, I.A., 1991. Oxygen in mammalian tissue: methods of measurement and affinities of various reactions. *Am. J. Physiol. - Cell Physiol.* 260. <https://doi.org/10.1152/ajpcell.1991.260.6.c1131>.
- Verberk, W.C.E.P., Bilton, D.T., Calosi, P., Spicer, J.I., 2011. Oxygen supply in aquatic ectotherms: partial pressure and solubility together explain biodiversity and size patterns. *Ecology* 92, 1565–1572. <https://doi.org/10.1890/10-2369.1>.
- Verberk, W.C.E.P., Atkinson, D., 2013. Why polar gigantism and Palaeozoic gigantism are not equivalent: effects of oxygen and temperature on the body size of ectotherms. *Funct. Ecol.* 27, 1275–1285. <https://doi.org/10.1111/1365-2435.12152>.
- Verberk, W.C.E.P., Bilton, D.T., 2013. Respiratory control in aquatic insects dictates their vulnerability to global warming. *Biol. Lett.* 9, 0–3. <https://doi.org/10.1098/rsbl.2013.0473>.
- Wang, L., Cui, S., Liu, Z., Ping, Y., Qiu, J., Geng, X., 2018. Inhibition of mitochondrial respiration under hypoxia and increased antioxidant activity after reoxygenation of *Tribolium castaneum*. *PLoS ONE* 13 (6), e0199056.
- Wigglesworth, V.B., 1983. The physiology of insect tracheoles. *Adv In Insect Phys* 17, 85–149.
- Wittenberg, B.A., Wittenberg, J.B., 1989. Transport of oxygen in muscle. *Annu. Rev. Physiol.* 51 (1), 857–878.
- Woods, H.A., 1999. Egg-mass size and cell size: effects of temperature on oxygen distribution. *Am. Zool.* 39, 244–252. <https://doi.org/10.1093/icb/39.2.244>.
- Woods, H.A., Moran, A.L., 2020. Reconsidering the oxygen-temperature hypothesis of polar gigantism: successes, failures, and nuance. *Integr. Comp. Biol.* 60, 1438–1453. <https://doi.org/10.1093/icb/icaa088>.
- Xie, L., Flippin, J.L., Deighton, N., Funk, D.H., Dickey, D.A., Buchwalter, D.B., 2009. Mercury(II) bioaccumulation and antioxidant physiology in four aquatic insects. *Environ. Sci. Technol.* 43, 934–940. <https://doi.org/10.1021/es802323r>.
- Zhang, Q., Yan, Q., Yang, H., Wei, W., 2019. Oxygen sensing and adaptability won the 2019 Nobel Prize in Physiology or medicine. *Genes Dis* 6, 328–332. <https://doi.org/10.1016/j.gendis.2019.10.006>.
- Zorov, D.B., Juhaszova, M., Sollott, S.J., 2014. Mitochondrial reactive oxygen species (ROS) and ROS-induced ROS release. *Physiol. Rev.* 94, 909–950. <https://doi.org/10.1152/physrev.00026.2013>.