

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/32521>

Please be advised that this information was generated on 2021-06-14 and may be subject to change.

FORMAL TOPOLOGY AND CONSTRUCTIVE MATHEMATICS: THE GELFAND AND STONE-YOSIDA REPRESENTATION THEOREMS

THIERRY COQUAND AND BAS SPITTERS

ABSTRACT. We present a constructive proof of the Stone-Yosida representation theorem for Riesz spaces motivated by considerations from formal topology. This theorem is used to derive a representation theorem for f -algebras. In turn, this theorem implies the Gelfand representation theorem for C^* -algebras of operators on Hilbert spaces as formulated by Bishop and Bridges. Our proof is shorter, clearer, and we avoid the use of approximate eigenvalues.

1. INTRODUCTION

This paper illustrates the relevance of locale theory for constructive mathematics. We present a constructive proof of the Stone-Yosida representation theorem for Riesz spaces motivated by considerations from formal topology. This theorem is used to derive a representation theorem for f -algebras. In turn, this theorem implies the Gelfand representation theorem for C^* -algebras of operators on Hilbert spaces as formulated by Bishop and Bridges [BB85]. Our proof is shorter, clearer, and we avoid the use of approximate eigenvalues.

The article is organized as follows. After dealing with some preliminaries we prove a pointfree Stone-Yosida representation theorem for Riesz spaces. In the next section this is used to obtain a representation theorem for f -algebras, which in turn is used to prove the Gelfand representation theorem. Next we discuss the similarity between Bishop's notion of compactness, i.e. complete and totally bounded, and compact overt spaces in formal topology. Finally we show that the axiom of dependent choice is needed to construct points in the formal spectrum.

We would like to stress that the present theory needs few foundational commitments, we work within Bishop-style mathematics. Moreover, the mathematics is predicative, even finitary, and we will not use the axiom of choice, even countable choice, unless explicitly stated.

2. RIESZ SPACES

We present a Stone-Yosida representation theorem for Riesz spaces. This theorem states that a Riesz space with a strong unit may be represented as a Riesz

2000 *Mathematics Subject Classification.* 03F60 Constructive and recursive analysis; 46S30 Constructive functional analysis; 06D22 Frames, locales.

Key words and phrases. Formal topology, constructive mathematics, Riesz space, f -algebra, axiom of choice.

This is an expanded version of our paper [CS05a]. For the convenience of the reader we have included more details and added a few clarifications. There are no new results. We are grateful to Bob Lubarsky and Fred Richman for suggesting improvements in the presentation.

Bas Spitters was supported by the Netherlands Organization for Scientific Research (NWO).

space of functions. In fact, one can even start with an l-group, or lattice ordered group, with a strong unit and construct a Riesz space from this, see [Coq05](sec. 3).

2.1. General definitions.

Definition 1. A Riesz space R is a \mathbb{Q} vector space with a compatible binary sup operation — that is, such that $a+(b\vee c) = (a+b)\vee(a+c)$ and, moreover, $\lambda a \geq 0$ and $-a \leq 0$, whenever a in R , λ in \mathbb{Q} , $a \geq 0$ and $\lambda \geq 0$. One can prove that A Riesz space (or vector lattice) is a partially ordered linear space which is a distributive lattice.

As usual we define $a^+ := a\vee 0$, $a^- := (-a)\vee 0$, $|a| := a^+ + a^-$ and $a \wedge b := -(-a \vee -b)$. One can prove that a Riesz space is a lattice and that the lattice operations are compatible with the vector space operations, see [Bir67][LZ71][Bou64].

Definition 2. A strong unit 1 in an ordered vector space R is a positive¹ element such that for all $a \in R$ there exists a natural number n such that $a \leq n \cdot 1$.

We will now consider a Riesz space R with a strong unit. When q is a rational number, we will often write $a \leq q$ to mean $a \leq q \cdot 1$.

Definition 3. A representation of R is a linear map $\sigma : R \rightarrow \mathbb{R}$ such that $\sigma(1) = 1$ and $\sigma(a \vee b) = \sigma(a) \vee \sigma(b)$.

Such a representation automatically preserves all the Riesz space structure.

Example 1. If X is a compact space, then $C(X)$, its space of continuous functions, is a Riesz space where the supremum is taken pointwise. Each point of x defines a representation $\sigma_x(f) := f(x)$.

In Example 2 we show that a complete commutative algebra of Hermitian operators on a Hilbert space is a Riesz space.

2.2. Spectrum. Given a Riesz space R , we will define a lattice that may be used to define the spectrum of R as a formal space, the points of which are then precisely the representations of R .

Let P denote the set of positive elements of a Riesz space R . For a, b in P we define $a \preceq b$ to mean that there exists n such that $a \leq nb$. We write $a \approx b$ for $a \preceq b$ and $a \succ b$. The following proposition is proved in [Coq05] and involves only elementary considerations on Riesz spaces.

Proposition 1. We write $L(R)$ for the quotient of P by \approx . Then $L(R)$ is a distributive lattice. In fact, if we define $D : R \rightarrow L(R)$ by $D(a) := [a^+]$, then $L(R)$ is the free lattice generated by $\{D(a) | a \in R\}$ subject to the following relations:

1. $D(a) = 0$, if $a \leq 0$;
2. $D(1) = 1$;
3. $D(a) \wedge D(-a) = 0$;
4. $D(a + b) \leq D(a) \vee D(b)$;
5. $D(a \vee b) = D(a) \vee D(b)$.

We have $D(a) \leq D(b)$ if and only if $a^+ \preceq b^+$ and $D(a) = 0$ if and only if $a \leq 0$.

For a in R and rational numbers p, q , we write $a \in (p, q) := (a - p) \wedge (q - a)$. Notice that this is an element of R by our convention that $(a - q)$ means $(a - q \cdot 1)$.

¹We follow the standard terminology using ‘positive’ to mean non-negative.

Lemma 1. $a \in (p, q) \preceq a \in (p, s) \vee a \in (t, q)$, whenever t, s are rational numbers such that $t < s$. Since our Riesz space has a strong unit, for each a , there exists p and q such that $p < q$ and $a \in (p, q) = 1$. Moreover, if I_0, \dots, I_n are open intervals covering (p, q) , then $\bigvee a \in I_i \approx 1$.

Proof. We may assume that $p < t < s < q$.

$$\begin{aligned} a \in (p, s) \vee a \in (t, q) &= ((a - p) \wedge (s - a)) \vee ((a - t) \wedge (q - a)) \\ &= ((a - p) \vee (a - t)) \wedge ((a - t) \vee (s - a)) \wedge \\ &\quad ((a - p) \vee (q - a)) \wedge ((q - a) \vee (s - a)) \\ &= (a - p) \wedge ((a - t) \vee (s - a)) \wedge \\ &\quad ((a - p) \vee (q - a)) \wedge (q - a) \\ &\geq ((a - p) \vee (q - a)) \wedge ((a - t) \vee (s - a)). \end{aligned}$$

We claim that $((a - t) \vee (s - a)) \geq \frac{t-s}{2}$. To show this we write $m := \frac{t+s}{2}$. Then

$$(a-s) \vee (t-a) = (a-m+(m-s)) \vee (t-m-(a-m)) = \frac{t-s}{2} + ((a-m) \vee -(a-m)) \geq \frac{t-s}{2}.$$

The proof is finished by the observation that $b \preceq c$ whenever for some ε , $\varepsilon \cdot 1 \wedge b \leq c$. \square

Lemma 2. If $D(b_1) \vee \dots \vee D(b_n) = 1$, then there exists $r > 0$ such that $D(b_1 - r) \vee \dots \vee D(b_n - r) = 1$.

Proof. Since $\frac{1}{N} \leq b_1^+ \vee \dots \vee b_m^+$ we see that

$$\begin{aligned} \frac{1}{2N} &\leq (b_1^+ \vee \dots \vee b_m^+) - \frac{1}{2N} \\ &= (b_1^+ - \frac{1}{2N}) \vee \dots \vee (b_m^+ - \frac{1}{2N}) \\ &\leq (b_1 - \frac{1}{2N})^+ \vee \dots \vee (b_m - \frac{1}{2N})^+. \end{aligned}$$

\square

The previous lemma is used to prove the following result, which can be found as Theorem 1.11 in [Coq05]. We will not need this, but only state it as a motivation.

Theorem 1. Define Σ , the spectrum of R , to be the locale generated by the elements $D(a)$ and the relations in Proposition 1 together with the relation $D(a) = \bigvee_{r>0} D(a - r)$. Then Σ is a compact completely regular locale.

Compact completely regular locales are the pointfree analogues of compact Hausdorff spaces. Moreover, the points of Σ can be identified with representations of R . In fact, if a representation σ is given, then $\sigma \in D(a)$ if and only if $\sigma(a) > 0$.

2.3. Normable elements. Dedekind cuts may be used to define real numbers, but it should be noted that constructively one needs to require that such cuts (L, U) are *located*, i.e. either $p \in L$ or $q \in U$, whenever $p < q$. Upper cuts in the rational numbers may be conveniently used to deal with certain objects that classically would also be real numbers, we call them *upper real* numbers, see [Ric98][Vic05]. In general, such a cut does not have a greatest lower bound in \mathbb{R} . If it does the upper real number is called *located* or simply a real number. Define the upper real $U(a) := \{q \in \mathbb{Q} \mid \exists q' < q. a \leq q' \cdot 1\}$ for each element of the Riesz space. If it is

located a is said to be *normable* and the greatest lower bound is denoted by $\sup a$. Then we have $\sup a < q$ if and only if $q \in U(a)$.

Proposition 2. *If all elements of R are normable, then the predicate $\text{Pos}(a) := \sup a > 0$ has the following properties:*

1. *If $\text{Pos}(a)$ and $D(a) \leq D(b)$, then $\text{Pos}(b)$;*
2. *If $\text{Pos}(a \vee b)$, then $\text{Pos}(a)$ or $\text{Pos}(b)$;*
3. *If r is a strictly positive rational number, then $\text{Pos}(a)$ or $D(a - r) = 0$.*

We note that $\text{Pos}(a)$ if and only if $\text{Pos}(a^+)$.

In fact, in localic terms this shows that Σ is *open*, or *overt*, see [Joh84], but we will not need this. In formal topology one would say that the formal space has a positivity predicate.

We remark that the standard terminologies from the two different fields seem to conflict. Clearly, it is not the case that $\text{Pos}(a)$ as soon as a is positive, i.e. $a \geq 0$. In particular, 0 is positive, but $\text{Pos}(0)$ does not hold.

In order to prove Proposition 2 we first need three lemmas.

Lemma 3. $\sup(a \vee b) = \sup a \vee \sup b$.

Proof. \geq : Suppose that $a \vee b \leq q' < q$, then $a, b \leq q'$, so both $\sup a \leq q'$ and $\sup b \leq q'$. Thus $q' \geq \sup a \vee \sup b$.

\leq : If $q' \geq a, b$, then $q' \geq a \vee b$ and hence $\sup a \vee \sup b \leq \sup a \vee \sup b$. □

Lemma 4. *Let r be a rational number. If $\sup b < r$ and $r < \sup(b \vee c)$, then $r < \sup c$.*

Proof. If $\sup b < r$, then $b \leq r' < r$, for some rational number r' . So, using Lemma 3

$$r < \sup(b \vee c) = \sup b \vee \sup c \leq r' \vee \sup c.$$

Consequently, $r < \sup c$. □

Lemma 5. *If $0 < \sup(b_1 \vee b_2)$, then $\sup b_1 > 0$ or $\sup b_2 > 0$.*

Proof. Suppose that $0 < r < \sup(b_1 \vee b_2)$. Either $\sup b_1 < r$ or $\sup b_1 > 0$. In latter case we are done. In the former case $\sup b_2 > r$ by Lemma 4. □

Proof. [of Proposition 2]

Property 1 is clear.

Property 2 is Lemma 5.

Finally, to prove property 3 we decide whether $\sup a > 0$ or $\sup a < r$. In the former case $\text{Pos}(a)$. In the latter case $D(a - r) = 0$. □

Corollary 1. *If $D(a_1) \vee \dots \vee D(a_n) = 1$, we can find $i_1 < \dots < i_k$ such that $D(a_{i_1}) \vee \dots \vee D(a_{i_k}) = 1$ and $\text{Pos}(a_{i_1}), \dots, \text{Pos}(a_{i_k})$.*

Proof. By Lemma 2 there exists $r > 0$ such that $D(a_1 - r) \vee \dots \vee D(a_n - r) = 1$. By Proposition 2 for all i , $D(a_i - r) = 0$ or $\text{Pos}(a_i)$. From this the result follows. □

Lemma 6. *Let $I := (p, q)$, $J := (r, s)$. Define $I + J := (p + r, q + s)$ and $I \vee J := (p \vee r, q \vee s)$. If $|a + b - c| \leq \varepsilon$ and $\text{Pos}(a \in I \wedge b \in J \wedge c \in K)$, then the distance between $I + J$ and K is bounded by ε . If $|a \vee b - c| \leq \varepsilon$ and $\text{Pos}(a \in I \wedge b \in J \wedge c \in K)$, then the distance between $I \vee J$ and K is bounded by ε .*

Finally, if $|a - b| \leq \varepsilon$ and $\text{Pos}(a \in I \wedge b \in J)$, then the distance between I and J is bounded by ε .

Proof. We only prove the last fact. If the distance between I and J is bigger than ε , then $r - q > \varepsilon$ or $p - s > \varepsilon$. Consequently, $D(q - a \wedge b - r) = 0$ or $D(a - p \wedge s - b) = 0$. Both cases imply that $D(a \in I \wedge b \in J) = 0$. \square

Definition 4. A Riesz space R is separable if there exists a sequence a_n such that for all a and $\varepsilon > 0$, there exists n such that $|a - a_n| \leq \varepsilon$.

Theorem 2. [DC] Let R be a separable Riesz space all elements of which are normable. Assume that $\text{Pos}(a)$, then there exists a representation σ such that $\sigma(a) > 0$.

Proof. We write $\varepsilon_n := 2^{-n}$. Using dependent choice and Lemma 1 we define a sequence (q_n) of rationals such that

$$\text{Pos}(a \in (\frac{\sup a}{2}, \sup a) \wedge a_0 \in (q_0 - \varepsilon_0, q_0 + \varepsilon_0) \wedge \dots \wedge a_n \in (q_n - \varepsilon_n, q_n + \varepsilon_n)).$$

If $b \in R$, we can find a sequence of elements a_{n_k} such that for any $\varepsilon > 0$ we have $|b - a_{n_k}| \leq \varepsilon$ when k is large enough. Then q_{n_k} is a Cauchy sequence and we define $\sigma(b) := \lim_k q_{n_k}$. By Lemma 6 this definition does not depend on the choice of the sequence a_{n_k} . The map σ is a representation such that $\sigma(a) > 0$ and $\sigma(a_n) \in (q_n - \varepsilon_n, q_n + \varepsilon_n)$ for all n . \square

A suggestive way to state that $\sigma(a) > 0$ is to say that σ is a *point* in $D(a)$.

Let Σ be the set of representations of R . We call Σ the *spectrum* of R . Each representation is a bounded linear functional. Each element a of R defines a pseudo norm $\rho_a(\varphi) := |\varphi(a)|$ on the space of bounded linear functionals. If R is separable and a_n is a dense sequence in $\{a \in R : |a| \leq 1\}$, we can collect, like Bishop, all the pseudo-norms into one norm $\rho(\varphi) := \sum_n 2^{-n} |\varphi(a_n)|$. Considering the restriction of these pseudo norm to the spectrum, which is not a linear space, we obtain a pseudo metric.

We have the following Stone-Yosida representation theorem, see [Sto41][Yos42].

Theorem 3. [DC] Let R is a separable Riesz space all elements of which are normable. The spectrum Σ is a complete totally bounded metric space. For a in R , we define $\hat{a} : \Sigma \rightarrow \mathbb{R}$ by $\hat{a}(\sigma) := \sigma(a)$ and $\|a\| := \sup(|a|)$. Then $\sup_\sigma |\hat{a}(\sigma)| = \|a\|$. Finally, the set of functions \hat{a} is dense in $C(\Sigma)$.

Proof. We define $U_{ars} := D(a \in (r, s))$ as an element of the lattice $L(R)$. Let $\varepsilon > 0$ and a_1, \dots, a_n in R . For each i we construct, using Lemma 1, finitely many s_{ij}, r_{ij} such that $s_{ij} - r_{ij} < \varepsilon$ and $\bigvee_j U_{a_i r_j s_j} = 1$ in the lattice $L(R)$. By Corollary 1 one can assume all these elements to be positive. Each of them contains a point σ_{ij} by Theorem 2. If τ in $U_{a_i r_j s_j}$, then $\rho_{a_i}(\tau, \sigma_{ij}) = |\tau(a) - \sigma_{ij}(a)| < s_{ij} - r_{ij}$. Since there are finitely many $U_{a_i r_j s_j}$ covering Σ , the collection of these points forms an ε -net for the pseudonorms ρ_{a_i} . Consequently, Σ is totally bounded.

It is straightforward to show that Σ is also complete as a uniform space.

For each $\sigma \in \Sigma$ we have $|\sigma(a)| \leq \|a\|$. To see this suppose that $\sigma(a) > \|a\|$. Then there exists $\varepsilon > 0$ such that $\sigma(a) - a \geq \varepsilon 1$, however $\sigma(\sigma(a) - a) = 0$. If $r < \|a\|$, then by Theorem 2, there exists σ such that $r < |\sigma(a)|$.

Finally, the density follows from Proposition 3.1 in [Coq05]. Its proof involves only elementary properties of Riesz spaces. \square

Notice the interplay between the *pointwise* and *pointfree* framework. From a formal covering, in the lattice $L(R)$, it is possible to deduce that Σ , a metric space, is totally bounded. This is remarkable since there are examples of Riesz spaces R such that in a recursive interpretation of Bishop's mathematics Σ does not have enough points. For instance, consider the Riesz space R of continuous real functions on Cantor space (2^ω) . The spectrum of R is precisely Cantor space and the representations are its points², which is known not to have enough points in a recursive interpretation. In particular, this means that we have a collection of open sets which covers all the recursive points, but does not allow a finite subcover. This is possible since this collection does not cover the space in the usual terminology of formal topology.

3. F-ALGEBRAS

In this section we apply the results of the previous section to f-algebras.

Definition 5. *An f-algebra is a Riesz space with a strong unit and a commutative³ multiplication such that $0 \leq ab$, whenever $0 \leq a$ and $0 \leq b$.*

3.1. f-algebra of operators.

Example 2. *If R is a complete commutative algebra of normable self-adjoint operators on a Hilbert space H , then R is a Riesz space with the order \leq defined by $0 \leq A$ if and only if $(Au, u) \geq 0$ for all u in H .*

In the rest of this subsection we prove that if $A, B \geq 0$, then $AB \geq 0$.

We have now defined two notions of boundedness on the algebra of operators. One as a bounded operator: A is bounded by a if for all x , $\|Ax\|^2 \leq a\|x\|^2$. The other from the ordering: A is bounded by a if $A \leq aI$, where I is the identity operator.

Lemma 7. *The two notions of boundedness coincide — that is, for all x , $(Ax, x) \leq (ax, x)$ if and only if for all x , $\|Ax\|^2 \leq a^2\|x\|^2$. Consequently, $\|A^2\| = \|A\|^2$.*

Proof. The usual proof, for instance in [Lan83] using the polarization identity, is constructive. \square

Since $(AB^2x, x) = (A(Bx), (Bx))$, we see that $AB^2 \geq 0$, whenever $A \geq 0$. This suffices to prove that R is an ordered ring.

Lemma 8. [Rie32] (p33, footnote 9) *Let R be as above. Then every positive element is the uniform limit of a sum of squares.*

Proof. We can assume that $0 \leq A \leq 1$. Define $A_0 := A$ and $A_{n+1} := A_n - A_n^2$. Then $0 \leq A_{n+1} \leq 1$, since $A_{n+1} = A_n(1 - A_n)^2 + (1 - A_n)A_n^2 \geq 0$ and $1 - A_{n+1} = 1 - A_n + A_n^2$. Moreover, $A_{n+1} = A_n - A_n^2 \leq A_n$. Since $A = A_1^2 + \dots + A_n^2 + A_{n+1}$, we have $A_n^2 \leq 1/n \rightarrow 0$. By Lemma 7 this implies that $A_n \rightarrow 0$. \square

Corollary 2. *$AB \geq 0$, whenever $A, B \geq 0$.*

²To see this, note that the characteristic function χ_u of any basic open u is continuous. So, given a representation σ , $\sigma(\chi_u \wedge (1 - \chi_u)) = 0$ and $\sigma(\chi_u \vee (1 - \chi_u)) = 1$. Consequently, $\sigma(\chi_u)$ is either 0 or 1. It follows that the representations can be identified with the points.

³One can prove classically that the commutativity requirement follows from the other properties of an f-algebra. We intend to provide a constructive proof of this separately using the pointfree description of the spectrum.

Proof. If $A, B \geq 0$, then $(ABx, x) = \sum (AB_n x, B_n x) \geq 0$, where B_n is a sequence such that $\sum B_n^2$ converges to B . \square

The following lemma shows that one can construct the square root⁴ when R is complete and thus one can define the absolute value as $|A| := \sqrt{A^2}$. From the absolute value one first defines $A^+ := (|A| + A)/2$ and then $A \vee B := A + (B - A)^+$. Consequently, the algebra is a Riesz space and an f-algebra.

Lemma 9. *For all $A \geq 0$ we can build a Cauchy sequence (A_n) of positive elements such that $A_n^2 \rightarrow A$.*

Proof. We can assume $0 \leq A \leq I$. We define the two sequences $A_n \in [0, I]$ and $r_n \in [0, 1]$ defined by $A_0 = 0$ and $r_0 = 0$ and

$$A_{n+1} = \frac{1}{2}(1 - A + A_n^2) \quad r_{n+1} = \frac{1}{2}(1 + r_n^2)$$

Clearly, we have $A_n \leq r_n$ for all n .

We claim that we have for all n

$$A_n \leq A_{n+1} \quad r_n \leq r_{n+1} \quad A_{n+1} - A_n \leq r_{n+1} - r_n$$

This is proved by induction from the equalities

$$A_{n+1} - A_n = \frac{1}{2}(A_n + A_{n-1})(A_n - A_{n-1}) \quad r_{n+1} - r_n = \frac{1}{2}(r_n + r_{n-1})(r_n - r_{n-1})$$

It follows that we have

$$(I - A_n)^2 - A = 2(A_{n+1} - A_n) \leq 2(r_{n+1} - r_n)$$

In order to conclude, all is left is to show that (r_n) has limit 1. We know that $0 \leq r_n \leq r_{n+1} \leq 1$ and we have

$$1 - r_{n+1} = \frac{1}{2}(1 - r_n^2) = (1 - r_n)\frac{1}{2}(1 + r_n) \leq (1 - r_n)(1 - \frac{\epsilon}{2})$$

if $r_n \leq 1 - \epsilon$. This shows that if $(1 - \frac{\epsilon}{2})^N \leq \epsilon$ we have $1 - r_n \leq \epsilon$ for all $n \geq N$. \square

3.2. Gelfand representation. Any f-algebra is a Riesz space so we have a Gelfand representation of the f-algebra qua Riesz space, see Theorem 3.

Theorem 4. *The Gelfand transform $\hat{\cdot}$ preserves multiplication.*

Proof. Since $2ab = (a + b)^2 - a^2 - b^2$. We need to prove that $\hat{\cdot}$ preserves squares — that is $\sigma(a^2) = \sigma(a)^2$. For this we first prove: $\sigma(ab) > 0$, whenever $\sigma(a), \sigma(b) > 0$.

If $\sigma(a) \geq r > 0$, then $\sigma(a - r) > 0$. By Lemma 6.3 in [Coq05], $(a - r)^+ \wedge b^+ \leq \frac{1}{r}(ab)^+$, so $\sigma(ab)^+ \geq r(\sigma(b)^+ \wedge \sigma(a - r)^+) > 0$, which was to be proved.

Suppose that $|\sigma(a)| < q$. Then $q > \sigma(a)$ and so $\sigma(q - a) > 0$. Similarly, $\sigma(q + a) > 0$. Consequently, $\sigma(q^2 - a^2) = \sigma((q - a)(q + a)) > 0$. By a similar argument we see that if $|\sigma(a)| < q$, then $|\sigma(a^2)| < q^2$. We conclude that $\sigma(a^2) = \sigma(a)^2$. \square

We have proved the following representation theorem for f-algebras which explains the name f-algebra: an f-algebra is an abstract function algebra.

⁴This is the usual lemma that R admits square root of positive elements if R is complete. Notice that the proof is directly constructive, and it corresponds to the usual Taylor expansion of $(1 - x)^{1/2}$.

Theorem 5. [DC] *Let \mathcal{A} be a separable f -algebra of normable elements, then the spectrum Σ is a compact metric space and there exists an f -algebra embedding of \mathcal{A} into $C(\Sigma)$.*

We now specialize this theorem to the f -algebra in Example 2 and obtain Bishop's version of the Gelfand representation theorem. In fact, like Bishop we first prove the theorem for Hermitian operators. As a corollary we obtain the Gelfand duality theorem for a separable Abelian C^* -algebra, exactly as stated by Bishop [BB85] Cor.8.28 by considering its self-adjoint part which is an f -algebra.

Corollary 3. [DC] *Let \mathcal{A} be a separable f -algebra of normable Hermitian operators on a Hilbert space, then the spectrum is a compact metric space and there exists an f -algebra embedding of \mathcal{A} into $C(\Sigma)$.*

Theorem 6. [DC] *Let R be an Abelian C^* -algebra of operators on a Hilbert space. Then there exists a C^* -algebra embedding φ of R into $C(\Sigma, \mathbb{C})$, where Σ is a compact metric space. Moreover, $\varphi(1) = 1$ and R is norm-dense.*

Bishop's Gelfand representation theorem states that for any commutative algebra of normable operators on a separable Hilbert space there exists a norm-preserving isomorphism to the algebra of continuous functions on its spectrum. To prove that this map is norm-preserving Bishop proves that certain ε eigenvectors can be computed. In fact, the computational information of the ε eigenvectors is used only to prove the non-computational statement that the map is norm-preserving. In contrast, we work directly on the approximations so that we can avoid these unused computational steps.

3.3. Peter-Weyl. For a typical application, we let G be a compact group and R be the algebra of operators over $L_2(G)$ generated by the unit operator and the operators $T(f)(g) := f * g$, where $*$ denotes the convolution product. Each operator $T(f)$ is compact and hence normable. The non-trivial representations of R are then exactly the characters of the group G . This gives a reduction of the Peter-Weyl theorem to the Gelfand representation theorem, see [CS05b].

4. COMPACT OVERT LOCALES

Bishop defines a metric space to be compact if it is complete and totally bounded and proves that all uniformly continuous functions defined on such a metric space are normable. In contrast, in the framework of locale theory it is not true in general that all functions on a compact regular locale are normable, i.e. the norm is only defined as an upper real which may not be a located.

However, the locale considered in Theorem 1 is not only compact completely regular, but also *overt* — that is, has a positivity predicate — as shown by Proposition 2. In this case all the continuous functions are normable. This is a general fact.

Theorem 7. *If X is a compact completely regular locale, then X is overt if and only if for any $f \in C(X)$ there exists $\sup f \in \mathbb{R}$ such that $\sup f < s$ if and only if $f^{-1}(-\infty, s) = X$.*

Proof. We prove only the 'only if' part. If X is overt and $f \in C(X)$, then an approximation of the supremum can be found by considering a finite covering of X by positive opens of the form $f^{-1}(r, s)$, where $s - r$ is small. \square

The previous facts suggest a similarity between Bishop’s compact metric spaces and the compact overt spaces in formal topology.

Bishop compact \Leftrightarrow compact overt

Clearly this requires further developments building on ideas in [ML70][Joh84]. However, we postpone this to further work.

It is interesting to note that Paul Taylor has independently found a similar relation between Bishop compact and compact overt in the context of his abstract Stone duality [Tay05]. He also introduced the term *overt*.

We would like to conclude this discussion with the following comparison between the three following frameworks: classical mathematics with the axiom of choice, Bishop’s mathematics and our framework, predicative constructive mathematics without dependent choice⁵. Using classical logic and the axiom of choice one can show that the spectrum defined in Theorem 1 has enough points [Joh82]. Thus in this setting the pointfree and pointwise description of the space coincide. In a recursive interpretation of Bishop’s framework these descriptions differ. However, using dependent choice, normability and separability assumptions, we have shown how to deduce that Σ is totally bounded from the pointfree description of Σ .

Our conclusion is that the best formulation of the representation theorem is the pointfree one, since, besides being neutral on the use of the axiom of choice and classical logic, it implies the usual formulations both in Bishop’s framework and in classical mathematics.

5. CHOICE

5.1. No points. As mentioned before, when all the elements of the algebra are normable, one can construct points in the spectrum using dependent choice. We claim that dependent choice is needed for this. In fact, it is known that there exist compact overt locales for which we need countable choice to construct a point. Thus it suffices to consider the space of continuous functions on such a locale. We think that a nice example can be extracted from [Ric00].

Richman [Ric00](p.5) gave an informal argument that indicates that one can not construct the zeroes of the complex polynomial $X^2 - a$ unless one knows whether $a = 0$ or not. To a in \mathbb{C} we can associate the locale Y_a of roots of $X^2 - a$. The existence of a point in Y_a requires dependent choice. On the other hand using results from [Vic05] it should be possible to show that Y_a is compact overt as an element of the completion of the metric space n -multisets in \mathbb{C} . The metric on this space is the usual Hausdorff metric on compact subsets of \mathbb{C} .

Finally, we remark that this leaves open the question whether it is possible to construct the points of the spectrum of a discrete countable Riesz space over the rationals, or more generally, to construct the points of the spectrum of a separable Riesz space.

5.2. Spreads. It is interesting to note that Richman [Ric02] proposes to use spreads to avoid dependent choice. We suggest to use formal spaces instead. One motivation of formal topology [ML70][Sam87] was precisely to give a direct treatment of Brouwer’s spreads by working with trees of finite sequences. Formal topology may be seen as a predicative and constructive version of locale theory. Johnstone [Joh82]

⁵This framework is related to Richman’s proposal to develop constructive mathematics without using countable choice, see [Ric00].

stresses that one may avoid the use of the axiom of choice in topology by using locale theory and dealing directly with the opens. In this light it may not be so surprising that Richman uses spreads to avoid choice.

Richman's definition of spread differs in two respects from Heyting's definition. The branching of the tree is arbitrary, i.e. not necessarily indexed by the natural numbers, and it is not decidable whether or not a branch can be continued. This may be compared to the present situation where we study the maximal spectrum Σ . When Σ has a countable base, we may define it as a finitely branching tree. When furthermore Σ is overt, every positive branch can be continued in a positive way. One difference between Richman's spreads and our approach is that Richman requires the infinite branches to be elements of a metric space.

6. CONCLUSION

We gave a constructive proof of the Stone-Yosida representation theorem for Riesz spaces. This theorem was used to prove a representation theorem for f -algebras, from which we derived the Gelfand representation theorem for commutative C^* -algebras of operators on a Hilbert space. This constructive theorem generalizes the one by Bishop and Bridges. In a similar way one may prove a generalization of Bishop's spectral theorem, see [Spi05b].

It should be noted that we have used normability and separability hypothesis in the statements of the main theorems and used the axiom of dependent choice. In fact, without these hypothesis we can still obtain the spectrum as a compact locale. The normability is necessary to show that the spectrum is overt. The separability hypothesis is used to obtain a metric space instead of a uniform space. Finally, the axiom of dependent choice is used in Theorem 2 to construct a point in each positive open, and thus obtain a metric space in the sense of Bishop.

In this context, we would like to mention a problem for both constructive versions of the Gelfand representation theorem: can it be applied to construct the Bohr compactification of, say, the real line, like Loomis [Loo53]? Considering that the Stone-ech compactification has been successfully treated in locale theory [Joh82], one would hope that a similar treatment is possible. Since the almost periodic functions do not form an algebra constructively, we may consider the f -algebra of functions generated by them. However, in this algebra not all elements are normable. Any element of the group determines a point in the spectrum. However, it is not possible to extend the group operation to the spectrum and obtain a localic group, since every compact localic group has a positivity predicate [Wra90] and since, moreover, the spectrum is compact this would imply that all the functions in the f -algebra are normable. This, as we stated before, is not the case. See Spitters [Spi05a] and the references therein for a constructive theory of almost periodic functions.

REFERENCES

- [BB85] Errett Bishop and Douglas Bridges. *Constructive analysis*, volume 279 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, 1985.
- [Bir67] Garrett Birkhoff. *Lattice theory*. Third edition. American Mathematical Society Colloquium Publications, Vol. XXV. American Mathematical Society, Providence, R.I., 1967.
- [Bou64] N. Bourbaki. *Algèbre, chapitre 6*. Hermann, 1964.
- [Coq05] Thierry Coquand. About Stone's notion of spectrum. *J. Pure Appl. Algebra*, 197(1-3):141–158, 2005.

- [CS05a] T. Coquand and B. Spitters. Formal Topology and Constructive Mathematics: the Gelfand and Stone-Yosida Representation Theorems. *Journal of Universal Computer Science*, 11(12):1932–1944, 2005. http://www.jucs.org/jucs_11_12/formal_topotoly_and_constructive.
- [CS05b] Thierry Coquand and Bas Spitters. A constructive proof of the Peter-Weyl theorem. *Mathematical Logic Quarterly*, 4:351–359, 2005.
- [Joh82] Peter T. Johnstone. *Stone Spaces*. Number 3 in Cambridge studies in advanced mathematics. Cambridge University press, 1982.
- [Joh84] Peter T. Johnstone. Open locales and exponentiation. In *Mathematical applications of category theory (Denver, Col., 1983)*, volume 30 of *Contemp. Math.*, pages 84–116. Amer. Math. Soc., Providence, RI, 1984.
- [Lan83] Serge Lang. *Real analysis*. Addison-Wesley Publishing Company Advanced Book Program, Reading, MA, second edition, 1983.
- [Loo53] Lynn H. Loomis. *An introduction to Abstract Harmonic Analysis*. University Series in Higher Mathematics. van Nostrand, New York, 1953.
- [LZ71] W. A. J. Luxemburg and A. C. Zaanen. *Riesz spaces. Vol. I*. North-Holland Publishing Co., Amsterdam, 1971. North-Holland Mathematical Library.
- [ML70] Per Martin-Löf. *Notes on constructive mathematics*. Almqvist & Wiksell, Stockholm, 1970.
- [Ric98] Fred Richman. Generalized real numbers in constructive mathematics. *Indagationes Mathematicae*, 9:595–606, 1998.
- [Ric00] Fred Richman. The fundamental theorem of algebra: a constructive development without choice. *Pacific Journal of Mathematics*, 196:213–230, 2000.
- [Ric02] Fred Richman. Spreads and choice in constructive mathematics. *Indagationes Mathematicae*, 13:259–267, 2002.
- [Rie32] F. Riesz. Ueber die linearen Transformationen des komplexen Hilbertschen Raumes. *Acta Sci. Math.*, 5:23–54, 1930–32.
- [Sam87] Giovanni Sambin. Intuitionistic formal spaces - a first communication. In D. Skordev, editor, *Mathematical logic and its Applications*, pages 187–204. Plenum, 1987.
- [Spi05a] Bas Spitters. Almost periodic functions, constructively. *Logical Methods in Computer Science*, 1(3:3):1–7, 2005.
- [Spi05b] Bas Spitters. Constructive algebraic integration theory without choice. *Dagstuhl proceedings*, 2005.
- [Sto41] M. H. Stone. A general theory of spectra. II. *Proc. Nat. Acad. Sci. U. S. A.*, 27:83–87, 1941.
- [Tay05] Paul Taylor. A lambda calculus for real analysis. In Tanja Grubba, Peter Hertling, Hideki Tsuiki, and Klaus Weihrauch, editors, *CCA*, volume 326-7/2005 of *Informatik Berichte*, pages 227–266. FernUniversität Hagen, Germany, 2005.
- [Vic05] Steven Vickers. Localic completion of generalized metric spaces I. *Theory and Applications of Categories*, 14:328–356, 2005.
- [Wra90] G. C. Wraith. Unsurprising results on localic groups. *J. Pure Appl. Algebra*, 67(1):95–100, 1990.
- [Yos42] Kôzaku Yosida. On the representation of the vector lattice. *Proc. Imp. Acad. Tokyo*, 18:339–342, 1942.

E-mail address: `coquand@cs.chalmers.se`

CHALMERS UNIVERSITY,, SWEDEN

E-mail address: `spitters@cs.ru.nl`

RADBOUD UNIVERSITY NIJMEGEN,, THE NETHERLANDS