

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/52122>

Please be advised that this information was generated on 2019-02-18 and may be subject to change.

# Heterozygous alpha-1 antitrypsin deficiency as a co-factor in the development of chronic liver disease: a review

K.F. Kok<sup>1\*</sup>, P.J. Wahab<sup>1</sup>, R.H.J. Houwen<sup>4</sup>, J.P.H. Drenth<sup>5</sup>, R.A. de Man<sup>6</sup>, B. van Hoek<sup>7</sup>, J.W.R. Meijer<sup>2</sup>, F.L.A. Willekens<sup>3</sup>, R.A. de Vries<sup>1</sup>

Departments of <sup>1</sup>Hepato-Gastroenterology, <sup>2</sup>Pathology and <sup>3</sup>Clinical Chemistry, Rijnstate Hospital, Arnhem, the Netherlands, <sup>4</sup>Department of Paediatric Gastroenterology, Wilhelmina Children's Hospital/University Medical Centre Utrecht, the Netherlands, <sup>5</sup>Department of Gastroenterology and Hepatology, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands, <sup>6</sup>Department of Hepato-Gastroenterology, Erasmus MC Rotterdam, the Netherlands, <sup>7</sup>Department of Gastroenterology and Hepatology, Leiden University Medical Centre, the Netherlands, \*corresponding author (presently at <sup>5</sup>): e-mail: karinkok@hetnet.nl

## ABSTRACT

Alpha-1 antitrypsin (A1AT) is an acute-phase protein that is produced in liver cells. A1AT deficiency is a hereditary disease which is defined by the hepatic production of an abnormal protein that can not be released into the plasma. This leads to deficiency of plasma A1AT and subsequently to an impaired protection against proteases, resulting in pulmonary disease. Accumulation of the abnormal protein in hepatocytes can lead to liver damage. Serum level measurement, phenotyping and liver biopsy can be used for establishing the diagnosis.

Homozygous A1AT deficiency can cause neonatal hepatitis; in adults end-stage liver disease, cirrhosis and hepatocellular carcinoma can develop. There are strong arguments to consider heterozygous A1AT deficiency as an important co-factor in the aetiology of chronic liver disease. Studies have shown that A1AT heterozygosity can be considered a modifier for hepatitis C virus, end-stage liver disease, cirrhosis and hepatocellular carcinoma. The accumulation of A1AT in the hepatocytes occurs more profoundly in a diseased liver, and as a consequence it affects the natural course of the liver disease. Therapeutic options include augmentation therapy (infusion of purified human plasma A1AT) in pulmonary disease; in end-stage liver disease liver transplantation is an option. For the future, other interventions such as gene therapy or strategies to inhibit polymerisation are promising.

## KEYWORDS

Alpha-1-antitrypsin deficiency, hepatocellular carcinoma, heterozygosity, liver disease

## INTRODUCTION

Alpha-1 antitrypsin (A1AT) is an acute-phase protein that is produced in liver cells. It is released into the plasma in response to an inflammatory stimulus. A1AT deficiency is a hereditary disease that is defined by the hepatic production of an abnormal protein that can not completely be released into the plasma. This leads to deficiency of plasma A1AT and subsequently to an impaired protection of the lungs against proteases. This results in pulmonary emphysema; hepatic accumulation of the abnormal protein can lead to chronic liver disease. This review gives an update of the present knowledge on partial A1AT deficiency in relation to various liver diseases.

## GENETICS AND (PATHO)PHYSIOLOGY

The A1AT molecule is a serum glycoprotein acting as an acute-phase protein. It is released during inflammatory processes from the hepatocyte, which results in increased plasma concentrations. The major physiological function

J.P.H. Drenth was not involved in the handling and review process of this paper.

of this protein is the inhibition of destructive neutrophil elastase, thus protecting against pulmonary damage.<sup>1</sup>

The A<sub>1</sub>AT protein is encoded by the protease inhibitor (Pi) locus located on chromosome 14q32.1. The Pi locus is highly polymorphic, resulting in different A<sub>1</sub>AT isotypes that can be detected by electrophoresis.

The most common allele is the M allele that results in a functionally normal protein with normal serum A<sub>1</sub>AT levels. The normal A<sub>1</sub>AT protein has a tertiary structure based on a large central  $\beta$ -sheet, surrounded by two other sheets and a reactive centre loop. The reactive centre loop can move in and out of the large  $\beta$ -sheet. At higher temperatures polymerisation can occur between molecules due to insertion of the loop of one molecule into the large  $\beta$ -sheet of the other.

The point mutation, found in the Z variant, destabilises the loop-sheet polymerisation of the A<sub>1</sub>AT molecule, resulting in chains of polymers that are retained in the hepatocytes. These polymers accumulate in the endoplasmic reticulum of the hepatocytes and may be recognised as PAS(+) inclusion bodies (figure 1). Only 15% of the Z variant of A<sub>1</sub>AT

can be secreted into the plasma, the other 85% accumulates in the liver. In Pi ZZ homozygous subjects, this results in a severe deficiency of serum A<sub>1</sub>AT and in accumulation of the abnormal protein in the endoplasmic reticulum of the hepatocyte, which can lead to chronic liver disease.

The S variant of the A<sub>1</sub>AT molecule has less effect on the loop-sheet polymerisation. Formation of S polymers is slower, resulting in less retention of protein in the hepatocytes compared with the Z variant. There is a mild reduction in serum A<sub>1</sub>AT levels. When an S and a Z variant are coinherited, the two interact with the formation of polymers within the hepatocytes; this can lead to reduction in the serum A<sub>1</sub>AT level, inclusion of the polymers, accumulation and subsequently development of cirrhosis. Less frequently found variants are null alleles resulting in undetectable A<sub>1</sub>AT levels due to intracellular degradation or intracellular accumulation of the protein; this is usually associated with severe pulmonary disease, but not with liver disease (table 1).<sup>2-7</sup>


A<sub>1</sub>AT deficiency is characterised by an imbalance between the protease neutrophil elastase and the protease inhibitor A<sub>1</sub>AT. It has been suggested that neutrophil elastase might promote the development of cancer.<sup>8</sup> The exact carcinogenic mechanism or sequence is not known.

In a recent paper, the hypothesis was given that in A<sub>1</sub>AT deficiency, the hepatocytes in which A<sub>1</sub>AT is accumulated are inhibited in their growth, but they do express regenerative signals. Relatively normal cells, without A<sub>1</sub>AT deposits, are thereby stimulated and this chronic stimulation of regeneration may lead to the formation of neoplasms.<sup>9</sup>

#### DIAGNOSTIC ASPECTS

To establish the diagnosis, various methods are available. The A<sub>1</sub>AT serum level can be determined by clinical chemistry. In homozygous A<sub>1</sub>AT deficiency the level is very low. However, in the heterozygous variant the A<sub>1</sub>AT level can be within the normal range, especially during an acute-phase reaction. Therefore, electrophoresis should be performed in any case of suspected A<sub>1</sub>AT deficiency in order to determine the phenotype. Liver biopsy is the

**Figure 1.** Liver biopsy: cirrhosis due to alcohol abuse and heterozygosity for  $\alpha_1$  antitrypsin (A<sub>1</sub>AT) deficiency<sup>33</sup>



A. HE staining, 20x: micro nodular liver cirrhosis.  
B. D-PAS staining, 40x: diastase resistant, Periodic-Schiff-Acid positive globuli, matching with A<sub>1</sub>AT accumulation.  
C. A<sub>1</sub>AT antibodies, 40x: confirmation of A<sub>1</sub>AT accumulation.

**Table 1.** Characteristics of the different A<sub>1</sub>AT alleles

Allele	Mutation	Cellular defect	Enzymatic activity
M	-	-	Normal
S	Glu264Val	Intracellular degradation	Intermediate
Z	Glu342Lys	Intracellular accumulation	Low
Null	Different mutations	Most mutations no mRNA	Null

gold standard for establishing A1AT accumulation and PAS-positive, diastase-resistant inclusions can be found. A specific immunohistochemical staining can confirm the diagnosis. It is also possible to determine the phenotype on paraffin-embedded liver slides.

## EPIDEMIOLOGY

Despite the fact that A1AT deficiency is a common disorder, it is poorly recognised in clinical practice. There are probably two main reasons for this; first in patients with liver disease, the diagnosis is not always considered and second not all subjects with a deficient phenotype develop liver disease, i.e. the penetrance is low.<sup>10</sup>

Given this well-known underdiagnosis, the real prevalence of A1AT deficiency has mainly been determined by epidemiological methods, either using a control cohort from an epidemiological study, or through neonatal screening. The Z allele is especially prevalent in Northern Europe, while the S allele is prevalent in Southern Europe (table 2).<sup>11-15</sup>

## CLINICAL ASPECTS

### Pulmonary disease

A1AT deficiency is associated with less inhibition of elastase resulting in pulmonary disease. A1AT homozygosity (Pi ZZ) results in a pulmonary phenotype with early onset of emphysema, asthma and bronchiectasia. A1AT deficiency results in panacinar pathology and disproportionate emphysematous involvement of the lung bases. Tobacco smoking is the most important additional risk factor for the development of pulmonary disease. Also subjects with the Pi MZ phenotype have an increased risk of developing pulmonary disease.<sup>7</sup>

### Neonatal/paediatric liver disease

A1AT deficiency is the most common genetic cause of liver disease in early childhood. The most common presentation is by prolonged jaundice. The stools generally contain no yellow or green pigment, indicating cholestasis and mimicking biliary atresia. All patients have hepatomegaly and about 50% also have splenomegaly. Approximately 5% of the patients present with an increased bleeding tendency. This is due to vitamin K deficiency caused by the cholestasis-induced malabsorption. Less commonly children present later in childhood with hepatosplenomegaly or with cirrhosis.<sup>16-19</sup>

In Sweden, between 1972 and 1974, 200,000 neonates were screened for A1AT deficiency: 120 Pi ZZ (0.06%), 2 Pi Z-, 54 Pi SZ and 1 Pi S- children were found. Only 14 of the Pi ZZ children had prolonged jaundice, nine of those had severe liver disease. All infants appeared healthy at six months of age. Infants with a Pi SZ phenotype had no signs of liver disease.

At the age of 16 years, elevated liver enzymes were found in 17% of Pi ZZ adolescents and in 8% of Pi SZ adolescents. The adults with liver disease in infancy were clinically healthy. At the age of 26 years, the Pi ZZ subjects were compared with Pi MM individuals. The Pi ZZ subjects had normal lung function; 4 to 9% of them had mild liver test abnormalities.<sup>11,20,21</sup> In the Province of Bozen in Northern Italy, Pi phenotyping in umbilical cord blood was performed as a routine neonatal screening. About 5% of Pi SZ children were affected by liver involvement with elevated liver enzymes and 7% of 833 Pi MZ heterozygotes had elevated liver enzymes in early childhood. At the age of 5 and 10 years, none had liver disease. The serum levels of A1AT were similar in the groups with and without liver test abnormalities; however these values had a wide range.<sup>22,23</sup>

Although these studies suggest a good prognosis for neonatal cholestasis due to A1AT deficiency, other studies have described children who developed severe liver disease.<sup>16,19</sup>

Table 2. Epidemiology<sup>10-15</sup>

Country	Frequency			Method	Year
	Pi ZZ	Pi MZ	Pi SZ		
Sweden	0.06%	-	0.03%	Neonates, phenotyping when serum A1AT <40%	1976
The Netherlands	Z allele 0.03%	-	S allele 0.04%	Neonates, phenotyping when A1AT/transferrine <1,2	1980
France	0.01%	2%	0.15%	Control cohort epidemiological study	2003
Spain	0.01%	-	0.2%	Control cohort epidemiological study	2003
Italy	0.02%	2.4%	0.06%	Control cohort epidemiological study	2003
Portugal	0.02%	2.3%	0.3%	Control cohort epidemiological study	2003
Australia	0.02%	2.5%	0.12%	Control cohort epidemiological study	2003
USA	0.02%	2.7%	0.09%	Control cohort epidemiological study	2003
New Zealand	0.05%	4.3%	0.17%	Control cohort epidemiological study	2003
Canada	0.02%	2.7%	0.11%	Control cohort epidemiological study	2003

### Liver disease in homozygous A1AT-deficient adults

Liver disease due to A1AT deficiency generally presents at adult age. One study reviewed adult patients with liver disease and A1AT deficiency; the mean age of the patients when liver disease became symptomatic was 58 years for the ZZ phenotype, 66 years for the SZ phenotype and 73 years for the MZ phenotype. At the time of diagnosis the liver disease was advanced, 42% of these patients died within two years.<sup>24</sup> A review of autopsy data on 94 Pi ZZ homozygous A1AT deficient patients showed that cirrhotic patients survived longer compared with noncirrhotic patients. The noncirrhotic patients had more severe lung disease and died earlier.<sup>25</sup>

A cohort of patients who are registered in the Alpha-1 Foundation Registry (a USA foundation providing increased research and improved health for A1AT deficiency), and who had reported liver disease or jaundice (165 of the 2175 participants in the registry) completed a questionnaire. Of these patients 71% were Pi ZZ and 18% were Pi MZ, the remainder did not know what their phenotype was. Mean age at diagnosis of liver disease was 31 years (range 0 to 68 years), 30% had undergone liver transplant or were on the waiting list. Male gender and obesity were risk factors for advanced liver disease, while white race, Pi phenotype, infant jaundice, diabetes or hypercholesterolaemia were not.<sup>26</sup> Although this survey is the largest cohort of A1AT deficiency and liver disease in the literature, the self-selected cohort runs a risk of inclusion bias.

The natural history of the disease is not completely known. The risk of cirrhosis in adults is difficult to estimate because most available data are retrospective and derived from patients known to have A1AT-deficient lung disease or cirrhosis.<sup>27,28</sup>

### Heterozygous A1AT deficiency and liver disease

Although the role of homozygous A1AT in liver disease is established, the association between heterozygous A1AT deficiency and chronic liver disease is still subject to ongoing investigation. Several studies, however, have shown an association between heterozygous A1AT deficiency and chronic liver disease.

In 1981 a study showed the association between Pi MZ and liver disease. About 1055 liver biopsies were screened for A1AT depositions in hepatocytes. A total of 34 patients with these inclusions were phenotyped; the prevalence of phenotype Pi MZ in the whole biopsy group was 2.4%. In liver cirrhosis, 9% had a Pi MZ phenotype. A percentage of 21% Pi MZ was found in cryptogenic cirrhosis and in chronic active hepatitis, this was significantly increased compared with other causes of cirrhosis. The prognosis of the Pi MZ cirrhotic patients was poor, most patients died within one year.<sup>29</sup>

More recently patients with end-stage liver disease, in work-up for liver transplantation, were investigated. Pi MZ was found in 7.3 to 8.2%, compared with 2.8% in the control population. A heterozygous phenotype was more prevalent in patients with hepatitis C, alcoholic liver disease, cryptogenic cirrhosis and hepatocellular carcinoma.<sup>30,31</sup>

In one study consecutive liver biopsies and autopsies were screened for Pi Z deposits. In the biopsy group 3.4% of cases were Pi MZ phenotyped, whereas in the autopsy group this was 1.8%. In biopsies from older people heterozygous for A1AT, more fibrosis and more Pi Z deposits were found; the liver involvement seems to be age-dependent. When there was another liver disease as well, the patients presented with more inflammation, more fibrosis and more Pi Z deposits than the biopsies without concomitant liver disease.<sup>32</sup>

We described three patients with alcoholic liver disease and a rapidly deteriorating clinical course, resulting in the patients' death. All three patients were found to be heterozygous for A1AT.<sup>33</sup>

To summarise, these studies showed that various liver diseases influence the A1AT accumulation and that the A1AT accumulation influences the course of the liver disease. The risk of developing liver cirrhosis is increased in patients with heterozygous A1AT deficiency, also without coexisting liver disease. The exact impact and involvement of liver disease by heterozygous A1AT deficiency are unknown. Further research is needed to give these data.

### Heterozygous A1AT deficiency and coexisting hepatitis C virus infection

The role of A1AT deficiency in the severity and the course of liver disease in chronic hepatitis C virus (HCV) infection is not clear, despite the fact that several studies have analysed the association of HCV-induced liver disease and A1AT deficiency.

In Austria, 1865 patients referred for the evaluation of chronic liver disease were analysed, 9% had a deficient phenotype. From these patients with cirrhosis, 62% were HCV positive, 33% had evidence of HBV infection, 41% abuse of alcohol and 12% had features of autoimmune liver disease. Out of 53 cirrhotic A1AT-deficient patients, only five had no coexisting liver disease. These authors concluded that the risk for chronic liver disease is increased in patients with the Pi Z gene, because they may have increased susceptibility to viral infection or additional factors, necessary to induce chronic liver disease.<sup>34</sup>

The same authors investigated the prognosis of patients with A1AT deficiency. Some 54 patients with A1AT deficiency had evidence of chronic liver disease, 78% showed positive viral markers (hepatitis B or hepatitis C); this was compared with 106 patients with A1AT deficiency without chronic liver disease, without signs of additional viral infection. Life expectancy in A1AT-deficient patients was significantly lower in patients with chronic liver disease in comparison with patients without chronic liver disease.<sup>35</sup>

Patients with end-stage liver disease, in work-up for liver transplantation, were also investigated. In the HCV patients Pi MZ was found in 10 to 13%, compared with 2.8% in the control population. This suggests that an abnormal heterozygous phenotype is a co-factor in the development of chronic liver disease in HCV.<sup>30,31</sup>

In contrast, other studies showed no association between hepatitis C infection and A1AT deficiency.<sup>36-38</sup>

To conclude, the results of these studies are controversial. Some studies show a higher incidence of A1AT deficiency in HCV infection and an increased susceptibility to viral infections in A1AT deficiency and other studies do not. Different methods to determine the A1AT state were used. Further research on the influence of A1AT deficiency in the course of HCV infection and vice versa is necessary.

### **Heterozygous A1AT deficiency and hepatocellular carcinoma**

Established risk factors for hepatocellular carcinoma include chronic hepatitis B, HCV infection and alcoholic liver cirrhosis. Several studies have investigated the correlation between A1AT deficiency and hepatocellular carcinoma.<sup>39,40</sup>

In 1986 it was suggested for the first time that men with A1AT deficiency may be at risk for cirrhosis and hepatocellular carcinoma (HCC). Autopsy was performed in 16 adult patients with A1AT deficiency. In five out of these 16 patients, an HCC was found.<sup>41,42</sup>

In 317 HCC patients, Pi Z deposits were found in 6% compared with 1.8% in the control group. In heterozygous A1AT deficiency, HCC had also developed in noncirrhotic livers and was frequently characterised by cholangio-cellular differentiation. In patients with A1AT deficiency bile duct lesions were frequently found. This might reflect a predisposition for the liver tissue for developing tumours with cholangiolar differentiation in A1AT deficiency.<sup>43,44</sup>

In contrast to these studies, others did not show an association between A1AT deficiency and hepatocellular carcinoma, although carcinomas in noncirrhotic livers were Pi MZ associated.<sup>45-48</sup>

To summarise, several studies have been performed to investigate the relation between A1AT deficiency and hepatocellular carcinoma. The outcomes are not uniform. In our opinion studies with large cohorts of patients with hepatocellular carcinoma are the most reliable; these studies did find an association.

### **A1AT deficiency and associations with other diseases**

A1AT deficiency is not only associated with liver and lung disease. Associations with panniculitis, nephrotic syndrome, intracranial aneurysm, hereditary haemochromatosis and celiac disease have been described.<sup>49-62</sup>

## **THERAPY**

Most therapeutic strategies in the treatment of A1AT deficiency are directed towards the pulmonary disease. Infusion of purified pooled human plasma A1AT is known as augmentation therapy. The goal of this treatment is to raise and maintain serum A1AT concentrations above

the protective threshold. Data from different studies suggest that intravenous augmentation therapy has a positive biochemical and clinical effect. The therapy is expensive (US \$ 28,075 to 65,973 per year).<sup>7</sup> Different concepts have been studied to prevent the polymerisation and accumulation of the A1AT protein. New peptides that block the polymerisation of the Z protein have been developed.<sup>63,64</sup> Gene therapy by injecting adeno-associated virus carrying the human A1AT gene is another promising concept.<sup>65</sup> Liver transplantation is used in end-stage liver disease and results in acquisition of the donor phenotype, a rise in serum levels of A1AT and prevention of associated diseases.<sup>66</sup>

## **CONCLUSION**

Alpha-1 antitrypsin deficiency is an autosomal recessive disorder that can lead to chronic pulmonary disease and liver disease. The liver disease is caused by accumulation of an abnormal, polymerised protein. Deficient phenotypes are present worldwide.

Homozygous A1AT deficiency in children can cause neonatal hepatitis. In adults homozygous patients are at risk for developing end-stage liver disease, cirrhosis and hepatocellular carcinoma. Heterozygous A1AT deficiency is probably an important co-factor in the aetiology of chronic liver disease, as several studies have shown associations with HCV, end-stage liver disease, cirrhosis and HCC.

Low-threshold screening for A1AT deficiency should therefore be carried out. A1AT serum levels may be used, but phenotyping is crucial, as serum levels may not reflect true deficiencies (as inflammation serum levels can be falsely normal). Especially in cryptogenic chronic liver disease and liver disease that deteriorates faster than may be expected, A1AT deficiency may be of clinical significance as a (co)-factor. Clinical research is needed in A1AT-related liver disease to investigate the association between heterozygous A1AT deficiency and the presentation and course of liver diseases.<sup>67</sup>

We believe that the current data are insufficient to decide on the pros and cons of screening on hepatocellular carcinoma in A1AT deficiency. Therapeutic options in A1AT deficiency include augmentation therapy in pulmonary disease; in end-stage liver disease liver transplantation is an option. For the future, gene therapy or strategies to inhibit polymerisation are promising.

## **ACKNOWLEDGEMENT**

Joost P.H. Drenth is a recipient of a VIDJ grant from the Netherlands Organisation for Health Research and Development (ZON-MW).

## REFERENCES

1. Teckmann JH, Qu D, Perlmutter DH. Molecular pathogenesis of liver disease in  $\alpha_1$ -antitrypsin deficiency. *Hepatology* 1996;24:1504-16.
2. Lomas DA. Loop-sheet polymerization: the mechanism of alpha<sub>1</sub>-antitrypsin deficiency. *Respir Med* 2000;94:S3-S6.
3. Dafforn TR, Mahadeva R, Elliott PR, Sivasothy P, Lomas DA. A kinetic mechanism for the polymerization of  $\alpha_1$ -antitrypsin. *J Biol Chem* 1999;274:9548-55.
4. Lomas DA, Evans DL, Finch JT, Carrell RW. The mechanism of Z  $\alpha_1$ -antitrypsin accumulation in the liver. *Nature* 1992;357:605-7.
5. DeMeo DL, Silverman EK.  $\alpha_1$ -Antitrypsin deficiency 2: genetic aspects of  $\alpha_1$ -antitrypsin deficiency: phenotypes and genetic modifiers of emphysema risk. *Thorax* 2004;59:259-64.
6. Carrell RW, Lomas DA. Alpha<sub>1</sub>-antitrypsin deficiency—a model for conformational diseases. *N Engl J Med* 2002;346:45-53.
7. Stoller JK, Aboussouan LS.  $\alpha_1$ -antitrypsin deficiency. *Lancet* 2005;365:2225-36.
8. Sun Z, Yang P. Role of imbalance between neutrophil elastase and  $\alpha_1$ -antitrypsin in cancer development and progression. *Lancet Oncol* 2004;5:182-90.
9. Perlmutter DH. Pathogenesis of chronic liver injury and hepatocellular carcinoma in alpha<sub>1</sub>-antitrypsin deficiency. *Resp Res* 2006;60:233-8.
10. Luisetti M, Seersholm N.  $\alpha_1$ -Antitrypsin deficiency 1: epidemiology of  $\alpha_1$ -antitrypsin deficiency. *Thorax* 2004;59:164-9.
11. Sveger T. Liver disease in  $\alpha_1$ -antitrypsin deficiency detected by screening of 200,000 infants. *N Engl J Med* 1976;294:1316-21.
12. Dijkman JH, Penders TJ, Kramps JA, Sonderkamp HJA, Broek WGM van den, Haar BGA ter. Epidemiology of Alpha<sub>1</sub>-antitrypsin deficiency in the Netherlands. *Hum Genet* 1980;53:409-13.
13. Serres FJ de, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. *Clin Genet* 2003;63:490-509.
14. Serres FJ de, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand, and the United States of America. *Clin Genet* 2003;64:382-97.
15. De Serres FJ. Worldwide racial and ethnic distribution of  $\alpha_1$ -antitrypsin deficiency: summary of an analysis of published genetic epidemiologic surveys. *Chest* 2002;122:1818-29.
16. Mowat AP. Alpha<sub>1</sub>-antitrypsin deficiency (PiZZ): features of liver involvement in childhood. *Acta Paediatr* 1994;393:13-7.
17. O'Brien ML, Buist NRM, Murphey WH. Neonatal screening for alpha<sub>1</sub>-antitrypsin deficiency. *J Pediatr* 1978;92:1006-10.
18. Cottrill K, Cook PJL. Neonatal hepatitis syndrome and alpha-1-antitrypsin deficiency: an epidemiological study in south-east England. *Postgrad Med J* 1974;50:376-80.
19. Grishan FK, Greene HL. Liver disease in children with PiZZ  $\alpha_1$ -antitrypsin deficiency. *Hepatology* 1988;8:307-10.
20. Sveger T, Eriksson S. The liver in adolescents with  $\alpha_1$ -antitrypsin deficiency. *Hepatology* 1995;22:514-7.
21. Pittulainen E, Carlson J, Ohlsson K, Sveger T.  $\alpha_1$ -Antitrypsin deficiency in 26-year-old subjects: lung, liver, and protease/protease inhibitor studies. *Chest* 2005;128:2076-81.
22. Pittschieler K. Liver disease and heterozygous alpha-1-antitrypsin deficiency. *Acta Paediatr Scand* 1991;80:323-7.
23. Pittschieler K. Liver involvement in alpha<sub>1</sub>-antitrypsin deficient phenotypes PiSZ and PiMZ. *Acta Paediatr* 2002;91:239-40.
24. Rakela J, Goldschmied M, Ludwig J. Late manifestation of chronic liver disease in adults with alpha-1 antitrypsin deficiency. *Dig Dis Sci* 1987;32:1358-62.
25. Eriksson S.  $\alpha_1$ -Antitrypsin deficiency and liver cirrhosis in adults. *Acta Med Scand* 1987;221:461-7.
26. Bowlus CL, Willner I, Zern MA, et al. Factors associated with advanced liver disease in adults with alpha<sub>1</sub>-antitrypsin deficiency. *Clin Gastroenterol Hepatol* 2005;3:390-6.
27. Von Schönfeld J, Breuer N, Zotz R, et al. Liver function in patients with pulmonary emphysema due to severe alpha-1-antitrypsin deficiency (Pi ZZ). *Digestion* 1996;57:165-9.
28. Larsson C, Eriksson S. Liver function in asymptomatic adult individuals with severe  $\alpha_1$ -antitrypsin deficiency (Pi Z). *Scand J Gastroent* 1977;12:543-6.
29. Hodges R, Millward-Sadler GH, Path MRC, Barbatis C, Wright R, Phil D. Heterozygous MZ alpha<sub>1</sub>-antitrypsin deficiency in adults with chronic active hepatitis and cryptogenic cirrhosis. *N Engl J Med* 1981;304:557-60.
30. Eigenbrodt ML, McCashland TM, Dy RM, Clark J, Galati J. Heterozygous  $\alpha_1$ -antitrypsin phenotypes in patients with end stage liver disease. *Am J Gastroenterol* 1997;92:602-7.
31. Graziadei IW, Joseph JJ, Wiesner RH, Therneau TM, Batts KP, Porayko MK. Increased risk of chronic liver failure in adults with heterozygous  $\alpha_1$ -antitrypsin deficiency. *Hepatology* 1998;28:1058-63.
32. Fischer HP, Ortiz-Pallardó ME, Ko Y, Esch C, Zhou H. Chronic liver disease in heterozygous  $\alpha_1$ -antitrypsin deficiency PiZ. *J Hepatol* 2000;33:883-92.
33. Kok KF, Wahab PJ, Vries RA. Heterozygosity for alpha<sub>1</sub>-antitrypsin deficiency as a cofactor in the development of chronic liver disease. *Ned Tijdschr Geneesk* 2005;149:2057-61.
34. Propst T, Propst A, Dietze O, Judmaier G, Braunsteiner H, Vogel W. High prevalence of viral infection in adults with homozygous and heterozygous alpha<sub>1</sub>-antitrypsin deficiency and chronic liver disease. *Ann Int Med* 1992;117:641-5.
35. Propst A, Propst T, Öfner D, Feichtinger H, Judmaier G, Vogel W. Prognosis and life expectancy in alpha-1-antitrypsin deficiency and chronic liver disease. *Scan J Gastroenterol* 1995;30:1108-12.
36. Serfaty L, Chazouillères O, Poujol-Robert A, et al. Risk factors for cirrhosis in patients with chronic hepatitis C virus infection: results of a case-control study. *Hepatology* 1997;26:776-9.
37. Elzouki AN, Verbaan H, Lindgren S, Widell A, Carlson J, Eriksson S. Serine protease inhibitors in patients with chronic viral hepatitis. *J Hepatol* 1997;27:42-8.
38. Scott BB, Egner W, on behalf of the Trent Hepatitis C Study Group. Does  $\alpha_1$ -antitrypsin phenotype PiMZ increase the risk of fibrosis in liver disease due to hepatitis C virus infection? *Eur J Gastroenterol Hepatol* 2006;18:521-3.
39. Palmer PE, Wolfe HJ.  $\alpha_1$ -antitrypsin deposition in primary hepatic carcinoma. *Arch Pathol Lab Med* 1976;100:232-6.
40. Reintoft I, Hägerstrand IE. Does the Z gene variant of alpha-1-antitrypsin predispose to hepatic cancer? *Hum Pathol* 1979;10:419-24.
41. Eriksson S, Carlson J, Velez R. Risk of cirrhosis and primary liver cancer in alpha<sub>1</sub>-antitrypsin deficiency. *N Engl J Med* 1986;314:736-9.
42. Elzouki AN, Eriksson S. Risk of hepatobiliary disease in adults with severe  $\alpha_1$ -antitrypsin deficiency (PiZZ): is chronic viral hepatitis B or C an additional risk factor for cirrhosis and hepatocellular carcinoma? *Eur J Gastroenterol Hepatol* 1996;8:989-94.
43. Zhou H, Ortiz-Pallardó ME, Ko Y, Fischer HP. Is heterozygous alpha-1-antitrypsin deficiency type PiZ a risk factor for primary liver cell carcinoma? *Cancer* 2000;88:2668-76.
44. Zhou H, Fischer HP. Liver carcinoma in PiZ alpha-1-antitrypsin deficiency. *Am J Pathol* 1998;22:742-8.
45. Propst T, Propst A, Dietze O, Judmaier G, Braunsteiner H, Vogel W. Prevalence of hepatocellular carcinoma in alpha-1-antitrypsin deficiency. *J Hepatol* 1994;21:1006-11.
46. Rabinovitz M, Gavalier JS, Kelly RH, Prieto M, van Thiel D. Lack of increase in heterozygous  $\alpha_1$ -antitrypsin deficiency phenotypes among patients with hepatocellular and bile duct carcinoma. *Hepatology* 1992;15:407-10.
47. Govindarajan S, Ashcaval M, Peters RL.  $\alpha_1$ -Antitrypsin deficiency phenotypes in hepatocellular carcinoma. *Hepatology* 1981;1:628-31.

48. Sparos L, Tountas Y, Chapuis-Cellier C, Theodoropoulos G, Trichopoulos D. Alpha<sub>1</sub>-antitrypsin levels and phenotypes and hepatitis B serology in liver cancer. *Br J Cancer* 1984;49:567-70.
49. Edmonds BK, Hodge JA, Rietschel RL. Alpha<sub>1</sub>-antitrypsin deficiency-associated panniculitis: case report and review of the literature. *Pediatr Dermatol* 1991;8:296-9.
50. O'Riordan K, Blei A, Rao MS, Abecassis M.  $\alpha_1$ -Antitrypsin deficiency-associated panniculitis: resolution with intravenous  $\alpha_1$ -antitrypsin administration and liver transplantation. *Transplantation* 1997;63:480-2.
51. Phelps RG, Shoji T. Update on panniculitis. *Mt Sinai J Med* 2001;68:262-7.
52. McBean J, Sable A, Maude J, Robinson-Bostom L.  $\alpha_1$ -Antitrypsin deficiency panniculitis. *Cutis* 2003;71:205-9.
53. Chowdhury MMU, Williams EJ, Morris JS, et al. Severe panniculitis caused by homozygous ZZ  $\alpha_1$ -antitrypsin deficiency treated successfully with human purified enzyme (Prolastin®). *Br J Dermatol* 2002;147:1258-61.
54. Elzouki AN, Lindgren S, Nilsson S, Veress B, Eriksson S. Severe  $\alpha_1$ -antitrypsin deficiency (PiZ homozygosity) with membranoproliferative glomerulonephritis and nephrotic syndrome, reversible after orthotopic liver transplantation. *J Hepatol* 1997;26:1403-7.
55. Elzouki AN, Eriksson S. Severe  $\alpha_1$ -antitrypsin deficiency and intracranial aneurysms. *Lancet* 1994;343:1037.
56. Elzouki AN, Hultcrantz R, Stål P, Befrits R, Eriksson S. Increased PiZ gene frequency for  $\alpha_1$ -antitrypsin in patients with genetic haemochromatosis. *Gut* 1995;36:922-6.
57. Rabinovitz M, Gavalier JS, Kelly RH, Thiel DH van. Association between heterozygous  $\alpha_1$ -antitrypsin deficiency and genetic hemochromatosis. *Hepatology* 1992;16:145-8.
58. Fargion S, Bissoli F, Francanzant AL, Suigo E, Sergi C, Taioli E. No association between genetic hemochromatosis and  $\alpha_1$ -antitrypsin deficiency. *Hepatology* 1996;24:1161-4.
59. Eriksson S, Lindmark B, Olsson S. Lack of association between hemochromatosis and  $\alpha$ -antitrypsin deficiency. *Acta Med Scand* 1986;219:291-4.
60. Walker-Smith J, Andrews J. Alpha-1 antitrypsin, autism, and coeliac disease. *Lancet* 1972;2:883-4.
61. Pons Romero F, Casafont F, Rodriguez de Lope C, San Miguel G, Artinano E, Cagigas J. Could alpha<sub>1</sub>-antitrypsin deficiency have any role in the development of celiac sprue after gastric operations? *J Clin Gastroenterol* 1986;8:559-61.
62. Klasen EC, Polanco I, Biemond I, Vazquez C, Pena AS.  $\alpha_1$ -Antitrypsin deficiency and celiac disease in Spain. *Gut* 1980;21:948-50.
63. Burrows JA, Willis LK, Perlmutter DH. Chemical chaperones mediate increased secretion of mutant  $\alpha_1$ -antitrypsin ( $\alpha_1$ -AT) Z: a potential pharmacological strategy for prevention of liver injury and emphysema in  $\alpha_1$ -AT deficiency. *Proc Natl Acad Sci USA* 2000;97:1796-801.
64. Lomas DA, Mahadeva R.  $\alpha_1$ -antitrypsin polymerization and the serpinopathies: pathobiology and prospects for therapy. *J Clin Invest* 2002;110:1585-90.
65. Stecenko AA, Brigham KL. Gene therapy progress and prospects: alpha-1 antitrypsin. *Gene Therapy* 2003;10:95-9.
66. Vennarecci G, Gunson BK, Ismail T, et al. Transplantation for end stage liver disease related to alpha 1 antitrypsin. *Transplantation* 1996;61:1488-95.
67. American Thoracic Society/European Respiratory Society Statement: Standards for the diagnosis and management of individuals with alpha-1 antitrypsin deficiency. *Am J Respir Crit Care Med* 2003;168:818-900.