

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/57032>

Please be advised that this information was generated on 2020-10-23 and may be subject to change.

NON - ARCHIMEDEAN DIFFERENTIATION

held on Tuesday, June 5, 1979 at the "VI Jornadas de Matemáticas Hispano-Lusas" organized by the University of SANTANDER,

by

W.H. Schikhof

§ 1. Introduction.

The subject is part of the so-called non-archimedean (or ultrametric) analysis. Roughly speaking, one may say that this is the analysis that one obtains when replacing in the "classical" analysis  $\mathbb{R}$  or  $\mathbb{C}$  by a non-archimedean valued field  $K$ .

A non-archimedean valued field is a (commutative) field  $K$ , together with a map  $|\cdot| : K \rightarrow \mathbb{R}$  (the valuation) satisfying

$$|a| \geq 0 \quad , \quad |a| = 0 \text{ iff } a = 0$$

$$|ab| = |a| |b|$$

$$|a+b| \leq \max(|a|, |b|) \quad (\text{the } \underline{\text{strong triangle inequality}})$$

for all  $a, b \in K$ .

We have the following remarks.

- (1) Apart from  $\mathbb{R}$  or  $\mathbb{C}$ , every complete valued field is non-archimedean.
- (2) If  $K$  is a non-archimedean valued field and if  $L \supset K$  is an overfield of  $K$  then the valuation on  $K$  can be extended to a non-archimedean valuation on  $L$ .
- (3) If  $K$  is a (non-archimedean) valued field then its completion  $\tilde{K}$  (with respect to the metric  $(x, y) \mapsto |x-y|$ ) can, in a natural

way, be given the structure of a non-archimedean valued field.

In the sequel we exclude the so-called trivial valuation given by

$$|x| := \begin{cases} 0 & \text{if } x = 0 \\ 1 & \text{if } x \neq 0. \end{cases}$$

The non-archimedean analysis has several branches, similar to the classical analysis. Thus we have non-archimedean functional analysis, harmonic analysis, theory of analytic functions in one or several variables, etc.

In this talk we consider a more elementary subject, namely infinitesimal calculus in  $K$ . More specifically, we want to see what remains of the so-called Fundamental Theorem of Calculus (in  $\mathbb{R}$ ) that states that the operations of differentiation and integration are in some sense each others inverses.

§ 2. Differentiation in  $K$ . Let  $X \subset K$  be a subset without isolated points. A function  $f : X \rightarrow K$  is called differentiable if for all  $a \in X$

$$f'(a) := \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

exists. The proof of the well known rules (sum-, product-, chain-rule) can formally be taken over from the classical theory. Thus, a rational function is differentiable if it has no poles on  $X$ . An analytic function  $x \mapsto \sum_n a_n x^n$  is differentiable on  $\{x : |x| < (\limsup_n |a_n|)^{-1}\}$ .

Deviations from the classical theory appear when we look at the functions whose derivative vanishes everywhere. For example,

let  $\varepsilon > 0$ ,  $a \in K$ . Then  $B(a, \varepsilon) := \{x \in K : |x-a| < \varepsilon\}$  is an open- and-closed subset of  $K$ , hence  $\xi_{B(a, \varepsilon)}$ , defined by

$$\xi_{B(a, \varepsilon)}(x) := \begin{cases} 1 & \text{if } x \in B(a, \varepsilon) \\ 0 & \text{elsewhere} \end{cases}$$

is differentiable and  $\xi'_{B(a, \varepsilon)} = 0$ .

Locally constant functions all have derivative zero. On the other hand they form a uniformly dense subset of  $C(X)$ , the space of all continuous functions:  $X \rightarrow K$ .

Even worse: let  $\mathbb{Q}_p$  the field of the  $p$ -adic numbers and let

$\mathbb{Z}_p := \{x \in \mathbb{Q}_p : |x| \leq 1\}$ . Then the function  $f : \mathbb{Z}_p \rightarrow \mathbb{Q}_p$  defined by

$$f(\sum_n a_n p^n) = \sum_n a_n p^{2n} \quad (\sum_n a_n p^n \in \mathbb{Z}_p)$$

satisfies  $|f(x) - f(y)| = |x - y|^2$  for all  $x, y \in \mathbb{Z}_p$ . So  $f' = 0$  but  $f$  is injective, hence not locally constant.

The above example shows also that a Mean Value Theorem is necessarily absent in our theory.

Notice that the difficulties encountered above also appear when we study differentiability of functions  $f : \mathbb{D} \rightarrow \mathbb{R}$ , where  $\mathbb{D} \subset [0, 1]$  is the Cantor set. So it is the domain of  $f$  that is responsible for the troubles rather than its range.

### § 3. Continuously differentiable functions.

If we follow naively the path of the classical analysis and define

$$C^1(X) := \{f : X \rightarrow K, f \text{ is differentiable, } f' \text{ is continuous}\}$$

then we run up against difficulties.

First of all, one can prove that  $C^1(\mathbb{Z}_p)$  (with the norm

$f \mapsto \max(\|f\|_\infty, \|f'\|_\infty)$  is not a Banach space. In fact one shows that for every pair of continuous functions  $f, g : \mathbb{Z}_p \rightarrow \mathbb{Q}_p$  there exists a sequence  $f_1, f_2, \dots$  in  $C^1(\mathbb{Z}_p)$  for which both  $f_n \rightarrow f$  and  $f'_n \rightarrow g$  uniformly.

What is worse, we have no local invertibility theorem for such  $C^1$ -functions.

In fact, let  $f : \mathbb{Z}_p \rightarrow \mathbb{Q}_p$  be defined by

$$f(x) = \begin{cases} x - p^{2n} & \text{if } |x - p^n| < p^{-2n} \quad (n \in \mathbb{N}) \\ x & \text{elsewhere} \end{cases}$$

Then  $f'(x) = 1$  for all  $x \in \mathbb{Z}_p$ . But  $f(p^n) = f(p^n - p^{2n})$  for all  $n \in \mathbb{N}$ , so  $f$  is not even locally injective at 0.

Therefore we are led to define:

Let  $f : X \rightarrow K$ . Put

$$\phi f(x, y) := \frac{f(x) - f(y)}{x - y} \quad (x, y \in X, x \neq y).$$

We say that  $f \in C^1(X)$  if  $\phi f$  can continuously be extended to a function  $\bar{\phi}f : X \times X \rightarrow K$ .

Then  $BC^1(X) := \{f \in C^1(X) : f \text{ and } \phi f \text{ are bounded}\}$  is a Banach space under  $f \mapsto \|f\|_1 := \max(\|f\|_\infty, \|\phi(f)\|_\infty)$ .

Further, if  $f \in C^1(X)$ ,  $f'(a) \neq 0$  for some  $a \in X$ , then  $f$  has a  $C^1$ -inverse, locally at  $a$ .

Theorem. Differentiation is a continuous surjection  $BC^1(X) \xrightarrow{D} BC(X)$ .

(here  $BC(X)$  is the space of all bounded continuous functions with the supremum norm)

#### § 4. "Integration".

Next, we want to define an "indefinite integral"  $P : BC(X) \rightarrow BC^1(X)$

(an analogue of  $(Pf)(x) := \int_0^x f(t)dt$  for real functions) such that  $DP$  is the identity on  $BC(X)$ .

A natural try is first to find an analogue of the Lebesgue measure in  $K$ . But this turns out to be a dead end road. For example if  $K = \mathbb{Q}_p$  there does not exist a nonzero translation invariant bounded additive  $\mathbb{Q}_p$ -valued function  $m$  defined on the compact open subsets of  $\mathbb{Z}_p$ . (By translation invariance  $|m(p^n \mathbb{Z}_p)| = p^n |m(\mathbb{Z}_p)| \rightarrow \infty$  if  $m(\mathbb{Z}_p) \neq 0$ ). For similar reasons it goes wrong for every local field  $K$ .

Following the ideas of Dieudonné, Treiber, we define for  $f \in BC(X)$

$$(Pf)(x) := \sum_{n=1}^{\infty} f(x_n) (x_{n+1} - x_n) \quad (x \in X)$$

Here the  $x_n$  are defined as follows. For each  $n \in \mathbb{N}$  the equivalence relation  $\sim_n$  defined by  $x \sim_n y$  if  $|x-y| < \frac{1}{n}$  yields a partition of  $X$  into balls. Choose a center in each ball and let  $R_n$  be the set of these centers.

For each  $x \in X$  and  $n \in \mathbb{N}$ ,  $x_n$  is defined by  $x_n \in R_n$ ,  $|x_n - x| < \frac{1}{n}$ .

Theorem. (A NON-ARCHIMEDEAN FORM OF THE FUNDAMENTAL THEOREM).

$P$  is a linear isometry of  $BC(X)$  into  $BC^1(X)$ .  $DP$  is the identity on  $BC(X)$ , whereas  $PD$  is a projection of  $BC^1(X)$  onto a complement of  $\{f \in BC^1(X) : f' = 0\}$ .

## § 5. Generalizations of the Fundamental Theorem.

We may ask whether there exists some form of the Fundamental Theorem for functions belonging to spaces, larger than  $BC(X)$ ,  $BC^1(X)$

respectively. (For example, compare the classical theorem on  $L^1$ -functions versus absolutely continuous functions).

We have the following striking fact that has no counterpart in classical analysis. We say that  $g : X \rightarrow K$  is of the first class of Baire if there exists a sequence  $g_1, g_2, \dots$  of continuous functions  $X \rightarrow K$  such that  $\lim g_n = g$  pointwise.

**THEOREM.** (a) Let  $f : X \rightarrow K$  be differentiable. Then  $f'$  is of the first class of Baire.

(b) Let  $g : X \rightarrow K$  be of the first class of Baire. Then  $g$  has an antiderivative.

Let  $B\mathbb{B}^1(X)$  be the Banach space consisting of all bounded functions  $X \rightarrow K$  of the first class of Baire with respect to the supremum norm.

Let  $BD(X)$  be the Banach space of all differentiable  $f : X \rightarrow K$  for which both  $f$  and  $\Phi f$  are bounded, with respect to the norm

$f \rightarrow \|f\|_\infty \vee \|\Phi f\|_\infty$ . Then we have

**THEOREM.** Differentiation is a quotient map  $BD(X) \xrightarrow{D} B\mathbb{B}^1(X)$ .

If  $K$  has discrete valuation then there exists a continuous linear  $P : B\mathbb{B}^1(X) \rightarrow BD(X)$  for which  $DP$  is the identity on  $B\mathbb{B}^1(X)$ .

### Notes.

1. The construction of the above  $P$  is awful and, contrary to § 4,  $P$  does not resemble an indefinite integral in any way.
2. If the valuation of  $K$  is dense the existence of such a  $P$  is still an open question.

§ 6. Restriction of the Fundamental Theorem.

In classical analysis, we have that if  $f \in C^n$  then  $x \mapsto \int_0^x f(t) dt$  is in  $C^{n+1}$ . In our situation we define for  $f : X \rightarrow K$ :  $f \in C^2(X)$  if the function  $\Phi_2 f$ , defined by

$$\Phi_2 f(x, y, z) = \frac{\Phi_1 f(x, z) - \Phi_1 f(y, z)}{x - y} \quad (x, y, z \in X, x \neq y, y \neq z, x \neq z)$$

can continuously be extended to  $\overline{\Phi}_2 f : X^3 \rightarrow K$ . Similarly, we define  $C^3(X), C^4(X), \dots$ . Let  $C^\infty(X) := \bigcap_{n=1} C^n(X)$ .

The map  $P$ , defined in § 4, does not always map  $C^1$ -functions into  $C^2$ -functions. But we have (notations as in § 4)

THEOREM. Let the characteristic of  $K$  be unequal to 2. Then the map  $P_2$  defined via

$$(P_2 f)(x) := \sum f(x_n)(x_{n+1} - x_n) + \frac{1}{2} \sum f'(x_n)(x_{n+1} - x_n)^2 \quad (x \in X)$$

maps  $C^1(X)$  into  $C^2(X)$  and  $(P_2 f)' = f$  for all  $f \in C^1(X)$ .

Similarly, one can define antiderivation maps  $P_n : C^{n-1}(X) \rightarrow C^n(X)$  (in case the characteristic of  $K$  is unequal to  $2, 3, \dots, n$ ).

OPEN QUESTION. Let  $K$  have characteristic 0. Does every  $f \in C^\infty(X)$  have a  $C^\infty$ -antiderivative?

W. Schikhof

Reference

Schikhof, W. : Non-archimedean calculus. (Lecture Notes). Report 7812, Mathematisch Instituut, Nijmegen, The Netherlands.