

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/57721>

Please be advised that this information was generated on 2020-11-01 and may be subject to change.

Regulation of the Epithelial Ca^{2+} Channel TRPV5 by the NHE Regulating Factor NHERF2 and the Serum and Glucocorticoid Inducible Kinase Isoforms SGK1 and SGK3 Expressed in *Xenopus* oocytes

Hamdy M. Embark¹, Iwan Setiawan¹, Susanne Poppendieck¹, Stan F.J. van de Graaf⁴, Christoph Boehmer¹, Monica Palmada¹, Thomas Wieder¹, Ruediger Gerstberger⁵, Philip Cohen², C. Chris Yun³, René J.M. Bindels⁴ and Florian Lang¹

¹Dept. of Physiology I, University of Tübingen, ²The MRC Protein Phosphorylation Unit, Division of Cell Signalling, MSI/WTB Complex, School of Life Sciences, University of Dundee, ³Dept. of Medicine, Div. of Digestive Diseases, Emory University, Atlanta, ⁴Department of Cell Physiology, University of Nijmegen, ⁵Dept. of Veterinary Physiology, University of Giessen

Key Words

1,25(OH)₂D₃ • TRPV5 • Calcium transport • Mineralisation • Kidney • Intestine • Bone • IGF1

Abstract

The epithelial Ca^{2+} channel TRPV5 (ECaC1) plays a key role in renal and intestinal Ca^{2+} (re)absorption and is thus regulated by 1,25(OH)₂D₃. The present study aims to explore whether TRPV5 is regulated by the serum and glucocorticoid inducible kinase SGK1, a kinase transcriptionally upregulated by 1,25(OH)₂D₃. To this end cRNA encoding TRPV5 has been injected into *Xenopus* oocytes with or without additional injection of SGK1, its isoforms SGK2 and SGK3, constitutively active S^{422D}SGK1, inactive K^{127N}SGK1, constitutively active T^{308D,S473D}PKB and/or the Na⁺/H⁺ exchanger regulating factor NHERF2. In *Xenopus laevis* oocytes expression of TRPV5 increases uptake of tracer Ca^{2+} and induces a Ca^{2+} current (I_{Ca}). In the

presence of Cl⁻, TRPV5 mediated Ca^{2+} entry leads to secondary activation of Ca^{2+} -sensitive Cl⁻ channels ($I_{\text{Cl}(\text{Ca})}$). Coexpression of TRPV5 with both S^{422D}SGK1 and NHERF2 stimulates tracer Ca^{2+} entry, I_{Ca} and $I_{\text{Cl}(\text{Ca})}$. The effect of S^{422D}SGK1 on TRPV5 and NHERF2 expressing oocytes is mimicked by SGK1 and SGK3, but not by SGK2, constitutively active T^{308D,S473D}PKB or inactive K^{127N}SGK1. The observations suggest that SGK1, SGK3 and NHERF2 regulate TRPV5 and are thus likely to participate in the regulation of calcium homeostasis.

Copyright © 2004 S. Karger AG, Basel

Introduction

Maintenance of Ca^{2+} homeostasis requires the fine tuning of intestinal Ca^{2+} absorption and renal Ca^{2+} excretion [1, 2]. The epithelial Ca^{2+} channel TRPV5 [3, 4] plays a key role in both, intestinal absorption and renal

reabsorption of Ca^{2+} . The channel accomplishes Ca^{2+} uptake across the apical membrane of epithelial cells. Subsequently, Ca^{2+} is extruded via the basolateral Ca^{2+} ATPase and $\text{Na}^{+}/\text{Ca}^{2+}$ exchanger [5]. As entry via TRPV5 is the rate limiting step in the transcellular transport of Ca^{2+} , TRPV5 activity is decisive for transepithelial transport regulation [2]. The C-terminal tail of TRPV5 contains a PDZ binding motif which may bind to the PDZ domains of NHE regulating factors NHERF1 or NHERF2 [6]. NHERF1 and NHERF2 modulate the targeting and trafficking of several proteins including TRPV4 into the plasma membrane [7-9]. TRPV5 [10] and NHERF2 [11] colocalize in principal cells. Thus, the possibility was considered that NHERF2 participates in the regulation of TRPV5 activity.

Most recent experiments disclosed a role of serum and glucocorticoid dependent kinase SGK1 in the interaction of NHE3 with NHERF2 [12, 13]. SGK1 has originally been cloned as a glucocorticoid sensitive gene from rat mammary tumor cells [14, 15]. The human isoform has been identified as a cell volume regulated gene [16]. Subsequent studies revealed the genomic regulation of SGK1 by aldosterone [17-21], $1,25(\text{OH})_2\text{D}_3$ [22], TGF β [23] and a variety of further cytokines [24].

SGK1 shares 80% homology with its isoforms SGK2 and SGK3 [25] and 60% homology with the protein kinase B [26]. All three isoforms SGK1-3 and protein kinase B are activated via a signaling cascade involving phosphatidylinositol 3 kinase and phosphoinositide dependent kinase PDK1 [25-28]. The activation of SGK1 requires phosphorylation at Ser422, the activation of PKB phosphorylation at Thr308 and Ser473. Replacement of those amino acids by aspartate leads to the respective constitutively active kinases, $\text{S}^{422\text{D}}$ SGK1 [27] and $\text{T}^{308\text{D},\text{S}^{473\text{D}}}$ PKB [26]. Destruction of the catalytic subunit by replacement of the lysine at position 127 with asparagine leads to the inactive mutant $\text{K}^{127\text{N}}$ SGK1 [27].

SGK1 [16], SGK2 [25], SGK3 [25] and protein kinase B [29] are expressed in a wide variety of human epithelial tissues including intestine, kidney and placenta. Thus, they may participate in the regulation of transport in those tissues. Similar to TRPV5 and NHERF2, SGK1 is expressed in principal cells of the distal nephron [30] and may well participate in the regulation of TRPV5.

The present study has been performed to test for a role of the SGK isoforms, protein kinase B and NHERF2 in the regulation of TRPV5 activity. To this end, cRNA encoding TRPV5 has been injected with or without NHERF2 and/or wild-type SGK1, SGK2, SGK3,

constitutively active $\text{S}^{422\text{D}}$ SGK1, inactive $\text{K}^{127\text{N}}$ SGK1, or constitutively active $\text{T}^{308\text{D},\text{S}^{473\text{D}}}$ PKB into *Xenopus* oocytes.

Materials and Methods

Expression in Xenopus laevis oocytes

cRNA encoding rabbit TRPV5 [3] which is highly homologous to human TRPV5, human NHERF2 [13], human SGK1 [16], human SGK2 [25], human SGK3 [25], constitutively active human $\text{S}^{422\text{D}}$ SGK1 [25], inactive human $\text{K}^{127\text{N}}$ SGK1 [25], and constitutively active human $\text{T}^{308\text{D},\text{S}^{473\text{D}}}$ PKB [26] have been synthesized as described [31]. Oocytes were injected with 2.5 ng TRPV5, 7.5 ng kinase and/or 5 ng NHERF2 or NHERF2 Δ P1 or NHERF2 Δ P2 cRNA or H_2O . All experiments were performed at room temperature 3 days after injection of the respective cRNAs.

Tracer Ca^{2+} uptake

Calcium uptake was measured similar to what has been described earlier [32-34] with slight modifications. In detail, uptake of $^{45}\text{Ca}^{2+}$ (ICN Biomedicals GmbH, Eschwege, Germany), delivered as CaCl_2 in aqueous solution (specific activity: 0.185-1.11 TBq/g Ca) was determined 3 days after injection of oocytes with the respective cRNAs by incubating 10 - 15 oocytes at 20 °C in uptake solution containing 96 mM NaCl, 2 mM KCl, 1 mM MgCl_2 , 0.1 mM CaCl_2 , 1 mM BaCl_2 , 10 μM methoxyverapamil, 5 mM HEPES-Tris, pH 7.4. Radioactive $^{45}\text{Ca}^{2+}$ was added to yield a final concentration of 7 $\mu\text{Ci/ml}$. After 30 min (within the linear range of $^{45}\text{Ca}^{2+}$ uptake) the oocytes were washed three times in stop buffer containing 96 mM NaCl, 1 mM MgCl_2 , 0.5 mM CaCl_2 , 1.5 LaCl_3 , 5 mM HEPES-Tris, pH 7.4. Single oocytes were then placed separately in scintillation vials and solubilized in 200 μl 10 % (w/v) sodium dodecyl sulfate. After addition of 3 ml scintillation fluid (Ultima Gold, Packard, Groningen, The Netherlands), radioactivity in the samples was measured in a liquid scintillation counter (Wallac, Freiburg, Germany).

Voltage-clamp analysis

In two-electrode voltage-clamp experiments currents were recorded during a 4 s linear voltage ramp from -150 mV to +50 mV. The intermediate holding potential between the voltage ramps was -50 mV. Data were filtered at 1000 Hz and recorded with MacLab digital to analog converter and software for data acquisition and analysis (ADInstruments, Castle Hill, Australia). The bath solution contained 96 mM NaCl, 2 mM KCl, 1 mM MgCl_2 , 1mM BaCl_2 , 10 μM methoxyverapamil, 5 mM HEPES, pH 7.4 with or without 10 mM CaCl_2 . Oocytes were kept in modified Barth's solution containing 88 mM NaCl, 1 mM KCl, 2.4 mM NaHCO_3 , 0.8 mM MgSO_4 , 0.3 mM $\text{Ca}(\text{NO}_3)_2$, 0.4 mM CaCl_2 and 5 mM HEPES, pH 7.4 supplemented with 25 $\mu\text{g/ml}$ gentamycin. The final solutions were titrated to the pH indicated using HCl or NaOH. The flow rate of the superfusion was 20 ml/min and a complete exchange of the bath solution was reached within about 10 s. For direct determination of Ca^{2+} currents (I_{Ca}), the experiments were performed with Cl^- depleted oocytes (bathed




Fig. 1. Stimulation of tracer Ca²⁺ entry by the combined expression of TRPV5, NHERF2 and SGK1 or SGK 3 but not of SGK2 or protein kinase B. *Xenopus laevis* oocytes were injected with either water or cRNA encoding TRPV5 with or without co-injection of SGK1, SGK2, SGK3, constitutively active T_{308D}, S_{473D}PKB or NHERF2 alone and combined with the mentioned kinases. ⁴⁵Ca²⁺ uptake was measured after 30 minutes of incubation (n = 8-46). * denotes significant difference (p < 0.05) between oocytes expressing TRPV5 together with either SGK1 or SGK3 and NHERF2 and oocytes expressing TRPV5 alone.

for 24 hours in Cl⁻ free medium) in the absence of extracellular Cl⁻ in the bath and KCl (3 M) filled agar bridges were used as reference electrodes to minimize liquid junction potentials. Under those experimental conditions, the currents were insensitive to the Cl⁻ channel blocker NPPB (100 μM). In the presence of Cl⁻ and absence of Cl⁻ channel inhibitors, the addition of 10 mM CaCl₂ induced an inward current (I_{Cl(Ca)}) which was created by entry of Ca²⁺ and subsequent activation of Ca²⁺ sensitive Cl⁻ channels [4]. The peak inward current was taken as a measure for TRPV5 activity. I_{Cl(Ca)} is triggered by the intracellular calcium concentration close to the cell membrane which is in turn a function of Ca²⁺ entry through the epithelial calcium channel TRPV5. Thus, I_{Cl(Ca)} activity mirrors activation and inactivation kinetics of TRPV5 [35]. I_{Cl(Ca)} is inhibited by NPPB. Where indicated, ruthenium red was added at a concentration of 5 μM to inhibit TRPV5 [3]. Ruthenium red further inhibits voltage-sensitive calcium channels [36], which in our experiments are already suppressed by 10 μM methoxyverapamil. Expression and currents may vary from batch




Fig. 2. Inhibitory effect of ruthenium red on TRPV5 activity. Ca²⁺ entry via TRPV5 with or without S^{422D}SGK1 and NHERF2 was inhibited by addition of 5 μM ruthenium red (n = 6-8). *Xenopus laevis* oocytes were injected with water alone or TRPV5 with or without cRNA encoding S^{422D}SGK1 and NHERF2. * denotes significant difference (p < 0.05) between oocytes expressing TRPV5 and oocytes injected with water. + denotes significant difference (p < 0.05) between oocytes expressing TRPV5 together with S^{422D}SGK1 and NHERF2 and oocytes expressing TRPV5 alone.

to batch. Thus, care was taken to make comparisons always within batches.

Detection of cell surface expression by chemiluminescence

Defolliculated oocytes were first injected with SGK1 cRNA (7.5 ng/oocyte) and/or NHERF2 cRNA (5 ng/oocyte), and one day later with TRPV5-HA cRNA which contains a HA epitope extracellularly between amino acid 376 and 377 (2.5 ng/oocyte). Oocytes were incubated with 1 μg/ml primary rat monoclonal anti-HA antibody (clone 3F10, Boehringer, Mannheim, Germany), and 2 μg/ml secondary, peroxidase-conjugated affinity-purified F(ab')₂ goat anti-rat IgG antibody (Jackson ImmunoResearch, West Grove, USA). Individual oocytes were placed in 20 μl of SuperSignal ELISA Femto Maximum Sensitivity Substrate (Pierce, Rockford, USA), and chemiluminescence was quantified in a luminometer by integrating the signal over a period of 1 s. Results are given in relative light units (RLU). RLU obtained with H₂O-injected




Fig. 3. Stimulation of Ca^{2+} currents by the combined expression of TRPV5, NHERF2 and $\text{S}^{422\text{D}}$ SGK1. **A** TRPV5 expressing oocytes but not water injected oocytes show an inward current upon addition of 10 mM Ca^{2+} to the bath. Coexpression of TRPV5 together with $\text{S}^{422\text{D}}$ SGK1 and NHERF2 leads to a significant increase of the current compared to oocytes expressing TRPV5 alone. $\text{S}^{422\text{D}}$ SGK1 and NHERF2 alone do not alter TRPV5 channel activity significantly ($n = 8-26$). Currents are normalized to the values obtained with oocytes expressing TRPV5 alone. * denotes significant difference ($p < 0.05$) between oocytes expressing TRPV5/ $\text{S}^{422\text{D}}$ SGK1 together with NHERF2 and

oocytes injected with TRPV5 alone. **B** The inward current into TRPV5 expressing oocytes in the absence of Cl^- induced by high extracellular Ca^{2+} is not sensitive to the chloride channel blocker NPPB. Currents are normalized to the values obtained with oocytes expressing TRPV5 alone in the absence of NPPB. * denotes significant difference ($p < 0.05$) between oocytes expressing TRPV5, $\text{S}^{422\text{D}}$ SGK1 together with NHERF2 and oocytes expressing TRPV5 alone. + denotes significant difference between oocytes expressing TRPV5, $\text{S}^{422\text{D}}$ SGK1 together with NHERF2 and oocytes expressing TRPV5 alone under continuous presence of 100 μM NPPB ($n = 5-12$).

oocytes (control oocytes) were subtracted from the values obtained with oocytes injected with TRPV5-HA alone or together with SGK1 and/or NHERF2.

Statistics


Data are provided as means \pm SEM, n represents the number of oocytes investigated. All experiments were repeated with at least 3 batches of oocytes from different frogs; in all repetitions qualitatively similar data were obtained. All data were tested for significance using ANOVA and only results with $P < 0.05$ were considered statistically significant.

Results

As evident from Fig. 1, uptake of radioactive Ca^{2+} ($^{45}\text{Ca}^{2+}$) into *Xenopus* oocytes was increased some 7 fold by expression of TRPV5. The increase of Ca^{2+} uptake reflects the Ca^{2+} transporting capacity of TRPV5.

Additional expression of either NHERF2, SGK1, SGK2, SGK3 or constitutively active $\text{T}^{308\text{D},\text{S}^{473\text{D}}}\text{PKB}$ alone did not further stimulate $^{45}\text{Ca}^{2+}$ entry. However, expression of TRPV5 together with both SGK1 and NHERF2 resulted in significant further stimulation of Ca^{2+} uptake. In the absence of TRPV5, expression of SGK1 and NHERF2 did not increase Ca^{2+} uptake (not shown). Thus, coexpression of SGK1 and NHERF2 increased Ca^{2+} uptake by stimulation of TRPV5. Similarly, the coexpression of TRPV5 together with SGK3 and NHERF2 increased Ca^{2+} uptake. Both kinases stimulated TRPV5, an effect requiring the presence of NHERF2. In contrast, the coexpression of neither SGK2 nor $\text{T}^{308\text{D},\text{S}^{473\text{D}}}\text{PKB}$ significantly modified the Ca^{2+} uptake into oocytes expressing TRPV5 and NHERF2. Thus, SGK1 and SGK3, but not SGK2 or $\text{T}^{308\text{D},\text{S}^{473\text{D}}}\text{PKB}$ stimulate TRPV5 in the presence of NHERF2.

Fig. 4. TRPV5 mediated calcium currents indirectly activate an endogenous chloride conductance ($I_{Cl(Ca)}$). **A - C** In TRPV5 expressing *Xenopus laevis* oocytes, hyperpolarization from -50 mV to -110 mV in the presence of 10 mM Ca^{2+} triggered a rapidly activating, slowly and partially inactivating inward current (A) while in water injected cells hyperpolarization does not increase any voltage-dependent current (B). Elevation of cytosolic Ca^{2+} by addition of the Ca^{2+} ionophore ionomycin (10 μ M) led to the stimulation of a non inactivating current (C). **D - E** $I_{Cl(Ca)}$ was stimulated by coexpression of TRPV5 together with NHERF2 and either S^{422D} SGK1 (D), SGK1 or SGK3 (E). Coexpression of TRPV5 with either NHERF2, S^{422D} SGK1, SGK1 or SGK3 alone did not significantly enhance $I_{Cl(Ca)}$ (D, E) ($n = 19-26$). **F** The ionomycin induced endogenous Cl⁻ current was not significantly enhanced by expression of S^{422D} SGK1, NHERF2 or both ($n = 6$). * denotes significant difference ($p < 0.05$) between oocytes expressing TRPV5 together with either S^{422D} SGK1, SGK1 or SGK3 and NHERF2 and oocytes expressing TRPV5 alone.



Previously, TRPV5 has been shown to be inhibited by ruthenium red [32], a polycationic dye. Thus, the effect of ruthenium red on Ca^{2+} uptake was tested. As shown in Fig. 2, 5 μ M ruthenium red completely inhibited Ca^{2+} uptake in TRPV5 expressing oocytes with or without additional expression of S^{422D} SGK1 and NHERF2.

Further studies have been performed to determine the influence of S^{422D} SGK1 on TRPV5 mediated currents. As reported earlier, the entry of Ca^{2+} triggers Ca^{2+} sensitive Cl⁻ channels [35]. To determine the Ca^{2+} current directly, the Cl⁻ current had to be suppressed. Therefore, the oocytes were bathed in Cl⁻ free extracellular solution




Fig. 5. Inhibition of the TRPV5 induced current $I_{Cl(Ca)}$ by ruthenium red and NPPB. *Xenopus laevis* oocytes were injected with water alone or TRPV5 with or without cRNA encoding S^{422D}SGK1 and NHERF2. **A** Inhibition of $I_{Cl(Ca)}$ by application of 5 µM ruthenium red (n = 7). **B** Inhibition of $I_{Cl(Ca)}$ by application of 100 µM NPPB (n = 5-8). Insert in A depicts dose-dependent inhibition of TRPV5 activity. * denotes significant difference (p < 0.05) between *Xenopus* oocytes expressing TRPV5 and oocytes injected with water. + denotes significant difference (p < 0.05) between *Xenopus* oocytes expressing TRPV5 together with S^{422D}SGK1 and NHERF2 and oocytes expressing TRPV5 alone.




Fig 6. TRPV5 currents and cell surface abundance are similarly increased by coexpression of S^{422D}SGK1 and NHERF2. *Xenopus laevis* oocytes were injected with water or cRNA encoding TRPV5 alone or with S^{422D}SGK1 and wild-type NHERF2. The combined expression of S^{422D}SGK1 and wild-type NHERF2 increases TRPV5 channel activity and cell surface abundance. * indicates significant difference (p < 0.05) between expression of TRPV5 alone and of TRPV5 coexpressed with S^{422D}SGK1 and NHERF2 (n = 20-29).

for 24 hours. As shown in figure 3, coexpression of TRPV5 together with S^{422D}SGK1 and NHERF2 led to a significant increase of the current induced by addition of 10 mM CaCl₂. In contrast, the current was not increased by coexpression of TRPV5 with either NHERF2 or S^{422D}SGK1 alone (Fig. 3A). I_{Ca} was insensitive to the Cl⁻ channel inhibitor NPPB (Fig. 3B).

In the presence of Cl⁻ the Ca²⁺ entry through TRPV5 stimulated Ca²⁺ sensitive Cl⁻ channels leading to the appearance of a large Cl⁻ current ($I_{Cl(Ca)}$). In TRPV5-expressing oocytes, hyperpolarization from -50 mV to -110 mV in the presence of 10 mM Ca²⁺ triggered a rapidly activating, slowly and partially inactivating inward current (Fig. 4A). The transiently activating current was only present in *Xenopus* oocytes expressing TRPV5. In water injected oocytes increase of cytosolic Ca²⁺ activity by addition of Ca²⁺ ionophore ionomycin (10 µM) led to the

stimulation of a non inactivating current (Fig. 4C). Fig. 4B depicts a recording from a non-injected oocyte. Similar to tracer the Ca^{2+} uptake and I_{Ca} , $I_{\text{Cl}(\text{Ca})}$ was stimulated by coexpression of TRPV5 together with NHERF2 and either, $\text{S}^{422\text{D}}$ SGK1 (Fig. 4D), SGK1 (Fig. 4E) or SGK3 (4E). Coexpression of TRPV5 with either NHERF2, $\text{S}^{422\text{D}}$ SGK1, SGK1 or SGK3 alone did not significantly enhance $I_{\text{Cl}(\text{Ca})}$. Moreover, neither SGK2 nor $\text{T}^{308\text{D},\text{S}^{473\text{D}}}$ PKB stimulated $I_{\text{Cl}(\text{Ca})}$, even upon NHERF2 coexpression (Fig. 4E). Thus, $I_{\text{Cl}(\text{Ca})}$ was enhanced by SGK1 and SGK3 only in the presence of NHERF2. In oocytes not expressing TRPV5, the Ca^{2+} sensitive Cl^- current could be activated by the Ca^{2+} ionophore ionomycin (10 μM). As shown in Fig. 4F, the ionomycin induced Cl^- current was not significantly enhanced by expression of $\text{S}^{422\text{D}}$ SGK1, NHERF2 or both. Thus, the increase of $I_{\text{Cl}(\text{Ca})}$ following coexpression of $\text{S}^{422\text{D}}$ SGK1 and NHERF2 was not due to upregulation of the Cl^- channel but due to upregulation of TRPV5.

As illustrated in Fig. 5, $I_{\text{Cl}(\text{Ca})}$ was abolished by ruthenium red (Fig. 5A) and by the Cl^- channel blocker NPPB (Fig. 5B).

To discriminate whether increased Ca^{2+} fluxes and Ca^{2+} currents upon coexpression of SGKs and NHERF2 are due to modulation of TRPV5 gating kinetics or induced transport of TRPV5 to the cell membrane, we quantified cell surface expression of the channel and the peak current voltage. As depicted in Fig. 6, TRPV5 currents and cell surface expression of the channel protein were similarly increased by coexpression of $\text{S}^{422\text{D}}$ SGK1 and NHERF2.

Discussion

The present observations clearly demonstrate that expression of TRPV5 induces Ca^{2+} influx allowing the cellular accumulation of tracer Ca^{2+} and generating a Ca^{2+} current. In the presence of Cl^- , Ca^{2+} entry through TRPV5 generates further currents [3] by activation of endogenous Ca^{2+} sensitive Cl^- channels [37].

More importantly, the present observations disclose a completely novel mechanism regulating TRPV5 activity, i.e. the regulation by two members of the serum and glucocorticoid inducible kinase family and the NHE regulating factor NHERF2. The effect of SGK1 depends on an intact catalytic subunit, as the inactive mutant $\text{K}^{127\text{N}}$ SGK1 does not influence TRPV5 even in the presence of NHERF2. Thus, the kinases are obviously effective through phosphorylation of target proteins. The

stimulating effect is shared by SGK1 and SGK3, but not by SGK2 or PKB. The TRPV5-stimulating capacity correlates with the ability of the two kinases to bind NHERF2 [6]. Expressed wild type SGK1 and SGK3 are not constitutively active but require activation. The kinases are activated through a signaling pathway involving phosphatidylinositol 3 kinase (PI3-K) and phosphoinositide dependent kinase (PDK) [27]. Obviously, the activation of the kinases is not limiting in oocytes, as expression of wild type SGK3 is similarly effective compared to $\text{S}^{422\text{D}}$ SGK1. In earlier studies, wild type SGK1 has similarly been shown to be effective [38]. In contrast to SGK1 and SGK3, SGK2 does not bind NHERF2 [6] and is, according to the present study, not able to stimulate TRPV5. PKB is similarly not able to stimulate TRPV5.

A well known function of SGK1 is its participation in the regulation of the epithelial Na^+ channel ENaC [17, 18, 20, 39-45]. SGK1 is effective by increasing the abundance of the ENaC protein within the cell membrane [30, 39, 45]. Most recently, SGK1 has been shown to upregulate the voltage gated K^+ channel Kv1.3 [46-48], the epithelial Na^+/H^+ exchanger NHE3 [6] and the renal epithelial K^+ channel ROMK1 [49]. Similar to SGK-dependent regulation of TRPV5, the regulation of ROMK1 by SGK1 has been shown to depend on NHERF2 [49].

In contrast to the regulation of TRPV5, NHE3 and ROMK1, the regulation of ENaC or Kv1.3 does not require the participation of NHERF2. SGK1 stimulates ENaC by phosphorylating the ubiquitin protein ligase Nedd4-2 in a PY motif-dependent manner [50, 51]. The phosphorylation impedes the binding of Nedd4-2 to ENaC [50]. The ENaC sequence does not include a PDZ motif, suggesting that NHERF2 can not directly interact with ENaC.

As NHERF2 regulates both TRPV5 and NHE3, it may participate in the link between TRPV5 activity and acid base balance. TRPV5 displays an exquisite H^+ sensitivity [1, 3, 22]. Its activity is markedly reduced by lowering of the ambient pH. The sensitivity of renal tubular Ca^{2+} transport to H^+ is of physiological significance, as on the one hand mineralization of bone depends on the deposition of highly alkaline Ca^{2+} salts, [52] and on the other, precipitation of Ca^{2+} phosphate salts is favoured by alkalization of urine [53]. Most recently, NHERF2 has been shown to direct the signaling of the PTH receptor [54]. Since PTH is a key regulator of TRPV5 [1], this effect may similarly contribute to the regulation of TRPV5. However, according to the present observations, NHERF2 alone is unable to activate TRPV5 but requires

the additional action of SGK1 or of one of its isoforms. SGK1 expression is highly variable, as it is upregulated by glucocorticoids [15], aldosterone [17-21], cell shrinkage [16] and a wide variety of additional factors [24]. Notably, SGK1 is under transcriptional control of $1,25(\text{OH})_2\text{D}_3$ [22]. It is thus tempting to speculate that SGK1 participates in the regulation of TRPV5 by $1,25(\text{OH})_2\text{D}_3$ and PTH. Activation of SGK1 can be accomplished by insulin and IGF1 through PI3 kinase and PDK1 [27, 28]. SGK1 integrates the signals coming from genomic regulators on the one hand and insulin or IGF1 on the other [55]. Thus, activation of SGK1 may contribute to the stimulating effect of both, $1,25(\text{OH})_2\text{D}_3$ and IGF1 on intestinal Ca^{2+} absorption [56].

In conclusion, TRPV5 is the target of a complex regulatory mechanism involving both, the NHE regulating factor NHERF2 and the serine/threonine kinases SGK1 and 3. The concerted action of NHERF2 and the kinases markedly up regulates the activity of this key channel in the regulation of Ca^{2+} homeostasis.

Abbreviations

SGK (serum and glucocorticoid inducible kinase); NHERF (sodium hydrogen exchanger regulating factor); Nedd (Neuronal cell expressed developmentally downregulated ubiquitin ligase).

Acknowledgements

The authors acknowledge the technical assistance of B. Noll and the meticulous preparation of the manuscript by Lejla Subasic and Tanja Loch. This study was supported by the Deutsche Forschungsgemeinschaft, Nr. La 315/4-4 and La 315/5-1, the Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (Center for Interdisciplinary Clinical Research) 01 KS 9602 and the National Institutes of Health Grant DK-44484. PC is supported by the UK Medical Research Council and The Royal Society. RB is supported by grants from the Dutch Organization of Scientific Research (Zon-Mw 016.006.001, Zon-Mw 902.18.298, NWO-ALW 810.38.004, NWO-ALW 805.09.042).

References

- Hoenderop JG, Nilius B, Bindels RJ: Molecular mechanism of active Ca^{2+} reabsorption in the distal nephron. *Annu Rev Physiol* 2002;64:529-549.
- Hoenderop JG, Muller D, Suzuki M, van Os CH, Bindels RJ: Epithelial calcium channel: gate-keeper of active calcium reabsorption. *Curr Opin Nephrol Hypertens* 2000;9:335-340.
- Hoenderop JG, Van Der Kemp AW, Hartog A, van Os CH, Willems PH, Bindels RJ: The epithelial calcium channel, ECaC, is activated by hyperpolarization and regulated by cytosolic calcium. *Biochem Biophys Res Commun* 1999;261:488-492.
- Hoenderop JG, Van Der Kemp AW, Hartog A, van de Graaf SF, van Os CH, Willems PH, Bindels RJ: Molecular identification of the apical Ca^{2+} channel in $1, 25$ -dihydroxyvitamin D_3 -responsive epithelia. *J Biol Chem* 1999;274:8375-8378.
- Hoenderop JG, Willems PH, Bindels RJ: Toward a comprehensive molecular model of active calcium reabsorption. *Am J Physiol Renal Physiol* 2000;278:F352-F360.
- Yun CC, Chen Y, Lang F: Glucocorticoid activation of $\text{Na}(+)/\text{H}(+)$ exchanger isoform 3 revisited. The roles of SGK1 and NHERF2. *J Biol Chem* 2002;277:7676-7683.
- Shenolikar S, Weinman EJ: NHERF: targeting and trafficking membrane proteins. *Am J Physiol Renal Physiol* 2001;280:F389-F395.
- Weinman EJ, Steplock D, Wang Y, Shenolikar S: Characterization of a protein cofactor that mediates protein kinase A regulation of the renal brush border membrane $\text{Na}(+)/\text{H}(+)$ exchanger. *J Clin Invest* 1995;95:2143-2149.
- Mery L, Strauss B, Dufour JF, Krause KH, Hoth M: The PDZ-interacting domain of TRPC4 controls its localization and surface expression in HEK293 cells. *J Cell Sci* 2002;115:3497-3508.
- Hoenderop JG, Hartog A, Stuver M, Doucet A, Willems PH, Bindels RJ: Localization of the epithelial $\text{Ca}(2+)$ channel in rabbit kidney and intestine. *J Am Soc Nephrol* 2000;11:1171-1178.
- Wade JB, Welling PA, Donowitz M, Shenolikar S, Weinman EJ: Differential renal distribution of NHERF isoforms and their colocalization with NHE3, ezrin, and ROMK. *Am J Physiol Cell Physiol* 2001;280:C192-C198.
- Yun CC: Concerted Roles of SGK1 and the Na^+/H^+ Exchanger Regulatory Factor 2 (NHERF2) in Regulation of NHE3. *Cell Physiol Biochem* 2003;13:029-040.
- Yun CH, Lamprecht G, Forster DV, Sidor A: NHE3 kinase A regulatory protein E3KARP binds the epithelial brush border Na^+/H^+ exchanger NHE3 and the cytoskeletal protein ezrin. *J Biol Chem* 1998;273:25856-25863.

- 14 Firestone GL, Giampaolo JR, O'Keefe BA: Stimulus-dependent regulation of the serum and glucocorticoid inducible protein kinase (Sgk) transcription, subcellular localization and enzymatic activity. *Cell Physiol Biochem* 2003;13:1-12.
- 15 Webster MK, Goya L, Ge Y, Maiyar AC, Firestone GL: Characterization of sgk, a novel member of the serine/threonine protein kinase gene family which is transcriptionally induced by glucocorticoids and serum. *Mol Cell Biol* 1993;13:2031-2040.
- 16 Waldegger S, Barth P, Raber G, Lang F: Cloning and characterization of a putative human serine/threonine protein kinase transcriptionally modified during anisotonic and isotonic alterations of cell volume. *Proc Natl Acad Sci U S A* 1997;94:4440-4445.
- 17 Chen SY, Bhargava A, Mastroberardino L, Meijer OC, Wang J, Buse P, Firestone GL, Verrey F, Pearce D: Epithelial sodium channel regulated by aldosterone-induced protein sgk. *Proc Natl Acad Sci U S A* 1999;96:2514-2519.
- 18 Naray-Fejes-Toth A, Canessa C, Cleveland ES, Aldrich G, Fejes-Toth G: sgk is an aldosterone-induced kinase in the renal collecting duct. Effects on epithelial Na⁺ channels. *J Biol Chem* 1999;274:16973-16978.
- 19 Brennan FE, Fuller PJ: Rapid upregulation of serum and glucocorticoid-regulated kinase (sgk) gene expression by corticosteroids in vivo. *Mol Cell Endocrinol* 2000;166:129-136.
- 20 Shigaev A, Asher C, Latter H, Garty H, Reuveny E: Regulation of sgk by aldosterone and its effects on the epithelial Na⁺ channel. *Am J Physiol Renal Physiol* 2000;278:F613-F619.
- 21 Cowling RT, Birnboim HC: Expression of serum- and glucocorticoid-regulated kinase (sgk) mRNA is up-regulated by GM-CSF and other proinflammatory mediators in human granulocytes. *J Leukoc Biol* 2000;67:240-248.
- 22 Akutsu N, Lin R, Bastien Y, Bestawros A, Enepekides DJ, Black MJ, White JH: Regulation of gene Expression by 1alpha,25-dihydroxyvitamin D3 and Its analog EB1089 under growth-inhibitory conditions in squamous carcinoma Cells. *Mol Endocrinol* 2001;15:1127-1139.
- 23 Waldegger S, Klingel K, Barth P, Sauter M, Rfer ML, Kandolf R, Lang F: h-sgk serine-threonine protein kinase gene as transcriptional target of transforming growth factor beta in human intestine. *Gastroenterology* 1999;116:1081-1088.
- 24 Lang F, Cohen P: Regulation and physiological roles of serum- and glucocorticoid-induced protein kinase isoforms. *Sci STKE* 2001;2001:RE17.
- 25 Kobayashi T, Deak M, Morrice N, Cohen P: Characterization of the structure and regulation of two novel isoforms of serum- and glucocorticoid-induced protein kinase. *Biochem J* 1999;344 Pt 1:189-197.
- 26 Alessi DR, Andjelkovic M, Caudwell B, Cron P, Morrice N, Cohen P, Hemmings BA: Mechanism of activation of protein kinase B by insulin and IGF-1. *EMBO J* 1996;15:6541-6551.
- 27 Kobayashi T, Cohen P: Activation of serum- and glucocorticoid-regulated protein kinase by agonists that activate phosphatidylinositide 3-kinase is mediated by 3-phosphoinositide-dependent protein kinase-1 (PDK1) and PDK2. *Biochem J* 1999;339 (Pt 2):319-328.
- 28 Park J, Leong ML, Buse P, Maiyar AC, Firestone GL, Hemmings BA: Serum and glucocorticoid-inducible kinase (SGK) is a target of the PI 3-kinase-stimulated signaling pathway. *EMBO J* 1999;18:3024-3033.
- 29 Meier R, Alessi DR, Cron P, Andjelkovic M, Hemmings BA: Mitogenic activation, phosphorylation, and nuclear translocation of protein kinase Bbeta. *J Biol Chem* 1997;272:30491-30497.
- 30 Loffing J, Zecevic M, Feraille E, Kaissling B, Asher C, Rossier BC, Firestone GL, Pearce D, Verrey F: Aldosterone induces rapid apical translocation of ENaC in early portion of renal collecting system: possible role of SGK. *Am J Physiol Renal Physiol* 2001;280:F675-F682.
- 31 Wagner CA, Friedrich B, Setiawan I, Lang F, Broer S: The use of *Xenopus laevis* oocytes for the functional characterization of heterologously expressed membrane proteins. *Cell Physiol Biochem* 2000;10:1-12.
- 32 Hoenderop JG, Vennekens R, Muller D, Prenen J, Droogmans G, Bindels RJ, Nilius B: Function and expression of the epithelial Ca²⁺ channel family: comparison of mammalian ECaC1 and 2. *J Physiol* 2001;537:747-761.
- 33 Peng JB, Chen XZ, Berger UV, Vassilev PM, Tsukaguchi H, Brown EM, Hediger MA: Molecular cloning and characterization of a channel-like transporter mediating intestinal calcium absorption. *J Biol Chem* 1999;274:22739-22746.
- 34 Peng JB, Chen XZ, Berger UV, Vassilev PM, Brown EM, Hediger MA: A rat kidney-specific calcium transporter in the distal nephron. *J Biol Chem* 2000;275:28186-28194.
- 35 Miledi R, Parker I: Chloride current induced by injection of calcium into *Xenopus* oocytes. *J Physiol* 1984;357:173-183.
- 36 Tapia R, Velasco I: Ruthenium red as a tool to study calcium channels, neuronal death and the function of neural pathways. *Neurochem Int* 1997;30:137-147.
- 37 Callamaras N, Parker I: Ca²⁺-dependent activation of Cl⁻ currents in *Xenopus* oocytes is modulated by voltage. *Am J Physiol Cell Physiol* 2000;278:C667-C675.
- 38 Boehmer C, Henke G, Schniepp R, Palmada M, Rothstein JD, Broer S, Lang F: Regulation of the glutamate transporter EAAT1 by the ubiquitin ligase Nedd4-2 and the serum and glucocorticoid-inducible kinase isoforms SGK1/3 and protein kinase B. *J Neurochem* 2003;86:1181-1188.
- 39 Alvarez de la Rosa D, Zhang P, Naray-Fejes-Toth A, Fejes-Toth G, Canessa CM: The serum and glucocorticoid kinase sgk increases the abundance of epithelial sodium channels in the plasma membrane of *Xenopus* oocytes. *J Biol Chem* 1999;274:37834-37839.
- 40 Böhrer C, Wagner CA, Beck S, Moschen I, Melzig J, Werner A, Lin JT, Lang F, Wehner F: The shrinkage-activated Na⁺ conductance of rat hepatocytes and its possible correlation to rENaC. *Cell Physiol Biochem* 2000;10:187-194.
- 41 Lang F, Klingel K, Wagner CA, Stegen C, Warntges S, Friedrich B, Lanzendorfer M, Melzig J, Moschen I, Steuer S, Waldegger S, Sauter M, Paulmichl M, Gerke V, Risler T, Gamba G, Capasso G, Kandolf R, Hebert SC, Massry SG, Broer S: Deranged transcriptional regulation of cell-volume-sensitive kinase hSGK in diabetic nephropathy. *Proc Natl Acad Sci U S A* 2000;97:8157-8162.
- 42 Lang F, Henke G, Embark HM, Waldegger S, Palmada M, Bohmer C, Vallon V: Regulation of channels by the serum and glucocorticoid-inducible kinase - implications for transport, excitability and cell proliferation. *Cell Physiol Biochem* 2003;13:41-50.
- 43 Pearce D: SGK1 Regulation of Epithelial Sodium Transport. *Cell Physiol Biochem* 2003;13:013-020.
- 44 Verrey F, Loffing J, Zecevic M, Heitzmann D, Staub O: SGK1: aldosterone-induced relay of Na⁺ transport regulation in distal kidney nephron cells. *Cell Physiol Biochem* 2003;13:021-028.
- 45 Wagner CA, Ott M, Klingel K, Beck S, Melzig J, Friedrich B, Wild KN, Broer S, Moschen I, Albers A, Waldegger S, Tummler B, Egan ME, Geibel JP, Kandolf R, Lang F: Effects of the serine/threonine kinase SGK1 on the epithelial Na⁺ channel (ENaC) and CFTR: implications for cystic fibrosis. *Cell Physiol Biochem* 2001;11:209-218.

- 46 Wärrntges S, Friedrich B, Henke G, Duranton C, Lang PA, Waldegger S, Meyermann R, Kuhl D, Speckmann EJ, Obermuller N, Witzgall R, Mack AF, Wagner HJ, Wagner A, Broer S, Lang F: Cerebral localization and regulation of the cell volume-sensitive serum- and glucocorticoid-dependent kinase SGK1. *Pflugers Arch* 2002;443:617-624.
- 47 Gamper N, Fillon S, Huber SM, Feng Y, Kobayashi T, Cohen P, Lang F: IGF-1 up-regulates K⁺ channels via PI3-kinase, PDK1 and SGK1. *Pflugers Arch* 2002;443:625-634.
- 48 Gamper N, Fillon S, Feng Y, Friedrich B, Lang PA, Henke G, Huber SM, Kobayashi T, Cohen P, Lang F: K⁽⁺⁾ channel activation by all three isoforms of serum- and glucocorticoid-dependent protein kinase SGK. *Pflugers Arch* 2002;445:60-66.
- 49 Yun CC, Palmada M, Embark HM, Fedorenko O, Feng Y, Henke G, Setiawan I, Boehmer C, Weinman EJ, Sandrasagra S, Korbmacher C, Cohen P, Pearce D, Lang F: The Serum and Glucocorticoid-Inducible Kinase SGK1 and the Na⁽⁺⁾/H⁽⁺⁾ Exchange Regulating Factor NHERF2 Synergize to Stimulate the Renal Outer Medullary K⁽⁺⁾ Channel ROMK1. *J Am Soc Nephrol* 2002;13:2823-2830.
- 50 Debonneville C, Flores SY, Kamynina E, Plant PJ, Tauxe C, Thomas MA, Munster C, Chraibi A, Pratt JH, Horisberger JD, Pearce D, Loffing J, Staub O: Phosphorylation of Nedd4-2 by Sgk1 regulates epithelial Na⁽⁺⁾ channel cell surface expression. *EMBO J* 2001;20:7052-7059.
- 51 Snyder PM, Olson DR, Thomas BC: Serum and glucocorticoid-regulated kinase modulates Nedd4-2-mediated inhibition of the epithelial Na⁺ channel. *J Biol Chem* 2002;277:5-8.
- 52 Bushinsky DA, Krieger NS: Role of the skeleton in calcium homeostasis.; in Seldin W, Giebisch G (eds): *The Kidney*. New York, Raven Press, 1992, pp 2395-2430.
- 53 Pak CYC: Pathophysiology of calcium nephrolithiasis.; in Seldin W, Giebisch G (eds): *The Kidney*. New York, Raven Press, 1992, pp 2461-2480.
- 54 Mahon MJ, Donowitz M, Yun CC, Segre GV: Na⁽⁺⁾/H⁽⁺⁾ exchanger regulatory factor 2 directs parathyroid hormone 1 receptor signalling. *Nature* 2002;417:858-861.
- 55 Wang J, Barbry P, Maiyar AC, Rozansky DJ, Bhargava A, Leong M, Firestone GL, Pearce D: SGK integrates insulin and mineralocorticoid regulation of epithelial sodium transport. *Am J Physiol Renal Physiol* 2001;280:F303-F313.
- 56 Bouillon R: Growth hormone and bone. *Horm Res* 1991;36 Suppl 1:49-55.