

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/60185>

Please be advised that this information was generated on 2018-10-20 and may be subject to change.

Anaerobic ammonium oxidation in an estuarine sediment

Nils Risgaard-Petersen^{1,*}, Rikke Louise Meyer², Markus Schmid³, Mike S. M. Jetten⁴, Alex Enrich-Prast⁵, Søren Rysgaard¹, Niels Peter Revsbech²

¹National Environmental Research Institute, Department of Marine Ecology, Vejlsøvej 25, 8600 Silkeborg, Denmark

²University of Aarhus, Institute of Biological Sciences, Department of Microbial Ecology, Ny Munkegade Building 540, 8000 Århus C, Denmark

³Delft University of Technology, Department of Biotechnology, Delft, The Netherlands

⁴KU Nijmegen, Department of Microbiology, Nijmegen, The Netherlands

⁵Depto. Ecologia, CCS, Ilha do Fundão, Cidade Universitária, Universidade Federal do Rio de Janeiro, Rio de Janeiro, 21951-590, Brasil

ABSTRACT: The occurrence and significance of the anammox (anaerobic ammonium oxidation) process relative to denitrification was studied in photosynthetically active sediment from 2 shallow-water estuaries: Randers Fjord and Norsminde Fjord, Denmark. Anammox accounted for 5 to 24 % of N₂ production in Randers Fjord sediment, whereas no indication was seen of the process in sediment from Norsminde Fjord. It is suggested that the presence of anammox in Randers Fjord and its absence from Norsminde Fjord is associated with differences in the availability of NO₃⁻ + NO₂⁻ (NO_x⁻) in the suboxic zone of the sediment. In Randers Fjord, NO_x⁻ is present in the water column throughout the year and NO_x⁻ porewater profiles showed that NO_x⁻ penetrates into the suboxic zone of the sediment. In Norsminde Fjord, NO_x⁻ is absent from the water column during the summer months and, via assimilation, benthic microalgae may prevent penetration of NO_x⁻ into the suboxic zone of the sediment. Volume-specific anammox rates in Randers Fjord were comparable with rates measured previously in Skagerrak sediment by other investigators, but denitrification rates were 10 to 15 times higher. Thus, anammox contributes less to N₂ production in Randers Fjord than in Skagerrak sediment. We propose that the lower contribution of anammox in Randers Fjord is linked to the higher availability of easily accessible carbon, which supports a higher population of denitrifying bacteria. Amplification of DNA extracted from the sediment samples from Randers Fjord using planctomycete-specific primers yielded 16S rRNA gene sequences closely related to candidatus *Scalindua sorokinii* found in the Black Sea by other investigators. The present study thus confirms the link between the presence of bacteria affiliated with candidatus *S. sorokinii* and the anammox reaction in marine environments. Anammox rates in sediment with intact chemical gradients were estimated using both ¹⁵N and microsensor techniques. Anammox rates estimated with microsensors were less than 22 % of the rates measured with isotopes. It is suggested that this discrepancy was due to the presence of fauna, because the applied ¹⁵N technique captures total N₂ production while the microsensor technique only captures diffusion-controlled N₂ production at the sediment surface. This hypothesis was verified by consistent agreement between the methods when applied to defaunated sediments.

KEY WORDS: Anammox · Denitrification · Planctomycetes · *Scalindula*

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

The biologically mediated reduction of NO₃⁻ + NO₂⁻ (NO_x⁻) to N₂ through denitrification is generally considered to be the major process responsible for

removal of nitrogen from the sea (Devol 1991). Until recently, it was believed that denitrification was based solely on the reduction of NO_x⁻ to N₂ in an oxygen-free environment by facultatively aerobic bacteria with an organotrophic metabolism (Zumft 1992). Within this

model, the major NO_x^- source for denitrification is nitrification, i.e. the bacterial oxidation of NH_4^+ or NO_2^- with O_2 . Nitrification and denitrification are associated with the sediment/water interface in marine sediments. Nitrification takes place in the upper oxic zone of the sediment and denitrification in the suboxic zone just below the oxic/suboxic interface (Jensen et al. 1993, 1994). According to the classical concept, coupled nitrification–denitrification is facilitated by the transport of NO_x^- from the oxic NO_x^- production zones to the suboxic NO_x^- consumption zones in the sediment. In bioturbated sediments, the transport of NO_x^- from oxic to suboxic zones may be further enhanced by fauna-mediated processes such as biodiffusion and irrigation (Aller 1982, Aller & Aller 1998, Kristensen 2000 and references therein). This enhanced transport results in elevated production of N_2 (e.g. Pelegri et al. 1994, Svensson et al. 2001, Newell et al. 2002).

The classical view has recently been challenged by the discovery of alternative pathways of combined nitrogen transformations. These alternative pathways include the anaerobic oxidation of NH_4^+ to NO_3^- or N_2 with manganese oxides (Luther et al. 1997, Hulth et al. 1999) and the anaerobic oxidation of NH_4^+ to N_2 with NO_2^- (Mulder et al. 1995). While direct experimental evidence for anaerobic NH_4^+ oxidation to N_2 with manganese oxides is still lacking (Thamdrup & Dalsgaard 2000), it is now well documented that NH_4^+ can be oxidized anaerobically to N_2 with NO_2^- both in marine sediments and in open waters (Dalsgaard & Thamdrup 2002, Thamdrup & Dalsgaard 2002, Dalsgaard et al. 2003, Kuypers et al. 2003, Trimmer et al. 2003). This biologically mediated process is called anammox (anaerobic ammonium oxidation) (Strous et al. 1999). Strictly viewed, anammox is equivalent to denitrification since denitrification is defined as the conversion of NO_2^- to gaseous N (Payne 1981). However, existing literature addressing anammox in natural environments (Dalsgaard & Thamdrup 2002, Thamdrup & Dalsgaard 2002, Trimmer et al. 2003) uses the term denitrification for the purely NO_x^- -based N_2 production ($\text{E-donor} + 2\text{NO}_x^- \rightarrow \text{N}_2$) and the term anammox for the production of N_2 from NH_4^+ and NO_2^- . For the sake of consistency, we choose to follow this nomenclature.

Little is known about the biogeography of the anammox process, its microbiology, and its importance as a source of N_2 production relative to denitrification. According to current knowledge, the anammox process is carried out by autotrophic, obligately anaerobic bacteria of the phylum *Planctomycetes* (Strous et al. 1999, Schmid et al. 2000, 2003, Kuypers et al. 2003). Knowledge of the microbiology of the process in marine environments is limited to a single study in the Black Sea (Kuypers et al. 2003), which, however, also linked the occurrence of the anammox process to the

occurrence of anammox bacteria of the phylum *Planctomycetes*. The few published data addressing the occurrence and significance of this newly discovered process in marine environments originate mainly from studies of deep-water, offshore sediments and anoxic water columns (see Dalsgaard & Thamdrup 2002, Thamdrup & Dalsgaard 2002, Dalsgaard et al. 2003, Kuypers et al. 2003). These studies suggested that the anammox reaction may account for 30 to 70% of oceanic N_2 production (Devol 2003), which poses a great challenge to our view on the control of marine N_2 production. Only very recently has anammox activity been located in estuarine sediment from the Thames estuary, UK (Trimmer et al. 2003), where it accounted for <10% of benthic N_2 production.

The present study addresses the occurrence and significance of the anammox process relative to denitrification in photosynthetically active sediments from 2 shallow-water estuaries: Randers Fjord and Norsminde Fjord, Denmark. We identified bacteria responsible for the process using anammox-specific FISH (fluorescence *in situ* hybridization) probes, 16S rRNA gene-targeted PCR amplification with anammox-specific primers, and subsequent sequencing and phylogenetic analysis (Schmid et al. 2000). We furthermore obtained quantitative data on the denitrification and anammox process rates. In order to compare the results of the present study with data from previous studies of the anammox process in marine sediments, we assessed anammox and denitrification activity in sediment slurries according to Thamdrup & Dalsgaard (2002). This technique estimates the volume-specific anammox and denitrification rates. To assess process rates in sediments with intact chemical gradients, we combined techniques that estimate benthic N_2 production in intact sediment cores (Nielsen 1992, Risgaard-Petersen et al. 2003) with the ^{15}N technique of Thamdrup & Dalsgaard (2002), which also estimates the contribution of anammox to total N_2 production. In addition, N_2 production was estimated from porewater profiles of NO_x^- , to address the diffusion-controlled activity directly linked to the sediment surface. Microsensor profiles of O_2 and NO_x^- were furthermore used to obtain a detailed picture of the distribution of oxic and suboxic zones as well as zones of NO_x^- production and consumption. As O_2 exposure and NO_x^- availability would be key factors in determining the distribution of anammox bacteria in the sediment, these detailed observations could explain some of the trends observed in anammox activity.

The first phase of this study took place in June 2001. During this phase we quantified anammox and denitrification rates in intact cores and in prepared cores from which fauna had been removed. The second phase took place in April, June and September 2003. In this

phase we revisited the sites to further validate the existence of the anammox process, and we furthermore analyzed occurrence of anammox bacteria using the molecular techniques described above.

MATERIALS AND METHODS

Study sites. The study was carried out in Randers Fjord and Norsminde Fjord, Denmark. Randers Fjord is a shallow eutrophic estuary, 27 km long. The study site, Mellerup, is situated 12 km from the mouth of the estuary and at this site salinity ranges from 3 to 15 psu. Water-column concentrations of $\text{NO}_3^- + \text{NO}_2^-$ (NO_x^-) range from 15 to 300 μM over the year and N_2 production from 2 to 6 $\text{mmol m}^{-2} \text{d}^{-1}$ (County of Aarhus 1999). The average water depth at the station is approximately 1 m. The sediment consists of very fine sand with a moderate content of organic C (4% ignition loss) (see Table 1). Benthic microalgae (mainly diatoms) were present at the sediment surface. Norsminde Fjord is a shallow eutrophic estuary, 5 km long. The study site Kysing is situated near the outlet, and at this site salinity ranges from 3 to 23.5 psu while NO_x^- concentrations range from 430 μM to below detection limit over the year, the minimum being found in the summer months (County of Aarhus 1994). N_2 production ranges from 0.2 to 1.6 $\text{mmol m}^{-2} \text{d}^{-1}$ (Nielsen et al. 1995). The average water depth at the station is approximately 0.5 m. The sediment consists of medium sand with a moderate content of organic C (4% ignition loss; see Table 1). Benthic microalgae (mainly diatoms) are present and periodically form dense mats.

Sediment sampling and handling. Sediment was sampled by hand in Plexiglas tubes at both of the sites. Cores for measurement of total N_2 production, NO_x^- and O_2 exchange rates anammox rates, and determination of sediment characteristics were collected in 300 mm tubes (inner diameter, i.d., 5.5 mm). Cores used for microsensor measurements of O_2 and NO_x^- porewater profiles were collected in 100 mm tubes (i.d. 55 mm).

Another batch of surface sediment was collected in Randers Fjord, sieved through a 1 mm mesh screen, and transferred to a plastic container. Sediment cores were then sampled from the container by core tubes of the type used for *in situ* sampling.

All cores were processed on return to the laboratory. The lengths of the sediment cores used for measurement of total N_2 production and exchange rates of dissolved inorganic nitrogen (DIN) and O_2 were adjusted to 10 cm, and magnetic stir bars were positioned about 5 cm above the sediment surface. The cores were then placed in a reservoir containing site water held at *in situ* temperature. An external magnetic rotor (ca. 50 rpm)

ensured stirring of the water inside the tubes. Cores for microsensor measurements were pushed upward in the Plexiglas cylinder until the surface of the sediment was flush with the cylinder edge, and then placed in an aerated reservoir. Measurements of process rates were initiated within 12 h. Cores containing sieved sediment were pre-incubated as described above in darkness for 1 wk in aerated seawater held at 17°C to allow microbial processes in the sediment to stabilize.

Anammox and denitrification in slurry incubations. In the first phase of the study we used a simplified version of the technique devised by Thamdrup & Dalsgaard (2002), which allowed us to address only the presence or absence of the anammox process and the contribution of anammox to benthic N_2 production. Slurries were prepared by transferring approximately 1 ml of homogenized sediment from the upper 0.5 cm of the intact Randers Fjord sediment, the sieved Randers Fjord sediment and the intact Norsminde Fjord sediment, to 12 ml gas-tight (Laughlin & Stevens 2003) glass vials (Exetainer, Labco). These vials were placed in an N_2 atmosphere inside a glove bag. The headspace of each vial was then purged with N_2 and capped. The samples were left to stand for 4 h to eliminate the background concentration of NO_x^- in the sediments. Test experiments showed that NO_x^- was absent after 2 h pre-incubation. N_2 -purged artificial seawater (Grasshoff et al. 1983) containing either 100 μM $^{15}\text{NO}_3^-$ (^{15}N at. %: 97.5), 100 μM $^{15}\text{NH}_4^+$ (^{15}N at. %: 99) or 100 μM $^{15}\text{NH}_4^+$ plus 100 μM $^{14}\text{NO}_3^-$ was then added to the vials ($n = 4$ for each combination). The vials were transferred to a gas-tight bag purged with N_2 and placed on a shaker tray. After 24 h, 200 μl of a 7 M ZnCl_2 solution were added to the slurries to stop bacterial activity.

In the second phase of the study, we used a slurry technique that allowed us to quantify volume-specific anammox and denitrification rates. The slurries were prepared as described above with the following modifications: Sediment of known weight and density was transferred to the glass vials together with N_2 -purged site water. The slurries were then pre-incubated for 18 h to remove NO_x^- in sediment and incubation media through denitrification and anammox. Control measurements of O_2 and NO_x^- confirmed that both O_2 and NO_x^- were depleted after this period. Subsequently, 100 μl of N_2 -purged stock solution of each isotopic mixture, i.e. (1) $^{15}\text{NO}_3^-$ (^{15}N at. %: 99), (2) $^{15}\text{NH}_4^+$ (^{15}N at. %: 99.6) and (3) $^{15}\text{NH}_4^+ + ^{14}\text{NO}_3^-$ was added with a Hamilton syringe resulting in a concentration of about 100 μM N (200 μM N in the $^{15}\text{NH}_4^+$ and $^{14}\text{NO}_3^-$ combination). Slurries based on $^{15}\text{NO}_2^-$ additions were prepared in a similar manner. Incubations of the slurries were stopped at 1 h intervals by adding 200 μl of a 7 M ZnCl_2 solution. The abundance

of $^{15}\text{N}_2$ -labeled gas ($^{29}\text{N}_2$ and $^{30}\text{N}_2$) in the samples was measured by combined gas chromatography/mass spectrometry (RoboPrep-G+ in line with Tracermass, Europa Scientific) as described by Risgaard-Petersen & Rysgaard (1995).

Production of $^{15}\text{N}_2$ gas in $^{15}\text{NH}_4^+$ + $^{14}\text{NO}_3^-$ -amended samples and absence of $^{15}\text{N}_2$ production from samples incubated only with $^{15}\text{NH}_4^+$ was interpreted as evidence of anammox activity (Thamdrup & Dalsgaard 2002). Anammox, denitrification and the contribution of anammox to N_2 production were calculated from the production of $^{29}\text{N}_2$ and $^{30}\text{N}_2$ in the samples amended with $^{15}\text{NO}_3^-$ or $^{15}\text{NO}_2^-$ using the expressions of Thamdrup & Dalsgaard (2002).

Total N_2 production rates and nutrient fluxes. For both the Randers Fjord and the Norsminde Fjord sediment, total N_2 production rates and O_2 and NO_x^- exchange rates were estimated in light and in darkness ($n = 5$ for each treatment). Light was provided by 400 W greenhouse lamps (HPI-T+ mercury, Philips). Irradiance at the sediment surface (photon flux density) was $200 \mu\text{mol m}^{-2} \text{s}^{-1}$. Process rates in the sieved Randers Fjord sediment were only determined in the dark ($n = 5$). N_2 production was measured with $^{15}\text{NO}_3^-$ as described by Risgaard-Petersen & Rysgaard (1995) and Dalsgaard et al. (2000). Exchange rates of O_2 and NO_x^- were likewise measured as described by Dalsgaard et al. (2000). Incubations were performed in 2 sessions: fluxes were measured first, and after an equilibrium period of 20 h the N_2 production measurements were performed. All measurements were initiated 4 h after a change in light regime. Abundance of $^{15}\text{N}_2$ ($^{29}\text{N}_2$ and $^{30}\text{N}_2$) gas in the $^{15}\text{NO}_3^-$ -amended cores was measured by combined gas chromatography/mass spectrometry as described above. Concentrations of NO_3^- and NO_2^- were determined by the vanadium chloride reduction method (Braman & Hendrix 1989) on an NO_x^- analyzer (Model 42c, Thermo Environmental Instruments). O_2 was measured by Winkler titration (Grasshoff et al. 1983).

Total ^{14}N - N_2 production rates were calculated from the production rates of $^{29}\text{N}_2$ and $^{30}\text{N}_2$ in the $^{15}\text{NO}_3^-$ -amended cores using the isotope pairing technique (IPT) (Nielsen 1992). This technique may, however, overestimate benthic ^{14}N - N_2 production, as the presence of anammox results in violation of central assumptions on which the IPT is based, i.e. independence between added $^{15}\text{NO}_3^-$ and ^{14}N - N_2 production and binomial distribution of produced $^{28}\text{N}_2$, $^{29}\text{N}_2$ and $^{30}\text{N}_2$ (Risgaard-Petersen et al. 2003). Therefore, we also used the procedure proposed by Risgaard-Petersen et al. (2003) for estimation of N_2 production in sediments where denitrification and anammox coexist. For a thorough discussion of the 2 calculation procedures see Risgaard-Petersen et al. (2003).

O_2 and NO_x^- porewater profiles. Diffusion-controlled N_2 production at the sediment surface and diffusive O_2 and NO_x^- uptake were estimated from porewater profiles of O_2 and NO_x^- in the Randers Fjord sediment, the Norsminde Fjord sediment and the sieved Randers Fjord sediment ($n = 5$). Profiles in sediment cores from Randers Fjord and Norsminde Fjord were measured in darkness and during illumination (irradiance: $200 \mu\text{mol photons m}^{-2} \text{s}^{-1}$). Light was provided by a halogen lamp. Profile measurements in cores of sieved sediment were performed only in darkness. A Clark-type O_2 sensor (Revsbech 1989), and a nitrate plus nitrite (NO_x^-) biosensor (Larsen et al. 1997) were used to measure concentration profiles of O_2 and NO_x^- , respectively. The measurements were performed as described by Meyer et al. (2001). Profiles of O_2 and NO_x^- production rates were obtained by modeling the experimental data using the numerical method described by Berg et al. (1998). The sediment diffusion coefficient (D_s) used in these calculations was estimated from the free-solution diffusion coefficient of O_2 and NO_3^- (Li & Gregory 1974) and sediment porosity (Boudreau 1997). Diffusive O_2 and NO_x^- uptake was calculated as the difference between the depth-integrated production and consumption estimated from the profiles of O_2 and NO_x^- production. Diffusive N_2 production was estimated from the depth-integrated NO_x^- consumption rates in the suboxic sediment strata according to the following rationale: N_2 production (p_{14}) is the sum of anammox and denitrification, and while 2 N_2 -N atoms are produced for every NO_2^- being reduced through anammox, only 1 N_2 -N atom is produced for every NO_x^- being reduced through denitrification. Thus:

$$\begin{aligned} p_{14} &= 2 \cdot \text{NO}_2^-_{\text{anammox}} + \text{NO}_x^-_{\text{denitrification}} \\ &= 2 \cdot \text{NO}_2^-_{\text{anammox}} + (\text{NO}_x^-_{\text{red}} - \text{NO}_2^-_{\text{anammox}}) \quad (1) \\ &= \text{NO}_2^-_{\text{anammox}} + \text{NO}_x^-_{\text{red}} \end{aligned}$$

where $\text{NO}_2^-_{\text{anammox}}$ is the rate of NO_2^- reduction via the anammox process, $\text{NO}_x^-_{\text{denitrification}}$ is NO_x^- reduction via denitrification and $\text{NO}_x^-_{\text{red}}$ is the total rate of NO_x^- reduction in the O_2 -free sediment strata. The rate of N - N_2 production via anammox can be expressed as follows:

$$2 \cdot \text{NO}_2^-_{\text{anammox}} = ra \cdot p_{14} = ra \cdot (\text{NO}_2^-_{\text{anammox}} + \text{NO}_x^-_{\text{red}}) \quad (2)$$

where ra is the contribution of anammox to N_2 production.

The rate of NO_2^- reduction via anammox can then be expressed as follows:

$$\text{NO}_2^-_{\text{anammox}} = \frac{ra \cdot \text{NO}_x^-_{\text{red}}}{2 - ra} \quad (3)$$

and p_{14} is thus equivalent to:

$$p_{14} = \text{NO}_x^-_{\text{red}} - \frac{ra \cdot \text{NO}_x^-_{\text{red}}}{2 - ra} \quad (4)$$

As in the total N_2 production measurements, we used the contribution of anammox to N_2 production estimated through slurry incubations in the first phase of this study as a proxy for *ra*.

Phylogenetic inference and fluorescence *in situ* hybridization (FISH). Analyses for anammox bacteria were performed on sediment sampled from the upper 0.5 cm of the Randers Fjord sediment. DNA extraction, cloning, sequencing, phylogenetic inference and FISH experiments were performed as reported by Schmid et al. (2003). Probes used in this study were S*-Amx-0368-a-A-18 (detecting all anammox organisms), S*-BS-0820-a-A-22 (detecting *Scalindua sorokinii* and candidate *S. wagneri*) and S-P-Planc-0046-a-A-18 (detecting bacteria in the phylum *Planctomycetes*). (For further probe details see www.probeBase.net; Loy et al. 2003.)

Sediment characteristics. Grain size distribution was determined on 3 pooled sediment cores from the sites as described by Berg et al. (2001). Sediment permeability was estimated from the porosity and the mean diameter of sediment particles using the Carman-Kozeny equation (Boudreau 1997). Chlorophyll *a* was determined on sediment subsamples collected from the upper 0.5 cm of the sediment ($n = 3$) using the method of Lorenzen (1967). Organic carbon content and porosity (vol/vol) were likewise determined on sediment subsamples from the upper 0.5 cm ($n = 3$), and estimated from loss on ignition and the water contents of known volumes of sediment. The fauna density was determined in 3 cores from the site. The sediment in these cores was sieved through a 0.5 mm sieve and animals found were sorted into groups and counted.

RESULTS

Sediment characteristics

More than 75 % of the sediment particles in Randers Fjord were $<125 \mu\text{m}$, and according to the Udden-Wenworth scheme (Fütterer 2000), the sediment from Randers could thus be characterized as very fine sand (Table 1). The sediment from Norsminde was somewhat coarser, with 63 % $<500 \mu\text{m}$, and could be classified as medium sand. Sediment permeability was $4 \times 10^{-15} \text{ m}^{-2}$ and $3 \times 10^{-14} \text{ m}^{-2}$ in the Randers Fjord and Norsminde Fjord sediment, respectively, and according to Glud et al. (1996) both sediments can be perceived as being impermeable. There was no major difference in chlorophyll *a* or organic C content between the Randers Fjord and Norsminde Fjord sediments. Intact sediment cores from both Randers Fjord and Norsminde Fjord were densely populated with polychaetes and *Corophium* sp. The density of *Corophium* sp. was 6734 ± 2105 and 3978 ± 1654 individuals m^{-2} and the density of polychaetes was 2525 ± 281 and 1473 ± 210 individuals m^{-2} in Randers Fjord and Norsminde Fjord, respectively.

Anammox and denitrification assessed through slurry incubations

In the case of slurries amended with $^{15}\text{NH}_4^+$ only, significant accumulation of $^{15}\text{N}_2$ -labeled gas was not seen in either the Randers Fjord or the Norsminde Fjord sediment (Table 2). When both $^{15}\text{NH}_4^+$ and $^{14}\text{NO}_3^-$ were present, $^{29}\text{N}_2$ accumulated in both the intact and

Table 1. Sediment characteristics of study sites in 2 shallow-water estuaries in Denmark. Values are means (SE), $n = 3$; nm: not measured; DW: dry weight

Site	Grain size distribution (%)						Porosity (vol/vol)	Organic C (% of DW)	Chl <i>a</i> (g m^{-2})
	$<63 \mu\text{m}$	$>63 \mu\text{m}$	$>125 \mu\text{m}$	$>250 \mu\text{m}$	$>500 \mu\text{m}$	$>1000 \mu\text{m}$			
Randers Fjord (intact)	19.1	56.1	14.3	1.1	0.5	8.9	0.72	4.2 (0.2)	5.08 (0.2)
Randers Fjord (sieved)	26.1	48.6	24.0	1.0	0.3	0.0	0.72	nm	nm
Norsminde Fjord	5.3	13.4	21.5	35.1	12.5	12.2	0.65	4.4(0.03)	4.2 (0.1)

Table 2. Concentrations (μM) of accumulated $^{15}\text{N}_2$ in slurries treated with either $^{15}\text{NO}_3^-$, $^{15}\text{NH}_4^+$ or $^{15}\text{NH}_4^+ + ^{14}\text{NO}_3^-$. The contribution of anammox to total N_2 production (*ra*) in the $^{15}\text{NO}_3^-$ -amended slurries is presented as % N_2 produced via anammox. Values are means (SE), $n = 4$. Data are from Phase 1 of study, June 2001

Site	$^{15}\text{NO}_3^-$		$^{15}\text{NH}_4^+$		$^{15}\text{NH}_4^+ + ^{14}\text{NO}_3^-$		<i>ra</i> (%)
	$^{29}\text{N}_2$	$^{30}\text{N}_2$	$^{29}\text{N}_2$	$^{30}\text{N}_2$	$^{29}\text{N}_2$	$^{30}\text{N}_2$	
Randers Fjord	5.39 (1.43)	41.190 (4.70)	0.05 (0.017)	0.00 (0.001)	2.48 (0.305)	0.03 (0.007)	6.2 (1)
Randers Fjord (sieved)	8.27 (0.629)	44.70 (0.899)	0.05 (0.013)	0.01 (0.005)	3.41 (0.722)	0.04 (0.008)	10.6 (1.2)
Norsminde Fjord	1.49 (0.15)	26.31 (1.49)	0.01 (0.004)	0.00 (0.001)	0.02 (0.003)	0.00 (0.001)	-0.5 (0.6)

the sieved sediment from Randers Fjord. However, there was no accumulation of $^{30}\text{N}_2$. This pattern was reproducible, as shown in the time-series experiments performed in the second phase of the study (Fig. 1). In the sediment from Norsminde Fjord, there was no accumulation of $^{29}\text{N}_2$ or $^{30}\text{N}_2$ in the $^{15}\text{NH}_4^+ + ^{14}\text{NO}_3^-$ -amended slurries (Table 2).

The contribution of anammox to total N_2 production was 6.2 and 10.6% in the intact and sieved Randers Fjord sediment, respectively, in the Phase 1 experiments. The contribution of anammox to N_2 production in Norsminde was below detection limit.

Volume-specific anammox activity estimated from slurry incubations of sediment from Randers Fjord was highest in April (ANOVA, $p = 0.01$), whereas no significant difference was seen between rates obtained in

Table 3. Volume-specific rates of anammox and denitrification estimated from slurry incubations with either $^{15}\text{NO}_3^-$ or $^{15}\text{NO}_2^-$, contribution of anammox to N_2 production (ra), and *in situ* temperature plus bottom-water concentrations of $\text{NO}_3^- + \text{NO}_2^-$ (NO_x^-). Values are means (SE), $n = 4$. Data are from Phase 2 of study (2003)

Month	Anammox ($\text{nmol N cm}^{-3} \text{ h}^{-1}$)	Denitrification ($\text{nmol N cm}^{-3} \text{ h}^{-1}$)	ra (%)	NO_x^- (μM)	T ($^\circ\text{C}$)
April	11 (0.2)	31 (1)	26.4	120	11
June				15	16
NO_3^-	5.2 (1.3)	137 (8)	3.7		
NO_2^-	5.9 (0.3)	131 (5.7)	4.3		
August				15	
NO_3^-	3.8 (0.6)	72 (8)	5.0		20
NO_2^-	4.1 (1.5)	69 (3.8)	5.5		

June and August (Table 3, ANOVA, $p = 0.6$). Volume-specific denitrification rates were highest in June and lowest in April (Table 3, ANOVA, $p = 0.01$). Accordingly, the contribution of anammox to N_2 production varied over the period investigated: anammox contributed approximately 26% in April, whereas the contribution from anammox in August was only 5%. There was no significant difference between rates obtained with NO_3^- or NO_2^- as substrate in either denitrification or anammox activities (ANOVA, $p > 0.5$; Table 3).

Phylogenetic analysis and detection of anammox bacteria

Phylogenetic analysis showed that the 16S rRNA sequence amplified from DNA extracted from the Randers Fjord sediment was affiliated with the anammox organism candidatus *Scalindua sorokinii* (Fig. 2). The overall sequence similarity to candidatus *S. sorokinii* was about 99%. FISH demonstrated that the organisms affiliated with candidatus *S. sorokinii*/candidatus *S. wagneri* were the only detectable anammox bacteria in the sample (Fig. 3). No other *Planctomycetes* were detected.

Porewater profiles

Average O_2 and NO_x^- concentration profiles and depth-specific rates of O_2 and NO_x^- production calculated from porewater profiles are shown in Figs. 4 & 5. In sediment cores from Randers Fjord, zones of NO_x^- production and consumption zones were separated at the oxic/suboxic interface. A very high production of oxygen by benthic microalgae in the top 0.3 mm of the sediment from Randers Fjord caused O_2 and NO_x^- penetration depths to increase by more than 1 mm during illumination. Despite these changes there was no sig-




Fig. 1. Examples of concentrations of $^{29}\text{N}_2$ and $^{30}\text{N}_2$ in samples incubated with either $^{15}\text{NO}_3^-$, $^{15}\text{NH}_4^+$ or $^{14}\text{NO}_3^-$ and $^{15}\text{NH}_4^+$ during Phase 2 of study (April, June, September 2003)




Fig. 2. Phylogenetic tree reflecting relationship of 16S rRNA sequence type amplified from Randers Fjord, other anammox organisms, other members of the order Planctomycetes and other reference organisms. Phylogenetic analyses were performed with maximum likelihood, neighbor-joining and maximum parsimony methods with 50% conservation filters for Bacteria and Planctomycetes. Triangles indicate phylogenetic groups. *S.*, *K.*, *B.* represent genus names *Scalindua*, *Kuenenia* and *Brocadia*, respectively

nificant difference in net NO_x^- consumption/production rates in light and darkness (Student's *t*-test, $p > 0.2$). In the sieved sediment, the distribution of NO_x^- production and consumption zones relative to the oxic/suboxic interface was similar to that found in natural cores from Randers Fjord. However, the rates of the NO_x^- transformation processes were higher (Fig. 5).




Fig. 3. Fluorescence *in situ* hybridization of Randers Fjord sediment; arrow indicates candidatus *Scalindua sorokinii*-related cells, which appear whitish yellow because of triple hybridization of S⁻-Amx-0368-a-A-18 (labeled with Cy3, red), S⁻-BS-0820-a-A-22 (labeled with Fluos green) and S-P-Planc-0046-a-A-18 (labeled with Cy5)




Fig. 4. Microprofiles of O_2 (○) and NO_x^- (●). Circles: means; error bars: SEs; lines: concentration profiles estimated from numerical simulations

Porewater profiles in dark-incubated cores from Norsminde Fjord showed distinct zones of NO_x^- production and consumption. Net NO_x^- production was lowest in light (Student's *t*-test, $p = 0.03$) whereas no significant difference was seen in net NO_x^- consumption rates measured in darkness and in light (Student's *t*-test, $p = 0.07$). In contrast to the other sediments, oxygen and NO_x^- were depleted at approximately the same depth (0.3 mm), and NO_x^- was consumed in the oxic zone well above the oxic/suboxic interface during both illumination and darkness (Fig. 4). A control experiment showed that N_2 was produced only in the absence of O_2 (data not shown), and NO_x^- consumption in the oxic zone was therefore not due to aerobic denitrification but most probably to microphytobenthic N-assimilation.

Total and diffusive N_2 production rates

Area-based rates of anammox and denitrification estimated from the porewater profiles of NO_x^- , the IPT and the revised IPT are shown in Table 4. N_2 produc-




Fig. 5. Vertical distribution of net consumption and production of O_2 (white area) and NO_x^- (shaded area) estimated from mean profiles

tion rates estimated with the IPT were not statistically different from the rates estimated with the revised technique. Likewise, no significant difference was seen between rates measured during illumination and

during darkness (Student's t -test, $p > 0.1$). On a diurnal scale, N_2 production rates estimated with ^{15}N isotopes in Norsminde Fjord were similar to the activity measured in Randers Fjord (Student's t -test, $p = 0.48$). N_2 production estimated from porewater profiles in the intact sediment from Randers Fjord was less than 20% of the activity measured with ^{15}N isotopes. The discrepancy was even more pronounced in the sediment from Norsminde Fjord, where porewater profiles showed that NO_x^- did not penetrate to the suboxic layers of the sediment, indicating absence of denitrification activity, although the IPT revealed high N_2 production rates. N_2 production rates estimated from the porewater profiles in the sieved and defaunated Randers Fjord sediment were not significantly different (Student's t -test, $p = 0.47\%$), however, from the estimates based on the revised IPT applied to the same sediment.

Exchange of O_2 and NO_x^- between sediment and water column

Exchange rates of O_2 and NO_x^- estimated from porewater profiles and from flux-core measurements are shown in Table 5. Whole-core measurements showed a consistent net uptake of O_2 in all cores at all times, whereas microsensor profiles indicated net efflux of oxygen during illumination. In darkness, the total O_2 uptake was about 7 (Randers) and about 20 (Norsminde) times higher than the diffusive uptake estimated from porewater profiles. A similar discrepancy was observed for NO_x^- fluxes, which also showed a consistent net uptake of NO_x^- in all undisturbed sediments when measured from whole cores. Porewater profiles, on the other hand, indicated a small efflux of NO_x^- from sediment cores from Randers Fjord in both light and darkness. In the sediment from Norsminde Fjord, porewater profiles suggested efflux of NO_x^-

Table 4. N_2 production rates ($\mu mol N m^{-2}$) estimated with NO_x^- microsensors (diffusive N_2 production) and from whole-core incubations with $^{15}NO_3^-$ (total N_2 production). Values are means (SE), $n = 5$. IPT: isotope pairing technique

Site	Diffusive N_2 production			Total N_2 production—IPT			Total N_2 production—revised IPT		
	Anaerobic NO_3^- reduction	Anammox	Denitrification	N_2 prod.	Anammox	Denitrification	N_2 prod.	Anammox	Denitrification
Randers Fjord									
Light	43.2 (1.8)	4.7 (0.2)	39.8 (1.7)	238 (43.0)	14.8 (2.7)	223 (40)	233 (42)	14.0 (2.5)	219 (40)
Dark	50.1 (3.9)	5.5 (0.4)	46.2 (3.6)	365 (49)	23 (3.1)	342 (46)	356 (48)	21 (2.9)	335 (46)
Randers Fjord (sieved)	77.8 (4.2)	8.7 (0.5)	73.4 (4.0)	97 (3.1)	10 (0.3)	87 (2.7)	91 (2.5)	9.6 (0.3)	81 (2.3)
Norsminde Fjord									
Light	0.0	0.0	0.0	236 (32)	0.0	236 (32)	236 (32)	0.0	236 (32)
Dark	0.0	0.0	0.0	191 (38)	0.0	191 (38)	191 (38)	0.0	191 (38)

Table 5. Oxygen and NO_x^- exchange rates ($\mu\text{mol N m}^{-2} \text{ h}^{-1}$) estimated with microsensors (Diffusive) and from whole-core incubations (Total). Values are means (SE), $n = 5$

Site	O_2		NO_x^-	
	Diffusive	Total	Diffusive	Total
Randers Fjord				
Light	1932 (112)	-3021 (623)	1.6 (1.4)	-47 (15)
Dark	-950 (89)	-7193 (399)	4.5 (2.3)	-68.0 (23.0)
Randers Fjord (sieved)	-788 (66)	-727 (39)	54 (3.8)	44 (4)
Norsminde Fjord				
Light	510 (113)	-2192 (2023)	-12 (1.9)	-120 (36)
Dark	-515 (38)	-11123 (497)	20 (4.7)	-51 (23)



in the dark and uptake in light at a rate more than 10 times lower than the uptake measured in whole cores. In contrast to the natural sediments, neither O_2 nor NO_x^- fluxes measured in whole cores prepared from sieved sediment were significantly different from equivalent parameters estimated from porewater profiles (Student's t -test, $p = 0.2$; Table 5).

DISCUSSION

Presence and absence of anammox in estuarine sediment

In our search for alternative N_2 -producing processes, we found evidence of anaerobic NH_4^+ oxidation in the presence of NO_3^- in slurries prepared with sediment from Randers Fjord (Table 1). The lack of ^{15}N - N_2 accumulation in samples from both fjords incubated with only $^{15}\text{NH}_4^+$ excludes the possibility of coupled nitrification–denitrification, which might have occurred if O_2 had been introduced into the slurries by mistake at the beginning of the experiment. Anaerobic oxidation of NH_4^+ to NO_x^- or N_2 with, for instance, MnO_2 (Luther et al. 1997, Hulth et al. 1999) can also be excluded. If this process was significant, NO_x^- produced through oxidation of NH_4^+ would undergo denitrification and result in accumulation of ^{15}N - N_2 gas in the slurries amended with $^{15}\text{NH}_4^+$ only. As mentioned above, no such accumulation was observed.

Accumulation of ^{15}N - N_2 in the $^{15}\text{NH}_4^+ + ^{14}\text{NO}_3^-$ -amended Randers Fjord sediments is the result of 1 of the following 2 reactions:



The lack of a 3:1 ratio between $^{29}\text{N}_2$ and $^{30}\text{N}_2$ production in the $^{15}\text{NH}_4^+ + ^{14}\text{NO}_3^-$ -amended slurries (Table 2,

Fig. 3) excludes Reaction 2 and points to Reaction 1, which is identical to the anammox reaction according to Strous et al. (1999). Thus, our data are strong proof of the presence of the anammox process in sediment from Randers Fjord. Furthermore, our data show that anaerobic NO_2^- generation rates were sufficiently high in this sediment to provide the bacteria with enough substrate for the reaction, as indicated by the fact that anammox rates measured with $^{15}\text{NO}_3^-$ and with $^{15}\text{NO}_2^-$ were similar (Table 3).

Autotrophic bacteria of the phylum *Planctomycetes* have been shown to be responsible for the anammox process (Strous et al. 1999, Schmid et al. 2000, 2003), and recently occurrence of anammox in the anoxic water column of the Black Sea was associated with the presence of a newly discovered *Planctomycetes* species candidatus *Scalindua sorokinii* (Kuypers et al. 2003). Phylogenetic analysis of 16S rRNA gene sequences amplified from DNA extracted from the Randers Fjord sediment showed a close relationship to candidatus *S. sorokinii* (Fig. 2). FISH analysis of sediment samples from Randers Fjord also showed the presence of small clusters of cells belonging to candidatus *S. sorokinii* (Fig. 3), confirming further that the bacteria belonging to the phylum *Planctomycetes* found in the Randers Fjord sediment were all affiliated with candidatus *S. sorokinii*. Since the FISH and the ^{15}N isotope data are consistent, the present study confirms the link between presence of bacteria affiliated with candidatus *S. sorokinii* and the anammox reaction in marine environments observed by Kuypers et al. (2003) in the Black Sea. It is remarkable that so closely related species are present in so distantly related environments as the anoxic water column of the Black Sea and the Randers Fjord sediment. However, this finding is in line with the classical view of Beijerinck (Brock 1961) that any bacterial species can occur anywhere, provided its environmental requirements are met, due to the enormous microbial population sizes that result in high dispersal probability and low probability of local extinction (Finlay & Clarke 1999, Fenchel 2003).

Our failure to demonstrate the existence of anammox in Norsminde Fjord (Table 2) indicates that the anammox process is not ubiquitous. We propose that the existence of anammox in the surface sediment of Randers Fjord and the absence of the process from Norsminde Fjord may be linked to differences in the availability of NO_x^- in the suboxic zone of the sediment. At the study site in Randers Fjord the water-column NO_x^- concentration is always above $15 \mu\text{M}$ (County of Aarhus 1999, Nielsen et al. 2001), and nitrification rates are high (Nielsen et al. 2001). Porewater profiles of NO_x^- measured in the surface sediment furthermore showed that NO_x^- penetrates into the suboxic zone, where it is consumed (Fig. 4). In contrast,

water-column NO_x^- is depleted in Norsminde Fjord during the summer months (Nielsen et al. 1995), and assimilation by benthic microalgae can prevent penetration of NO_x^- into the suboxic zone of the sediment (Fig. 4), probably because the supply of NO_x^- from the water column is low (see for instance Meyer et al. 2001). According to current knowledge, anammox bacteria are slow-growing obligate anaerobes and base their energy production solely on NO_2^- and NH_4^+ conversion (Strous et al. 1999). In contrast, most denitrifying bacteria are organotrophic organisms capable of using O_2 as an electron acceptor (Zumft 1992). Thus, denitrifying bacteria seem better adapted to the fluctuating availability of O_2 and NO_x^- imposed by microalgae. Experimental studies of anammox and denitrification in sediments with and without microphyto-benthic activity confirm this hypothesis (R. L. Meyer & N. Risgaard-Petersen unpubl.). Nitrite porewater profiles furthermore indicate that net NO_2^- production takes place mainly in the suboxic zone of the sediment as a result of NO_3^- reduction (Stief et al. 2002). This may indicate that anammox bacteria are dependent on release of NO_2^- from anaerobic NO_3^- reducers such as denitrifying bacteria. If NO_3^- availability is low, as in Norsminde Fjord during the summer months, the loss of NO_2^- from denitrifiers would probably be insignificant and insufficient to support a population of anammox bacteria.

Importance of anammox as an N_2 source in Randers Fjord

In the surface sediment of Randers Fjord we observed the highest volume-specific anammox rates in April and the lowest in September 2003, suggesting that the activity of anammox bacteria decreases during the course of the summer period. Denitrification did not follow a similar trend, and as a consequence the contribution of anammox to N_2 production decreased from 26 to 5%. This indicates seasonal variations in the abundance of the respective bacterial groups and that denitrifying bacteria and anammox bacteria are controlled in different ways. The observed indications of seasonal fluctuations in the contribution of anammox to N_2 production probably reflect differences in the availability of organic carbon and NO_x^- over the season. In the summer months, O_2 consumption rates in Randers Fjord are generally higher than during spring and winter (Nielsen et al. 2001). This may indicate that the availability of organic carbon is higher and that conditions are more favorable for the organotrophic denitrifying bacteria than for the lithotrophic anammox bacteria in the sediments during the summer period.

The volume-specific anammox activity measured in Randers Fjord was close to the activity measured in the Skagerrak ($5 \text{ nmol N cm}^{-3} \text{ h}^{-1}$) at Stn S9 at the temperature applied in the present study (16°C) (see Dalsgaard & Thamdrup 2002) and within the range reported by Trimmer et al. (2003) for the Thames Estuary (0.2 to $10 \text{ nmol N cm}^{-3} \text{ h}^{-1}$). This may be an indication that population densities of anammox bacteria in Randers Fjord, Skagerrak and Thames sediments are comparable. Despite these similarities, anammox in Randers Fjord and the Thames is relatively less important as a source of N_2 production ($ra = 5$ to 25% in Randers Fjord [Tables 2 & 3] and 1 to 8% in the Thames Estuary) than in the Skagerrak sediment, where the process accounts for ca. 70% of N_2 production (Thamdrup & Dalsgaard 2002). Volume-specific denitrification rates in Randers Fjord exceeded the rates in the Skagerrak by a factor of 15 in April and a factor of 30 in August. The Randers Fjord sediment thus seemed to be a more favorable habitat for denitrifying bacteria than the Skagerrak sediment, which explains the difference in contribution of anammox to N_2 production. This is not surprising, as it is well known that benthic carbon mineralization rates decrease with increasing water depth (Canfield 1993) due to a decrease in easily degradable carbon. The Skagerrak sediment (water depth 695 m) would thus be expected to have a much lower heterotrophic microbial activity than the shallow Randers Fjord (water depth ca. 1 m), which is furthermore heavily eutrophic. The difference in overall heterotrophic activity between Randers Fjord and the Skagerrak is reflected both in the O_2 penetration depth and in the sediment O_2 uptake rate. The O_2 penetration depth in the Skagerrak sediment is 1.5 cm and the sediment O_2 consumption rate $4 \text{ mmol m}^{-2} \text{ d}^{-1}$ (Rysgaard et al. 2001), which is 10 times higher and 150 times lower, respectively, than the equivalent parameters measured in darkness in Randers Fjord in the present study.

Anammox and denitrification in cores with natural substrate gradients

Sediment anammox rates presented in the literature (Dalsgaard & Thamdrup 2002, Thamdrup & Dalsgaard 2002, Trimmer et al. 2003) are at best potential rates, being estimated with methods that disrupt the natural substrate gradients in the sediment. In the present study, we applied ^{15}N and microsensor techniques to estimate anammox activity in sediments with intact stratification. The applied ^{15}N calculation procedures—the IPT (Nielsen 1992) and the revised IPT (Risgaard-Petersen et al. 2003)—yielded similar esti-

mates of anammox and denitrification rates (Table 4) because the contribution of anammox to N_2 production was too low to seriously affect the assumptions underlying the IPT (Risgaard-Petersen et al. 2003). Using the revised IPT on $^{15}N_2$ raw data from Rysgaard et al. (2001), Risgaard-Petersen et al. (2003) estimated an anammox rate of $4 \mu\text{mol N m}^{-2} \text{ h}^{-1}$ in the Skagerrak sediment studied by Thamdrup & Dalsgaard (2002). This is between 18 and 29% of the rates estimated in Randers Fjord with ^{15}N isotopes (Table 4). The anammox process is thus quantitatively more important in the Randers Fjord sediment than in the Skagerrak.

We observed a large difference between the estimates of denitrification and anammox obtained with microsensors and those obtained with ^{15}N isotopes (Table 4). N_2 production rates estimated from interpretation of porewater profiles were 22 and 0% of the N_2 production rates obtained with ^{15}N isotopes in the Randers Fjord and the Norsminde Fjord sediments, respectively. A similar difference was observed when total and diffusive O_2 and NO_x^- uptake rates were compared (Table 5). This discrepancy was probably due to the presence of fauna, because the applied ^{15}N technique captures total N_2 production, while the microsensor technique only captures the diffusion-controlled N_2 production at the sediment surface. This hypothesis is supported by the consistent agreement found between the methods regarding all parameters measured in the sieved and defaunated sediment from Randers Fjord (Tables 4 & 5). Furthermore, the hypothesis is fully in line with conclusions from previous studies comparing diffusive O_2 or N_2O fluxes estimated from microsensor profiles and total exchange rates in impermeable sediments (Andersen & Helder 1987, Binnerup et al. 1992, Glud et al. 1994, Berg et al. 2001). The mechanisms responsible for this fauna-mediated stimulation of biogeochemical processes include enhanced porewater transport caused by biodiffusion and irrigation as well as a several-fold increase in the area of the oxic/suboxic interface in the presence of polychaete burrows (Kristensen 2000 & references therein). Several experimental studies addressing the impact of benthic animals on the N-cycle processes have shown that via these mechanisms benthic fauna may significantly stimulate benthic N_2 production (e.g. Binnerup et al. 1992, Pelegri et al. 1994, Svensson et al. 2001, Newell et al. 2002).

Acknowledgements. We thank Anna Haxen, Kitte Gerlich, Egon Frandsen and Marlene Jessen for assistance in the laboratory. This study was in part supported by grants from the Danish Natural Science Research Councils (Contract No. 51-00-0458 and Contract No. 51-00-0320) and in part by the ICON project under the European Union 5th Framework Programme, Project No. EVK1-CT2000-00054.

LITERATURE CITED

- Aller RC (1982) The effects of macrobenthos on chemical properties of marine sediment and overlying water. In: McCall PL, Tevesz PJS (eds) *Animal-sediment relations*. Plenum, New York, p 53–102
- Aller RC, Aller JY (1998) The effect of biogenic irrigation intensity and solute exchange on diagenetic reaction rates in marine sediments. *J Mar Res* 56:905–936
- Andersen FØ, Helder W (1987) Comparison of oxygen microgradients, oxygen flux rates and electron transport system activity in coastal marine sediments. *Mar Ecol Prog Ser* 37: 259–264
- Berg P, Risgaard-Petersen N, Rysgaard S (1998) Interpretation of measured concentration profiles in sediment pore water. *Limnol Oceanogr* 43:1500–1510
- Berg P, Rysgaard S, Funch P, Sejr MK (2001) Effects of bioturbation on solutes and solids in marine sediments. *Aquat Microb Ecol* 26:81–94
- Binnerup SJ, Jensen K, Revsbech NP, Jensen MH, Sørensen J (1992) Denitrification, dissimilatory reduction of nitrate to ammonia and nitrification in a bioturbated estuarine sediment as measured with ^{15}N and microsensor techniques. *Appl Environ Microbiol* 58:303–313
- Boudreau BP (1997) *Diagenetic models and their implementation*. Springer-Verlag, Berlin
- Braman RS, Hendrix SA (1989) Nanogram nitrite and nitrate determination in environmental and biological materials by vanadium (III) reduction with chemiluminescence detection. *Analyt Chem* 61:2715–2718
- Brock TD (1961) *Milestones in microbiology*. Prentice-Hall, Englewood Cliffs, NJ
- Canfield D (1993) Organic matter oxidation in marine sediments. In: Wollast R, Mackenzie FT, Chou L (eds) *Interactions of C, N, P and S biogeochemical cycles*. Springer-Verlag, Berlin, p 333–363
- County of Aarhus (1994) *Norsminde Fjord 1992*. Natur og Miljø Århus Amt, Århus
- County of Aarhus (1999) *Randers Fjord status 1997*. Natur og Miljø Århus Amt, Århus
- Dalsgaard T, Thamdrup B (2002) Factors controlling anaerobic ammonium oxidation with nitrite in marine sediments. *Appl Environ Microbiol* 68:3802–3808
- Dalsgaard T, Nielsen LP, Brotas V, Viaroli P and 10 others (2000) *Protocol handbook for NICE—nitrogen cycling in estuaries: a project under the EU research programme: marine science and technology (MAST III)*. National Environmental Research Institute, Silkeborg
- Devol AH (1991) Direct measurements of nitrogen gas fluxes from continental shelf sediments. *Nature* 349:319–321
- Devol AH (2003) Nitrogen cycle—solution to a marine mystery. *Nature* 422:575–576
- Fenchel TB (2003) Biogeography for bacteria. *Science* 301: 925–926
- Finlay BJ, Clarke KL (1999) Ubiquitous dispersal of microbial species. *Nature* 400:828
- Fütterer DF (2000) The solid phase of marine sediments. In: Schulz HD, Zabel M (eds) *Marine geochemistry*. Springer-Verlag, Berlin, p 1–22
- Glud RN, Gundersen JK, Jørgensen BB, Revsbech NP, Schulz HD (1994) Diffusive and total oxygen uptake of deep-sea sediments in the eastern South Atlantic Ocean: in situ and laboratory measurements. *Deep-Sea Res* 41:1767–1788
- Glud RN, Forster S, Huettel M (1996) Influence of radial pressure gradients on solute exchange in stirred benthic chambers. *Mar Ecol Prog Ser* 141:303–311
- Grasshoff K, Erhardt M, Kremling K (1983) *Methods of sea-*

- water analysis. Verlag Chemie, Weinheim
- Hulth S, Aller RC, Gilbert F (1999) Coupled anoxic nitrification manganese reduction in marine sediments. *Geochim Cosmochim Acta* 63:49–66
- Jensen K, Revsbech NP, Nielsen LP (1993) Microscale distribution of nitrification activity in sediment determined with a shielded microsensor for nitrate. *Appl Environ Microbiol* 59:3287–3296
- Jensen K, Sloth NP, Risgaard-Petersen N, Rysgaard S, Revsbech NP (1994) Estimation of nitrification and denitrification from microprofiles of oxygen and nitrate in model sediment systems. *Appl Environ Microbiol* 60:2094–2100
- Kristensen E (2000) Organic matter diagenesis at the oxic/anoxic interface in coastal marine sediments, with emphasis on the role of burrowing animals. *Hydrobiologia* 426: 1–24
- Kuypers MMM, Sliekers AO, Lavik G, Schmid M and 5 others (2003) Anaerobic ammonium oxidation by anammox bacteria in the Black Sea. *Nature* 422:608–611
- Larsen LH, Kjær T, Revsbech NP (1997) A microscale NO₃-biosensor for environmental applications. *Analyt Chem* 69:3527–3531
- Laughlin RJ, Stevens RJ (2003) Changes in composition of nitrogen-15-labeled gases during storage in septum-capped vials. *Soil Sci Soc Am J* 67:540–543
- Li YH, Gregory S (1974) Diffusions of ions in sea water and in deep-sea sediments. *Geochim Cosmochim Acta* 38: 703–714
- Lorenzen CJ (1967) Determination of chlorophyll and pheopigments—spectrophotometric equations. *Limnol Oceanogr* 12:343–346
- Loy A, Horn M, Wagner M (2003) probeBase—an online resource for rRNA-targeted oligonucleotide probes. *Nucleic Acids Res* 31:514–516
- Luther GW, Sundby B, Lewis BL, Brendel PJ, Silverberg N (1997) Interactions of manganese with the nitrogen cycle: alternative pathways to dinitrogen. *Geochim Cosmochim Acta* 61:4043–4052
- Meyer RL, Kjær T, Revsbech NP (2001) Use of NO_x⁻ microsensors to estimate the activity of sediment nitrification and NO_x-consumption along an estuarine salinity, nitrate, and light gradient. *Aquat Microb Ecol* 26:181–193
- Mulder A, van de Graaf AA, Robinson LA, Kuenen JG (1995) Anaerobic ammonium oxidation in a denitrifying fluidized bed reactor. *FEMS Microbiol Ecol* 16:177–184
- Nielsen K, Nielsen LP, Rasmussen P (1995) Estuarine nitrogen retention independently estimated by the denitrification rate and mass balance methods: a study of Norsminde Fjord, Denmark. *Mar Ecol Prog Ser* 119:275–283
- Nielsen K, Risgaard-Petersen N, Somod B, Rysgaard S, Bergo T (2001) Nitrogen and phosphorus retention estimated independently by flux measurements and dynamic modelling in the estuary, Randers Fjord, Denmark. *Mar Ecol Prog Ser* 219:25–40
- Nielsen LP (1992) Denitrification in sediments determined from nitrogen isotope pairing. *FEMS Microbiol Ecol* 86: 357–362
- Newell RIE, Cornwell JC, Owens MS (2002) Influence of simulated bivalve biodeposition and microphytobenthos on sediment nitrogen dynamics: a laboratory study. *Limnol Oceanogr* 47:1367–1379
- Payne JW (1981) Denitrification. John Wiley & Sons, New York
- Pelegri SP, Nielsen LP, Blackburn TH (1994) Denitrification in estuarine sediment stimulated by irrigation activity of the amphipod *Corophium volutator*. *Mar Ecol Prog Ser* 105:285–290
- Revsbech NP (1989) An oxygen microelectrode with a guard cathode. *Limnol Oceanogr* 34:474–478
- Risgaard-Petersen N, Rysgaard S (1995) Nitrate reduction in sediments and waterlogged soils measured by ¹⁵N techniques. In: Alef K, Nannipieri P (eds) *Methods in applied soil microbiology*. Academic Press, New York, p 287–296
- Risgaard-Petersen N, Nielsen LP, Rysgaard S, Dalsgaard T, Meyer RL (2003) Application of the isotope pairing technique in sediments where anammox and denitrification coexist. *Limnol Oceanogr: Methods* 1:63–73
- Rysgaard S, Fossing H, Jensen MM (2001) Organic matter degradation through oxygen respiration, denitrification, and manganese, iron, and sulfate reduction in marine sediments (the Kattegat and the Skagerrak). *Ophelia* 55:77–91
- Schmid M, Twachtman U, Klein M, Strous and 5 others (2000) Molecular evidence for genus level diversity of bacteria capable of catalyzing anaerobic ammonium oxidation. *Syst Appl Microbiol* 23:93–106
- Schmid M, Walsh K, Webb R, Rijpstra WIC and 11 others (2003) Candidatus '*Scalindua brodae*', sp. nov., candidatus '*Scalindua wagneri*', sp. nov., two new species of anaerobic ammonium oxidizing bacteria. *Syst Appl Microbiol* 26: 529–538
- Stief P, De Beer D, Neumann D (2002) Small-scale distribution of interstitial nitrite in freshwater sediment microcosms: the role of nitrate and oxygen availability, and sediment permeability. *Microb Ecol* 43:367–378
- Strous M, Fuerst JA, Kramer EHM, Logemann S and 5 others (1999) Missing lithotroph identified as new planctomycete. *Nature* 400:446–449
- Svensson JM, Enrich-Prast A, Leonardson L (2001) Nitrification and denitrification in a eutrophic lake sediment bioturbated by oligochaetes. *Aquat Microb Ecol* 23:177–186
- Thamdrup B, Dalsgaard T (2000) The fate of ammonium in anoxic manganese oxide-rich marine sediment. *Geochim Cosmochim Acta* 64:4157–4164
- Thamdrup B, Dalsgaard T (2002) Production of N₂ through anaerobic ammonium oxidation coupled to nitrate reduction in marine sediments. *Appl Environ Microbiol* 68: 1312–1318
- Trimmer M, Nicholls JC, Deflandre B (2003) Anaerobic ammonium oxidation measured in sediments along the Thames Estuary, United Kingdom. *Appl Environ Microbiol* 69:6447–6454
- Zumft WG (1992) The denitrifying prokaryotes. In: Barlows A (ed) *The prokaryotes: a handbook on the biology of bacteria: ecophysiology, isolation, identification, applications*. Springer-Verlag, Berlin, p 554–582