

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/73075>

Please be advised that this information was generated on 2019-11-21 and may be subject to change.

LETTER TO THE EDITOR

Reply to: can CBT substantially change grey matter volume in chronic fatigue syndrome?

Floris P. de Lange,¹ Anda Koers,¹ Joke S. Kalkman,² Gijs Bleijenberg,² Peter Hagoort,^{1,3}
Jos W. M. van der Meer⁴ and Ivan Toni^{1,3}

¹F. C. Donders Centre for Cognitive Neuroimaging, Radboud University Nijmegen, The Netherlands, ²Expert Center Chronic Fatigue, Radboud University Nijmegen Medical Center, The Netherlands, ³Nijmegen Institute for Cognition and Information, Radboud University Nijmegen, The Netherlands and ⁴Department of General Internal Medicine, Radboud University Nijmegen Medical Center, The Netherlands

Correspondence to: Floris P. de Lange, PhD, F. C. Donders Centre for Cognitive Neuroimaging, Radboud University Nijmegen, Kapittelweg 29, 6500 HB Nijmegen, The Netherlands
E-mail: floris.delange@fcdonders.ru.nl
doi: 10.1093/brain/awn208

Received and Accepted August 1, 2008

Sir, We thank Dr Bramsen for her interest in our recently published article (de Lange *et al.* 2008). In this article, we report on an increase in prefrontal grey matter volume following cognitive behavioural therapy (CBT) in a group of patients suffering from the chronic fatigue syndrome (CFS).

Dr Bramsen raises two points of criticism. First, she questions the lack of a control group of patients receiving no treatment or a different treatment. While this limitation was already emphasized in the discussion section of de Lange *et al.* (2008), it remains unclear how the observed behavioral and cerebral changes could be accounted for by generic and CBT-independent factors. Dr Bramsen wonders whether the reported behavioural improvements are really the result of CBT, or whether they could be due to natural course or lifestyle changes, while the underlying disease process is not influenced by CBT. A large body of evidence suggests that this is not the case: there is overwhelming evidence in the literature from several carefully conducted randomized controlled trials (RCT) that demonstrate the specific efficacy of CBT for CFS (e.g. Sharpe *et al.* 1996; Deale *et al.* 1997; Prins *et al.* 2001). It was beyond the scope of this study to disprove that treatments other than CBT would have resulted in the same—or even better—results, simply because this possibility has already been tested and rejected (Whiting *et al.* 2001; Chambers *et al.* 2006).

The second point of criticism raised is that the structural changes in the brain are small, comprising <1% of the total brain volume. Dr Bramsen argues that, although statistically significant, these changes are unlikely to be clinically relevant. This argument is more an intuition than a scientific argument. Given the limited knowledge of

structural plasticity in the adult human brain, it seems arbitrary to set an *a priori* threshold for clinical relevance. More importantly however, Dr Bramsen neglects the fact that the reported increase in grey matter volume was largely regionally specific, confined to left and right lateral prefrontal cortex [see Fig. 3 of de Lange *et al.* (2008)]. The increase in grey matter volume might appear numerically modest when expressed in terms of *whole-brain* grey matter volume. Yet, when relating the change to a more specific anatomical structure (the lateral prefrontal cortex), the increase in grey matter volume was on average 8%. Nevertheless, we prefer to focus on whole-brain grey matter volume as a dependent measure, as it is a reliable, sensitive and unbiased measure of overall cerebral morphology (Good *et al.* 2001; Chard *et al.* 2002).

In summary, while our study did not compare the efficacy of CBT in bringing about the behavioural and brain changes with other treatments or no treatment, there is an overwhelming amount of evidence from the literature that shows the specific efficacy of CBT. The size of the structural changes should naturally be interpreted in relation to the dependent measure used. Finally, we agree with Dr Bramsen that our results do open a wide array of new questions, like: ‘What is the long-term time course of these structural brain changes?’ and ‘Can successful CBT also lead to structural brain changes in other psychopathologies, e.g. depression?’ These exciting research questions will hopefully be the topic of future studies.

References

Chambers D, Bagnall AM, Hempel S, Forbes C. Interventions for the treatment, management and rehabilitation of patients with chronic

- fatigue syndrome/myalgic encephalomyelitis: an updated systematic review 1. *J R Soc Med* 2006; 99: 506–20.
- Chard DT, Parker GJ, Griffin CM, Thompson AJ, Miller DH. The reproducibility and sensitivity of brain tissue volume measurements derived from an SPM-based segmentation methodology. *J Magn Reson Imaging* 2002; 15: 259–67.
- de Lange FP, Koers A, Kalkman JS, Bleijenberg G, Hagoort P, van de Meer JW, et al. Increase in prefrontal cortical volume following cognitive behavioural therapy in patients with chronic fatigue syndrome. *Brain* 2008; 131: 2172–80.
- Deale A, Chalder T, Marks I, Wessely S. Cognitive behavior therapy for chronic fatigue syndrome: a randomized controlled trial. *Am J Psychiatry* 1997; 154: 408–14.
- Good CD, Johnsrude IS, Ashburner J, Henson RNA, Friston KJ, Frackowiak RSJ. A voxel-based morphometric study of ageing in 465 normal adult human brains. *NeuroImage* 2001; 14: 21–36.
- Prins JB, Bleijenberg G, Bazelmans E, Elving LD, de Boo TM, Severens JL, et al. Cognitive behaviour therapy for chronic fatigue syndrome: a multicentre randomised controlled trial. *The Lancet* 2001; 357: 841–7.
- Sharpe M, Hawton K, Simkin S, Surawy C, Hackmann A, Klimes I, et al. Cognitive behaviour therapy for the chronic fatigue syndrome: a randomized controlled trial. *BMJ* 1996; 312: 22–6.
- Whiting P, Bagnall AM, Sowden AJ, Cornell JE, Mulrow CD, Ramirez G. Interventions for the treatment and management of chronic fatigue syndrome: a systematic review. *JAMA* 2001; 286: 1360–8.