

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The version of the following full text has not yet been defined or was untraceable and may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/79883>

Please be advised that this information was generated on 2019-02-15 and may be subject to change.

# Circulating Lipoproteins Are a Crucial Component of Host Defense against Invasive *Salmonella typhimurium* Infection

Mihai G. Netea<sup>1,2</sup>, Leo A. B. Joosten<sup>1,2</sup>, Monique Keuter<sup>1,2</sup>, Frank Wagener<sup>3</sup>, Anton F. H. Stalenhoef<sup>1</sup>, Jos W. M. van der Meer<sup>1,2</sup>, Bart Jan Kullberg<sup>1,2\*</sup>

**1** Division of General Internal Medicine, Department of Medicine, Radboud University Nijmegen Medical Center, Nijmegen, The Netherlands, **2** Nijmegen Institute of Infection, Inflammation and Immunity (N4i), Nijmegen, The Netherlands, **3** Department of Pharmacology and Toxicology, Nijmegen Center for Molecular Life Sciences, Radboud University Nijmegen Medical Center, Nijmegen, The Netherlands

## Abstract

**Background:** Circulating lipoproteins improve the outcome of severe Gram-negative infections through neutralizing lipopolysaccharides (LPS), thus inhibiting the release of proinflammatory cytokines.

**Methods/Principal Findings:** Low density lipoprotein receptor deficient (LDLR<sup>-/-</sup>) mice, with a 7-fold increase in LDL, are resistant against infection with *Salmonella typhimurium* (survival 100% vs 5%,  $p < 0.001$ ), and 100 to 1000-fold lower bacterial burden in the organs, compared with LDLR<sup>+/+</sup> mice. Protection was not due to differences in cytokine production, phagocytosis, and killing of *Salmonella* organisms. The differences were caused by the excess of lipoproteins, as hyperlipoproteinemic ApoE<sup>-/-</sup> mice were also highly resistant to *Salmonella* infection. Lipoproteins protect against infection by interfering with the binding of *Salmonella* to host cells, and preventing organ invasion. This leads to an altered biodistribution of the microorganisms during the first hours of infection: after intravenous injection of *Salmonella* into LDLR<sup>+/+</sup> mice, the bacteria invaded the liver and spleen within 30 minutes of infection. In contrast, in LDLR<sup>-/-</sup> mice, *Salmonella* remained constrained to the circulation from where they were efficiently cleared, with decreased organ invasion.

**Conclusions:** plasma lipoproteins are a potent host defense mechanism against invasive *Salmonella* infection, by blocking adhesion of *Salmonella* to the host cells and subsequent tissue invasion.

**Citation:** Netea MG, Joosten LAB, Keuter M, Wagener F, Stalenhoef AFH, et al. (2009) Circulating Lipoproteins Are a Crucial Component of Host Defense against Invasive *Salmonella typhimurium* Infection. PLoS ONE 4(1): e4237. doi:10.1371/journal.pone.0004237

**Editor:** David M. Ojcius, University of California Merced, United States of America

**Received:** November 10, 2008; **Accepted:** November 21, 2008; **Published:** January 21, 2009

**Copyright:** © 2009 Netea et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The authors have no support or funding to report.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: b.kullberg@aig.umcn.nl

## Introduction

*Salmonella* infections are a significant cause of morbidity and mortality, despite preventive measures and the availability of antibiotics. A major virulence factor of *Salmonella* is lipopolysaccharide (LPS) [1,2], and *Salmonella* strains with a reduced LPS expression have a poor growth under stress conditions and are less virulent [2]. In addition, LPS induces proinflammatory cytokines and is essential for internalization of *Salmonella* by host cells [3]. Interaction of LPS with cellular receptors is essential and therefore, strategies aimed at blocking this interaction may have a therapeutic potential in invasive infections.

Lipoproteins bind and neutralize bacterial LPS, and they prevent the induction of potentially harmful proinflammatory cytokines such as IL-1 $\beta$  and TNF $\alpha$  [4]. In experimental models, administration of lipoproteins protects against endotoxic shock [5–7]. Low density lipoprotein receptor deficient (LDLR<sup>-/-</sup>) mice have a 7 times higher LDL-cholesterol level than control mice. We have shown previously that LDLR<sup>-/-</sup> mice survive longer and have lower proinflammatory cytokine concentrations than control mice after LPS challenge, as well as after infection with *Klebsiella*

*pneumoniae* [8]. In addition, LDL administration can protect against Gram-negative microorganisms through its neutralizing effects on LPS [9].

Although *Salmonella* is a Gram-negative organism, the clinical picture and inflammatory response in systemic *Salmonella* infections (e.g., typhoid fever) differs from that in other Gram-negative sepsis [10]. This is most likely due to the behaviour of *Salmonella* as facultative intracellular pathogens, and to the fact that the pattern of cytokine induction differs from other Gram-negative infections. In mice, TNF $\alpha$  is undetectable in the circulation until several days after *S. typhimurium* infection, whereas TNF $\alpha$  rises at 1 hour after extracellular Gram-negative infection [11]. The level of cytokine during *Salmonella* infections does not reach the toxic levels seen in endotoxic shock, and inhibition of TNF $\alpha$  during *Salmonella* infection worsens the outcome [12,13].

Considering these differences in the pathogenesis of *Salmonella* and extracellular Gram-negative infections, one would envisage either beneficial effects of lipoproteins on the host resistance to *Salmonella* through blockade of cellular internalization, or deleterious effects by blocking the induction of cytokines by *Salmonella* LPS that are required for the activation of host defense. In the

present study, we investigated the effect of lipoproteins on the outcome of *Salmonella* infection.

## Methods

### Animals

Homozygous C57Bl/6J mice lacking low density lipoprotein receptors (LDLR<sup>-/-</sup>) and their wild-type littermates (C57Bl/6J LDLR<sup>+/+</sup>) were obtained from Jackson Laboratory (Bar Harbour, ME) [8]. Homozygous apolipoprotein E (ApoE)-deficient mice on a C57Bl/6 background were obtained from the Transgenic Facility of Leiden University Medical Center, Leiden, The Netherlands [14]. Six to eight weeks old littermate LDLR<sup>+/+</sup> and LDLR<sup>-/-</sup> mice were used, weighing 20–25 grams. The animals were fed standard laboratory chow and housed under specific pathogen free conditions. The experiments were approved by the Ethics Committee for animal experiments at the Radboud University Nijmegen.

### *Salmonella typhimurium* infection

A serum-resistant strain of *S. typhimurium* (phage type 510) was grown by overnight incubation at 37°C in nutrient broth (BHI Oxoid). Mice were injected i.v. or i.p. with  $1 \times 10^8$  cfu of *S. typhimurium*. Survival was assessed daily for 21 days in groups of at least 20 animals. On day 1, 3 and 7 after infection, mice were killed by cervical dislocation and blood for cytokines or organs for outgrowth of the microorganisms were collected. For this purpose, the liver and spleen were removed aseptically, and bone marrow was flushed from the femur aseptically with 1 ml of sterile saline. The number of viable *Salmonella* organisms was determined by plating several dilutions on Brilliant Green agar (BGA) plates. The results were expressed as log cfu per gram of tissue.

### Distribution of *Salmonella* cfu

In a separate experiment, *Salmonella* cfu ( $10^5$ /mouse) were injected i.v. Distribution was determined in blood, liver and spleen after 30, 60, 120 and 360 minutes by plating serial dilutions of blood and homogenized tissue samples on BGA plates. Groups of 5 mice were used for each time point.

### Intracellular killing of *S. typhimurium* by peritoneal phagocytes

Phagocytosis and intracellular killing of *S. typhimurium* was assessed in vitro using peritoneal macrophages and PMN of LRLR<sup>+/+</sup> and LDLR<sup>-/-</sup> mice. Exudate peritoneal neutrophils (PMN) were harvested 4 h after an i.p. injection of 10% proteose peptone, and exudate macrophages 72 h after i.p. injection of proteose peptone.  $5 \times 10^5$  cells in 100  $\mu$ L of RPMI were dispensed into 96-well flat bottom plates (Costar) and incubated at 37°C and 5% CO<sub>2</sub>. To assess phagocytosis,  $1 \times 10^5$  *Salmonella* organisms/mL were incubated on the phagocyte monolayers at 37°C in RPMI with 10% serum. After 15 min, supernatants were aspirated and the monolayers were gently washed with medium to remove uningested bacteria. The supernatants were plated on BGA agar (the non-phagocytosed fraction). To assess intracellular killing, the wells containing the cells with phagocytosed bacteria were scraped with a plastic paddle and washed with 200  $\mu$ L distilled H<sub>2</sub>O to lyse the phagocytes. The number of viable bacteria was determined by plating serial dilutions on BGA plates.

### In vitro cytokine production

Resident peritoneal macrophages were harvested from peritoneal cavity, and cells were resuspended in RPMI 1640 containing

1 mM pyruvate, 2 mM L-glutamine and 100  $\mu$ g gentamicin per ml, and incubated ( $10^5$ /well) in 96-wells microtiter plates (Costar). Heat-killed (30 min, 100°C) *S. typhimurium* ( $10^6$  microorganisms in 100  $\mu$ L of RPMI) were added to peritoneal macrophages and incubated at 37°C in 5% CO<sub>2</sub>. After 24 h, the supernatants were collected and stored at -70°C until assayed. To the macrophages in the monolayer, 200  $\mu$ L of RPMI was added and the cells were disrupted by three freeze-thaw cycles to determine the cell-associated cytokine contents.

### Cytokine measurements

TNF $\alpha$ , IL-1 $\alpha$  and IL-1 $\beta$  concentrations were determined using specific radioimmunoassays (RIA), as previously described [8]. To assess cytokine mRNA expression, total RNA from spleen cells 24 hours after infection was isolated as described [15]. The following primers were used for the PCR reactions: GAPDH, sense, 5'-AACTCCCTCAAGATTGTCAGCA-3', and antisense, 5'-TCC-ACCACCCTGTTGCTGTA-3'; TNF $\alpha$ , sense, 5'-TCTCAT-CAGTTCTATGGCCC-3', and antisense, 5'-GGGAGTAGA-CAAGGTACAAC-3'; IL-1 $\alpha$ , sense, 5'-CAGTTCCTGCCATT-GACCATC-3', and antisense, 5'-TCTCACTGAACTCAG-CCGT-3', IL-1 $\beta$ , sense, 5'-TTGACGGACCCAAAAGATG-3', and antisense, 5'-AGAAGGTGCTCATGTCTCCTCA-3' (Eurogentec, Seraing, Belgium). After checking the reactions to be in the log phase, thirty PCR cycles were performed with sets at 92°C for 30 sec., 55°C for 30 sec., and 72°C for 90 sec., using a Mastercycler 5330 (Eppendorf). PCR products were run on 2% agarose gels stained with ethidium bromide. The gels were scanned on a densitometer (GS-670, Bio-Rad) and analyzed using Molecular Analyst software (Bio-Rad). The relative amount of TNF $\alpha$ , IL-1 $\alpha$  and IL-1 $\beta$  mRNA in a sample was expressed as a ratio versus the amount of mRNA for the housekeeping gene GAPDH.


### Growth of *Salmonella* in vitro

To investigate the effect of lipoproteins on microbial growth in vitro,  $0.5 \times 10^3$  cfu *S. typhimurium* in 0.5 mL BHI were incubated with 0.5 mL of plasma obtained from control C57Bl/6J mice, or from LDLR<sup>-/-</sup> mice and ApoE<sup>-/-</sup> mice. After 2, 7, 12 and 24 hours, aliquots of 0.1 mL were removed, serial dilutions were plated on BGA agar, and cfu were counted after overnight incubation at 37°C.

### Effect of lipoproteins on interaction of *Salmonella* with monocytes and endothelial cells

To assess the effect of lipoproteins on the production of cytokines, *S. typhimurium* LPS (10 ng/mL; Sigma) and heat-killed (30 min, 100°C) *S. typhimurium* ( $10^7$  organisms/mL) were preincubated with lipoprotein-depleted plasma (LPDP) or isolated LDL [16] at various concentrations for 60 min, before being added to the macrophages of LDLR<sup>+/+</sup> mice ( $10^5$ /well). The production of TNF $\alpha$  after 24 h stimulation was measured as described above, and expressed as relative TNF production compared to controls in LPDP.

To assess the effect of lipoproteins on the attachment of *Salmonella* to vascular endothelial cells, *S. typhimurium* were resuspended to  $6 \times 10^8$ /ml in 0.01 mg/ml FITC (Fluka) in 0.05 M carbonate-bicarbonate buffer (pH 9.5). After incubation for 15 min at room temperature in the dark, FITC-labeled *Salmonella* cells were washed twice in PBS containing 1% BSA and subsequently incubated with isolated 1.1 mmol/L LDL for 4 hours, or with LPDP as a negative control. The human endothelial cell line (HMEC-1) (CDC Atlanta, GA) was cultured in MCDB131 medium supplemented with 10% fetal calf serum,



**Figure 1. LDLR<sup>-/-</sup> mice are more resistant to *S. typhimurium* infection.** Survival of LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> C57Bl/6J mice after i.v. injection of  $10^2$  *S. typhimurium*. n=20/group. doi:10.1371/journal.pone.0004237.g001

EGF (10 ng/ml), hydrocortison (1  $\mu$ g/ml), and glutamine at 37°C and 5% CO<sub>2</sub>.  $1 \times 10^5$  HMEC-1 cells were trypsinized and incubated for 1 hour with  $3 \times 10^9$  FITC-labeled *S. typhimurium*, which were preincubated with either LDL or LPDP. After incubation, the non-bound *Salmonella* was thoroughly washed off, after which the cells were fixated with 2% paraformaldehyde in PBS and analyzed for binding of FITC-labeled *Salmonella* by flow cytometry using the FACScalibur (BD Biosciences).

#### Statistical analysis

Survival of groups of mice was compared by the Kaplan-Meier log-rank test. Differences in concentrations of cytokines and in organ counts of the microorganisms were analyzed by the Mann-Whitney U test. Differences were considered significant at  $P < .05$ . All the experiments were at least performed in duplicate.

## Results

### Outcome of *Salmonella* infection in LDLR<sup>-/-</sup> mice

The total cholesterol concentrations were significantly higher in the uninfected LDLR<sup>-/-</sup> mice than in their wild-type littermates


( $9.6 \pm 1.1$  mmol/L vs.  $2.3 \pm 0.5$  mmol/L). After i.v. infection with  $10^2$  cfu of *S. typhimurium*, only 5% of the LDLR<sup>-/-</sup> mice died, whereas the mortality of control LDLR<sup>+/+</sup> was 100% within 12 days of infection ( $P < .001$ ; Fig. 1). A similar difference in mortality was apparent when mice were infected intraperitoneally with *S. typhimurium* (10% mortality in LDLR<sup>-/-</sup> mice, vs. 100% mortality in control LDLR<sup>+/+</sup> mice,  $p < 0.01$ ). The reduced mortality to infection in LDLR<sup>-/-</sup> mice was accompanied by a markedly reduced bacterial load in the organs (Fig. 2). On day 7 of infection, the differences between the control and LDLR<sup>-/-</sup> mice approached 10,000-fold ( $P < .001$ ; Fig. 2).

### Intracellular killing of *Salmonella* by cells from LDLR<sup>+/+</sup> and LDLR<sup>-/-</sup> mice in vitro


The numbers of *Salmonella* CFU phagocytized by neutrophils and macrophages of LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice were similar (Fig. 3A). In addition, the intracellular killing assay demonstrated that neutrophils and macrophages of LDLR<sup>-/-</sup> mice and LDLR<sup>+/+</sup> mice did not differ in their ability to kill *S. typhimurium* intracellularly (Fig. 3B). The killing rate did not differ when lipoprotein-rich serum of LDLR<sup>-/-</sup> mice was coincubated with LDLR<sup>+/+</sup> control macrophages, and likewise, serum from LDLR<sup>+/+</sup> mice did not affect the killing of *Salmonella* by LDLR<sup>-/-</sup> macrophages (not shown).

### Circulating cytokines during *Salmonella* infection

On day 1, cytokine concentrations in all samples were under the detection limit. No detectable concentrations of IL-1 $\beta$  (<20 pg/ml) were found at any time point during the infection. On day 3, IL-1 $\alpha$  and TNF $\alpha$  were under the detection limit in LDLR<sup>-/-</sup> mice, while TNF $\alpha$  concentrations tended to be slightly higher ( $45 \pm 10$  pg/ml) in LDLR<sup>+/+</sup> mice (n.s.). On day 7, circulating concentrations of IL-1 $\alpha$  and TNF $\alpha$  were significantly higher in LDLR<sup>+/+</sup> than in LDLR<sup>-/-</sup> mice:  $95 \pm 63$  pg/ml vs  $30 \pm 10$  pg/ml for IL-1 $\alpha$  ( $P < 0.02$ ) and  $1140 \pm 290$  pg/ml vs  $43 \pm 6$  pg/ml for TNF $\alpha$  ( $P < 0.01$ ) (Fig. 4A). These differences were most likely due to the greater amounts of *Salmonella* in the LDLR<sup>+/+</sup> mice, leading to increased cytokine stimulation.



**Figure 2. Outgrowth of *S. typhimurium* in the organs of LDLR<sup>+/+</sup> and LDLR<sup>-/-</sup> mice.** Outgrowth of *S. typhimurium* in the liver, spleen and bone marrow of LDLR<sup>-/-</sup> and control (LDLR<sup>+/+</sup>) C57Bl/6J mice after i.v. injection of  $10^2$  cfu. Each point represents the mean  $\pm$  SD for at least 10 animals. Significant differences between LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice are indicated (\*,  $P < .01$ ; \*\*,  $P < .001$ ; Mann-Whitney U test). doi:10.1371/journal.pone.0004237.g002



**Figure 3. Phagocytosis and killing of *S. typhimurium* by neutrophils and macrophages of LDLR<sup>-/-</sup> mice.** Phagocytosis of *S. typhimurium* after 15 min incubation, and intracellular killing of *S. typhimurium* after 4 h, by proteose peptone-elicited peritoneal neutrophils and macrophages. Data are expressed as percentage of the initial number of microorganisms. No significant differences between LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> were found. doi:10.1371/journal.pone.0004237.g003

Importantly, the differences in cytokine concentrations were not secondary to an intrinsically deficient cytokine production in the organs of the LDLR<sup>-/-</sup> mice: the amounts of TNF and IL-1 $\beta$  mRNA in the spleens of LDLR<sup>+/+</sup> and LDLR<sup>-/-</sup> mice on day 1 of infection was similar (Fig. 4B). No differences in the expression of IL-1 $\alpha$  mRNA were observed either (not shown). In line with this, peritoneal macrophages of LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice stimulated with heat-killed *Salmonella* produced similar amounts of TNF $\alpha$  in vitro: unstimulated macrophages, 194 $\pm$ 91 pg/ml and 186 $\pm$ 55 pg/ml, macrophages coincubated with heat-killed *Salmonella*, 550 $\pm$ 166 pg/ml and 520 $\pm$ 135 pg/ml for LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice, respectively ( $P>.05$ ).

#### Resistance of ApoE<sup>-/-</sup> mice to *Salmonella* infection

To investigate whether the increased lipoprotein concentrations in another model of hyperlipoproteinemia can also protect against salmonellosis, hyperlipoproteinemic ApoE<sup>-/-</sup> mice (total serum cholesterol, 16.1 $\pm$ 3.7 vs. 1.9 $\pm$ 0.2 mmol/L [14]) were infected i.v. with 10<sup>2</sup> cfu of *S. typhimurium*. Whereas little differences were apparent on day 1 of infection, the outgrowth of *Salmonella* on day 3 and 7 after the infection was 100 to 1000-fold less in the liver and

spleen of ApoE<sup>-/-</sup> mice compared to that in ApoE<sup>+/+</sup> control animals (Fig. 5).

#### Effect of lipoproteins on the growth of *S. typhimurium* in vitro

To test whether lipoproteins have a direct inhibitory effect on the growth of *Salmonella*, plasma isolated from control mice (cholesterol concentration 2.3 mmol/L) and hyperlipoproteinemic plasma from either LDLR<sup>-/-</sup> or ApoE<sup>-/-</sup> mice (cholesterol concentrations, 9.6 and 16.1 mmol/L) was added to the culture. The growth curves of *Salmonella* were similar in broth with plasma obtained from all mouse strains (Fig. 6A).

#### Inhibition of the interaction of *Salmonella* with monocytes and endothelial cells by lipoproteins

Because lipoproteins are known to bind and neutralize LPS, we preincubated *Salmonella* LPS and heat-killed whole *Salmonella* bacteria with LDL at various concentrations, and subsequently stimulated normal (LDLR<sup>+/+</sup>) macrophages for cytokine production. As shown in Fig. 6B, TNF $\alpha$  production induced by *S. typhimurium* is reduced in the presence of elevated LDL concentrations. To assess the effect of lipoproteins on the attachment of *Salmonella* to endothelial cells, FITC-labelled *S. typhimurium* was preincubated with LDL. Their attachment to endothelial cells was significantly reduced compared to that of *Salmonella* preincubated with lipoprotein-free plasma, as shown by both reduction of the percentage of cells binding *Salmonella* (97% cells bound LDPD-*Salmonella*, whereas only 82% cells bound LDL-*Salmonella*), as well as the mean fluorescence intensity per cell (41% reduction, from 56 to 33 conventional units) (see also Fig. 6C).



#### Protection against organ invasion by *Salmonella* in LDLR<sup>-/-</sup> mice

To investigate whether hyperlipoproteinemia influences the early organ invasion from the bloodstream by *Salmonella*, we determined the early distribution of the microorganisms after i.v. injection of 10<sup>5</sup> *S. typhimurium* cfu. Blood from LDLR<sup>-/-</sup> mice contained significantly more *Salmonella* cfu than that of LDLR<sup>+/+</sup> mice 30 minutes after injection ( $P<0.01$ ), but an efficient elimination of the microorganisms occurred during the next 6 hours (Fig. 7A). The numbers of cfu in the liver and spleen were 70–80% lower in LDLR<sup>-/-</sup> mice than those in LDLR<sup>+/+</sup> mice at 30 minutes after injection, and remained lower throughout the experiment, the difference between mouse strains being significant at 30, 60, 120 and 360 minutes for the liver (Fig. 7B,  $P<0.05$ ) and at 30 and 60 minutes for the spleen (Fig. 7C,  $P<0.05$ ).

#### Discussion

In the present study, we demonstrate that hyperlipoproteinemic mice are resistant against *S. typhimurium* infection. The protection was not due to the absence of the LDLR in the knock-out mouse strain, but to a direct effect of hyperlipoproteinemia. The beneficial effect of lipoproteins was exerted by blocking the interaction of *Salmonella* with host cells, including endothelial cells and monocytes, which led to inhibition of organ invasion. This resulted in an altered distribution of the microorganism to the organs of the host, and increased survival.


It has previously been shown that lipoproteins bind and neutralize LPS, with beneficial effects in Gram-negative infections [5–8]. As *S. typhimurium* is an LPS-containing Gram-negative bacterium, and *Salmonella* LPS plays a crucial role in cytokine stimulation [17] and induction of mortality in vivo [18], an






**Figure 4. In-vivo cytokine production in LDLR<sup>+/+</sup> and LDLR<sup>-/-</sup> mice infected with *S. typhimurium*.** LDLR<sup>-/-</sup> and control (LDLR<sup>+/+</sup>) C57Bl/6J mice were infected i.v. with  $10^2$  cfu of *S. typhimurium*. After 1, 3 or 7 days, groups of 5 mice/group were sacrificed, and circulating concentrations of cytokines in the plasma were measured by specific ELISA. In a separate experiment, TNF and IL-1 $\beta$  mRNA in the spleens of LDLR<sup>+/+</sup> and LDLR<sup>-/-</sup> mice was assessed by semi-quantitative RT-PCR, and expressed as ratio to GAPDH mRNA expression. Data are presented as means $\pm$ SD, \* $p$ <0.05.  
doi:10.1371/journal.pone.0004237.g004

improved survival of LDLR<sup>-/-</sup> mice during systemic *S. typhimurium* infection would have been expected. Indeed, LDLR<sup>-/-</sup> mice were less susceptible to *S. typhimurium* infection, but this was not due to blunted cytokine production, as we observed undetectable or low cytokine circulating concentrations during the infection in the LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice. In addition, the expression of cytokine mRNA was similar in the organs of LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice. Thus, the cytokine response is not responsible for the mortality due to systemic *S. typhimurium* infection in this model, and the major difference is the almost complete absence of *Salmonella* in the organs of the LDLR<sup>-/-</sup> mice.


The low circulating cytokine response during systemic *S. typhimurium* infection may be attributed to the facultative intracellular nature of the organism. It should be noted that at later time points during the infection, the LDLR<sup>+/+</sup> mice



**Figure 5. Outgrowth of *S. typhimurium* in the organs of ApoE<sup>+/+</sup> and ApoE<sup>-/-</sup> mice.** Outgrowth of *S. typhimurium* in the liver and spleen of ApoE<sup>-/-</sup> and control (ApoE<sup>+/+</sup>) C57Bl/6J mice after i.v. injection of  $10^2$  cfu. Each point represents the mean $\pm$ SD for at least 10 animals. Significant differences between ApoE<sup>-/-</sup> and ApoE<sup>+/+</sup> mice were found for both liver and spleen on days 3 and 7 ( $p$ <0.01).  
doi:10.1371/journal.pone.0004237.g005



**Figure 6. The effect of LDL on the interaction between *S. typhimurium* and the host cells.** The growth of *S. typhimurium* was identical in plasma harvested from control C57Bl/6, LDLR<sup>-/-</sup> or ApoE<sup>-/-</sup> mice (upper panel). Preincubation of *Salmonella* LPS or heat-killed *S. typhimurium* with various concentrations of LDL led to a significantly diminished stimulation of TNF when added to normal macrophages (middle panel). Similarly, preincubation of FITC-labelled *S. typhimurium* with LDL led to a diminished adhesion of the microorganisms to vascular endothelial cells, when compared with the lipoprotein-deficient serum (LPDS).  
doi:10.1371/journal.pone.0004237.g006



**Figure 7. Hyperlipoproteinemia inhibits organ invasion by *S. typhimurium*.** Distribution of *S. typhimurium* to the blood (cfu/ml), the liver (cfu/organ), and the spleen (cfu/organ) at various time points after i.v. injection of  $10^5$  *Salmonella* cfu. Each point represents the mean  $\pm$  SD for at least 5 animals. Significant differences between LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice are indicated (\* $p < 0.05$ ; \*\* $p < 0.01$ ; Mann-Whitney U test). doi:10.1371/journal.pone.0004237.g007

exhibited a greater cytokine response than did LDLR<sup>-/-</sup> mice, and this is likely due to the 1000-fold greater bacterial burden in the control mice, leading to substantial stimulation of the host response.

We hypothesized that the decreased bacterial growth in the LDLR<sup>-/-</sup> mice may be a result of enhanced phagocytosis and intracellular killing of *Salmonella* organisms in these mice. However, this proved not to be the case: phagocytosis and subsequent intracellular killing of the organisms by both neutrophils and macrophages did not differ between LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice. Another reason for protection could have been the absence of the LDL receptor in the knock-out mice. *Toxoplasma gondii* is known to use the host LDLR for cholesterol acquisition [19], whereas penetration of cells by *Pseudomonas* exotoxin A is mediated through LDLR-related protein [20,21]. Thus, use of the LDLR by *Salmonella* during organ invasion could be envisaged, but was ruled out by the observation that hyperlipoproteinemic ApoE<sup>-/-</sup> mice, which have an intact LDLR, also were resistant to *Salmonella* infection. In addition, LDL affected the adhesion of *S. typhimurium* to LDLR-bearing endothelial cells. This demonstrates that elevated lipoprotein concentrations, and not the lack of LDLR itself, are responsible from the resistance of mice against *Salmonella* infection.

Theoretically, there are several mechanisms that could account for the beneficial effects of the lipoproteins on *Salmonella* infection. Firstly, a direct effect of lipoproteins on the growth of *Salmonella* could be envisaged. In this respect, it is of interest that HDL has been found to be cidal against *Trypanosoma cruzi* [20]. However, the growth of *Salmonella* was similar in the plasma of LDLR<sup>-/-</sup>, ApoE<sup>-/-</sup> and control mice, excluding a direct antimicrobial effect of LDL. Secondly, lipoproteins may interact with *Salmonella*, putatively with its LPS component, and thus block bacterial binding and internalization by host cells. As LPS is crucial for the internalization of *Salmonella* [22], blocking the interaction between LPS and host cells may prevent subsequent tissue invasion. Indeed, preincubation of *Salmonella* with LDL led to reduced cytokine production, demonstrating that lipoproteins are able to inhibit the interaction of *Salmonella* with monocytes. Even more relevant for tissue invasion, preincubation of *Salmonella* with LDL significantly reduced its attachment to endothelial cells. This protective mechanism in which lipoproteins block *Salmonella* interaction with endothelial cells by their blockade of LPS represents the same type of mechanism as previously shown by the blockade of MSCRAMMs (microbial surface components recognizing adhesive matrix molecules) of *Staphylococcus* by naturally occurring antibodies, resulting in the inhibition of staphylococcal adhesion to endothelial cell and reduced tissue invasion [23].

Interaction of *Salmonella* with host cells likely is an important early step in the pathogenesis of invasive infection. The ability to infect tissue macrophages has been described as an invasive trait of intracellular bacteria such as *Salmonella* spp. [24], and attachment to endothelial cells is the first step in organ invasion by *Salmonella* from the bloodstream. The hypothesis that lipoproteins are able to directly modify organ invasion by *Salmonella*, was tested by

assessing early distribution of *Salmonella* organisms in LDLR<sup>-/-</sup> and LDLR<sup>+/+</sup> mice after i.v. injection of a large bacterial load. Indeed, we observed that invasion of *Salmonella* organisms into the liver and spleen of LDLR<sup>-/-</sup> mice was markedly lower than that in control mice. The difference in clearance of bacteria between the two strains of mice was already apparent within 30 minutes after injection. In this early phase of infection, the numerical balance between bacterial burden and host defense mechanisms in the tissues will determine the outcome of infection, and the LDLR<sup>-/-</sup> mice start with a substantial advantage over control animals. In contrast to the LDLR<sup>+/+</sup> mice, the vast majority of *Salmonella* organisms in the LDLR<sup>-/-</sup> remain sequestered in the circulation, where are eliminated by complement and neutrophils, known to efficiently clear *Salmonella* from the bloodstream [25]. Blocking tissue invasion by lipoproteins is not unique for *Salmonella*, as others have reported that VLDL inhibits liver invasion by *Plasmodium* sporozoites, leading to protection against malaria [26].

The precise molecular interaction between *Salmonella* and lipoproteins remains to be elucidated, but the bacterial LPS is the most likely candidate to be involved. *Salmonella* is known to interact with host cell Toll-like receptor (TLR)-4 through its LPS component [17,27], and this signaling mechanism is likely blocked by binding of lipoproteins to the LPS. The exact nature of the lipoprotein particle responsible for interaction with *Salmonella* has yet to be identified. Phospholipids have been shown to mediate LPS neutralization, but protein components, such as apolipoprotein E, have also been reported to bind LPS [28]. Our findings in the apoE- and LDLR-deficient mice, which are also protected against salmonellosis, however, point to a binding site other than apoE. In addition to TLR4, the macrophage scavenger receptors [29], and the cystic fibrosis transmembrane conductance regulator protein [30] are probably involved in the entry of *Salmonella* into cells. Whether the interaction of these receptors with *Salmonella* is also influenced by lipoproteins remains to be elucidated.

In conclusion, plasma lipoproteins appear to be an important host defense mechanism against invasive *Salmonella* infection. A direct and rapid interaction between lipoproteins and *Salmonella* in the bloodstream occurs, preventing the invasion of the microorganisms from the bloodstream into the organs. These new insights improve the understanding of the pathogenesis of *Salmonella* infection, and could ultimately lead to the design of new therapeutic strategies.

## Acknowledgments

The authors thank Prof. Jaap van Dissel (Leiden University Medical Center) and Dr Pierre Demacker for advice, and Ineke Verschuere for assistance with the animal experiments and cytokine assays.

## Author Contributions

Conceived and designed the experiments: MGN LAJ MK JWMVdM BJK. Performed the experiments: MGN MK FW AFHS. Analyzed the data: MGN LAJ FW. Contributed reagents/materials/analysis tools: AFHS. Wrote the paper: MGN LAJ MK AFHS JWMVdM BJK.

## References

- Khan SA, Everest P, Servos S, Foxwell N, Zahringer U, et al. (1998) A lethal role for lipid A in *Salmonella* infections. *Mol Microbiol* 29: 571–579.
- Thomsen LE, Chadfield MS, Bispham J, Wallis TS, Olsen JE, et al. (2003) Reduced amounts of LPS affect both stress tolerance and virulence of *Salmonella enterica* serovar Dublin. *FEMS Microbiol Lett* 228: 225–231.
- Lyczak JB, Zaidi TS, Grout M, Bitner M, Contreras I, et al. (2001) Epithelial cell contact-induced alterations in *Salmonella enterica* serovar Typhi lipopolysaccharide are critical for bacterial internalization. *Cell Microbiol* 3: 763–772.
- Flegel WA, Wolpl A, Mannel DN, Northoff N (1989) Inhibition of endotoxin-induced activation of human monocytes by human lipoproteins. *Infect Immun* 57: 2237–2245.
- Read TE, Grunfeld C, Kumwenda Z, Calhoun MC, Kane JP, et al. (1995) Triglyceride-rich lipoproteins prevent septic death in rats. *J Exp Med* 182: 267–272.
- Harris HW, Grunfeld C, Feingold KR, Rapp JH (1990) Human very low density lipoproteins and chylomicrons can protect against endotoxin-induced death in mice. *J Clin Invest* 86: 696–702.

7. Pajkrt D, Doran JE, Koster F, Lerch PG, Arnet B, et al. (1996) Antiinflammatory effects of reconstituted high-density lipoprotein during human endotoxemia. *J Exp Med* 184: 1601–1608.
8. Netea MG, Demacker PNM, Kullberg BJ, Boerman OC, Verschuereen I, et al. (1996) Low-density-lipoprotein receptor deficient mice are protected against lethal endotoxemia and severe Gram-negative infections. *J Clin Invest* 97: 1366–1372.
9. Park KH, Kim JS, Lee YR, Moon YJ, Hur H, et al. (2007) Low-density lipoprotein protects *Vibrio vulnificus*-induced lethality through blocking lipopolysaccharide action. *Exp Mol Med* 39: 673–678.
10. Keuter M, Dharmana E, Gasem MH, Van der Ven-Jongekrijg J, Djokomocljanto R, et al. (1994) Patterns of proinflammatory cytokines and inhibitors during typhoid fever. *J Infect Dis* 169: 1306–1311.
11. Jotwani R, Tanaka Y, Watanabe K, Tanaka K, Kato N, et al. (1995) Cytokine stimulation during *Salmonella typhimurium* sepsis in Itys mice. *J Med Microbiol* 42: 348–352.
12. Nauciel C, Espinasse-Maes F (1992) Role of gamma-interferon and tumor necrosis factor alpha in resistance to *Salmonella typhimurium* infection. *Infect Immun* 60: 450–454.
13. Mastroeni P, Skepper JN, Hormacche CE (1995) Effect of anti-tumor necrosis factor alpha antibodies on histopathology of primary *Salmonella* infections. *Infect Immun* 63: 3674–3682.
14. Vonk AG, De Bont N, Netea MG, Demacker PN, van der Meer JW, et al. (2004) Apolipoprotein-E-deficient mice exhibit an increased susceptibility to disseminated candidiasis. *Med Mycol* 42: 341–348.
15. Chomczynski P, Sacchi N (1987) Single step method of RNA isolation by acid guanidinium phenol chloroform extraction. *Anal Biochem* 162: 156–159.
16. Demacker PNM, Vos-Janssen HE, Jansen AP, Van't Laar A (1977) Evaluation of the dual precipitation method by comparison with the ultracentrifugation method for the measurement of lipoproteins in serum. *Clin Chem* 23: 1238–1244.
17. Tapping RI, Akashi S, Miyake K, Godowski PJ, Tobias PS (2000) Toll-like receptor 4, but not toll-like receptor 2, is a signaling receptor for *Escherichia* and *Salmonella* lipopolysaccharides. *J Immunol* 165: 5780–5787.
18. Freudenberg MA, Tchaptchet S, Keck S, Fejer G, Huber M, et al. (2008) Lipopolysaccharide sensing an important factor in the innate immune response to Gram-negative bacterial infections: benefits and hazards of LPS hypersensitivity. *Immunobiology* 213: 193–203.
19. Coppens I, Sinai AP, Joiner KA (2000) *Toxoplasma gondii* exploits host low-density lipoprotein receptor-mediated endocytosis for cholesterol acquisition. *J Cell Biol* 149: 167–180.
20. Smith AB, Esko JD, Hajduk SL (1995) Killing of trypanosomes by the human haptoglobin-related protein. *Science* 268: 284–286.
21. Laithwaite JE, Benn SJ, Marshall WS, FitzGerald DJ, LaMarre J (2001) Divergent *Pseudomonas* exotoxin A sensitivity in normal and transformed liver cells is correlated with low-density lipoprotein receptor-related protein expression. *Toxicol* 39: 1283–1290.
22. Freudenberg MA, Merlin T, Gumenscheimer M, Kalis C, Landmann R, et al. (2001) Role of lipopolysaccharide susceptibility in the innate immune response to *Salmonella typhimurium* infection: LPS, a primary target for recognition of Gram-negative bacteria. *Microbes Infect* 3: 1213–1222.
23. Patti JM, Allen BL, McGavin MJ, Hook M (1994) MSCRAMM-mediated adherence of microorganisms to host tissues. *Annu Rev Microbiol* 48: 585–617.
24. Wijburg OL, Simmons CP, van Rooijen N, Strugnell RA (2000) Dual role for macrophages in vivo in pathogenesis and control of murine *Salmonella enterica* var. *Typhimurium* infections. *Eur J Immunol* 30: 944–953.
25. Fierer J (2001) Polymorphonuclear leukocytes and innate immunity to *Salmonella* infections in mice. *Microbes Infect* 3: 1233–1237.
26. Sinnis P, Willnow TE, Briones MR, Herz J, Nussenzweig V (1996) Remnant lipoproteins inhibit malaria sporozoite invasion of hepatocytes. *J Exp Med* 184: 945–954.
27. Royle MC, Totemeyer S, Alldridge LC, Maskell DJ, Bryant CE (2003) Stimulation of Toll-like receptor 4 by lipopolysaccharide during cellular invasion by live *Salmonella typhimurium* is a critical but not exclusive event leading to macrophage responses. *J Immunol* 170: 5445–5454.
28. Van Oosten M, Rensen PC, Van Amersfoort ES, Van Eck M, Van Dam AM, et al. (2001) Apolipoprotein E protects against bacterial lipopolysaccharide-induced lethality. A new therapeutic approach to treat gram-negative sepsis. *J Biol Chem* 276: 8820–8824.
29. van Oosten M, van Amersfoort ES, van Berkel TJ, Kuiper J (2001) Scavenger receptor-like receptors for the binding of lipopolysaccharide and lipoteichoic acid to liver endothelial and Kupffer cells. *J Endotoxin Res* 7: 381–384.
30. Lyczak JB, Pier GB (2002) *Salmonella enterica* serovar typhi modulates cell surface expression of its receptor, the cystic fibrosis transmembrane conductance regulator, on the intestinal epithelium. *Infect Immun* 70: 6416–6423.