

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/89306>

Please be advised that this information was generated on 2019-06-27 and may be subject to change.

Sequence variants of the *DFNB31* gene among Usher syndrome patients of diverse origin

Elena Aller,^{1,2} Teresa Jaijo,^{1,2} Erwin van Wijk,^{3,4,5,6} Inga Ebermann,⁷ Ferry Kersten,^{3,4,5,6,8}
 Gema García-García,¹ Krysta Voesenek,⁴ María José Aparisi,¹ Lies Hoefsloot,⁴ Cor Cremers,^{3,6}
 Manuel Díaz-Llopis,^{9,10} Ronald Pennings,^{3,6} Hanno J. Bolz,⁷ Hannie Kremer,^{3,5,6} José M. Millán^{1,2}

(The first two authors contributed equally to this work)

¹Unidad de Genética, Hospital Universitario La Fe, Valencia, Spain; ²CIBER de Enfermedades Raras (CIBERER), Valencia, Spain; ³Dept Otorhinolaryngology, Head and Neck Surgery, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands; ⁴Dept Human Genetics, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands; ⁵Nijmegen Centre for Molecular Life Sciences, Radboud University Nijmegen, Nijmegen, The Netherlands; ⁶Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen, Nijmegen, The Netherlands; ⁷Institute of Human Genetics, University Hospital of Cologne, Cologne, Germany; ⁸Dept Ophthalmology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands; ⁹Servicio de Oftalmología, Hospital Universitario La Fe, Valencia, Spain; ¹⁰Departamento de Oftalmología, Facultad de Medicina, Universidad de Valencia, Valencia, Spain

Purpose: It has been demonstrated that mutations in deafness, autosomal recessive 31 (*DFNB31*), the gene encoding whirlin, is responsible for nonsyndromic hearing loss (NSHL; *DFNB31*) and Usher syndrome type II (*USH2D*). We screened *DFNB31* in a large cohort of patients with different clinical subtypes of Usher syndrome (*USH*) to determine the prevalence of *DFNB31* mutations among *USH* patients.

Methods: *DFNB31* was screened in 149 *USH2*, 29 *USH1*, six atypical *USH*, and 11 unclassified *USH* patients from diverse ethnic backgrounds. Mutation detection was performed by direct sequencing of all coding exons.

Results: We identified 38 different variants among 195 patients. Most variants were clearly polymorphic, but at least two out of the 15 nonsynonymous variants (p.R350W and p.R882S) are predicted to impair whirlin structure and function, suggesting eventual pathogenicity. No putatively pathogenic mutation was found in the second allele of patients with these mutations.

Conclusions: *DFNB31* is not a major cause of *USH*.

Usher syndrome (OMIM 276900–2; OMIM 276905; OMIM 605472) is the most common cause of genetic deafblindness. This syndrome follows an autosomal recessive pattern of inheritance and is characterized by retinitis pigmentosa (RP), sensorineural hearing impairment, and in some cases vestibular dysfunction. Three clinical types can be distinguished [1]. Patients with Usher syndrome type I (*USH1*) show severe to profound congenital hearing loss, early onset RP, and vestibular areflexia. Patients with type II (*USH2*) suffer from moderate to severe congenital hearing loss, onset of RP around puberty or in adulthood, but normal vestibular function. Usher type III (*USH3*) patients present with progressive hearing loss, RP, and variable vestibular function. Furthermore, some Usher syndrome patients cannot be classified into any of these three subtypes and are considered atypical Usher (*USHA*) patients. To date, nine genes are known to be involved in *USH*: *MYO7A* (*USH1B*),

USH1C, *CDH23* (*USH1D*), *PCDH15* (*USH1F*), and *USH1G* for *USH1*; *USH2A*, *GPR98* (*USH2C*), and *DFNB31* (*USH2D*) for *USH2*; and *USH3A* for *USH3* [2–14]. The eight genes responsible for *USH1* and *USH2* encode proteins that interact in a functional network.

The *DFNB31* gene was initially found to be responsible for recessive nonsyndromic sensorineural hearing impairment (*NSHI*), underlying the *DFNB31* locus [15]. *DFNB31* comprises 12 exons and encodes the whirlin protein. Two isoforms of whirlin are known. The short isoform is encoded by exons 6–12 and contains one PDZ domain (PDZ3) and a proline-rich region. The long isoform is encoded by all 12 *DFNB31* exons and contains three PDZ domains (PDZ1, PDZ2, and PDZ3) and a proline-rich region [15]. PDZ1 and PDZ2 have been shown to interact with usherin, myosin VIIa, G-Protein coupled Receptor 98 (*GPR98*), Scaffold protein containing Ankyrin repeats and SAM domain (*SANS*), and myosin XVa, while PDZ3 interacts only with myosin XVa [16–18]. All these findings made *DFNB31* an excellent candidate for Usher syndrome. More recently, Ebermann et al. [14] described a novel genetic subtype for Usher syndrome

Correspondence to: Jose Millan, Hospital Universitario La Fe and CIBERER, Genetics Avda. Campanar, 21, Valencia, Spain 46009; Phone: +34 96 197 3153; FAX: +34 96 3089; email: millan_jos@gva.es

TABLE 1. CLINICAL CLASSIFICATION OF USH PATIENTS.

Origin	USH1	USH2	USHA	USH NC*	Total
The Netherlands	0	27	2	10	39
Spain	21	51	2	0	74
Germany	2	44	0	1	47
Hungary	2	13	2	0	17
Canada	1	4	0	0	5
Russia	1	2	0	0	3
Arabia	0	1	0	0	1
Egypt	2	2	0	0	4
Turkey	0	3	0	0	3
Unknown	0	2	0	0	2
Total	29	149	6	11	195

*USH NC: Clinical data not available to classify the patients into any subtype.

type II (USH2D) caused by truncating mutations in the long isoform of whirlin.

The protein whirlin is thought to play an important role in the elongation of stereocilia and development of cochlear hair cells and has been demonstrated to be the major scaffold protein of an Usher protein network in the ankle link complex of inner hair cells [15,17,19]. In the retina, whirlin is a key partner of another USH interactome in the periciliary region of photoreceptor cells. This Usher protein network is hypothesized to be essential for the regulation of cargo transfer from the photoreceptor inner segment to the outer segment by the ciliary transport system [18].

The recent finding that mutations in *DFNB31* cause USH2 prompted us to screen the 12 exons of this gene in 195 patients from diverse origins who have USH1, USH2, and USHA. Our aim was to elucidate the prevalence of *DFNB31* mutations in the pathogenesis of USH.

METHODS

Patients: We included 149 patients with USH2, 29 patients with USH1, and six patients with USHA in this study. Furthermore, 11 unclassified USH patients were included from whom clinical data were not available. Of the 195 unrelated patients, 74 were of Spanish origin, 47 of German, 39 of Dutch, 17 of Hungarian, and 13 of other different origins (see Table 1).

Most of these patients were selected for *DFNB31* screening after discarding the presence of pathologic mutations in the most prevalent USH genes. Of the 195 patients, 116 patients correspond to those briefly mentioned in previous studies [13,16]. Clinical diagnosis of Usher syndrome patients was performed in different hospitals according to standard ophthalmologic, otorhinolaryngologic, and electroretinographic procedures. Written informed consent was obtained for each participant, and the study was approved by the review board of the Ethics Committee of each institution that participated in this work.

Mutation screening: Venous blood samples were obtained and genomic DNA was extracted by standard protocols [7]. For mutation analysis of *DFNB31*, the 12 coding exons were PCR amplified using primers located in flanking introns and untranslated region (UTR) sequences (Table 2). PCR products were directly sequenced using standard procedures (Table 2). The obtained sequences were compared with the consensus sequence [NM_015404](#).

Isocoding and missense changes found in a low frequency were analyzed with the program [RESCUE-ESE](#) (Christopher Burge Laboratory, Massachusetts Institute of Technology; Cambridge, MA). This program predicts the creation or elimination of exonic splicing enhancer sites (ESEs).

To assess the effect of the amino acid substitutions on the protein and to study the degree of interspecies amino acid conservation we used the software [Alamut v1.5 Interactive Biosoftware](#) (Rouen, France).

Intronic, isocoding, and missense changes were also analyzed using the programs [NNSPLICE](#) (Berkeley Drosophila Genome Project [BDGP] University of California Berkeley; Berkeley, CA) and [Splice View](#) (Institute of Biomedical Technologies, Milano, Italy) to predict if those changes could be affecting, creating, or eliminating donor/acceptor splice sites.

RESULTS

Mutation screening of *DFNB31* in 195 USH patients of diverse origins identified 38 different sequence variants: eight synonymous, 15 missense, and 15 intronic variants. Most of these changes are clearly polymorphic as they were detected in high frequencies and represent known single nucleotide polymorphisms (SNPs). Twelve sequence variants were found only once in our series, five changes were found twice, one variant was identified in three alleles, and another one in four (see Table 3). None of these 19 rare changes were found in the homozygous state or in trans to another putatively pathogenic variation.

TABLE 2. PRIMER SEQUENCES FOR THE AMPLIFICATION OF THE CODING REGION OF *DFNB31*.

Primer name	Sequence (5'-3')	Annealing temperature
Exon 1	F: CAGCAGCCAACTCTTGTGTC R: CCAGAAAGGCCAAGTGATTC	55 °C
Exon 2	F: ACTCCCCAAATCCAAGTTCC R: CAGAACCAGCCTCTTCTTGC	58 °C
Exon 3	F: CTCCTTGCCAGTCGGATAAG R: GAGTGCTGATTGCTCTGCTG	55 °C
Exon 4	F: ATAAGGGGACCCCTTGGAAATG R: TCCCCACTTTTTGGATGAAG	55 °C
Exon 5	F: GTCCGGAGTTTCCTTTACCC R: TGGTCTGCTCTGTTCAATTGC	55 °C
Exon 6	F: TGGCAATGAACAGAGCAGAC R: GGAGGGCTTGTGAAGATGAC	58 °C
Exon 7	F: GACAGGGAAGCAGGAGTGAG R: GATTCGAACTCAGGCTGGTC	58 °C
Exon 8	F: CAGCATCTCTGGCAGTTCAG R: GGCTGTCATGGAGAGGAGAG	58 °C
Exon 9	F: GTGACAAGCTCTGGCTGATG R: TTCAAACCTGGGGTCTCCAAC	58 °C
Exon 10	F: GGTCTGGTTGAAAGGACAGG R: GGCCTCCAGATTCCTAATCC	58 °C
Exon 11	F: GAGGCTGAGATTGGTCTTGG R: CCTAGGTCTGCCCTTGAGTG	55 °C
Exon 12	F: CCCTTTCTCAGCATCTCCAG R: GTCTGCCTTGTCTCTCTCTC	58 °C

Amplification conditions were 95 °C 5 min followed by 35 cycles of 30 s at 95 °C, 30 s at an annealing temperature specific for each exon (column 3) and 30 s at 72 °C.

Isocoding changes: Only three out of the eight silent changes were found with an allele frequency ≤ 0.01 (see Table 3 for more details). Variant c.1486C>T was only found in one USH2 patient from uncertain origin, and c.2307C>T was found in one USH1 patient from Spain and in one USH2 patient from Germany. Variant c.1455G>A was detected in four alleles of patients from the Netherlands and Germany. These three rare isocoding variants were analyzed using the [RESCUE-ESE](#) program, but none were found to create or eliminate any ESEs.

These three changes were also analyzed with NNSPLICE and Splice View. Both programs predicted the creation of one new donor site when introducing the change c.2307C>T (NNSPLICE score of 0.66 [0.00–1.00] and Splice View score of 86 [0–100]).

Missense changes: From the 15 missense variants detected, only p.T77S, p.R350W, p.T383N, p.T383S, p.D447H, p.S628R, p.S648Y, p.M723I, and p.R882S showed a frequency ≤ 0.01 (see Table 3).

These nine variants were analyzed using the [Alamut program](#). For two of these changes (p.R350W and p.R882S),

this program predicted possible implications in protein structure and function. The amino acid substitution p.R350W (c.1048C>T) affected a highly conserved nucleotide at the cDNA level with a score of 0.9 (0–1). The amino acid conservation among 13 species was moderate, and the Grantham distance between amino acids Arg and Trp was 101 (0–215). Furthermore, this variation was found to be located in the PDZ2 protein domain. The change p.R882S (c.2644C>A at the cDNA level) affects a highly conserved nucleotide (score 1.0 [0–1]) and an amino acid highly conserved up to the fruit fly (considering 13 species). The physicochemical difference between Arg and Ser was found to be moderate since a Grantham distance of 110 (0–215) was obtained. Also this variation was located in the PDZ3 domain.

These nine rare missense variants were further evaluated for an effect on exonic splice enhancers by using the [RESCUE-ESE](#) program. The change c.1148C>A (p.T383N), found in two unrelated USH2 patients from the Netherlands, creates four new putative ESE sites, whereas c.1339G>A (p.D447H), detected in one USH2 patient from Hungary, suppresses one existing ESE sequence and creates two new

TABLE 3. VARIANTS DETECTED IN THIS STUDY

Nucleotide change	SNP	Exon	Amino acid change	Protein location	Number of alleles	Allele frequency	Homozygotes	Heterozygotes	Origin
Isocoding									
c.117G>A	rs2297815	1	None	LC1-PDZ1	111	0.383	20	71	N, S, G, H, C, R, T, A, E
c.1353T>Ca	rs4979387	6	None	PDZ2-LC2	263	0.674	96	71	N, S, G, H, C, R, T, E
c.1455G>A		7	None	PDZ2-LC2	4	0.01	0	4	N, G
c.1486C>T		7	None	PDZ2-LC2	1	0.003	0	1	Unknown
c.1515G>A	rs34252199	7	None	PDZ2-LC2	10	0.026	0	10	S, G, H
c.1886G>A		9	None	LC3	18	0.046	6	6	S, N
c.2283C>T	rs34963246	10	None	LC3-PDZ3	30	0.077	3	24	S, G, H, S, C, T
c.2307C>T		10	None	LC3-PDZ3	2	0.005	0	2	S, G
Missense									
c.229A>T	rs56204273	1	p.T7TS	Between LC1-PDZ1	1	0.003	0	1	N
c.1048C>T		4	p.R350W	PDZ2	1	0.003	0	1	E
c.1148C>A		4	p.T383N	Between PDZ2-LC2	2	0.005	0	2	N
c.1148C>G		4	p.T383S	Between PDZ2-LC2	1	0.003	0	1	S
c.1309G>A		6	p.A437T	Between PDZ2-LC2	6	0.015	0	6	S
c.1318G>A	rs4978584	6	p.A440T	Between PDZ2-LC2	88	0.226	16	55	N, S, G, C, R, T, E
c.1339G>A		p.D447H	Between PDZ2-LC2	1	0.003	0	1	H	
c.1684C>G	rs12339210	8	p.P562A	Between LC2-LC3	39	0.1	2	35	N, S, G, H, C, R, T, E
c.1838T>Ca	rs942519	9	p.M613T	Between LC2-LC3	179	0.459	45	89	N, S, G, H, C, R, T, E
c.1884C>A		p.S628R	LC2	1	0.003	0	1	1	E
c.1943C>A		p.S648Y	Between LC3-PDZ3	1	0.003	0	1	1	S
c.2169G>A		p.M723I	Between LC3-PDZ3	1	0.003	0	1	1	H
c.2348T>C*	rs2274159	10	p.V783A	Between LC3-PDZ3	153	0.392	30	93	N, S, G, H, C, R, T, E
c.2388C>A	rs2274158	10	p.N796K	Between LC3-PDZ3	85	0.218	11	63	N, S, G, H, C, R, T, E
c.2644C>A		12	p.R882S	PDZ3	1	0.003	0	1	S
Intronic									
c.619-41A>G		2	None		1	0.003	0	1	H
c.837+41A>G		2	None		2	0.005	0	2	E, unknown
c.964-21A>G	rs2274163	4	None		114	0.292	14	86	N, S, G, H, C, C, R, T, E
c.1203+114C>T		5	None		2	0.005	0	2	S
c.1416+151A>G	rs4979385	6	None		19	0.049	2	15	S
c.1416+62delC		6	None		1	0.003	0	1	S
c.1416+22A>T	rs4979386	6	None		77	0.197	8	61	N, S, G, H, C, T, E
c.1627-12G>A	rs2274160	8	None		68	0.174	5	19	N, S, G, H, C, R, T, A, E
c.2236+84G>T	rs10982200	9	None		8	0.021	0	8	G, H
c.2237-44C>T	rs766835	10	None		3	0.008	0	3	N
c.2418+142A>G	rs10739410	10	None		29	0.074	5	19	N, S
c.2419-199A>G		11	None		18	0.046	5	8	N, S
c.2419-118G>A	rs55833018	11	None		12	0.031	0	12	S
c.2419-16T>C		11	None		1	0.003	0	1	G
c.2644-157C>T		12	None		2	0.005	0	2	G, S

The asterisk indicates previously reported as polymorphism by Tili et al. [20]. Abbreviations: N represents The Netherlands, S represents Spain, G represents Germany, H represents Hungary, C represents Canada, R represents Russia, T represents Turkey, A represents Arabia, E represents Egypt, LC represents Low complexity protein domain.

putative ESE sites. The remaining variants were not found to affect any ESE site.

These changes were also analyzed with NNSPLICE and Splice View, but neither was found to affect, create, or eliminate any donor or acceptor splice site.

Intronic changes: All intronic variants were detected in frequencies ≥ 0.01 , except for c.619–41A>G, c.837+41A>G, c.1203+114C>T, c.1416+62delC, c.2237–44C>T (rs766835), c.2419–16T>C, and c.2644–157C>T (Table 3). None of these changes were predicted to affect or create splice donor or splice acceptor sites in the analysis with the NNSPLICE and Splice View programs.

DISCUSSION

To date only one Usher family with clearly pathogenic, truncating, *DFNB31* mutations has been described [14]. Here, we describe additional sequence variants in USH patients tested for mutations in *DFNB31* and discuss their putative pathologic effects.

A total of 38 different variants were found in the *DFNB31* sequence, but no variant was a frameshift or a nonsense change. Out of these, 19 rare variants were found with allelic frequency ≤ 0.01 ; two of them were located in PDZ protein domains (p.R350W and p.R882S), and four were predicted to create or abrogate splice sites (c.2307C>T, p.T383N, p.D447H, and p.R882S).

By the introduction of p.R350W and p.R882S, uncharged amino acids may affect the three-dimensional structure of the protein, which is important for the adequate function of interacting domains. Furthermore, both the amino acids R350 and R882 were well conserved throughout evolution, indicating these residues have an important role in protein structure and function. All these factors point to a possible pathologic effect for these two variants.

Segregation analysis for the variants p.R350W, p.G769G, p.T383N, and p.D447H could not be performed. In some cases, we only had a DNA sample from a patient without any family history. In other cases, the patient was a sporadic case and segregation analysis did not reveal any information about the pathogenicity of the variant (e.g., in the case of p.R882S). We found variant p.R882S in a sporadic Spanish USH2 patient together with the variant p.S648Y. Segregation analysis for these changes was performed in healthy relatives and revealed that both variants were located on the same allele. In addition, p.R882S and p.S648Y were present in the patient, in his healthy brother and sister, and in his two healthy children.

To explore the possibility of yet unidentified exons of *DFNB31*, which may contain mutations in the second allele of patients with a potentially pathogenic mutation on one allele, we performed an in silico search by using Geneid (Genome Bioinformatics Research Lab; Center for Genomic Regulation, Barcelona, Spain), N-SCAN (Computational

Genomics Laboratory; Washington University of St. Louis, St. Louis; MO) and we also searched in the UCSC Genome Browser (University of California Santa Cruz Genome Bioinformatics Web Site, Santa Cruz, CA), but no additional exons were predicted in the *DFNB31* locus.

In summary, this study did not reveal evidence for the involvement of *DFNB31* mutations in 195 unrelated USH patients. However, total or partial gene deletions and duplications can escape the screening method applied herein and therefore cannot be excluded. Our results indicate a minor causative role for *DFNB31* in Usher syndrome. However, modifying effects of the variants we detected might contribute to the phenotype of Usher syndrome.

Regarding the implication of this gene in nonsyndromic hearing loss, the studies performed so far also indicate that *DFNB31* is a rare form of deafness [15,20]. We previously hypothesized that *DFNB31* mutations may also be causative for nonsyndromic retinal degenerations. This remains to be confirmed.

ACKNOWLEDGMENTS

The authors are grateful to the participating patients and their relatives. Supported by grants BO 2954/1–1 (Deutsche Forschungsgemeinschaft), Koeln Fortune Program, grant 113/2004 (Faculty of Medicine, University of Cologne) and Forschung contra Blindheit – Initiative Usher Syndrom e.V. to H.J.B and Research Projects PI07/0558 and PI08/90311 (Institute of Health Carlos III, Spanish Ministry of Science), grant GR552 of the BRPS to H.K. and by the Forschung contra Blindheit – Initiative Usher Syndrome.V to H.K. Elena Aller, Teresa Jaijo and Jose M. Millán are members of the CIBER for Rare Diseases (CIBERER), an initiative of the Institute of Health Carlos III from the Spanish Ministry of Science and Innovation.

REFERENCES

1. Kimberling WJ, Möller C. Clinical and molecular genetics of Usher syndrome. *J Am Acad Audiol* 1995; 6:63-72. [PMID: 7696679]
2. Weil D, Blanchard S, Kaplan J, Guilford P, Gibson F, Walsh J, Mburu P, Varela A, Levilliers J, Weston MD, Kelley PM, Kimberling WJ, Wagenaar M, Levi-Acobas F, Larget-Piet D, Munnich A, Steel KP, Brown SDM, Petit C. Defective myosin VIIA gene responsible for Usher syndrome type 1B. *Nature* 1995; 374:60-1. [PMID: 7870171]
3. Eudy JD, Weston MD, Yao S, Hoover DM, Rehm HL, Ma-Edmonds M, Yan D, Ahmad I, Cheng JJ, Ayuso C, Cremers C, Davenport S, Moller C, Talmadge CB, Beisel KW, Tamayo M, Morton CC, Swaroop A, Kimberling WJ, Sumegi J. Mutation of a gene encoding a protein with extracellular matrix motifs in Usher syndrome type IIa. *Science* 1998; 280:1753-7. [PMID: 9624053]
4. Bitner-Glindzicz M, Lindley KJ, Rutland P, Blaydon D, Smith VV, Milla PJ, Hussain K, Furth-Lavi J, Cosgrove KE, Shepherd RM, Barnes PD, O'Brien RE, Farndon PA, Sowden J, Liu XZ, Scanlan MJ, Malcolm S, Dunne MJ, Aynsley-

- Green A, Glaser B. A recessive contiguous gene deletion causing infantile hyperinsulinism, enteropathy and deafness identifies the Usher type 1C gene. *Nat Genet* 2000; 26:56-60. [PMID: 10973248]
5. Verpy E, Leibovici M, Zwaenepoel I, Liu XZ, Gal A, Salem N, Mansour A, Blanchard S, Kobayashi I, Keats BJ, Slim R, Petit C. A defect in harmonin, a PDZ domain-containing protein expressed in the inner ear sensory hair cells, underlies Usher syndrome type 1C. *Nat Genet* 2000; 26:51-5. [PMID: 10973247]
 6. Ahmed ZM, Riazuddin S, Bernstein SL, Ahmed Z, Khan S, Griffith AJ, Morell RJ, Friedman TB, Riazuddin S, Wilcox ER. Mutations of the protocadherin gene PCDH15 cause Usher syndrome type 1F. *Am J Hum Genet* 2001; 69:25-34. [PMID: 11398101]
 7. Alagramam KN, Yuan H, Kuehn MH, Murcia CL, Wayne S, Srisailpathy CR, Lowry RB, Knaus R, Van Laer L, Bernier FP, Schwartz S, Lee C, Morton CC, Mullins RF, Ramesh A, Van Camp G, Hageman GS, Woychik RP, Smith RJ. Mutations in the novel protocadherin PCDH15 cause Usher syndrome type 1F. *Hum Mol Genet* 2001; 10:1709-18. [PMID: 11487575] Erratum in: *Hum Mol Genet* 2001;10:2603
 8. Bolz H, von Brederlow B, Ramirez A, Bryda EC, Kutsche K, Nothwang HG, Seeliger M, del C-Salcedo Cabrera M, Vila MC, Molina OP, Gal A, Kubisch C. Mutation of CDH23, encoding a new member of the cadherin gene family, causes Usher syndrome type 1D. *Nat Genet* 2001; 27:108-12. [PMID: 11138009]
 9. Bork JM, Peters LM, Riazuddin S, Bernstein SL, Ahmed ZM, Ness SL, Polomeno R, Ramesh A, Schloss M, Srisailpathy CR, Wayne S, Bellman S, Desmukh D, Ahmed Z, Khan SN, Kaloustian VM, Li XC, Lalwani A, Bitner-Glindzicz M, Nance WE, Liu XZ, Wistow G, Smith RJ, Griffith AJ, Wilcox ER, Friedman TB, Morell RJ. Usher syndrome 1D and nonsyndromic autosomal recessive deafness DFNB12 are caused by allelic mutations of the novel cadherin-like gene CDH23. *Am J Hum Genet* 2001; 68:26-37. [PMID: 11090341]
 10. Adato A, Vreugde S, Joensuu T, Avidan N, Hamalainen R, Belenkiy O, Olender T, Bonne-Tamir B, Ben-Asher E, Espinos C, Millan JM, Lehesjoki AE, Flannery JG, Avraham KB, Pietrokovski S, Sankila EM, Beckmann JS, Lancet D. USH3A transcripts encode clarin-1, a four-transmembrane-domain protein with a possible role in sensory synapses. *Eur J Hum Genet* 2002; 10:339-50. [PMID: 12080385]
 11. Weil D, El-Amraoui A, Masmoudi S, Mustapha M, Kikkawa Y, Laine S, Delmaghani S, Adato A, Nadifi S, Zina ZB, Hamel C, Gal A, Ayadi H, Yonekawa H, Petit C. Usher syndrome type I G (USH1G) is caused by mutations in the gene encoding SANS, a protein that associates with the USH1C protein, harmonin. *Hum Mol Genet* 2003; 12:463-71. [PMID: 12588794]
 12. Weston MD, Lujendijk MW, Humphrey KD, Moller C, Kimberling WJ. Mutations in the VLGR1 gene implicate G-protein signaling in the pathogenesis of Usher syndrome type II. *Am J Hum Genet* 2004; 74:357-66. [PMID: 14740321]
 13. van Wijk E, Pennings RJ, te Brinke H, Claassen A, Yntema HG, Hoefsloot LH, Cremers FP, Cremers CW, Kremer H. Identification of 51 novel exons of the Usher syndrome type 2A (USH2A) gene that encode multiple conserved functional domains and that are mutated in patients with Usher syndrome type II. *Am J Hum Genet* 2004; 74:738-44. [PMID: 15015129]
 14. Ebermann I, Scholl HP, Charbel Issa P, Becirovic E, Lamprecht J, Jurklics B, Millán JM, Aller E, Mitter D, Bolz H. A novel gene for Usher syndrome type 2: mutations in the long isoform of whirlin are associated with retinitis pigmentosa and sensorineural hearing loss. *Hum Genet* 2007; 121:203-11. [PMID: 17171570]
 15. Mburu P, Mustapha M, Varela A, Weil D, El-Amraoui A, Holme RH, Rump A, Hardisty RE, Blanchard S, Coimbra RS, Perfettini I, Parkinson N, Mallon AM, Glenister P, Rogers MJ, Paige AJ, Moir L, Clay J, Rosenthal A, Liu XZ, Blanco G, Steel KP, Petit C, Brown SD. Defects in whirlin, a PDZ domain molecule involved in stereocilia elongation, cause deafness in the whirler mouse and families with DFNB31. *Nat Genet* 2003; 34:421-8. [PMID: 12833159]
 16. van Wijk E, van der Zwaag B, Peters T, Zimmermann U, Te Brinke H, Kersten FF, Märker T, Aller E, Hoefsloot LH, Cremers CW, Cremers FP, Wolfrum U, Knipper M, Roepman R, Kremer H. The DFNB31 gene product whirlin connects to the Usher protein network in the cochlea and retina by direct association with USH2A and VLGR1. *Hum Mol Genet* 2006; 15:751-65. [PMID: 16434480]
 17. Michalski N, Michel V, Bahloul A, Lefèvre G, Barral J, Yagi H, Chardenoux S, Weil D, Martin P, Hardelin JP, Sato M, Petit C. Molecular characterization of the ankle-link complex in cochlear hair cells and its role in the hair bundle functioning. *J Neurosci* 2007; 27:6478-88. [PMID: 17567809]
 18. Maerker T, van Wijk E, Overlack N, Kersten FF, McGee J, Goldmann T, Sehn E, Roepman R, Walsh EJ, Kremer H, Wolfrum U. A novel Usher protein network at the periciliary reloading point between molecular transport machineries in vertebrate photoreceptor cells. *Hum Mol Genet* 2008; 17:71-86. [PMID: 17906286]
 19. Belyantseva IA, Boger ET, Naz S, Frolenkov GI, Sellers JR, Ahmed ZM, Griffith AJ, Friedman TB. Myosin-XVa is required for tip localization of whirlin and differential elongation of hair-cell stereocilia. *Nat Cell Biol* 2005; 7:148-56. [PMID: 15654330]
 20. Tlili A, Charfedine I, Lahmar I, Benzina Z, Mohamed BA, Weil D, Idriss N, Drira M, Masmoudi S, Ayadi H. Identification of a novel frameshift mutation in the DFNB31/WHRN gene in a Tunisian consanguineous family with hereditary non-syndromic recessive hearing loss. *Hum Mutat* 2005; 25:503. [PMID: 15841483]

The print version of this article was created on 23 March 2010. This reflects all typographical corrections and errata to the article through that date. Details of any changes may be found in the online version of the article.