

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/97014>

Please be advised that this information was generated on 2021-02-26 and may be subject to change.

Immunovirological Response to Triple Nucleotide Reverse-Transcriptase Inhibitors and Ritonavir-Boosted Protease Inhibitors in Treatment-Naïve HIV-2–Infected Patients: The ACHIEV_{2E} Collaboration Study Group

Antoine Benard,¹ Ard van Sighem,⁶ Audrey Taieb,¹ Emilia Valadas,⁷ Jean Ruelle,⁸ Vicente Soriano,⁹ Alexandra Calmy,¹⁰ Claudia Balotta,¹¹ Florence Damond,^{3,4} Françoise Brun-Vezinet,^{3,4} Geneviève Chene,^{1,2} Sophie Matheron,^{4,5} and the ACHIEV_{2E} Collaboration Study Group*

¹INSERM, U897; ²University Bordeaux Segalen, Bordeaux; ³APHP, Hôpital Bichat – Claude Bernard, Laboratoire de Virologie; ⁴Paris VII Denis Diderot University; ⁵APHP, Hôpital Bichat-Claude Bernard, Service de Maladies infectieuses et Tropicales, Paris, France; ⁶Stichting HIV Monitoring, Amsterdam, the Netherlands; ⁷Hospital de Santa Maria, Clinica Universitaria de Doenças Infecciosas, Lisbon, Portugal; ⁸Université Catholique de Louvain, AIDS Reference Laboratory, Brussels, Belgium; ⁹Hospital Carlos III, Department of Infectious Diseases, Madrid, Spain; ¹⁰Hôpitaux Universitaires de Genève, Service de Maladies Infectieuses, Unite VIH/SIDA, Geneva, Switzerland; and ¹¹University of Milan, Department of Clinical Sciences "L. Sacco", Section of Infectious Diseases, Milan, Italy

Background. Triple nucleoside reverse-transcriptase inhibitors (NRTIs) are recommended by the World Health Organization as first-line regimen in treatment-naïve HIV-2–infected patients. However, ritonavir-boosted protease inhibitor (PI/r)–containing regimens are frequently prescribed. In the absence of previous randomized trials, we retrospectively compared these regimens in observational cohorts.

Methods. HIV-2–infected patients from 7 European cohorts who started triple NRTI or PI/r since January 1998 were included. Piecewise linear models were used to estimate CD4 cell count and plasma HIV-2 RNA level slopes, differentiating an early phase (until end of month 3) and a second phase (months 4–12). On-treatment analyses censored data at major treatment modification and systematically at month 12.

Results. Forty-four patients started triple NRTI therapy and 126 started PI/r therapy. Overall, the median CD4 cell count was 191 cells/mm³ and the median plasma HIV-2 RNA level was ≥ 2.7 log₁₀ copies/ml in 61% of the patients at combination antiretroviral therapy (cART) initiation; the median duration of the first cART was 20 months, not differing between groups. PI/r regimens were associated with better CD4 cell count and HIV-2 RNA level outcomes, compared with NRTI regimens. Estimated CD4 cell count slopes were +6 and +12 cells/mm³/month during the early phase ($P = .22$), and -60 cells/mm³/year versus $+76$ cells/mm³/year during the second phase ($P = .002$), for triple NRTI and PI/r, respectively. Estimated mean HIV-2 RNA levels at month 12 in patients with detectable viremia at cART initiation were 4.0 and 2.2 log₁₀ copies/ml, respectively ($P = .005$).

Conclusions. In this observational study, PI/r-containing regimens showed superior efficacy over triple NRTI regimens as first-line therapy in HIV-2–infected patients.

Received 8 October 2010; accepted 3 February 2011.

* The ACHIEV_{2E} Collaboration Study Group is listed in full in the Acknowledgments. Correspondence: Antoine Benard, MD, INSERM Research Centre U897, Université Bordeaux Segalen, Bâtiment ISPED-Case 11, 146 rue Léo Saignat, 33076 Bordeaux Cedex, France (antoine.benard@isped.u-bordeaux2.fr).

Clinical Infectious Diseases 2011;52(10):1257–1266

© The Author 2011. Published by Oxford University Press on behalf of the Infectious Diseases Society of America. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.
1058-4838/2011/5210-0009\$14.00
DOI: 10.1093/cid/cir123

Although ultimately leading to AIDS and death, HIV-2 is associated with a slower T CD4⁺ lymphocyte depletion [1, 2], a lower viral load at comparable CD4 cell counts [1, 3, 4], and a poorer CD4 cell recovery after treatment initiation in naïve patients [1, 5], compared with HIV-1.

Because of the limitation of the epidemic and the fewer treatment options because of natural resistance to enfuvirtide and nonnucleoside reverse-transcriptase

inhibitors (NNRTI) [6, 7], no randomized clinical trial has assessed the efficacy of specific combination antiretroviral therapies (cART) in treatment-naïve HIV-2-infected patients. Available data regarding nucleoside reverse-transcriptase inhibitors (NRTIs) and protease inhibitors (PIs) are provided by small observational studies, evaluating first-generation antiretrovirals and showing no difference between drugs [1, 5, 8–10].

Current World Health Organization (WHO) treatment guidelines for HIV-2 infection recommend triple NRTI regimens as first-line cART [11]. PI in an initial treatment regimen would essentially rule out second-line options in areas with limited access to cART. This is important when considering that most HIV-2-infected individuals are living in sub-Saharan African countries, where tuberculosis is highly prevalent, with available rifamycin curative therapy limited to rifampicin, which interacts with PIs.

However, recent noncomparative studies have suggested better immunological and virological responses to ritonavir-boosted PI-containing cART in antiretroviral-naïve HIV-2-infected patients [12–14]. Furthermore, in vitro phenotypic susceptibility studies of HIV-2 to PI have shown similar half maximal inhibitory concentration, compared with those reported with HIV-1 for ritonavir-boosted darunavir, lopinavir, and saquinavir [15]. On the basis of these observations, we investigated whether PI/r-containing cART was associated with better immunological and virological responses, compared with triple NRTI regimens, as first-line therapy in HIV-2-infected patients, regardless of the level of immunodeficiency at treatment initiation, in a large European collaboration.

METHODS

Study Population

ACHIEV_{2E} (<http://etudes.isped.u-bordeaux2.fr/achiev2e/>) was established in 2005 as a collaboration of 15 observational cohort studies in 10 European countries, Gambia, and North America.

Adult patients included in the analysis had a confirmed HIV-2 infection (HIV-1 and HIV-2 dually infected patients were not included) and started triple NRTI or PI/r (defined as ≥ 3 antiretrovirals) as their first-line regimen from 1 January 1998 through 20 June 2008 (when data were merged). For each individual, follow-up began on the date of cART initiation and ended on the date of the last recorded CD4 cell count.

Seven cohorts participated in the present analysis: the Belgium and Luxemburg HIV-2 Database ($n = 16$); the ANRS CO5 HIV-2 cohort, France ($n = 145$); a cohort from the Section of Infectious Diseases at the “L. Sacco” Hospital in Milan, Italy ($n = 3$); the ATHENA cohort in the Netherlands ($n = 35$); the Santa Maria HIV₂ Cohort in Portugal ($n = 29$); the Spanish

HIV-2 cohort ($n = 9$); and the Swiss HIV Cohort Study ($n = 5$). Each cohort submitted information, using a standardized data format (ie, the HIV Collaboration Data Exchange Protocol) [16], to the coordinating data center at the Bordeaux School of Public Health, France. Data collected included patient demographic characteristics, ART, CD4 cell counts and percentages, HIV-2 RNA level, AIDS, and deaths. The coordinating data center ensured adherence to strict quality-assurance guidelines and performed data quality checks.

Markers and End Point Definition

In each of the 7 cohorts, CD4 cell counts were measured using flow cytometry, but different plasma HIV-2 RNA quantification assays were used [17]. Because each assay had a different threshold of detectability (1.7 – $2.7 \log_{10}$ copies/mL), the highest threshold was taken into account. Furthermore, the best reproducibility was achieved for plasma RNA values above this threshold [17].

All CD4 cell count and plasma HIV-2 RNA level measurements between treatment initiation and month 12 were taken into account to estimate the immunovirological response. In addition, treatment success at month 12 (± 1.5) was defined as an increase in CD4 cell count of ≥ 50 cells/mm³ from treatment initiation, in conjunction with undetectable plasma RNA in the absence of progression to AIDS, death, or major treatment modification (ie, switch from triple NRTI or PI/r to another cART).

Statistical Analysis

The changes in CD4 cell counts and plasma HIV-2 RNA level after cART initiation were studied using 2-phase linear mixed models in which data were censored for major treatment modification or after 12 months of treatment, whichever came first. The date of first cART initiation was considered as baseline. Plasma HIV-2 RNA level changes were estimated in patients with detectable values at baseline. Trends in the evolution of markers were fitted using 2 slopes: one for the early change (0–3 months, in unit/month) and a second for the long-term trend (4–12 months, in unit/year). The correlation between individual baseline value(s) and the subsequent slope(s) was handled through the unstructured covariance matrix of random effects. We performed a secondary analysis stratified by baseline CD4 cell count with use of a threshold of 200 cells/mm³. Left-censoring of plasma viral load because of undetectable values was taken into account by imputing half the value of the assay’s threshold of detectability. A sensitivity analysis was conducted among patients treated with currently recommended regimens: 3 NRTI (lamivudine [3TC], zidovudine [AZT], and abacavir or tenofovir) or PI/r (lopinavir, saquinavir, or darunavir) [11, 18, 19].

Comparisons of proportions were performed using Fisher’s exact tests. Data analyses were conducted using SAS, version 9.1 (SAS Institute).

RESULTS

Overall, 242 HIV-2-infected adults were included in the database. Of these, 72 patients were excluded from subsequent analyses for the following reasons: 56 patients received a non-boosted PI-containing regimen, 15 received NNRTI-based cART, and 1 patient was treated with an enfuvirtide-containing regimen. Of the 170 patients included in the analysis, 44 (26%) received a triple NRTI regimen and 126 (74%) were treated with PI/r. The vast majority (72%) of patients treated with 3 NRTIs received a combination of abacavir, AZT and 3TC. In patients treated with PI/r, 61% received lopinavir, 14% received indinavir, and 13% received saquinavir (Table 1). Backbone regimens were a combination of AZT and 3TC in 79 patients (63%), whereas tenofovir was prescribed in 21 patients (17%; in association with emtricitabine in 11 patients [9%]). As shown in Table 1, 33 patients (75%) treated with 3 NRTIs and 93 (74%) treated with PI/r were selected for the sensitivity analysis including only currently recommended regimens.

Patients treated with 3 NRTIs did not differ from those receiving PI/r with regard to sex, age, mode of infection, and history of AIDS (Table 2). The proportion of patients originating from Africa was higher among those treated with PI/r than those treated with a triple NRTI regimen and patients treated with PI/r tended to have a more advanced infection at treatment initiation, as indicated by a higher proportion of plasma HIV-2 RNA values >2.7 log₁₀ copies/mL and a lower median CD4 cell count, although these differences were not statistically significant.

Table 1. Description of the First-Line cART Prescribed in Treatment-Naive HIV-2-Infected Patients: The ACHIEVE Collaboration, 1998–2008

cART	N	(%)
3 NRTI		
abacavir + lamivudine + zidovudine	32	(73)
tenofovir + lamivudine + zidovudine	1	(2)
didanosine + lamivudine + zidovudine	3	(7)
didanosine + lamivudine + stavudine	3	(7)
abacavir + didanosine + stavudine	2	(5)
tenofovir + lamivudine + abacavir	1	(2)
tenofovir + lamivudine + stavudine	1	(2)
didanosine + lamivudine + tenofovir	1	(2)
Ritonavir-boosted PI		
Lopinavir	76	(61)
Saquinavir	16	(13)
Darunavir	1	(1)
Indinavir	18	(14)
Atazanavir	8	(6)
fos-amprenavir	7	(5)

NOTE. Bold: treatment recommended in antiretroviral-naive HIV-2-infected patients in current guidelines.

The median duration of first-line cART was 19 months (interquartile interval [IQR], 7–40 months) in patients receiving 3 NRTIs and 20 months (IQR, 11–34 months) in patients treated with PI/r. A major treatment modification during the first 12 months of treatment was reported in 10 patients (23%) treated with 3 NRTIs and in 13 patients (10%) treated with PI/r. Reasons were unknown except for 2 patients treated with a triple NRTI regimen that experienced virological failure (threshold variable across participating centers) and 3 patients treated with PI/r, 2 of whom experienced toxicity issues and another who became pregnant.

Virological Response

Sixty-seven patients with detectable plasma RNA values at baseline were included in the estimations of plasma HIV-2 RNA level changes (Figure 1). A total of 229 plasma HIV-2 RNA measurements were available, with a median number of 3 (IQR, 2–3) for patients treated with 3 NRTIs and 4 (IQR, 3–5) for patients treated with PI/r. During the first 3 months of treatment, the estimated decrease in HIV-2 RNA values did not differ in patients treated with 3 NRTIs and those treated with PI/r ($P = .77$). From month 4 through month 12, plasma RNA values remained low in patients treated with PI/r (-0.002 log₁₀ copies/mL/year) and increased in patients treated with 3 NRTIs ($+1.6$ log₁₀ copies/mL/year), although the difference between slopes was not statistically significant ($P = .12$). These changes resulted in estimated plasma HIV-2 RNA values at month 12 being higher in patients treated with 3 NRTIs than in those treated with PI/r (4.0 vs 2.2 log₁₀ copies/mL; $P = .005$).

Only 9 patients treated with 3 NRTIs and 38 treated with PI/r in the subset with detectable plasma RNA values at baseline could be included in the sensitivity analysis restricted to patients given recommended regimens only, and too few RNA measurements were available to use a 2-phase linear mixed model. However, on the basis of observed data, 1 patient (11%) had sustained undetectable RNA values during months 3–12 among those who received 3 NRTIs, compared with 30 (79%) among those who received PI/r.

In patients with undetectable baseline plasma HIV-2 RNA, 1 (8%) of 12 treated with a triple NRTI regimen and 1 (3%) of 31 patients treated with PI/r had at least 1 subsequent detectable HIV-2 RNA value within the first 12 months of treatment.

Immunological Response

Overall, 158 patients were included in the estimation of CD4 cell count changes (Figure 2); the other 12 patients had no CD4 cell count measurements available. A total of 669 CD4 cell count measurements were available, with a median number of 6 (IQR, 4–8) for patients treated with 3 NRTIs and 6 (IQR, 4–10) for patients treated with PI/r. During the first 3 months of treatment, the estimated CD4 cell count change did not differ significantly

Table 2. Main Characteristics of Treatment-Naive HIV-2-Infected Patients Starting a First-Line Triple-NRTI Regimen or PI/r-Containing cART: The ACHIEVE Collaboration, 1998–2008

		Overall (n = 170)		PI/r (n = 126)		3 NRTI (n = 44)		p ^b
At treatment initiation								
Male gender	N (%)	87	(51)	64	(51)	23	(52)	0.67
Age ≥ 45 years	N (%)	90	(53)	64	(51)	26	(59)	0.38
	Median [IQR]	45.6	[38.5–52.2]	45.3	[37.5–51.8]	46.9	[40.4–54.6]	0.45
Infection through heterosexual contact	N (%)	139	(82)	104	(83)	35	(80)	0.65
Originating from Africa	N (%)	128	(75)	100	(79)	28	(64)	0.04
AIDS before cART initiation	N (%)	36	(21)	28	(22)	8	(18)	0.67
Plasma HIV-2 RNA (log ₁₀ copies/ml)	N (%) ≥ 2.7 ^a	67	(61)	56	(64)	11	(48)	0.16
(n = 110 with available data)	Median [IQR] for detectable values	4.0	[3.4–4.6]	4.0	[3.4–4.6]	4.0	[2.9–4.6]	0.64
CD4 count (cells/mm ³)	Median [IQR]	191	[90–275]	170	[72–275]	216	[150–287]	0.21
(n = 134 with available data)								
During follow-up								
Duration of first line cART	Median [IQR]	20	[8–36]	20	[11–34]	19	[7–40]	
Major treatment modifications	N (%)	23	(14)	15	(12)	8	(18)	

NOTE. IQR: Inter-quartile range.

^a The highest threshold of detectability was taken into account to define detectable and undetectable RNA values.

^b Comparison of patients treated with ritonavir-boosted PI based regimen and those receiving a 3-NRTI cART.

between patients treated with 3 NRTIs and those treated with PI/r ($P = .24$). Beyond 3 months of treatment, the estimated CD4 cell count decreased in patients treated with 3 NRTIs and increased in those treated with PI/r (-60 vs $+76$ cells/mm³/year; $P = .002$). These changes resulted in estimated CD4 cell counts at month 12 being lower in patients treated with 3 NRTIs than in patients treated with PI/r (191 vs 327 cells/mm³; $P = .001$).

The difference in estimated CD4 cell counts at month 12 between patients treated with 3 NRTIs and those treated with PI/r remained statistically significant after adjustment for geographical origin ($P = .0009$) or for baseline HIV-2 RNA level ($P = .05$). Among patients originating from Africa, the estimated CD4 cell count at 12 months was 320 cells/mm³ (3 NRTIs), compared with 176 cells/mm³ (PI/r), and among patients with undetectable RNA value at treatment initiation, it was 354 cells/mm³ (3 NRTIs), compared with 242 cells/mm³ (PI/r).

Immunological response appeared to be different after 3 months, regardless of baseline CD4 cell counts. In patients with a baseline CD4 cell count ≥ 200 cells/mm³ (Figure 3.A), the estimated change was -99 cells/mm³/year (95% confidence interval [CI], -201 to 4) among patients receiving a triple NRTI regimen, compared with $+52$ cells/mm³/year in patients receiving PI/r (95% CI, -11 to 119 ; $P = .02$). The same trends were observed in patients with a baseline CD4 cell count < 200 cells/mm³ (Figure 3.B), although the difference according to treatment regimens was not statistically significant ($P = .56$ for the months 0–3 and $P = .26$ for months 4–12).

When the analysis was restricted to recommended regimens only (sensitivity analysis), the baseline estimated CD4 cell count did not differ between patients treated with 3 NRTIs (227 cells/mm³; 95% CI, 163 – 291 cells/mm³) and those treated with PI/r (238 cells/mm³; 95% CI, 201 – 275 cells/mm³; $P = .77$). During the first 3 months of treatment, the estimated CD4 cell count change did not differ significantly between patients treated with 3 NRTIs ($+3$ cells/mm³/month; 95% CI, -7 to 13 cells/mm³/month) and patients treated with PI/r ($+13$ cells/mm³/month; 95% CI, 7 – 19 cells/mm³/month; $P = .09$). Beyond 3 months of treatment, the estimated CD4 cell count slope was -122 cells/mm³/year (95% CI, -139 to 51 cells/mm³/year) in patients treated with 3 NRTIs and $+88$ cells/mm³/year (95% CI, 43 – 134 cells/mm³/year) in those treated with PI/r ($P = .01$). This evolution resulted in lower estimated CD4 cell counts at month 12 in patients treated with 3 NRTIs, compared with those treated with PI/r: 344 cells/mm³ (95% CI, 298 – 390 cells/mm³) versus 204 cells/mm³ (95% CI, 118 – 290 cells/mm³; $P = .005$).

Only 106 patients (62%; 21 treated with 3 NRTIs and 85 with PI/r) had available data at month 12 (± 1.5). The observed success rate was 10% among patients treated with 3 NRTIs and 55% among those treated with PI/r ($P < .001$). Five patients (26%) treated with 3 NRTIs and 50 (67%) treated with PI/r experienced an increase in CD4 cell count of at least 50 cells/mm³ together with undetectable plasma RNA at month 12 ($P = .003$).

None of the patients died during the first 12 months of treatment. One patient (2%) receiving a triple-NRTI regimen experienced progression to AIDS (tuberculosis) 5 months after




Figure 1. Estimated HIV-2 RNA changes with 95% confidence interval in treatment-naive HIV-2-infected patients starting a first-line triple-NRTI regimen or PI/r-containing cART, with detectable values at treatment initiation ($n = 67$). The ACHIEV_{2E} collaboration, 1998–2008. 1st slope (M0–M3): $-0.3 \log_{10}$ copies/mL/month in patients treated with a PI/r-containing cART; $-0.2 \log_{10}$ copies/mL/month in those treated with 3 NRTIs ($P = .77$). 2nd slope (M3–M12): $-0.002 \log_{10}$ copies/mL/month in patients treated with a PI/r-containing cART; $+1.6 \log_{10}$ copies/mL/month in those treated with 3 NRTIs ($P = .12$). M12 estimates: $2.2 \log_{10}$ copies/mL in patients treated with a PI/r-containing cART; $4.0 \log_{10}$ copies/mL in those treated with 3 NRTIs ($P = .005$).




Figure 2. Estimated CD4 cell count changes with 95% confidence interval in treatment-naive HIV-2-infected patients starting a first-line triple-NRTI regimen or PI/r-containing cART ($n = 158$). The ACHIEV_{2E} collaboration, 1998–2008. 1st slope (M0–M3): $+12$ cells/mm³/month in patients treated with a PI/r-containing cART; $+6$ cells/mm³/month in those treated with 3 NRTIs ($P = .24$). 2nd slope (M3–M12): -60 cells/mm³/year in patients treated with a PI/r-containing cART; $+76$ cells/mm³/month in those treated with 3 NRTIs ($P = .002$). M12 estimates: 327 cells/mm³ in patients treated with a PI/r-containing cART; 191 cells/mm³ in those treated with 3 NRTIs ($P = .001$).




Figure 3. Estimated CD4 cell count changes with 95% confidence interval in treatment-naive HIV-2-infected patients starting a first-line triple-NRTI regimen or PI/r-containing cART, with ≥ 200 CD4 cell/mm³ at treatment initiation (A, $n = 71$) or with < 200 CD4 cell/mm³ at treatment initiation (B, $n = 63$). The ACHIEVE collaboration, 1998–2008. A, ≥ 200 CD4 cell/mm³ at treatment initiation 1st slope (M0–M3): +9 cells/mm³/month in patients treated with a PI/r-containing cART; +5 cells/mm³/month in those treated with 3 NRTIs ($P = .45$). 2nd slope (M3–M12): +52 cells/mm³/year in patients treated with a PI/r-containing cART; -99 cells/mm³/month in those treated with 3 NRTIs ($P = .02$). M12 estimates: 451 cells/mm³ in patients treated with a PI/r-containing cART; 236 cells/mm³ in those treated with 3 NRTIs ($P < .001$). B, < 200 CD4 cell/mm³ at treatment initiation 1st slope (M0–M3): +14 cells/mm³/month in patients treated with a PI/r-containing cART; +9 cells/mm³/month in those treated with 3 NRTIs ($P = .56$). 2nd slope (M3–M12): +80 cells/mm³/year in patients treated with a PI/r-containing cART; -12 cells/mm³/month in those treated with 3 NRTIs ($P = .26$). M12 estimates: 228 cells/mm³ in patients treated with a PI/r-containing cART; 156 cells/mm³ in those treated with 3 NRTIs ($P = .2$).

treatment initiation. Patients who progressed to AIDS were 9 (7%) in those treated with PI/r (cytomegalovirus infections [2], recurrent bacterial pneumonia [1], candidiasis [1], toxoplasmosis [1], cryptococcosis [1], pneumocystosis [1], HIV wasting syndrome [1], and unknown [1]) within a median delay of 2 months (range, 0.5–7.5 months) after treatment initiation.

DISCUSSION

In this large collaborative analysis comparing the immunological and virological response to PI/r with that to triple NRTI regimens among ART-naive HIV-2-infected patients followed up in developed countries, we showed better viral suppression and higher CD4 cell recovery associated with PI/r than with a triple NRTI regimen. This result was observed regardless of geographical origin or baseline HIV-2 RNA values and even in patients with baseline CD4 cell counts >200 cells/mm³. A subgroup analysis including only patients treated with currently recommended cART (126; 74%) yielded the same favorable trend for those initially treated with PI/r regimens. A combined end point reflecting successful clinical, therapeutic, virological, and immunological measurements at month 12 showed superiority of PI/r over triple NRTI regimens.

There are several limitations to our analysis. Patients were not randomized to receive one regimen or the other; thus, there were differences between the 2 patient groups at baseline. Although we adjusted for geographical origin and baseline plasma HIV-2 RNA level, a bias due to unmeasured confounding might still remain. Nevertheless, because patients receiving a PI/r regimen had markers of more advanced disease at baseline, this bias would lead to underestimate rather than overestimate the difference between both regimens. No data on adverse events were available. Reasons for major treatment modifications could have provided reasonable information on this matter, but we were not able to collect this information from the participating cohorts. However, major treatment modifications were rather uncommon during the 12-month study period, and we believe that the benefit-to-risk ratio remained in favor of PI/r over 3 NRTI regimens in HIV-2-infected patients. No data on adherence were available in this retrospective study. A higher pill burden (>10 per day) has been associated with a poorer adherence [20, 21]. Because PI/r regimens involve a higher pill burden than triple NRTI regimens used in our study, lack of adherence is expected to occur more frequently with PI/r and to jeopardize the response mainly in that group, again reinforcing our conclusion. Of note, none of the antiretroviral regimens considered in our study necessitated >10 pills per day. Another limitation is the lack of a longer follow-up period and sufficient power to investigate clinical outcomes. However, CD4 cell counts and plasma HIV-2 RNA levels are recognized as major predictors of clinical progression in HIV-2 infection

[3, 22, 23], and we may rely on our conclusion showing a superiority of PI/r over triple NRTI regimens based on these surrogate markers.

To our knowledge, our study was the first to evaluate the efficacy of PI/r in comparison with triple NRTI regimens, since the latter regimens have been recommended by 2010 WHO guidelines as first-line cART in HIV-2-infected patients [11]. Our results are in line with previous noncomparative observational studies in treatment-naive HIV-2-infected patients: poor immunological and virological responses to triple NRTI regimens [8, 14] and good immunological and virological responses to PI/r-containing cART [12, 14].

A large proportion of HIV-2-infected patients in our study had undetectable plasma RNA at treatment initiation, even with low CD4 cell counts. This highlights the importance of taking into account CD4 cell count changes and plasma HIV-2 RNA levels to adequately evaluate treatment responses in HIV-2-infected patients [24]. In our study, the coherence between virological and immunological responses is in favor of a higher effectiveness of PI/r, compared with triple NRTI regimens, in treatment-naive HIV-2-infected patients. Indeed, in patients receiving a triple NRTI regimen, the poor immunological response beyond 3 months of treatment was often observed together with an increase in plasma HIV-2 RNA values during the same period. In contrast, in patients treated with PI/r, the sustained CD4 cell count increase was generally supported by sustained viral suppression.

In vitro phenotypic susceptibility studies have reported a full activity of all NRTIs (including zidovudine, lamivudine, and abacavir) against wild-type ROD and EHO HIV-2 isolates [25]. The difference in viro-immunological response observed between patients receiving a triple-NRTI regimen and those treated with PI/r might be explained by the resistance mutation profile of HIV-2. Indeed, HIV-2 displays NRTI-resistance pathways different from those in HIV-1 [26, 27]. Mutations Q151M (+/–V111I) and K65R develop more frequently in HIV-2 than in HIV-1 and are the main NRTI resistance pathways [28–30]. The Q151M mutation, together with K65R or M184V, is sufficient to confer high-level resistance to both lamivudine and zidovudine, the most frequently prescribed NRTIs in our study. Furthermore, the combination of K65R, Q151M, and M184V mutations confers classwide NRTI resistance [27]. M184V/I appears at treatment failure in patients treated with lamivudine-emtricitabine and has been reported to occur in vitro within 6 weeks [31]. Our results are consistent with these observations even if no resistance data are available yet to establish the role of resistance mutations in response to both triple NRTI regimens and PI/r. A recent study has also increased concerns about the risk of transmission of drug-resistant HIV-2 strains [14], further emphasizing the need for prescribing the most potent first-line ART in HIV-2-infected patients [32].

The vast majority (61%) of patients treated with a PI/r-containing cART in our study received lopinavir, and very few were treated with other potent PI/r regimens that have shown promising results in vitro against HIV-2, such as saquinavir or darunavir [15]. Further research is needed to evaluate the optimal cART regimen for treatment-naïve patients with earlier HIV-2-infection, at best through a randomized clinical trial [24, 33].

In the mean time, our results represent the best evidence to date for the treatment of ART-naïve HIV-2-infected patients and suggest that PI/r-containing cART should be considered as first-line ART, even when CD4 cell counts are >200 cells/mm³.

Acknowledgments

Financial support. The ANRS CO5 HIV-2 Cohort is funded by the French Agency for research on AIDS and viral hepatitis (ANRS). The Swiss HIV Cohort Study is supported by the Swiss National Science Foundation. The ATHENA database is supported by a grant from the Dutch Health Minister. The Hospital Carlos III, Madrid, was funded by FIPSE (US Department of Education), NEAT (European AIDS Treatment Network), and Fundacion Investigacion y Education en Sida. The AIDS Reference Laboratory in Brussels received research grants from Merck Sharp and Dohme and travel grants from Janssen and Viiv Healthcare. The virology laboratory of Bichat-Claude Bernard Hospital, Paris, received grants from the ANRS. The INSERM U897, Bordeaux, received a research grant from Gilead through the EuroCOORD-CHAIN collaboration.

Potential conflicts of interest. A. B. received payment for development of educational presentations from Abbott. G. C. received payment for development of educational presentations from Boehringer Ingelheim and made punctual consultancy for Roche. J. R. received payment for lectures, including service on speakers, bureaus from Siemens Healthcare diagnostics. V. S. received payment for development of educational presentations from BMS, Gilead, and Merck Sharp and Dohme. F. B.-V. received payment for lectures, including service on speakers bureaus from Tibotec, and is board member at Gen Probe. S. M. made punctual consultancy for Boehringer Ingelheim, Abbott, and Gilead and received payment for development of educational presentations from MSD, Gilead, Abbott, Glaxo, Boringher Ingelheim, and BMS. All other authors: no conflicts.

The ACHIEVE Collaboration Study Group

Clinical centres.

France: Clinical centres from the ANRS CO5 HIV-2 Cohort: Bichat-Claude Bernard Hospital, Paris (Sophie Matheron); Pitié-Salpêtrière Hospital, Paris (Roland Tubiana); Saint-Antoine Hospital, Paris (Marie-Caroline Meyohas); Cochin Hospital (Cornélia Bernasconi, Nicolas Dupin); Tenon Hospital, Paris (Laurence Slama); Saint-Louis Hospital, Paris (Diane Ponscarne, Caroline Lascoux-Combe, Françoise-Julie Timsit); Delafontaine Hospital, Saint-Denis (Marie-Aude Khuong); Lariboisière Hospital, Paris (Agathe Rami); Paul Brousse Hospital, Villejuif (Elina Teichner); Villeneuve Saint Georges Hospital (Caroline Semaille); Bicêtre Hospital, Le Kremlin Bicêtre (Yann Quertainmont); Louis Mourier Hospital, Colombes (Martine Bloch); Lagny Hospital, Marne la Vallée (Eric Froguel); Victor Dupouy Hospital, Argenteuil (Philippe Genet); Simone Veil Hospital, Eaubonne (Annie Leprêtre); Foch Hospital, Suresnes (David Zucman); Georges Pompidou Hospital, Paris (Marina Karmochkine); René Dubos Hospital, Pontoise (Laurent Blum); Gilles de Corbeil Hospital,

Corbeil Essones (Pierre Chevojon); Ambroise Paré Hospital, Boulogne Billancourt (Cyril Olivier); Robert Ballanger Hospital, Aulnay sous Bois (Jean-Luc Delassus); Montsouris Hospital, Paris (Loïc Bodard); Bégin Hospital, Saint Mandé (Patrick Imbert); Antoine Bécclère Hospital, Clamart (François Boué); Hôtel-Dieu Hospital, Nantes (Eric Billaud); Saint-Jacques Hospital, Besançon (Christine Drobacheff-Thiébaud); Hôtel-Dieu Hospital, Lyon (Laurent Cotte); Pays d'Aix Hospital, Aix en Provence (Thierry Allègre); Côte de Nacre Hospital, Caen (Claude Bazin); Bretonneau Hospital, Tours (Pascale Nau); Charles Nicolle Hospital, Rouen (Yasmine Debab); Michallon Hospital, Grenoble (Pascale Leclercq); Pontchaillou Hospital, Rennes (Cédric Arvieux); Intercommunal Hospital, Toulon-La Seyne sur Mer (Alain Lafeuillade); Hôpital Pellegrin Hospital, Bordeaux (Jean-Marie Ragnaud, Hervé Dutronc); La Roche sur Yon Hospital (Philippe Perré); Cannes Hospital (Nathalie Montagne); Gui de Chauliac Hospital, Montpellier (Jacques Reynes); Hôtel Dieu Hospital, Clermont Ferrand (Christiane Jacomet); Archet Hospital, Nice (Frédéric Sanderson); Civil Hospital, Strasbourg (David Rey); Saint André Hospital, Bordeaux (Maïté Longy-Boursier); Angers Hospital (Jean-Marie Chennebault); Digne les Bains Hospital (Patricia Granet).

Netherlands: The ATHENA database was set up and is maintained by the Stichting HIV Monitoring. The physicians and data analysts include (*site coordinating physicians): Prof. dr. F. de Wolf (director), Dr D. O. Bezemer, Drs L. A. J. Gras, Drs A. M. Kesselring, Dr A. I. van Sighem, Dr C. Smit, Drs S. Zhang (data analysis group), Drs S. Zaheri (data collection); Academisch Medisch Centrum bij de Universiteit van Amsterdam, Amsterdam: Prof. dr. J. M. Prins*, Prof. dr T. W. Kuijpers, Dr H. J. Scherpbier, Dr K. Boer, Dr J. T. M. van der Meer, Dr F. W. M. N. Wit, Dr M. H. Godfried, Prof. dr. P. Reiss, Drs M. E. Haverkort, Prof. dr. T. van der Poll, Dr F. J. B. Nellen, Prof. dr. J. M. A. Lange, Dr S. E. Geerlings, Dr M. van Vugt, Drs S. M. E. Vrouwenraets, Drs D. Pajkrt, Drs J. C. Bos, Drs M. van der Valk; Academisch Ziekenhuis Maastricht, Maastricht: Dr G. Schreij*, Dr S. Lowe, Dr A. Oude Lashof; Catharina Ziekenhuis, Eindhoven: Drs M. J. H. Pronk*, Dr B. Bravenboer; Erasmus Medisch Centrum, Dr M. E. van der Ende*, Drs T. E. M. S. de Vries-Sluijs, Dr C. A. M. Schurink, Drs M. van der Feltz, Dr J. L. Nouwen, Dr L. B. S. Gelinck, Dr A. Verbon, Drs B. J. A. Rijnders, Drs E. D. van de Ven-de Ruiter, Dr L. Slobbe; Erasmus Medisch Centrum-Sophia, Rotterdam: Dr N. G. Hartwig, Dr G. J. A. Driessen; Flevoziekenhuis, Almere: Dr J. Branger*. HagaZiekenhuis, Den Haag: Dr R. H. Kauffmann*, Dr E. F. Schippers; Isala Klinieken, Zwolle: Dr P. H. P. Groeneveld*, Dr M. A. Alleman, Drs J. W. Bouwhuis; Kennemer Gasthuis, Haarlem: Prof. dr. R. W. ten Kate*, Dr R. Soetekouw; Leids Universitair Medisch Centrum, Leiden: Dr F. P. Kroon*, Prof. dr. P. J. van den Broek, Prof. dr. J. T. van Dissel, Dr S. M. Arend, Drs C. van Nieuwkoop, Drs M. G. J. de Boer, Drs H. Jolink; Maasstad Ziekenhuis, Rotterdam: Dr J. G. den Hollander*, Dr K. Pogany; Medisch Centrum Alkmaar, Alkmaar: Drs W. Kortmann, Drs G. van Twillert*; Medisch Centrum Haaglanden, Den Haag: Dr R. Vriesendorp*, Dr E. M. S. Leyten; Medisch Spectrum Twente, Enschede: Dr C. H. H. ten Napel*, Drs G. J. Kootstra. Onze Lieve Vrouwe Gasthuis, Amsterdam:

Prof. dr. K. Brinkman*, Dr W. L. Blok, Dr P. H. J. Frissen, Drs W. E. M. Schouten, Drs G. E. L. van den Berk; St. Elisabeth Ziekenhuis, Tilburg; Dr J. R. Juttman*, Dr M. E. E. van Kasteren, Drs A. E. Brouwer; Sint Lucas Andreas Ziekenhuis, Amsterdam; Dr J. Veenstra*, Dr K. D. Lettinga; Slotervaart Ziekenhuis, Amsterdam; Dr J. W. Mulder*, Dr E. C. M. van Gorp, Drs P. M. Smit, S. Weijer; Stichting Medisch Centrum Jan van Goyen, Amsterdam; Drs A. van Eeden*, Dr D. W. M. Verhagen*; Universitair Medisch Centrum Groningen, Groningen; Dr H. G. Sprenger*, Dr R. Doedens, Dr E. H. Scholvinck, Drs S. van Assen, C. J. Stek; Universitair Medisch Centrum Sint Radboud, Nijmegen; Dr P. P. Koopmans*, Prof. dr. R. de Groot, Dr M. Keuter, Dr A. J. A. M. van der Ven, Dr H. J. M. ter Hofstede, Dr M. van der Flier, Drs A. M. Brouwer, Dr A. S. M. Dofferhoff; Universitair Medisch Centrum Utrecht, Utrecht; Prof. dr. A. I. M. Hoepelman*, Dr T. Mudrikova, Dr M. M. E. Schneider, Drs C. A. J. J. Jaspers, Dr P. M. Ellerbroek, Dr E. J. G. Peters, Dr L. J. Maarschalk-Ellebroek, Dr J. J. Oosterheert, Dr J. E. Arends, Dr M. W. M. Wassenberg, Dr J. C. H. van der Hilst. VU Medisch Centrum, Amsterdam; Prof. dr. S. A. Danner*, Dr M. A. van Agtmael, Drs J. de Vocht, Dr R. M. Perenboom, Drs F. A. P. Claessen, Drs W. F. W. Bierman, Drs E. V. de Jong, Drs E. A. bij de Vaate; Wilhelmina Kinderziekenhuis, Utrecht; Dr S. P. M. Geelen, Dr T. F. W. Wolfs; Ziekenhuis Rijnstate, Arnhem; Dr C. Richter*, Dr J. P. van der Berg, Dr E. H. Gisolf; Admiraal De Ruyter Ziekenhuis, Vlissingen; Drs M. van den Berge*, Drs A. Stegeman; Medisch Centrum Leeuwarden, Leeuwarden; Drs D. P. F. van Houte*, Dr M. B. Polée, Dr M. G. A. van Vonderen; Sint Elisabeth Hospitaal, Willemstad, Curaçao; Dr C. Winkel, Prof. dr. A. J. Duits.

Portugal: Clinica Universitaria de Doenças Infecciosas, Lisbon (Francisco Antunes, Luis França, Kamal Mansinho, Emilia Valadas).

Spain: Hospital Carlos III, Madrid (Vicente Soriano, Ana Trevino, Berta Rodes).

Switzerland: Swiss HIV Cohort Study. The members of the Swiss HIV Cohort Study are Barth J, Battegay M, Bernasconi E, Böni J, Bucher HC, Bürgisser P, Burton-Jeangros C, Calmy A, Cavassini M, Dubs R, Egger M, Elzi L, Fehr J, Flepp M, Francioli P (President of the SHCS), Furrer H (Chairman of the Clinical and Laboratory Committee), Fux CA, Gorgievski M, Günthard H (Chairman of the Scientific Board), Hasse B, Hirsch HH, Hirschel B, Hösli I, Kahlert C, Kaiser L, Keiser O, Kind C, Klimkait T, Kovari H, Ledergerber B, Martinetti G, Martinez de Tejada B, Müller N, Nadal D, Pantaleo G, Rauch A, Regenass S, Rickenbach M (Head of Data Center), Rudin C (Chairman of the Mother & Child Substudy), Schmid P, Schultze D, Schöni-Affolter F, Schüpbach J, Speck R, Taffé P, Telenti A, Trkola A, Vernazza P, von Wyl V, Weber R, Yerly S.

Laboratories.

Belgium: AIDS Reference Laboratory, Université Catholique de Louvain, AIDS Reference Laboratory, Brussels (Patrick Goubau, Jean Ruelle).

France: Cellular immunology laboratory, Pitié-Salpêtrière Hospital, Paris (Brigitte Autran); virology laboratory, Bichat-Claude Bernard Hospital, Paris (Françoise Brun-Vezinet,

Florence Damond, Diane Descamps), Saint-Louis Hospital, Paris (François Simon).

Italy: University of Milan, Department of Clinical Sciences “L. Sacco” (Claudia Balotta).

Portugal: Hospital Egas Moniz, Lisbon (Ricardo Camacho, Perpetua Gomes).

Spain: Laboratory of Molecular Biology, Infectious Diseases Department, Hospital Carlos III, Madrid (Ana Treviño & Vincent Soriano).

Switzerland: Laboratories of the Swiss HIV Cohort Study (resp. Jürg Böni).

Coordinating center.

France: ANRS Clinical Trials Unit INSERM U897 (Antoine Bénard, Marie Bertoncello, Geneviève Chêne, Audrey Taieb).

References

1. Drylewicz J, Matheron S, Lazaro E, et al. Comparison of viro-immunological marker changes between HIV-1 and HIV-2-infected patients in France. *AIDS* **2008**; 22:457–68.
2. Jaffar S, Wilkins A, Ngom PT, et al. Rate of decline of percentage CD4+ cells is faster in HIV-1 than in HIV-2 infection. *J Acquir Immune Defic Syndr Hum Retrovirol* **1997**; 16:327–32.
3. Alabi AS, Jaffar S, Ariyoshi K, et al. Plasma viral load, CD4 cell percentage, HLA and survival of HIV-1, HIV-2, and dually infected Gambian patients. *AIDS* **2003**; 17:1513–20.
4. MacNeil A, Sarr AD, Sankale JL, Meloni ST, Mboup S, Kanki P. Direct evidence of lower viral replication rates in vivo in human immunodeficiency virus type 2 (HIV-2) infection than in HIV-1 infection. *J Virol* **2007**; 81:5325–30.
5. Matheron S, Damond F, Benard A, et al. CD4 cell recovery in treated HIV-2-infected adults is lower than expected: results from the French ANRS CO5 HIV-2 cohort. *AIDS* **2006**; 20:459–62.
6. Poveda E, Rodes B, Toro C, Soriano V. Are fusion inhibitors active against all HIV variants? *AIDS Res Hum Retroviruses* **2004**; 20:347–8.
7. Tuaille E, Gueudin M, Lemee V, et al. Phenotypic susceptibility to nonnucleoside inhibitors of virion-associated reverse transcriptase from different HIV types and groups. *J Acquir Immune Defic Syndr* **2004**; 37:1543–9.
8. Adje-Toure CA, Cheingsong R, Garcia-Lerma JG, et al. Antiretroviral therapy in HIV-2-infected patients: changes in plasma viral load, CD4+ cell counts, and drug resistance profiles of patients treated in Abidjan, Cote d'Ivoire. *AIDS* **2003**; 17(Suppl 3):S49–54.
9. Mullins C, Eisen G, Popper S, et al. Highly active antiretroviral therapy and viral response in HIV type 2 infection. *Clin Infect Dis* **2004**; 38:1771–9.
10. van der Ende ME, Prins JM, Brinkman K, et al. Clinical, immunological and virological response to different antiretroviral regimens in a cohort of HIV-2-infected patients. *AIDS* **2003**; 17(Suppl 3):S55–61.
11. World Health Organisation. World Health Organisation Antiretroviral therapy in adults and adolescents—recommendations for a public health approach—2010 revision. Available at: <http://www.who.int/hiv/pub/guidelines/artadultguidelines.pdf>. Accessed 6 January 2011.
12. Benard A, Damond F, Campa P, et al. Good response to lopinavir/ritonavir-containing antiretroviral regimens in antiretroviral-naïve HIV-2-infected patients. *AIDS* **2009**; 23:1171–3.
13. Drylewicz J, Eholie S, Maiga M, et al. First-year lymphocyte T CD4+ response to antiretroviral therapy according to the HIV type in the IeDEA West Africa collaboration. *AIDS* **2010**; 24:1043–50.
14. Ruelle J, Roman F, Vandenbroucke AT, et al. Transmitted drug resistance, selection of resistance mutations and moderate antiretroviral

- efficacy in HIV-2: analysis of the HIV-2 Belgium and Luxembourg database. *BMC Infect Dis* **2008**; 8:21.
15. Desbois D, Roquebert B, Peytavin G, et al. In vitro phenotypic susceptibility of human immunodeficiency virus type 2 clinical isolates to protease inhibitors. *Antimicrob Agents Chemother* **2008**; 52:1545–8.
 16. Kjaer J, Ledergerber B. HIV cohort collaborations: proposal for harmonization of data exchange. *Antivir Ther* **2004**; 9:631–3.
 17. Damond F, Benard A, Ruelle J, et al. Quality control assessment of human immunodeficiency virus type 2 (HIV-2) viral load quantification assays: results from an international collaboration on HIV-2 infection in 2006. *J Clin Microbiol* **2008**; 46:2088–91.
 18. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents. Department of Health and Human Services, 2009; 1–161. Available at: <http://www.aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf>. Accessed 6 January 2011.
 19. Yeni P. Prise en charge medicale des personnes infectees par le VIH: rapport 2008. Available at: <http://www.ladocumentationfrancaise.fr/rapports-publics/084000593/>. Accessed 6 January 2011.
 20. Atkinson MJ, Petrozzino JJ. An evidence-based review of treatment-related determinants of patients' nonadherence to HIV medications. *AIDS Patient Care STDS* **2009**; 23:903–14.
 21. Mills EJ, Nachega JB, Buchan I, et al. Adherence to antiretroviral therapy in sub-Saharan Africa and North America: a meta-analysis. *JAMA* **2006**; 296:679–90.
 22. Ariyoshi K, Jaffar S, Alabi AS, et al. Plasma RNA viral load predicts the rate of CD4 T cell decline and death in HIV-2-infected patients in West Africa. *AIDS* **2000**; 14:339–44.
 23. Matheron S, Pueyo S, Damond F, et al. Factors associated with clinical progression in HIV-2 infected-patients: the French ANRS cohort. *AIDS* **2003**; 17:2593–601.
 24. Matheron S. HIV-2 infection: a call for controlled trials. *AIDS* **2008**; 22:2073–4.
 25. Witvrouw M, Pannecouque C, Switzer WM, Folks TM, De Clercq E, Heneine W. Susceptibility of HIV-2, SIV and SHIV to various anti-HIV-1 compounds: implications for treatment and postexposure prophylaxis. *Antivir Ther* **2004**; 9:57–65.
 26. Boyer PL, Sarafianos SG, Clark PK, Arnold E, Hughes SH. Why do HIV-1 and HIV-2 use different pathways to develop AZT resistance? *PLoS Pathog* **2006**; 2:e10.
 27. Smith RA, Anderson DJ, Pyrak CL, Preston BD, Gottlieb GS. Antiretroviral drug resistance in HIV-2: three amino acid changes are sufficient for classwide nucleoside analogue resistance. *J Infect Dis* **2009**; 199:1323–6.
 28. Damond F, Matheron S, Peytavin G, et al. Selection of K65R mutation in HIV-2-infected patients receiving tenofovir-containing regimen. *Antivir Ther* **2004**; 9:635–6.
 29. Descamps D, Damond F, Matheron S, et al. High frequency of selection of K65R and Q151M mutations in HIV-2 infected patients receiving nucleoside reverse transcriptase inhibitors containing regimen. *J Med Virol* **2004**; 74:197–201.
 30. Rodes B, Holguin A, Soriano V, et al. Emergence of drug resistance mutations in human immunodeficiency virus type 2-infected subjects undergoing antiretroviral therapy. *J Clin Microbiol* **2000**; 38:1370–4.
 31. Ntemgwa ML, Toni T, Brenner BG, et al. Nucleoside and nucleotide analogs select in culture for different patterns of drug resistance in human immunodeficiency virus types 1 and 2. *Antimicrob Agents Chemother* **2009**; 53:708–15.
 32. Gottlieb GS, Badiane NM, Hawes SE, et al. Emergence of multiclass drug-resistance in HIV-2 in antiretroviral-treated individuals in Senegal: implications for HIV-2 treatment in resource-limited West Africa. *Clin Infect Dis* **2009**; 48:476–83.
 33. Gottlieb GS, Eholie SP, Nkengasong JN, et al. A call for randomized controlled trials of antiretroviral therapy for HIV-2 infection in West Africa. *AIDS* **2008**; 22:2069–72.