

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/26147>

Please be advised that this information was generated on 2021-02-25 and may be subject to change.

# Isolation and Characterization of Single Anti-U1A-specific B Cells from Autoimmune Patients

RUUD M. T. DE WILDT,<sup>a</sup> FRANK H. J. VAN DEN HOOGEN,<sup>b</sup>  
WALTHER J. VAN VENROOIJ,<sup>a</sup> AND RENÉ M. A. HOET<sup>a,c</sup>

<sup>a</sup>*Department of Biochemistry  
University of Nijmegen*

<sup>b</sup>*Department of Rheumatology  
University Hospital Nijmegen  
6500 HB Nijmegen, the Netherlands*


Patients with systemic lupus erythematosus (SLE) or SLE-overlap syndromes often produce autoantibodies directed to U1RNA-binding proteins.<sup>1</sup> Previously, we isolated and characterized autoantigen-binding Ab fragments directed to the U1RNA-associated A protein (U1A) from several variable (V) gene combinatorial phage libraries.<sup>2</sup> Although these libraries have proven to be very successful, heavy (V<sub>H</sub>) and light (V<sub>L</sub>) chains are randomly combined during construction of such libraries. Because we were interested in the utilization of the original V<sub>H</sub>/V<sub>L</sub> pairings of the encoding autoantibodies, we developed a single cell culture system for B cells using mouse thymoma EL-4 B5 cells.<sup>3</sup> In combination with a preselection via recombinant antigen and single cell sorting using fluorescence-activated cell sorting (FACS), we were able to enrich for autoantigen-specific B cells. V<sub>H</sub> and V<sub>L</sub> genes originating from single U1A-specific B cells were cloned in a phage display vector for expression of single-chain variable fragments (scFv).

Human mononuclear cells were isolated from SLE patients and selected against U1A-coated plates or biotinylated U1A coated to streptavidin-coupled superparamagnetic microbeads (MACS). Adhering cells were collected from the plates via trypsin treatment or were directly used in the case of U1A-bound MACS (FIGURE 1). The selected lymphocytes were plated as single CD19/CD20-positive cells using FACS with an automatic cell deposit unit. Single selected B cells were seeded on 96-well plates containing 20,000 irradiated EL-4 B5 cells and 10% supernatant of phorbol myristate acetate (PMA)- and phytohemagglutinin (PHA)-activated human T cells (FIGURE 2). Cultures were grown for 10–11 days and tested in ELISA for antigen (Ag)-specific antibodies and total Ig production. Cell cultures that were positive in the Ag-specific ELISA screening were used for RNA isolation. An RT reaction was performed with an oligo-dT primer and the cDNA was used in a first PCR with V<sub>H</sub> family-specific primers or a mixture of V-kappa or V-lambda primers (in the case of the light chains) as in Marks *et al.*<sup>4</sup> These first PCR products were used in a second

<sup>c</sup>To whom all correspondence should be addressed.


PCR with nested joining-region primers to introduce the appropriate restriction sites for cloning into pHENIX.<sup>2</sup>

Typical percentages of Ig-positive cells determined by ELISA after 10–11 days of culturing on EL-4 B5 cells in the presence of 10% human T cell supernatant varied between 50% and 70%. The frequency of Ag-specific B cell clones varied between 1% and 2.5% as a percentage of Ig-positive wells. Assuming that the frequency of Ag-specific cells in the periphery varies between  $10^{-5}$  and  $10^{-4}$ , this would indicate an enrichment factor of 100–1000. Distributions of IgG, IgM, and IgG/IgM double-



**FIGURE 1.** Antigen selection on peripheral blood mononuclear cells. Human mononuclear cells were isolated from autoimmune patients (SLE) and used to select on U1A-coated plates or biotinylated U1A coated to streptavidin-coupled MACS. Selected cells were collected from the plates via trypsin treatment or were directly used in the case of U1A-bound MACS. Magnetically sorted cells (MACS) can be processed using FACS.

positive isotypes in Ig-producing single B cell cultures were 3:3:1. After culture in the EL-4 B5 system, the B cells obtained a plasmablast-like phenotype expressing  $CD38^{\text{HIGH}}$  and  $SYNDECAN-1^{\text{MOD}}$  (plasma cell marker). PCR products using cDNA from a single B cell clone amplified with separate  $V_H$  family-specific primers and constant region primers (IgG- or IgM-specific) gave rise to only one  $V_H$  product, which indicates clonality. In the case of the light chains, a product with either V-kappa or V-lambda primers was obtained. Five IgM and 2 IgG anti-U1A-specific clones have been isolated. So far, we have sequenced only one clone with V regions derived from



**FIGURE 2.** The selected lymphocytes were plated as single cells using FACS with an automatic cell deposit unit in 96-well plates containing 20,000 irradiated EL-4 B5 cells and 10% supernatant of phorbol myristate acetate (PMA)- and phytohemagglutinin (PHA)-activated human T cells. Cultures were grown for 10-11 days and tested in ELISA for Ag-specific antibodies and total Ig production.

DP-49 ( $V_H3$ ) for the heavy chain and lv318 for the lambda light chain. We have been able to show binding of the cloned V regions to the U1A protein in ELISA. Further studies are currently being performed to isolate functional recombinant antibodies containing original  $V_H/V_L$  pairings.

## REFERENCES

1. VAN VENROOIJ, W. J. & G. J. M. PRUIJN. 1995. *Curr. Opin. Immunol.* 7: 819-824.
2. DE WILDT, R. M. T., R. FINNERN, W. H. OUWEHAND, A. D. GRIFFITHS, W. J. VAN VENROOIJ & R. M. A. HOET. 1996. *Eur. J. Immunol.* 26: 629-639.
3. STEENBAKKERS, P. G. A., H. A. J. M. HUBERS & A. W. M. RIJNDERS. 1994. *Mol. Biol. Rep.* 19: 125-134.
4. MARKS, J. D., H. R. HOOGENBOOM, T. P. BONNERT, J. MCCAFFERTY, A. D. GRIFFITHS & G. WINTER. 1991. *J. Mol. Biol.* 222: 581-597.